



Title	Random parallel transport on surfaces of finite type, and relations to homotopy
Author(s)	Albeverio,S.; Iwata,K.; Kolsrud,T.
Citation	Hokkaido University Preprint Series in Mathematics, 162, 1-8
Issue Date	1992-08
DOI	10.14943/83306
Doc URL	http://hdl.handle.net/2115/68908
Type	bulletin (article)
File Information	pre162.pdf



[Instructions for use](#)

**Random Parallel Transport on Surfaces
of Finite Type, and Relations to Homotopy**

S. Albeverio, K. Iwata, T. Kolsrud

Series #162. August 1992

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 133: Y. Okabe, Applications of the theory of KM_2O -Langevin equations to the linear prediction problem for the multi-dimensional weakly stationary time series, 22 pages. 1992.
- # 134: P. Aviles, Y. Giga and N. Komuro, Duality formulas and variational integrals, 22 pages. 1992.
- # 135: S. Izumiya, The Clairaut type equation, 6 pages. 1992.
- # 136: S. Izumiya, Singular solutions of first order differential equations, 6 pages. 1992.
- # 137: S. Izumiya, W.L. Marar, The Euler characteristic of a generic wave front in a 3-manifold, 6 pages. 1992.
- # 138: S. Izumiya, W.L. Marar, The Euler characteristic of the image of a stable mapping from a closed n -manifold to a $(2n - 1)$ -manifold, 5 pages. 1992.
- # 139: Y. Giga, Z. Yoshida, A bound for the pressure integral in a plasma equilibrium, 20 pages. 1992.
- # 140: S. Izumiya, What is the Clairaut equation ?, 13 pages. 1992.
- # 141: H. Takamura, Weighted deformation theorem for normal currents, 27 pages. 1992.
- # 142: T. Morimoto, Geometric structures on filtered manifolds, 104 pages. 1992.
- # 143: G. Ishikawa, T. Ohmoto, Local invariants of singular surfaces in an almost complex four-manifold, 9 pages. 1992.
- # 144: K. Kubota, K. Mochizuki, On small data scattering for 2-dimensional semilinear wave equations, 22 pages. 1992.
- # 145: T. Nakazi, K. Takahashi, Hyponormal Toeplitz operators and extremal problems of Hardy spaces, 30 pages. 1992.
- # 146: N. Hayashi, T. Ozawa, Remarks on nonlinear Schrödinger equations in one space dimension, 10 pages. 1992.
- # 147: M. Sato, Interface evolution with Neumann boundary condition, 16 pages. 1992.
- # 148: Y. Okabe, Langevin equations and causal analysis, 49 pages. 1992.
- # 149: Y. Giga, S. Takahashi, On global weak solutions of the nonstationary two-phase Stokes flow, 25 pages. 1992.
- # 150: G. Ishikawa, Determinacy of envelope of the osculating hyperplanes to a curve, 9 pages. 1992.
- # 151: G. Ishikawa, Developable of a curve and determinacy relative to osculation-type, 15 pages. 1992.
- # 152: H. Kubo, Global existence of solutions of semilinear wave equations with data of non compact support in odd space dimensions, 25 pages. 1992.
- # 153: Y. Watatani, Lattices of intermediate subfactors, 33 pages. 1992.
- # 154: T. Ozawa, On critical cases of Sobolev inequalities, 11 pages. 1992.
- # 155: M. Ohnuma, M. Sato, Singular degenerate parabolic equations with applications to geometric evolutions, 20 pages. 1992.
- # 156: S. Izumiya, Perestroikas of optical wave fronts and graphlike Legendrian unfoldings, 13 pages. 1992.
- # 157: A. Arai, Momentum operators with gauge potentials, local quantization of magnetic flux, and representation of canonical commutation relations, 11 pages. 1992.
- # 158: S. Izumiya, W.L. Marar, The Euler number of a topologically stable singular surface in a 3-manifold, 11 pages. 1992.
- # 159: T. Hibi, Cohen-Macaulay types of Cohen-Macaulay complexes, 26 pages. 1992.
- # 160: A. Arai, Properties of the Dirac-Weyl operator with a strongly singular gauge potential, 26 pages. 1992.
- # 161: A. Arai, Dirac operators in Boson-Fermion Fock spaces and supersymmetric quantum field theory, 30 pages. 1992.

Random Parallel Transport on Surfaces of Finite Type, and Relations to Homotopy

by

S. Albeverio^{*,**}, K. Iwata^{*}, T. Kolsrud[‡]

* Fakultät für Mathematik, Ruhr-Universität, D-4630 Bochum 1 (Germany)

SFB237 Essen-Bochum-Düsseldorf

BiBoS Research Centre, D-4800 Bielefeld 1

** CERFIM, Locarno, (Switzerland)

‡ Department of Mathematics, Royal Institute of Technology, S10044 Stockholm (Sweden)

Current address for K.I.: Department of Mathematics, Hokkaido University, Sapporo 060 (Japan)

Abstract

For general surfaces of finite type, probability measures for parallel transport are constructed. Relations to the topology of the surface are pointed out. We also discuss possible loop invariants.

1. Introduction.

In recent years there has been an increasing interest in stochastic differential geometry. One reason for this is the construction of quantized Yang-Mills fields, in Euclidean space-time. In the physics literature, see e.g. [DM] and [Sei], one studies lattice gauge theory, and let the lattice spacing tend to zero to obtain the continuum limit.

In [AH-KH], for a two dimensional case, the continuum limit is constructed directly as a particular case of Markov cosurfaces. The theory was expanded in [K], and similar ideas later appeared in [D], [GKS], [S1,2]. In [Be], [Br] and [F1,2] expectations involving Wilson loops have been studied. In [W], using some ideas in Atiyah's lecture notes [A], heuristic arguments were employed to point out certain connections with topological invariants.

In the present paper we shall only consider the two dimensional case, presenting the "basic construction" initiated in [AH-KH] from a different, simplicial, point of view. (Simplicial approximations to construct Markov random fields on surfaces were considered in [AZ].) This permits us to relate our work with the results of [W] and others.

2. Random Parallel Transport

Let G be a Lie group and let M be a manifold. The action of a connection (one form) on the product bundle $M \times G$ is infinitesimally given by the associated parallel transport. This can be described by a G -valued function, defined for curves in M , and satisfying

$$m(c_1 c_2) = m(c_1) m(c_2), \quad m(c^{-1}) = m(c)^{-1},$$

where $c_1 c_2$ designates that the curve c_2 is followed by c_1 and c^{-1} denotes c with opposite orientation.

To obtain quantised field theories, one way is to construct random fields satisfying, sample wise, these two conditions. Below we shall sketch a general construction for surfaces M of finite type.

The particularity of gauge theory in 2 dimensions makes this possible. This fact has certainly been known among physicists for quite some time. The mathematical formulation and construction goes back to [A, H-K, H], and was given a detailed treatment in Kaufmann [K]. Some of these ideas have appeared later in several papers [D], [F], [G,K,S], [S]. There, the starting point, and the principal object, is the Yang-Mills action functional, and the Feynman path integral formalism. It also appears, among several other subjects, in the recent paper by Witten [W]. In this connection, see also Atiyah's Lincei Lecture Notes [A].

The construction presented below is based on certain inner invariant, or class, functions, and partitionings of M . One also needs to "measure" the plaquettes. Except for this feature, the theory is really "topological" (combinatorial; cf. the comment about the "Hauptvermutung of topology" in Sengupta [S], and [W]), rather than based on the differentiable structure of the manifold. Essentially, this is lattice gauge theory, but the particular family of class functions (convolution semigroups), yields invariance under subdivision. Hence, in physics terminology, the continuum limit is in fact realised.

We remark, finally, that in our 2-dimensional case, if in addition G is simply connected, then $E = M \times G$ is the general principal G -bundle over M . In the general case, one must take into account the topology of the bundle. As remarked in [S], [W], one has to sum over all topologically inequivalent principal G -bundles E over M . For further discussions on this point, refer to the latter article.

Basic construction Let G be a compact topological group, and let $G/\text{Ad } G$ denote the orbits under the inner automorphisms $x \rightarrow g^{-1} x g$, $g \in G$. Functions that are invariant under inner automorphisms (class functions) are called inner invariant. For such functions, the convolution is commutative.

Let now M be an oriented surface which admits a finite triangulation K . The sets of (non-oriented) vertices, bonds and plaquettes are denoted by $\mathcal{V}(K)$, $\mathcal{B}(K)$, and $\mathcal{P}(K)$, respectively. It is understood that the plaquettes are polygons (rather than triangles), hence simply connected. Denote by σ an assignment of orientation to each bond.

We start from a probability space $(\Omega, \mathcal{F}, P_0)$ (which may depend on the oriented triangulation (K, σ)) upon which a collection of random variables $\{X^{(K, \sigma)}(c), c \in \mathcal{B}(K)\}$ are i.i.d. with respect to the Haar measure on G . Each sample $\omega \in \Omega$ describes a connection on the product bundle $M \times G$ and $X^{(K, \sigma)}(c, \omega)$ describes the parallel transport along the oriented bond (c, σ) with respect to the connection ω . (We may choose Ω as the product

of $n = \#\mathcal{B}(K)$ copies of G , P_0 as the n -fold product of Haar measure and realise the X s canonically.)

The gauge group $G^{\#\mathcal{V}(K)}$ is supposed to act on Ω by a measure preserving transformation as

$$(2.1) \quad X^{(K,\sigma)}(c, \omega \cdot g) = g(c(1))^{-1} X^{(K,\sigma)}(c, \omega) g(c(0)),$$

where $c(0)$ and $c(1)$ denote the initial and endpoint of c as given by σ , and where $g(c(0))$ and $g(c(1))$ are the elements of the $\#\mathcal{V}(K)$ -tuple g that correspond to these vertices.

Referring to the orientation of M , we form the oriented boundary $\partial\Delta$ for $\Delta \in \mathcal{P}(K)$. We write, for the appropriate bonds $c_1, \dots, c_k \in \mathcal{B}(K)$,

$$\partial\Delta = \langle c_1 \cdots c_k \rangle$$

to indicate that each bond c_j is given the orientation inherited from $\partial\Delta$, and that no specific initial vertex is preferred. We then consider the $G/\text{Ad } G$ -valued random variable

$$[(X^{(K,\sigma)}(c_1))^{\sigma(c_1)} \cdots (X^{(K,\sigma)}(c_k))^{\sigma(c_k)}].$$

Here, $\sigma(c_j) = 1$ if the orientations of c_j given by σ and that of $\partial\Delta$ coincide, whereas $\sigma(c_j) = -1$ otherwise. Furthermore, the brackets indicate the conjugacy class. Since it is independent of the specific starting curve, it is denoted by $Y^{(K,\sigma)}(\Delta)$. Due to (2.1), it is gauge invariant:

$$Y^{(K,\sigma)}(\Delta, \omega \cdot g) = Y^{(K,\sigma)}(\Delta, \omega), \quad \Delta \in \mathcal{P}(K), g \in G^{\#\mathcal{V}(K)}.$$

To each plaquette we assign its measure, written $|\Delta|$. This could be the area obtained from a Riemannian metric on M , but this is not at all necessary. We shall assume however, that the measure of the closure and the interior of each plaquette coincide.

Let now $(\mu_t)_{t>0}$ denote a convolution semigroup of inner invariant probability measures on G , absolutely continuous w.r.t. Haar measure, and with continuous densities. The same notation for the densities, which are looked upon as class functions, will be used. We shall make use of the following consequence of our assumptions:

$$(2.2) \quad \int_G \mu_s(xyz) \mu_t(x'y^{-1}z') dy = \mu_{s+t}(zz'x').$$

We define the following measure on our probability space:

$$P = P^{(K,\sigma)} = \frac{1}{Z(K,\sigma)} \prod_{\Delta \in \mathcal{P}(K)} \mu_{|\Delta|}(Y^{(K,\sigma)}(\Delta)) P_0,$$

where $Z(K,\sigma)$ is a normalisation constant, and we put $\mu_{|\Delta|} = 1$ if $|\Delta|$ is infinite.

Remarks 1. If the μ_t are symmetric, i.e. invariant under $x \rightarrow x^{-1}$, we can allow non-oriented surfaces M . Then the Y s above take their values in $G/\text{Ad } G/\mathbb{Z}_2$. Abusing notation, this means that $\partial\Delta$ and $(\partial\Delta)^{-1}$ are identified.

2. Consider the case where M is a compact oriented surface of genus $p \geq 1$ from which we have removed s holes. Then

$$Z(K, \sigma) = \int_{G^{2p+s}} \mu_{|M|}(x_1 y_1 x_1^{-1} y_1^{-1} \cdots x_p y_p x_p^{-1} y_p^{-1} z_1 \cdots z_s) dx_1 dy_1 \cdots dx_p dy_p dz_1 \cdots dz_s.$$

Except for the area of M , this is a topological invariant. It is identically one if $s \geq 1$, i.e. when M has a boundary, but not otherwise.

3. The construction is obviously invariant under area preserving homeomorphisms $M \rightarrow M$.

Consistency under subdivision The system is stable under refinements. To explain this, we have to elaborate a bit more. Suppose that K and K' are two triangulations. We write $K \preceq K'$ if each bond in K is a product of bonds in K' . For such ordered pairs, let σ' be an orientation for the bonds of K' . We can construct two systems (Ω, P) and (Ω', P') , one for K and one for K' , together with a map $\pi : \Omega' \rightarrow \Omega$ so that P' and π induce the measure P , i.e. $P = P' \circ \pi^{-1}$. Assume that the sample spaces are realised as products of G , and the random variables X, X' are realised canonically. Enlargement of the triangulation K can be done inductively by the following two kinds of operations.

First, the number of vertices is unchanged, but we add a bond between two vertices. If the variables in Ω' are $\omega' = (\omega_1, \dots, \omega_n, \omega_{n+1})$, where ω_{n+1} refers to the new bond, we simply cancel it, so that $\pi(\omega') = (\omega_1, \dots, \omega_n)$. This is the case if the other bonds have the same orientation in both cases. Otherwise, we map those components ω_j corresponding to bonds where the orientations σ and σ' are opposite to ω_j^{-1} .

The second case is when a bond, say c_1 , corresponding to ω_1 , is divided into two new ones: $c_1 = c_{12}c_{11}$. Then, assuming the orientations σ' at c_{12} and c_{11} are inherited from σ at c_1 (it should be clear how to handle the general case), we map $(\omega_{11}, \omega_{12}, \omega_2, \dots, \omega_n)$ to $(\omega_{12}\omega_{11}, \omega_2, \dots, \omega_n)$. This way, $X(c_1) \circ \pi = X'(c_{12})X'(c_{11})$, i.e. the π respects multiplication of curves.

In both cases one will find, due to (2.2), that π pushes out P' to P .

Relations to homotopy Simple examples show that in general the distributions defined above are not homotopy invariant. A general question is to what extent the distribution of let's say $(X(c_1), \dots, X(c_n))$ depends on c_1, \dots, c_n . We are very far from an answer to this, but already in simple cases very interesting features appear, and we shall look at some examples on the torus $M = T^2$. Let α and β be the homotopy generators. Consider two immersed images of S^1 , a and b , representatives of α and β , and the law of $(X(a), X(b))$.

If a is a simple loop and b a simple loop onto which a further simple loop is added, it turns out that we get a contribution when the orientation of the extra loop is such that the total curve has to cross itself (Fig 1.a). When the orientation is reversed (Fig. 1.b) we do not get any contribution. In the first, but not in the second, case a "tie" is formed, so the constructions separates untied formations from tied ones. (The curves studied are "plats" in the sense of Jones [J].) To be more precise, we should exclude abelian, to obtain effects of this kind. The situation, when b has one tie whose small loop is oriented counterclockwise and a is simple, is described by Fig. 2.a, which is equivalent to the graph in Fig. 2.b, provided that the small loops inscribe the same area.

In this case

$$E[f(X(a), X(b))] = \frac{1}{Z} \int f(x, y) \mu_t(z) \mu_{|M|-t}(xyz^{-2}x^{-1}zy^{-1}) dx dy dz,$$

and the word $(t; z)(|M| - t; xyz^{-2}x^{-1}zy^{-1})$ corresponds to Fig. 2. If we put the small loop going around clockwise on the other side so that the loop b is tied in the different way (Figs. 3) another word appears, namely, $(t; z^{-1})(|M| - t; xyz^{-1}x^{-1}z^2y^{-1})$. This does not give the same distribution as above, so there is also a difference between left and right ties. We give some further examples with two ties. Figures 4.a, b correspond to the word $(t_1; z_1)(t_2; z_2)(|M| - t_1 - t_2; xyz_2^{-1}z_1^{-2}z_2^{-1}x^{-1}z_1z_2y^{-1})$, in that

$$E[f(X(a), X(b))] = \frac{1}{Z} \int f(x, y) \mu_{t_1}(z_1) \mu_{t_2}(z_2) \\ \times \mu_{|M|-t_1-t_2}(xyz_2^{-1}z_1^{-2}z_2^{-1}x^{-1}z_1z_2y^{-1}) dx dy dz_1 dz_2.$$

If instead a left tie and a right tie are simultaneously on the loop b , as in Fig. 5, we get the non-equivalent word $(t_1; z_1)(t_2; z_2^{-1})(|M| - t_1 - t_2; xyz_2^{-1}z_1^{-2}x^{-1}z_2z_1z_2y^{-1})$. On the other hand if both a and b have ties as in Fig. 6, we obtain the word $(t_1; z_1)(t_2; z_2)(|M| - t_1 - t_2; xz_1^{-2}yz_2^{-2}z_1x^{-1}z_2y^{-1})$.

We can also form figures looking like the number eight in various ways. As an example, Fig. 7 corresponds to the word $(t_1; z_1^{-1})(t_2; z_2)(|M| - t_1 - t_2; xyz_2^{-1}z_1^{-1}z_2^{-1}z_1x^{-1}z_1z_2y^{-1})$ which is equivalent to $(t_1; z_1^{-1})(t_2; z_2)(|M| - t_1 - t_2; xyz_1^{-1}z_2^{-1}z_1z_2^{-1}x^{-1}z_2z_1y^{-1})$. Finally, the nested loops in Fig. 8, where the inner one gives the area t_1 , and the area between the inner and the outer loop is t_2 , is associated with the word $(t_1; z_1)(t_2; z_1^{-1}z_2)(|M| - t_1 - t_2; xyz_1^{-1}z_2^{-2}x^{-1}z_2z_1y^{-1})$.

In general, it seems that we can simplify to certain types of graphs and corresponding words. To find more invariant expressions, it would be desirable to get rid of the area parameters above. We therefore propose to study the corresponding infinitesimal expressions, i.e. to differentiate w.r.t. the area parameters, and put them all equal to zero, to obtain infinitesimal Schwinger functions. Already in the case of $SU(2)$, this leads to complicated expressions involving the generator of the semigroup, and products of at least three group characters. In principle, though, combinatorial invariants can be read off from the Wigner-Racah $6j$ symbol (cf. Witten [W]).

Acknowledgements

We gratefully acknowledge very stimulating discussions with C. Becker, Z. Haba, J. Schäfer and B. Zegarliniski.

References

- [A] M. F. Atiyah: The geometry and physics of knots. Cambridge Univ. Press 1990.
- [AHKH] S. Albeverio, R. Høegh-Krohn, H. Holden: Markov cosurfaces and gauge fields. Acta Phys. Austr. Suppl. XXVI, 211-231 (1984)

- [AZ] S. Albeverio, B. Zegarlinski: Construction of convergent simplicial approximations of quantum fields on Riemannian manifolds. *Comm. Math. Phys.* 132, 39-72 (1990).
- [Be] C. Becker: Wilson loops in two-dimensional space-time regarded as generalized white noise. SFB 237 Preprint September 1991.
- [Br] N. E. Bralic: Exact computation of loop averages in two-dimensional Yang-Mills theory. *Phys. Rev. D* 22, 3090-3103 (1980).
- [D] B. Driver: YM_2 : Continuum expectation, lattice convergence and lassos; *Comm. Math. Phys.* 123, 576-616 (1989)
- [DM] H. G. Dosch, V.F. Müller: Lattice gauge theory in two spacetime dimensions. *Fort. Phys.* 27, 547-559 (1979).
- [F1] D.S.Fine, Quantum Yang-Mills theory on the two sphere. *Commun. Math. Phys.* 134, 273-292 (1990).
- [F2] D.S.Fine, Quantum Yang-Mills on a Riemann surface. *Commun. Math. Phys.* 140, 321-338 (1991).
- [GKS] L.Gross, C.King, A.Sengupta: Two dimensional Yang-Mills theory via stochastic differential equations; *Ann. Phys.* 194, 65-112 (1989)
- [J] V. Jones: Hecke algebra representations of braid groups and link polynomials. *Ann. Math.* 126, 335-388 (1987).
- [K] A. Kaufmann: Stetigkeit von Gruppenwertigen Stochastischen Koflächen. Diplomarbeit, Bochum 1986.
- [S1] A. Sengupta: The Yang-Mills measure for S^2 . To appear in *J. Funct. Anal.*
- [S2] A. Sengupta: Quantum gauge theory on compact surfaces. Princeton University Preprint 1991.
- [Sei] E. Seiler: Gauge theories as a problem of constructive quantum field theory and statistical mechanics; *Lect. Notes Phys.* 159, Springer 1982
- [W] E. Witten: On quantum gauge theories in two dimensions. *Comm. Math. Phys.* 141, 153-209 (1991).




Fig. 1.a Tied loop




Fig. 1.b Untied loop




Fig. 2.a




Fig. 2.b




Fig. 3.a




Fig. 3.b




Fig. 4.a




Fig. 4.b




Fig. 5




Fig. 6




Fig. 7




Fig. 8