



Title	Vanishing theorem for the tempered distributions
Author(s)	Honda, N.
Citation	Hokkaido University Preprint Series in Mathematics, 205, 1-11
Issue Date	1993-06
DOI	10.14943/83349
Doc URL	<a href="http://hdl.handle.net/2115/68951">http://hdl.handle.net/2115/68951</a>
Type	bulletin (article)
File Information	pre205.pdf



[Instructions for use](#)

**VANISHING THEOREM FOR THE  
TEMPERED DISTRIBUTIONS**

**Naofumi Honda**

**Series #205. June 1993**

HOKKAIDO UNIVERSITY  
PREPRINT SERIES IN MATHEMATICS

- # 178: T. Nakazi, Toeplitz operators and weighted norm inequalities, 17 pages. 1993.
- # 179: O. Ogurusu, Existence and structure of infinitely degenerate zero-energy ground states of a Wess-Zumino type model in supersymmetric quantum mechanics, 26 pages. 1993.
- # 180: O. Ogurusu, Ground state of a spin  $1/2$  charged particle in an even dimensional magnetic field, 9 pages. 1993.
- # 181: K. Sugano, Note on H-separable Galois extension, 6 pages. 1993.
- # 182: M. Yamada, Distance formulas of asymptotic Toeplitz and Hankel operators, 13 pages. 1993
- # 183: G. Ishikawa, T. Ozawa, The genus of a connected compact real algebraic surface in the affine three space, 11 pages. 1993.
- # 184: T. Hibi, Canonical modules and Cohen-Macaulay types of partially ordered sets, 6 pages. 1993.
- # 185: Y. Giga, K. Yama-uchi, On a lower bound for the extinction time of surfaces moved by mean curvature, 16 pages. 1993.
- # 186: Y. Kurokawa, On functional moduli for first order ordinary differential equations, 9 pages. 1993.
- # 187: O. Ogurusu, Unitary equivalence between a spin  $1/2$  charged particle in a two-dimensional magnetic field and a spin  $1/2$  neutral particle with an anomalous magnetic moment in a two-dimensional electric field, 4 pages. 1993.
- # 188: A. Jensen, T. Ozawa, Existence and non-existence results for wave operators for perturbations of the laplacian, 30 pages. 1993.
- # 189: T. Nakazi, Multipliers of invariant subspaces in the bidisc, 12 pages. 1993.
- # 190: S. Izumiya, G.T. Kossioris, Semi-local classification of geometric singularities for Hamilton-Jacobi equations, 24 pages. 1993.
- # 191: N. Hayashi, T. Ozawa, Finite energy solutions of nonlinear Schrödinger equations of derivative type, 21 pages. 1993.
- # 192: J. Seade, T. Suwa, A residue formula for the index of a holomorphic flow, 22 pages. 1993.
- # 193: H. Kubo, Blow-up of solutions to semilinear wave equations with initial data of slow decay in low space dimensions, 8 pages. 1993.
- # 194: F. Hiroshima, Scaling limit of a model of quantum electrodynamics, 52 pages. 1993.
- # 195: T. Ozawa, Y. Tsutsumi, Global existence and asymptotic behavior of solutions for the Zakharov equations in three space dimensions, 34 pages. 1993.
- # 196: H. Kubo, Asymptotic behaviors of solutions to semilinear wave equations with initial data of slow decay, 25 pages. 1993.
- # 197: Y. Giga, Motion of a graph by convexified energy, 32 pages. 1993.
- # 198: T. Ozawa, Local decay estimates for Schrödinger operators with long range potentials, 17 pages. 1993.
- # 199: A. Arai, N. Tominaga, Quantization of angle-variables, 31 pages. 1993.
- # 200: S. Izumiya, Y. Kurokawa, Holonomic systems of Clairaut type, 17 pages. 1993.
- # 201: K.-S. Saito, Y. Watatani, Subdiagonal algebras for subfactors, 7 pages. 1993.
- # 202: K. Iwata, On Markov properties of Gaussian generalized random fields, 7 pages. 1993.
- # 203: A. Arai, Characterization of anticommutativity of self-adjoint operators in connection with Clifford algebra and applications, 13 pages. 1993.
- # 204: J. Wierzbicki, An estimation of the depth from an intermediate subfactor, 7 pages. 1993.

# VANISHING THEOREM FOR THE TEMPERED DISTRIBUTIONS

NAOFUMI HONDA

University of Hokkaido

## 0. Introduction

The functor  $\text{RH}(\cdot)$  which gives the Riemann-Hilbert correspondence was constructed and deeply studied by Kashiwara [K 3]. This functor plays an important role in the construction of the tempered micro-localization functor due to Andronikof ([A 1, A 2]). The tempered micro-localization functor in Gevrey classes is also constructed by the author of this paper [H 2] using  $\text{RH}^{(s)}(\cdot)$ . It is an important problem to study the vanishing of the higher cohomology groups of  $\text{RH}(\mathbb{C}_\Omega)$  ( $\text{RH}^{(s)}(\mathbb{C}_\Omega)$ ) with  $\Omega$  a pseudo-convex open set, because it is deeply connected with the concentration of cohomology groups of the tempered micro-localization functor (see section 3). We also need the vanishing of cohomology groups to study the tempered (Gevrey class) micro-localization functor with Čech representation.

Let  $\Omega \subset X = \mathbb{C}^n$  be an open set and  $s \in (1, \infty)$  or  $\phi$ . We denote by  $\mathbb{C}_\Omega$  a constructible sheaf whose stalk is  $\mathbb{C}$  in  $\Omega$  and 0 outside of  $\Omega$ .

**Theorem 0.1.** *Assume  $\Omega$  is a relative compact pseudo-convex open set. Then the complex of sheaves  $\text{RH}^{(s)}(\mathbb{C}_\Omega)$  is concentrated degree 0, and we have*

$$H^k(\mathbb{R}\Gamma(X, \text{RH}^{(s)}(\mathbb{C}_\Omega))) = 0 \quad k \neq 0.$$

We give the definitions and notations of Theorem 0.1 in the next section, and some applications in section 3. The proof of the main theorem is given in section 2.

### 1. Notations

Let  $M = \mathbb{R}^n$  with a coordinate  $(x_i)$  and  $s \in (1, \infty)$  or  $\phi$ . We denote by  $\mathcal{E}(\mathcal{D})$  the sheaf of infinitesimal differentiable functions (with compact support). We set

$$\|f\|_K^{l,s} = \sup_{x \in K, \alpha} \frac{|D^\alpha f|}{|\alpha|^{l(s)} l^{|\alpha|}} \quad (s \neq \phi)$$

for a compact set  $K \subset M$  and  $l > 0$ .

We define the sheaf  $\mathcal{E}^{(s)}$  ( $\mathcal{D}^{(s)}$ ) of ultradifferentiable functions of Beurling class  $(s)$  as follows. For an open set  $U$  and  $s \neq \phi$ ,

$$\mathcal{E}^{l,s}(U) := \{f \in \mathcal{E} : \|f\|_K^{l,s} < \infty \text{ for any } K \subset\subset U\},$$

$$\mathcal{D}^{l,s}(U) := \mathcal{E}^{l,s} \cap \mathcal{D}(U),$$

and

$$\mathcal{E}^{(s)} := \varinjlim \mathcal{E}^{l,s},$$

$$\mathcal{D}^{(s)} := \mathcal{E}^{(s)} \cap \mathcal{D}.$$

We set  $\mathcal{E}^{(\phi)} := \mathcal{E}$  and so on.

The sheaf of ultradistributions  $\mathcal{D}_b^{(s)}$  of Beurling class  $(s)$  is the topological dual of  $\mathcal{D}^{(s)}$ . When  $s = \phi$ , this is nothing but the sheaf of distributions by definition. For the details, refer to Komatsu [Ko 2, Ko 3].

Let  $U$  be an open set of  $M$  and  $\mathbb{C}_U$  a constructible sheaf whose stalk is  $\mathbb{C}$  in  $U$  and 0 outside of  $U$ . We define the sheaf  $\text{TH}^{(s)}(\mathbb{C}_U)$  of tempered ultradistributions as follows. For any open set  $V \subset M$

$$\text{TH}^{(s)}(\mathbb{C}_U)(V) := \{u \in \mathcal{D}_b^{(s)}(U \cap V) : \text{there exists a } u' \in \mathcal{D}_b^{(s)}(V) \text{ satisfying } u = u'|_{U \cap V}\}.$$

We remark that  $\text{TH}(\cdot) := \text{TH}^{(\phi)}(\cdot)$  was constructed by Kashiwara [K 3] for general constructible sheaves.

**Proposition 1.1** (cf. [K 3; Lemma 3.3]). *Let  $U \subset M$  be a relative compact open set. Then we have the following equivalent conditions (1), (2) and (3) for any  $s \in (1, \infty)$  and  $u \in \Gamma(U, \mathcal{D}_b^{(s)})$ .*

- (1) *There exists a  $v \in \Gamma(M, \mathcal{D}_b^{(s)})$  satisfying  $u = v$  in  $U$ . (i.e.  $u \in \Gamma(M, \text{TH}^{(s)}(\mathbb{C}_U))$ ).*
- (2) *There exist  $C, l > 0$  satisfying*

$$|\langle u, \psi \rangle| \leq C \| \psi \|_U^{l,s}$$

for any  $\psi \in \mathcal{D}_b^{l,(s)}(U)$ .

- (3) *There exist positive constants  $C, l$  and  $L$  satisfying*

$$|\langle u, \psi \rangle| \leq C \sup_{x \in U, \alpha} \frac{\exp(L \text{dist}(CU, x)^{\frac{-1}{s-1}}) |D^\alpha f|}{|\alpha|^{l(s)} l^{|\alpha|}}$$

for any  $\psi \in \mathcal{D}_b^{l,(s)}(U)$ .

In the case of  $s = \phi$ , we have the same proposition for the tempered distributions [K 3; Lemma 3.3].

Finally we introduce the functor  $\text{RH}^{(s)}(\cdot)$ . Let  $X$  be a complex manifold. We denote by  $\bar{X}$  the complex conjugate space of  $X$ .

$$\text{RH}^{(s)}(\mathbb{C}_U) := \mathbb{R}\text{Hom}_{\mathcal{D}_{\bar{X}}}(\mathcal{O}_{\bar{X}}, \text{TH}^{(s)}(\mathbb{C}_U))$$

where  $\mathcal{D}_{\bar{X}}$  is the sheaf of linear differential operators with holomorphic coefficients in  $\bar{X}$  and  $\mathcal{O}_{\bar{X}}$  is the sheaf of holomorphic functions in  $\bar{X}$  (i.e. Cauchy-Riemann system of  $X^{\mathbb{R}}$  as  $\mathcal{D}_{X \times \bar{X}}$  module).

## 2. Proof of the main theorem

In this section, we give the proof of Theorem 0.1 in the case of  $s \neq \phi$ . Since the proof of the vanishing of  $\text{RH}^{\phi}(\mathbb{C}_{\Omega})$  goes in the similar way, we omit it.

We make a several preparation. Let  $s > 1$  and  $Q_l(\zeta)$  ( $l > 0$ ) entire functions in  $\mathbb{C} = (\zeta)$  satisfying the following conditions. There exists a function  $l' : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  with  $l'(l) \rightarrow \infty$  ( $l \rightarrow \infty$ ), and there exist positive constants  $B_l$  and  $C_l$ , we have estimates

$$|Q_l(\zeta)| \leq B_l \exp(l|\zeta|^{\frac{1}{2s}}) \quad \zeta \in \mathbb{C},$$

and

$$|Q_l(\zeta)| \geq C_l \exp(l'(l)|\zeta|^{\frac{1}{2s}}) \quad \text{Re } \zeta \geq 0.$$

We can concretely construct a family of entire functions satisfying the above conditions ( see [K 1; Theorem 10.1]). Our first step is to construct a differential operator of infinite order which possesses an elliptic property. Let  $X = \mathbb{C}^n = (\zeta_1, \dots, \zeta_n)$  with a coordinate  $(\xi_1 + i\eta_1, \dots, \xi_n + i\eta_n)$ . We set

$$P_l(\zeta) := Q_l(\zeta_1^2 + \dots + \zeta_n^2).$$

Then  $P_l(\zeta)$  satisfies the following estimations.

$$|P_l(\zeta)| \leq B_l \exp(l|\zeta|^{\frac{1}{2}}) \quad \zeta \in \mathbb{C}^n,$$

and

$$|P_l(\zeta)| \geq C_l \exp(l'(l)|\zeta|^{\frac{1}{2}}) \quad \frac{1}{\sqrt{2}}|\zeta| \geq |\eta|.$$

We consider a kernel function in  $X = \mathbb{C}^n = (z_1 = x_1 + iy_1, \dots, z_n = x_n + iy_n)$  as

$$K_l(z) := \int_{\mathbb{R}^n} \frac{e^{-iz\zeta}}{P_l(\zeta)} d\zeta \quad z \in \mathbb{R}^n.$$

It is easy to see  $K_l(x)$  is an ultradifferentiable function of Gevery order ( $s$ ) in  $\mathbb{R}^n$ . Moreover we have the following lemma.

**Lemma 2.1.** *There exist  $\sigma > 0$  and a function  $l' : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  with  $l'(l) \rightarrow \infty$  ( $l \rightarrow \infty$ ),  $K_l(z)$  is holomorphic in  $T_\sigma := \{z; \sigma|x| > |y|\}$  and satisfies the estimate*

$$\sup_{z \in T_\sigma} \left| \frac{l'(l)^{|\alpha+\beta|} D_x^\alpha D_y^\beta K_l(z)}{|\alpha + \beta|!^s} \right| < \infty.$$

*Proof.* The first assertion of the lemma is easily shown by changing the path of the integral as

$$\eta = \frac{-x}{\sqrt{2}|x|} |\xi|.$$

To obtain the estimate of the lemma, we remark the inequality

$$t^n \exp(-lst^{\frac{1}{s}}) \leq l^{-sn} n!^s,$$

for any  $t > 0$  and  $n \in \mathbb{N}$ . Since  $K_l(z)$  is holomorphic in  $T_\sigma$ , it is enough to show the estimates for  $(\frac{\partial}{\partial z})^\alpha K_l(z)$ , and this is a direct calculation.

We remark that  $P_l(\frac{\partial}{\partial z})$  is a partial differential operator of infinite order with Gevrey growth ( $s$ ) and satisfies

$$P_l\left(\frac{\partial}{\partial x}\right)K_l(x) = \delta(x).$$

Let  $U \subset \mathbb{R}^n$  be a relative compact open set and  $u \in \mathcal{D}_b^{(s)}(\mathbb{R}^n)$  with  $\text{supp}(u) \subset \partial U$ . We define an ultradistribution in  $\mathbb{R}^n$  as

$$g(x) := K_l * u \in \mathcal{D}_b^{(s)}(\mathbb{R}^n).$$

**Lemma 2.2.** *If we take a sufficiently large  $l$ , we have*

- (1)  $g(x)$  is an ultradifferentiable function of Gevrey class ( $s$ ) in  $\mathbb{R}^n$ ,
- (2) and,  $g$  is an analytic function outside of  $\partial U$  and can be holomorphically extended over  $W_\sigma = \{z \in X : |y| < \text{dist}(\partial U, x)\}$  for some  $\sigma > 0$ . Moreover any derivative of  $g(z)$  is bounded in  $W_\sigma$ .

*Proof.* On account of the estimation of  $K_l$  in Lemma 2.1, we easily obtain the first assertion of the lemma. Since for any point  $(x, y) \in W_\sigma$

$$\text{supp}(u) - (x, y) \subset \subset T_\sigma,$$

and  $K_l(z)$  is holomorphic in  $T_\sigma$ ,  $g$  can be holomorphically extended over  $W_\sigma$ . Moreover  $K_l(z)$  can be extended over  $\mathbb{C}^n$  as an ultradifferentiable function preserving Gevrey growth order. It is easy to see  $g$  is bounded in  $W_\sigma$ .

Let  $\Omega$  a relative compact pseudo-convex open set and  $\phi = |z|^2$ . We denote by  $L_{(p,q)}^2(\Omega)$  (resp.  $C_{(p,q)}^\infty(\Omega)$ ,  $\text{TH}_{(p,q)}^{(s)}(\mathbb{C}_\Omega)(X)$ ) the space of  $(p, q)$  forms with coefficients in  $L^2(\Omega)$  (resp.  $C^\infty(\Omega)$ ,  $\text{TH}^{(s)}(\mathbb{C}_\Omega)(X)$ ).  $L_{(p,q)}^2(\Omega)$  is Hilbert space equipped with a norm

$$\|f\|_{L^2(\Omega)}^2 := \sum_{I, J} \int_{\mathbb{C}^n} |f_{I, J}|^2 dz \wedge d\bar{z}.$$

We denote by  $\theta : C_{(p,q+1)}^\infty(\Omega) \rightarrow C_{(p,q)}^\infty(\Omega)$  the formal adjoint operator of  $\bar{\partial}$  which has the concrete expression as

$$\theta f := \sum_{I, J} \sum_{j \in [1, n]} \frac{\partial}{\partial z_j} f_{I, J, k} dz^I \wedge d\bar{z}^J,$$

for any  $f \in C_{(p,q+1)}^\infty(\Omega)$ . From now on, we fix  $\phi = |z|^2$ .

**Proposition 2.3.** Let  $f \in C_{(p,q+1)}^\infty(\Omega)$  and  $u \in C_{(p,q)}^\infty(\Omega)$ . We assume  $u$  satisfies

$$\bar{\partial}u = f,$$

and

$$\theta(e^{-\phi}u) = 0.$$

Then we have the following estimate.

$$\|\delta(z)^i \sum_{|\alpha|=i} D^\alpha u\|_{L^2(\Omega)}^2 \leq C^i (\|u\|_{L^2(\Omega)}^2 + \sum_{|\alpha|=i-1} \|\delta(z)^{|\alpha|} D^\alpha f\|_{L^2(\Omega)}^2) \quad (i \geq 1)$$

where  $C$  is a constant which depends only on  $\Omega$  and  $\delta(z) := \text{Min}(\text{dist}(C\Omega, z), 1)$ .

*Proof.* We assume  $q \geq 1$ . In the case of  $q = 0$ , we can prove the proposition in the similar way as the following proof. Set

$$U_k = \{z \in \Omega; \text{dist}(C\Omega, z) \geq \frac{1}{2^k}\} \quad k \geq 0.$$

We choose  $C^1$  functions  $\{\chi_k(z)\}_{k \geq 0}$  satisfying

$$0 \leq \chi_k \leq 1,$$

$$\chi_k(z) = 1 \quad z \in U_k,$$

$$\text{supp}(\chi_k) \subset U_{k+1}^-$$

and, there exists a positive constant  $C$

$$\sum_{i=0}^n (|\frac{\partial}{\partial z_i} \chi_k| + |\frac{\partial}{\partial \bar{z}_i} \chi_k|) \delta(z) \leq C$$

for any  $k$ . By the assumption, we have

$$\bar{\partial}u = f,$$

$$\theta u = au$$

where  $a$  is a differential operator of order 0 whose coefficients are polynomials of degree 1. We have for any multi index  $\alpha$

$$\bar{\partial}(\chi_k D^\alpha u) = \bar{\partial}\chi_k \wedge D^\alpha u + \chi_k D^\alpha f,$$

$$\theta(\chi_k D^\alpha u) = b D^\alpha u + \chi_k D^\alpha (au)$$

where  $b$  is a differential operator of degree 0. Remark that

$$\delta(z)b : L_{(p,q+1)}^2(\Omega) \rightarrow L_{(p,q)}^2(\Omega)$$



is a bounded operator. We have for any multi index  $\alpha$  ( $|\alpha| \geq 1$ ),

$$\begin{aligned} \|\delta^{|\alpha|+1} \bar{\partial} \chi_k D^{\alpha+1} u\|_{L^2(\Omega)}^2 &\leq \|\delta^{|\alpha|+1} \bar{\partial} \chi_k \wedge D^\alpha u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|+1} \chi_k D^\alpha f\|_{L^2(\Omega)}^2 \\ &\leq C_1 (\|\delta^{|\alpha|} D^\alpha u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|} D^\alpha f\|_{L^2(\Omega)}^2). \end{aligned}$$

Similary we have

$$\begin{aligned} \|\delta^{|\alpha|+1} \theta(\chi_k D^\alpha u)\|_{L^2(\Omega)}^2 &\leq C_2 (\|\delta^{|\alpha|} D^\alpha u\|_{L^2(\Omega)}^2 \\ &\quad + \sum_{|\beta|=|\alpha|-1, \beta \prec \alpha} \|\delta^{|\beta|} D^\beta u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|} D^\alpha u\|_{L^2(\Omega)}^2). \end{aligned}$$

Therefore we obtain the following estimation.

$$\begin{aligned} &\sum_{1 \leq i \leq n} (\|\delta^{|\alpha|+1} \frac{\partial}{\partial z_i} \chi_k D^{\alpha+1} u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|+1} \frac{\partial}{\partial \bar{z}_i} \chi_k D^{\alpha+1} U\|_{L^2(\Omega)}^2) \\ &\leq 4 (\|\delta^{|\alpha|+1} \bar{\partial} \chi_k D^{\alpha+1} u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|+1} \bar{\partial} \chi_k D^{\alpha+1} u\|_{L^2(\Omega)}^2) \\ &\leq C_3 (\|\delta^{|\alpha|} D^\alpha u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|} D^\alpha f\|_{L^2(\Omega)}^2 \\ &\quad + \sum_{|\beta|=|\alpha|-1, \beta \prec \alpha} \|\delta^{|\beta|} D^\beta u\|_{L^2(\Omega)}^2) \end{aligned}$$

where  $C_3$  is a positive constat depending only on  $\Omega$ . Taking limit  $k \rightarrow \infty$ , we have for any multi index  $\alpha$

$$\begin{aligned} &\sum_{1 \leq i \leq n} (\|\delta^{|\alpha|+1} \frac{\partial}{\partial x_i} D^{\alpha+1} u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|+1} \frac{\partial}{\partial \bar{x}_i} D^{\alpha+1} u\|_{L^2(\Omega)}^2) \\ &\leq C_4 (\|\delta^{|\alpha|} D^\alpha u\|_{L^2(\Omega)}^2 + \|\delta^{|\alpha|} D^\alpha f\|_{L^2(\Omega)}^2 \\ &\quad + \sum_{|\beta|=|\alpha|-1, \beta \prec \alpha} \|\delta^{|\beta|} D^\beta u\|_{L^2(\Omega)}^2). \end{aligned}$$

By induction and the above inequality, we obtain the disired estimation.

Now we show the proof of the main theorem.

*proof.* Consider the complex

$$\mathrm{TH}_{(p,q)}^{(s)}(\mathbb{C}_\Omega)(X) \xrightarrow{\bar{\partial}} \mathrm{TH}_{(p,q+1)}^{(s)}(\mathbb{C}_\Omega)(X) \xrightarrow{\bar{\partial}} \mathrm{TH}_{(p,q+2)}^{(s)}(\mathbb{C}_\Omega)(X).$$

Given  $f \in \mathrm{TH}_{(p,q+1)}^{(s)}(\mathbb{C}_\Omega)(X)$  with  $\bar{\partial} f = 0$ . We take  $f' \in \mathcal{D}_b^{(s)}(X)$  satisfying

$$f'|_U = f|_U,$$

$$\mathrm{supp}(f') \subset \bar{U},$$

and

$$\mathrm{supp}(\bar{\partial} f') \subset \partial U.$$

For a sufficiently large  $l > 0$ , we set

$$h := K_l * \bar{\partial} f',$$

where  $K(z, \bar{z})$  is a kernel function associated with  $P_l(\zeta, \bar{\zeta})$  constructed in the beginning of this section. Remark that, on account of Lemma 2.1,  $K_l(z, \bar{z})$  is a real analytic function in the underlying space  $X^{\mathbb{R}} = \mathbb{R}^{2n}$  except for the origin and can be holomorphically extended over the domain  $T_\sigma \subset X \times \bar{X}$  described in the same lemma. Therefore  $h$  satisfies the following.

- (1)  $\bar{\partial} h = k_l * \bar{\partial} \bar{\partial} f' = 0$  in  $X^{\mathbb{R}}$ .
- (2)  $h$  is a  $(p, q + 2)$  form with coefficients in ultradifferentiable functions of Gevrey order  $(s)$  in  $X^{\mathbb{R}}$ .
- (3) There exists a positive constant  $C$ , we have the estimate

$$\left| \frac{\partial^\alpha}{\partial z^\alpha} \frac{\partial^\beta}{\partial \bar{z}^\beta} h \right| \leq C \frac{|\alpha + \beta|!}{\text{dist}(C\Omega, z)^{|\alpha + \beta|}} \quad (z, \bar{z}) \in X^{\mathbb{R}} \subset X \times \bar{X}$$

on account of Lemma 2.2.

We fix a relative compact convex open set  $W \subset X^{\mathbb{R}}$  with  $\Omega \subset\subset W$ . By the theorem of Hörmander [Hö; chapter 4], we can find  $g \in C_{p, q+1}^\infty(W)$  satisfying

$$\bar{\partial} g = h \quad \text{in } W,$$

and

$$\theta(e^{-\phi} u)|_U = 0$$

where  $\phi = |z|^2$ . Then by proposition 2.3, we have

$$\|\delta(z)^{|\alpha + \beta|} \frac{\partial^\alpha}{\partial z^\alpha} \frac{\partial^\beta}{\partial \bar{z}^\beta} g\|_{L^2(\Omega)} \leq C^{|\alpha + \beta|} |\alpha + \beta|! \quad (z \in \Omega)$$

with a positive constant  $C$ . Since

$$P_l\left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}}\right) = \Sigma C_{\alpha\beta} \frac{\partial^\alpha}{\partial z^\alpha} \frac{\partial^\beta}{\partial \bar{z}^\beta}$$

is a differential operator of infinite order with Gevrey class  $(s)$ , there exists a positive constants  $l$  satisfying

$$|C_{\alpha\beta}| \leq \frac{l^{|\alpha + \beta|}}{|\alpha + \beta|!^s}.$$

Therefore we obtain

(2.1)

$$\begin{aligned} \left\| \left( P_l\left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}}\right) g \right) \exp(-l' \delta(z)^{\frac{-1}{s-1}}) \right\|_{L^2(\Omega)} &\leq \Sigma \frac{(Cl)^{|\alpha + \beta|} |\alpha + \beta|!}{|\alpha + \beta|!^s} \sup_{z \in \Omega} |\delta(z)^{-|\alpha + \beta|} \exp(-l' \delta(z)^{\frac{-1}{s-1}})| \\ &\leq C_1 \Sigma \frac{(Cl)^{|\alpha + \beta|}}{|\alpha + \beta|!^{(s-1)}} \frac{|\alpha + \beta|!^{(s-1)}}{(l')^{|\alpha + \beta|}} \\ &\leq C_2 \Sigma \left(\frac{Cl}{l'}\right)^{|\alpha + \beta|} < \infty \end{aligned}$$

for a sufficiently large  $l'$ .

We set  $f''(z) := P_l(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}})g$ . Since  $W$  is convex and

$$\bar{\partial}(f' - f'') = \bar{\partial}f' - P_l \bar{\partial}g = \bar{\partial}f' - P_l h = \bar{\partial}f' - P_l(K_l * \bar{\partial}f') = 0$$

on account of Palamndov [P; p.300 Theorem 1], we can find  $u' \in \mathcal{D}_b^{(s)}(W)$  with  $\bar{\partial}u' = f' - f''$ . On the other hand, since  $\bar{\partial}f''|_\Omega = \bar{\partial}f'|_\Omega = 0$  and  $f''$  satisfies the esitimation (2.1), we can find  $u'' \in L_{(p,q)}^{loc}(\Omega)$  satisfying

$$f'' = \bar{\partial}u'' \quad \text{in } \Omega$$

and the estimation

$$\|u'' \frac{\exp(-l'\delta(z)^{\frac{-1}{s-1}})}{(1+|z|^2)}\|_{L^2(\Omega)} < \infty$$

due to [Hö; Theorem 4.4.2]. The above inequality means, in particular,  $u'' \in \text{TH}_{(p,q)}^{(s)}(\mathbb{C}_\Omega)(X)$ . Thus  $u = u' + u'' \in \text{TH}_{(p,q)}^{(s)}(\mathbb{C}_\Omega)(X)$  satisfies

$$\bar{\partial}(u' + u'')|_U = ((f' - f'') + f'')|_U = f|_U.$$

### 3. The application of the main theorem

One of the application of the main theorem, we give the vanishing of distribution solutions along the submanifolds which are micro-locally weakly pseudo-convex.

Let  $M$  be a real analytic submanifold in a complex manifold  $X$ , and  $\pi : T^*X \rightarrow X$  (resp.  $\pi : T_M^*X \rightarrow X$ ) a cotangent bundle of  $X$  (resp. conormal bundle of  $M$ ). We set for  $p \in T^*X$ ,

$$\begin{aligned} E(p) &= T_p(T^*X), \\ \lambda_M(p) &= T_p(T_M^*X), \\ \lambda_0(p) &= T_p(\pi^{-1}\pi(p)), \\ \rho_M(p) &= \lambda_M(p) \cap i\lambda_M(p), \\ \delta(p) &= \dim_{\mathbb{C}}(\lambda_M(p) \cap i\lambda_M(p) \cap \lambda_0(p)). \end{aligned}$$

The vector space  $(E(p), \sigma)$  has a natural complex symplectic structure. For any real Lagrangian vector space  $\lambda \in E(p)^{\mathbb{R}}$ , we denote by  $\lambda^{\rho_M}$  the quotient space

$$((\lambda \cap \rho(p))_M^\perp + \rho_M(p)) / \rho_M(p).$$

We also define the bilinear form  $\gamma_{\lambda_M}$  on  $\lambda_0^{\rho_M}(p)$  by setting for  $(u, v) \in \lambda_0^{\rho_M}(p) \times \lambda_0^{\rho_M}(p)$

$$\gamma_{\lambda_M}(u, v) := \sigma_{\rho_M}(u, \bar{v})$$

where  $\bar{v}$  is the complex conjugate of  $v$  with respect to  $\lambda_M^{\rho_M}(p)$ . We denote by  $S_+(M, p)$  (resp  $S_-(M, p)$ ) the number of positive (resp. negative) eigen values of the hermitian form  $\gamma_{\lambda_M}$ , and define  $S(M, p) = S_+(M, p) - S_-(M, p)$ .

**Theorem 3.1** (cf. [K-S 1; Theorem 11.3.5]). *Let  $p_0 \in T_M^*X \setminus T_X^*X$ . We assume the following conditions.*

- (1)  $\dim_{\mathbb{R}}(\lambda_M(p_0) \cap \nu(p_0)) = 1$  where  $\nu(p_0)$  is a complex line generated by the Euler vector field.
- (2)  $S_-(M, p) - \delta(p)$  is locally constant in a neighborhood of  $p_0$ .

*Then the complex  $T\text{-}\mu_M(\mathcal{O}_X)_{p_0}$  is concentrated degree  $k = \text{codim } M + S_-(M, p_0) - \delta(p_0)$ , and the canonical morphism*

$$H^k T\text{-}\mu_M(\mathcal{O}_X)_{p_0} \rightarrow H^k \mu_M(\mathcal{O}_X)_{p_0}$$

*is injective.*

When  $X$  is the complexification of a real analytic manifold  $M$ , the above theorem has been shown by Maltineau [Ma 1]. Andronikof, in his papers [A 1, A 2], has obtained the edge of the wedge with tempered growth order, and proved the same results in the complex case (i.e.  $M$  is a complex manifold). The first assertion replaced the functor  $T\text{-}\mu_M$  with a micro-localization functor  $\mu_M$  was established by Kashiwara-Schapira [K-S 1]. They proved this to construct a quantized contact transformation cohomologically and reduce the problem to the case  $M$  is a weakly pseudo-convex hypersurface. Therefore the important point is to construct a quantized contact transformation cohomologically for the tempered micro-localization functor. We first study how to construct the quantized contact transformation. Let  $X, Y$  be complex manifolds and  $N \subset Y$ ,  $M \subset X$  real analytic submanifolds,  $\Phi : T^*X \rightarrow T^*Y$  a complex contact transformation which interchange  $(T_M^*X, p_0)$  and  $(T_N^*Y, \Phi(p_0))$  in a neighborhood of  $p_0 \in T_M^*X \setminus T_X^*X$ . We denote by  $Q_1, Q_2$  the first and the second projection from  $X \times Y$  respectively and by  $P_1, P_2$  the first and the second projection from  $T^*(X \times Y)$ . We set  $P_1^a = P_1 \cdot a$  where  $a$  is the anti-podal map on  $T^*X$ . Let  $\Lambda \subset T^*(X \times Y)$  be a complex conic Lagrangian submanifold in a neighborhood of  $(p_0, -\Phi(p_0))$  obtained by taking the image of graph of  $\Phi$  by the anti-podal map on  $T^*X$ . Moreover we may assume  $Z = \pi(\Lambda) \subset X \times Y$  is a smooth complex hypersurface. We denote by  $K$  a pure sheaf  $\mathbb{C}_Z[-1]$ . And if necessarily, we cut the support of  $K$  in a neighborhood of  $\pi(p, -\Phi(p))$  satisfying that  $Q_2^a|_{\text{supp}(K)}$  is proper over  $\text{supp}(\mathbb{C}_N)$ . This does not affect on the following arguments because we consider the problem in the micro-local point of view. To construct the quantized contact transformation, we use direct image and restriction of the tempered micro-localization functor due to Andronikof [A 1, A 2]. We consider the following diagram.

$$\begin{array}{ccc} X & \xleftarrow{Q_1} & X \times (X \times Y) \\ & & \uparrow i \\ & & \Delta_X \times Y \simeq X \times Y \xrightarrow{Q_2} Y. \end{array}$$

$$\begin{aligned}
& (\mathrm{T}\text{-}\mu\mathrm{hom}(K, \mathcal{O}_{X \times Y}) \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/Y})_{(-p_0, \Phi(p_0))} \boxtimes \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_X)_{p_0} \\
& \rightarrow \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M \boxtimes K, \mathcal{O}_{X \times X \times Y})_{(p_0, -p_0, \Phi(p_0))} \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/Y} \\
& \rightarrow \mathcal{D}_{X \times Y \rightarrow X \times X \times Y} \overset{\mathbb{L}}{\otimes}_{\mathcal{D}_{X \times X \times Y}} \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M \boxtimes K, \mathcal{O}_{X \times X \times Y})_{(p_0, -p_0, \Phi(p_0))} \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/Y} \\
& \xrightarrow{i^*} \mathrm{T}\text{-}\mu\mathrm{hom}(i^{-1}(\mathbb{C}_M \boxtimes K), \mathcal{O}_{X \times Y})_{(-p_0, \Phi(p_0))} \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/Y} \\
& \rightarrow \mathcal{D}_{Y \rightarrow X \times Y} \overset{\mathbb{L}}{\otimes}_{\mathcal{D}_{X \times Y}} \mathrm{T}\text{-}\mu\mathrm{hom}(i^{-1}(\mathbb{C}_M \boxtimes K), \mathcal{O}_{X \times Y})_{(-p_0, \Phi(p_0))} \\
& \xrightarrow{f_!} \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{R}Q_{2*}i^{-1}(\mathbb{C}_M \boxtimes K), \mathcal{O}_Y)_{\Phi(p_0)}[-\dim_{\mathbb{C}} X].
\end{aligned}$$

The second map is given by choosing a canonical section  $1_{X \times Y \rightarrow X \times X \times Y} \in \mathcal{D}_{X \times Y \rightarrow X \times X \times Y}$ . Since

$$\mathrm{T}\text{-}\mu\mathrm{hom}(K, \mathcal{O}_{X \times Y}) = \mathcal{E}_{X \times Y}^{\mathbb{R}, f} \otimes_{\mathcal{D}_{X \times Y}} \mathcal{B}_{Z|X \times Y}$$

and  $\mathcal{B}_{Z|X \times Y}$  is a simple holonomic system, we can take a section  $s \in \mathcal{E}_{Z|X \times Y}^{\mathbb{R}, f}$  which generates  $\mathcal{E}_{Z|X \times Y}^{\mathbb{R}, f}$  over  $\mathcal{E}_{X \times Y}^{\mathbb{R}, f}$ . Then we have the morphism

$$\mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_X)_{p_0} \rightarrow (\mathrm{T}\text{-}\mu\mathrm{hom}(K, \mathcal{O}_{X \times Y}) \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/Y})_{(-p_0, \Phi(p_0))} \boxtimes \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_X)_{p_0},$$

by  $u \rightarrow s \boxtimes u$ . Thus we have the cohomological quantized contact transformation map

$$\mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_{X \times Y})_{p_0} \rightarrow \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{R}Q_{2*}i^{-1}(\mathbb{C}_M \boxtimes K), \mathcal{O}_Y)_{\Phi(p_0)}[-\dim_{\mathbb{C}} X].$$

Since  $\mathbb{R}Q_{2*}i^{-1}(\mathbb{C}_M \boxtimes K)$  is a simple sheaf along  $T_N^*Y$  with the shift  $-\frac{1}{2} \dim M - \frac{1}{2}(S(N, p) - S(M, \Phi(p)))$  in a neighborhood  $p_0$ , we have

$$\mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_{X \times Y})_{p_0} \rightarrow \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_N, \mathcal{O}_Y)_{\Phi(p_0)}[d]$$

where  $d := \frac{1}{2}(\dim M - \dim N + S(M, p_0) - S(N, \Phi(p_0)))$ . We can easily check the map constructed above satisfies the composition law, and is isomorphism. Moreover we have a commutative diagram

$$\begin{array}{ccc}
\mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_X)_{p_0} & \longrightarrow & \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_N, \mathcal{O}_Y)_{\Phi(p_0)}[d] \\
\downarrow & & \downarrow \\
\mathbb{R}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_X)_{p_0} & \longrightarrow & \mathbb{R}\mu\mathrm{hom}(\mathbb{C}_N, \mathcal{O}_Y)_{\Phi(p_0)}[d].
\end{array}$$

*proof of Theorem 3.1.* In the same way of the proof of [K-S 1, Theorem 11.3.5], we can find a hypersurface  $N$  which is the boundary of a pseudo-convex open

set  $j : \Omega \rightarrow X$  (in particular,  $S_-(N, q) = 0$ ) and a contact transformation  $\phi$  interchanging  $(T_M^*X, p_0)$  and  $(T_N^*Y, q_0)$  in a neighborhood of  $p_0$ . We obtain

$$\mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_M, \mathcal{O}_X)_{p_0} [\mathrm{codim} M + S_-(M, p_0) - \delta(p_0)] = \mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_N, \mathcal{O}_Y)_{q_0} [1].$$

Since there is a triangle

$$\mathrm{T}\text{-}\mu\mathrm{hom}(\mathbb{C}_N, \mathcal{O}_Y)_{q_0} \rightarrow \mathcal{O}_{\pi(q_0)} \rightarrow \mathrm{RH}(\mathbb{C}_\Omega)_{\pi(q_0)} \xrightarrow{+1},$$

we obtain the first assertion of Theorem 3.1 due to Theorem 0.1. Moreover

$$\mathrm{H}^0(\mathrm{RH}(\mathbb{C}_\Omega))/\mathcal{O} \rightarrow j_*\mathcal{O}/\mathcal{O}$$

is clearly injective, we have injectivity of  $\mathrm{H}^k \mathrm{T}\text{-}\mu_M(\mathcal{O}_X) \rightarrow \mathrm{H}^k \mu_M(\mathcal{O}_X)$ .

#### REFERENCES

- [A 1] E. Andronikof, *Microlocalisation tempérée des distributions et des fonctions holomorphes I*, C.R.Acad.Sci 303, 347-350 (1986).
- [A 2] E. Andronikof, *Microlocalisation tempérée des distributions et des fonctions holomorphes II*, C.R.Acad.Sci 304, 511-514 (1987).
- [A 3] E. Andronikof, *On the  $C^\infty$ -singularities of regular holonomic distributions*, Ann. Inst. Fourier, 42, 695-704 (1992).
- [Hö 1] Hörmander, *An introduction to complex analysis in several variables.*, Van Nostrand, Princeton, (1966).
- [Ho 1] N.Honda, *On the reconstruction theorem of holonomic modules in Gevrey classes*, Publ. RIMS, 27, Kyoto Univ., 923-943 (1991).
- [Ho 2] N.Honda, *Microlocalization in Gevrey classes*, in preparation.
- [Ho 3] N.Honda, *Regularity theorems for holonomic modules*, in preparation.
- [K 1] M.Kashiwara, *On the maximally overdetermined systems of linear differential equations, I*, Publ. RIMS, 10, Kyoto Univ., 563-579 (1975).
- [K 2] M.Kashiwara, *On the holonomic systems of linear differential equations II*, Inventiones Math., 49, 121-135 (1978).
- [K 3] M.Kashiwara, *The Riemann-Hilbert problem for holonomic systems*, Publ. RIMS, 20, Kyoto Univ., 319-365 (1984).
- [K-K 1] M.Kashiwara, T.Kawai, *On the holonomic systems of microdifferential equations III*, Publ. RIMS, 17, Kyoto Univ., 813-979 (1981).
- [K-K 2] M.Kashiwara, T.Kawai, *Second microlocalization and asymptotic expansions*, Lect. Notes Phys. 126, 21-76 (1980).
- [K-S 1] M.Kashiwara, P.Schapira, *Microlocal study of sheaves*, Astérisque, 128 (1985).
- [K-S 2] M.Kashiwara, P.Schapira, *Sheaves on manifolds*, Grundlehren der Math., 292, Springer-Verlag (1990).
- [Ko 1] H.Komatsu, *Untradistributions, I, Structure theorems and a characterizations*, J. Fac. Sci. Univ. Tokyo, Sec. IA, 20, 25-105 (1973).
- [Ko 2] H.Komatsu, *On the regularity of hyperfunction solutions of linear ordinary differential equations with real analytic coefficients*, J. Fac. Sci. Univ. Tokyo, Sec. IA, 20, 107-119 (1973).
- [Ko 3] H.Komatsu, *Untradistributions, II, The kernel theorems and ultradistributions with compact support in a submanifold*, J. Fac. Sci. Univ. Tokyo, Sec. IA, 24, 607-628 (1977).
- [Ma 1] A. Martineau, *Oeuvre de Andre Martineau*, Publ. C.N.R.S., (1977).
- [P 1] V.P.Palamodov, *Linear differential operators with constant coefficients*, Grundlehren der Math., 168, Springer-Verlag, (1970).
- [S-K-K] M.Sato, T.Kawai, M.Kashiwara, *Hyperfunctions and pseudodifferential equations*, Lecture Notes in Math., 287, Springer-Verlag, 265-529 (1973).