



Title	Note on non-commutative local field
Author(s)	Sugano, K.
Citation	Hokkaido University Preprint Series in Mathematics, 213, 1-3
Issue Date	1993-08
DOI	10.14943/83357
Doc URL	http://hdl.handle.net/2115/68959
Type	bulletin (article)
File Information	pre213.pdf



[Instructions for use](#)

Note on non-commutative local field

Kozo Sugano

Series #213. August 1993

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 186: Y. Kurokawa, On functional moduli for first order ordinary differential equations, 9 pages. 1993.
- # 187: O. Ogurusu, Unitary equivalence between a spin $1/2$ charged particle in a two-dimensional magnetic field and a spin $1/2$ neutral particle with an anomalous magnetic moment in a two-dimensional electric field, 4 pages. 1993.
- # 188: A. Jensen, T. Ozawa, Existence and non-existence results for wave operators for perturbations of the laplacian, 30 pages. 1993.
- # 189: T. Nakazi, Multipliers of invariant subspaces in the bidisc, 12 pages. 1993.
- # 190: S. Izumiya, G.T. Kossioris, Semi-local classification of geometric singularities for Hamilton-Jacobi equations, 24 pages. 1993.
- # 191: N. Hayashi, T. Ozawa, Finite energy solutions of nonlinear Schrödinger equations of derivative type, 21 pages. 1993.
- # 192: J. Seade, T. Suwa, A residue formula for the index of a holomorphic flow, 22 pages. 1993.
- # 193: H. Kubo, Blow-up of solutions to semilinear wave equations with initial data of slow decay in low space dimensions, 8 pages. 1993.
- # 194: F. Hiroshima, Scaling limit of a model of quantum electrodynamics, 52 pages. 1993.
- # 195: T. Ozawa, Y. Tsutsumi, Global existence and asymptotic behavior of solutions for the Zakharov equations in three space dimensions, 34 pages. 1993.
- # 196: H. Kubo, Asymptotic behaviors of solutions to semilinear wave equations with initial data of slow decay, 25 pages. 1993.
- # 197: Y. Giga, Motion of a graph by convexified energy, 32 pages. 1993.
- # 198: T. Ozawa, Local decay estimates for Schrödinger operators with long range potentials, 17 pages. 1993.
- # 199: A. Arai, N. Tominaga, Quantization of angle-variables, 31 pages. 1993.
- # 200: S. Izumiya, Y. Kurokawa, Holonomic systems of Clairaut type, 17 pages. 1993.
- # 201: K.-S. Saito, Y. Watatani, Subdiagonal algebras for subfactors, 7 pages. 1993.
- # 202: K. Iwata, On Markov properties of Gaussian generalized random fields, 7 pages. 1993.
- # 203: A. Arai, Characterization of anticommutativity of self-adjoint operators in connection with Clifford algebra and applications, 13 pages. 1993.
- # 204: J. Wierzbicki, An estimation of the depth from an intermediate subfactor, 7 pages. 1993.
- # 205: N. Honda, Vanishing theorem for the tempered distributions, 11 pages. 1993.
- # 206: T. Hibi, Betti number sequences of simplicial complexes, Cohen-Macaulay types and Möbius functions of partially ordered sets, and related topics, 25 pages. 1993.
- # 207: A. Inoue, Regularly varying correlations, 23 pages. 1993.
- # 208: S. Izumiya, B. Li, Overdetermined systems of first order partial differential equations with singular solution, 9 pages. 1993.
- # 209: T. Hibi, Hochster's formula on Betti numbers and Buchsbaum complexes, 7 pages. 1993.
- # 210: T. Hibi, Star-shaped complexes and Ehrhart polynomials, 5 pages. 1993.
- # 211: S. Izumiya, G. T. Kossioris, Geometric singularities for solutions of single conservation laws, 28 pages. 1993.
- # 212: A. Arai, On self-adjointness of Dirac operators in Boson-Fermion Fock spaces, 43 pages. 1993.

Throughout this paper A will be a division ring with non-trivial valuation v , and C will be the center of A . A has the completion with respect to the v -topology, which is also a division ring. We will denote it by A^* . For each division subring B of A the closure of B in A^* with respect to the v -topology will be also denoted by B^* . B^* is isomorphic to the completion of B as topological ring.

The aim of this paper is to show that B^* coincides with the double centralizer of B in A^* for each division C -subalgebra B of A , in the case where A is finite over C and A^* is a local field with respect to v .

In this paper we will use the same terminology as [4] and [6]. In particular for each division subring B of A we write

$$O(B) = \{x \in B : v(x) \leq 1\}, \quad P(B) = \{x \in B : v(x) < 1\} \text{ and}$$

Since v is non-trivial, A is locally compact if and only if $O(A)$ is compact (See § 17.5 [4]). In this case A is complete (See the proof of Proposition 17.5 b [4]).

Let v be non-archimedean and B an arbitrary division subring of A . $O(B)$ is a local ring with the maximal ideal $P(B)$. Hence $O(B)/P(B)$ is a division ring, which will be denoted by $E(B)$. Write $E(B) = K$ and $E(A) = E$. Then K is a division subring of E . We will write $f_r(A/B) = [E:K]_r$, $f_l(A/B) = [E:K]_l$ and $e(A/B) = [v(A^\circ) : v(B^\circ)]$, where A° and B° are the unit groups of A and B , respectively. In the case where $[E:K]_l = [E:K]_r$, we will write $f(A/B)$ in stead of $f_r(A/B)$ or $f_l(A/B)$. The next lemma is well known in the case where A is a commutative field.

Lemma 1. Let A, B and v be as above, and assume that both $e(A/B)$ and $f_r(A/B)$ are finite. Then we have $e(A/B)f_r(A/B) \leq [A:B]_r$

Proof. Since $v(ab) = v(a)v(b) = v(b)v(a) = v(ba)$ for any $a, b \in A$, we can follow the same lines as the proof of Theorem 4.5 Chap. 2 [2].

Now suppose that A is finite dimensional over C . Then for any division C -subalgebra B of A we have $B = V_A(V_A(B))$, and $V_A(B)$ is finite dimensional division C -subalgebra (See e.g., Theorem 12.7 [4]). Therefore A is an H -separable extension of every division C -subalgebra B of A by Theorem 1 [5]. Moreover $(A^*/A, B^*/B)$ have the centralizer property in the sense of [6], and we have $[A^*:B^*] = [A^*:B^*]$ and $B^* \supset B^*$, where $B^* = V_A(V_A(B))$ by Theorem 1.3 (4) [6]. In particular C^* is contained in the center of A^* . C is an infinite field, otherwise A must be commutative by Wedderburn's Theorem. Therefore, if furthermore v is non-archimedean and $O(A^*)$ is compact in the v -topology, for any division C -subalgebra B of A we have $e(B^*/C^*)f(B^*/C^*) = [B^*:C^*] < \infty$ by Proposition 17.7 [4], since $O(B^*)$ is also compact and C^* is infinite and closed in B^* .

Now we are ready to have our main theorem.

Theorem 1. Let A be a division ring with non-trivial non-archimedean valuation v . Assume A is finite dimensional over its center C , and $O(A^*)$ is compact in the v -topology. Then we have

(1) The closure C^* of C in A^* coincides with the center of A^* , and we have $[A^*:C^*] = [A:C]$

(2) For any division C -subalgebra B of A , we have $B^* = V_A(V_A(B))$, $A^* = B^*A = AB^*$ and $B^* \cap A = B$.

Proof. (1). Write $C^* = V_A(V_A(C))$. C^* coincides with the center of A^* by Proposition 1.3 [6]. Since v is non-archimedean, we have

$$e(C^*/C) = e(A^*/A) = 1 \text{ and } f(A^*/A) = f(C^*/C) = 1$$

(See § 17.4 [4]). Since

$$e(A^*/C) = e(A^*/C^*)e(C^*/C) = e(A^*/A)e(A/C),$$

we have $e(A/C) = e(A^*/C^*)$. Similarly we have $f(A^*/C^*) = f(A/C)$. Then by Proposition 17.7 (1) [4] and Lemma 1 we have

$$[A^*:C^*] = e(A^*/C^*)f(A^*/C^*) = e(A/C)f(A/C) \leq [A:C].$$

But by Theorem 1.7 (3) [6] we have $[A:C] = [A^*:C^*] \leq [A^*:C^*]$. Hence we have $[A^*:C^*] = [A^*:C^*]$ and $C^* = C^*$.

(2). Put $B^* = \bigvee_{A^*} (V_{A^*} \cdot (B))$. B^* is complete, and $[A^* : B^*] = [A : B]$ by Theorem 1.7 (3) and (4) [6]. On the other hand since $e(B^*/B) = f(B^*/B) = 1$, we have

$$e(A^*/B^*)f(A^*/B^*) = e(A/B)f(A/B) \leq [A : B]$$

$$e(B^*/C^*)f(B^*/C^*) = e(B/C)f(B/C) \leq [B : C]$$

where the inequalities \leq 's are due to Lemma 1. Then we have

$$[A^* : C^*] = e(A^*/C^*)f(A^*/C^*) = e(A^*/B^*)e(B^*/C^*)f(A^*/B^*)f(B^*/C^*)$$

$$\leq [A : B][B : C] = [A : C] = [A^* : C^*]$$

This means that $e(A/B)f(A/B) = [A : B]$ and $e(B/C)f(B/C) = [B : C]$. The latter equality shows $[B : C] = e(B^*/C^*)f(B^*/C^*) = [B^* : C^*]$. Then $[A^* : B^*] = [A : B] = [A^* : B^*]$ and we have $B^* = B$. The proof of the remaining part is due to Theorem 1.7 [6].

References

- [1] E. Artin and G. Whaples, The theory of simple rings, Amer. J. Math., 65 (1943), 87-107.
- [2] S. Iyanaga: Number Theory (in Japanese), Iwanami (1969).
- [3] K. Hirata: Some types of separable extensions of rings, Nagoya Math. J., 33 (1968), 107-115.
- [4] R. Pierce: Associative Algebras, Springer GTM 88 (1980).
- [5] K. Sugano: On H-separable extensions of two-sided simple rings, Hokkaido Math. J., 11 (1982), 246-252.
- [6] K. Sugano: On bicommutators of modules over H-separable extension rings III, to appear in Hokkaido Math. J..

Hokkaido University