



Title	Dilation Method and Smoothing Effect of the Schrödinger Evolution Group
Author(s)	Hayashi, N.; Kato, K.; Ozawa, T.
Citation	Hokkaido University Preprint Series in Mathematics, 243, 2-10
Issue Date	1994-5-1
DOI	10.14943/83390
Doc URL	http://hdl.handle.net/2115/68994
Type	bulletin (article)
File Information	pre243.pdf



[Instructions for use](#)

**DILATION METHOD AND
SMOOTHING EFFECT OF
THE SCHRÖDINGER
EVOLUTION GROUP**

N. Hayashi, K. Kato and T. Ozawa

Series #243. May 1994

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 216: Y. Giga, N. Mizoguchi, Existence of periodic solutions for equations of evolving curves, 45 pages. 1993.
- # 217: T. Suwa, Indices holomorphic vector fields relative to invariant curves, 10 pages. 1993.
- # 218: S. Izumiya, G. T. Kossioris, Realization theorems of geometric singularities for Hamilton-Jacobi equations, 14 pages. 1993.
- # 219: Y. Giga, K. Yama-uchi, On instability of evolving hypersurfaces, 14 pages. 1993.
- # 220: W. Bruns, T. Hibi, Cohen-Macaulay partially ordered sets with pure resolutions, 11 pages. 1993.
- # 221: S. Jimbo, Y. Morita, Ginzburg Landau equation and stable solutions in a rotational domain, 32 pages. 1993.
- # 222: T. Miyake, Y. Maeda, On a property of Fourier coefficients of cusp forms of half-integral weight, 12 pages. 1993.
- # 223: I. Nakai, Notes on versal deformation of first order PDE and web structure, 34 pages. 1993.
- # 224: I. Tsuda, Can stochastic renewal of maps be a model for cerebral cortex?, 30 pages. 1993.
- # 225: H. Kubo, K. Kubota, Asymptotic behaviors of radial solutions to semilinear wave equations in odd space dimensions, 47 pages. 1994.
- # 226: T. Nakazi, K. Takahashi, Two dimensional representations of uniform algebras, 7 pages. 1994.
- # 227: N. Hayashi, T. Ozawa, Global, small radially symmetric solutions to nonlinear Schrödinger equations and a gauge transformation, 16 pages. 1994.
- # 228: S. Izumiya, Characteristic vector fields for first order partial differential equations, 9 pages. 1994.
- # 229: K. Tsutaya, Lower bounds for the life span of solutions of semilinear wave equations with data of non compact support, 14 pages. 1994.
- # 230: H. Okuda, I. Tsuda, A coupled chaotic system with different time scales: Toward the implication of observation with dynamical systems, 31 pages. 1994.
- # 231: A. Hoshiga, The asymptotic behaviour of radial solutions near the blow-up point to quasi-linear wave equations in two space dimensions, 8 pages. 1994.
- # 232: Y. Giga, N. Mizoguchi, Existence of periodically evolving convex curves moved by anisotropic curvature, 12 pages. 1994.
- # 233: C. Dohmen, Y. Giga, N. Mizoguchi, Existence of selfsimilar shrinking curves for anisotropic curvature flow equations, 13 pages. 1994.
- # 234: T. Nakazi, M. Yamada, Invertible Toeplitz operators and uniform algebras, 14 pages. 1994.
- # 235: C. Dohmen, A Pohozaev-type inequality for a quasilinear Haraux-Weissler equation, 6 pages. 1994.
- # 236: T. Mikami, Large deviations for the first exit time on small random perturbations of dynamical systems, 32 pages. 1994.
- # 237: T. Ozawa, Characterization of Trudinger's inequality, 7 pages. 1994.
- # 238: T. Hibi, Buchsbaum complexes with linear resolutions, 12 pages. 1994.
- # 239: Y. Giga, N. Mizoguchi, On time periodic solutions of the Dirichlet problem for degenerate parabolic equations of nondivergence type, 23 pages. 1994.
- # 240: C. Dohmen, Existence of Fast Decaying Solutions to a Haraux-Weissler Equation With a Prescribed Number of Zeroes, 12 pages. 1994.
- # 241: K. Sugano, Note on H-separable Frobenius extensions, 8 pages. 1994.
- # 242: J. Zhai, Some Estimates For The Blowing up Solutions of Semilinear Heat Equations, 11 pages. 1994.

DILATION METHOD AND SMOOTHING EFFECT OF THE SCHRÖDINGER EVOLUTION GROUP

N. HAYASHI*, K. KATO** AND T. OZAWA***

*Department of Mathematics, Faculty of
Engineering, Gunma University, Kiryu 376, Japan

**Department of Mathematics, Osaka University, Toyonaka 560, Japan

***Department of Mathematics, Hokkaido University, Sapporo 060, Japan

ABSTRACT. We reexamine the mechanism of smoothing effects of the Schrödinger evolution group in the weighted Sobolev spaces by using the generator of space-time dilations instead of Galilei transformations.

1. Introduction

In this paper we prove smoothing effects of the Schrödinger evolution groups in the weighted Sobolev spaces on the basis of the commutation relations for the generator of space-time dilations instead of Galilei transformations. Let $H = -(1/2)\Delta + V$ be a Schrödinger operator in $L^2 = L^2(\mathbb{R}^n)$ with potential V , the multiplication operator by the real-valued function V on \mathbb{R}^n . Our main attention is concentrated on the smoothing effects of the unitary propagator e^{-itH} in L^2 . Looking through an extensive literature on the smoothing effects for Schrödinger evolution equations [1-4, 7, 9-13, 15, 16, 20-27, 29-31, 33-36, 38-42], we find that the methods are classified into three categories. The first method is related to the Strichartz type estimate, which is formulated in the L^p setting and depends essentially on the mapping properties of the Fourier transform with respect to the space variables. It implies in particular that for any data in L^2 the corresponding solution belongs for almost all times to $L^2 \cap L^p$ for all p with $2 < p < 2n/(n-2)$. The second method originates in the Kato smoothness estimate, which is formulated in the L^2 setting and depends essentially on the mapping properties of the Fourier transform in the time variable. It implies in particular that for any data in L^2 the corresponding solution belongs for almost all times to the local Sobolev space $H_{loc}^{1/2}$. Roughly speaking, both results above ensure that for almost all times propagators for Schrödinger evolution equations improve the spatial local regularity of the initial data. As for the time variable, we remark here that the restriction of almost everywhere convergence cannot be reduced to the pointwise convergence. In contrast, the third method holds pointwisely in time and depends exclusively on the commutation relations between the unperturbed operator $L = i\partial_t + (1/2)\Delta$

and the generators of transformations which leave the associated Lagrangean invariant. To be specific the only operator that has been used for that purpose is the generator $J = x + it\nabla$ of Galilei transformations, which commutes with L and provides the spatial local regularity through the formula $(x + it\nabla)U(t) = U(t)x$, where $U(t) = \exp(i(t/2)\Delta)$. It implies in particular that for any data $\phi \in L^2$ with $|x|^m\phi \in L^2$, m being a positive integer, the corresponding solution belongs to the local Sobolev space H_{loc}^m away from the initial time. In other words the last method reveals the connection between the spatial decay at infinity of the data measured by the position operator x and the regularity of the corresponding solutions measured by the order of local Sobolev spaces. Despite the fact that smoothing effects of pointwise times have been recognized as implications of the spatial decay at infinity, it does not logically mean that the decay at infinity is the only source of the local regularity of solutions. In fact the purpose of this paper is to prove that this is the case. We prove in particular that for any data $\phi \in L^2$ with $(x \cdot \nabla)^m\phi \in L^2$, the corresponding solution belongs to H_{loc}^{2m} away from the initial time. This reveals the connection between the regularity of the data with respect to the self-adjoint operator $A = (1/2i)(x \cdot \nabla + \nabla \cdot x)$ and that of the corresponding solutions measured by the order of local Sobolev spaces. As a corollary, it turns out that the propagator e^{-itH} maps C^∞ -vectors for A into smooth functions for any $t \in \mathbf{R} \setminus \{0\}$. Our argument exploits the generator $P = x \cdot \nabla + 2t\partial_t$ of space-time dilations, which satisfies the commutation relation $[L, P] = 2L$. In this sense our method falls into the third category. We derive the local regularity from the time dilation part of P in the associated a priori estimates. To this end we prove that in the L^2 space with weight of negative index the effect of the space dilation part of P amounts to an infinitesimally small perturbation with respect to the time dilation part. As is expected by equation, the time derivative deserves the second derivatives in space, so that actually we prove that $\langle x \rangle^{-2}x \cdot \nabla$ is infinitesimally small with respect to $\langle x \rangle^{-2}\Delta$. The description of the resulting smoothing effect is therefore efficiently given by the use of the weighted Sobolev spaces, which traces back to Jensen[20]. The operator P has recently been used in the study of the nonlinear Schrödinger equation[5, 8, 9, 12]. To state our result precisely, we introduce several notations.

Notation. $\partial_t = \partial/\partial t$, $\partial_j = \partial/\partial x_j$, $\nabla = (\partial_1, \dots, \partial_n)$, $x = (x_1, \dots, x_n)$; $\partial^\alpha = \prod_{j=1}^n \partial_j^{\alpha_j}$, $x^\alpha = \prod_{j=1}^n x_j^{\alpha_j}$; L^p denotes the usual Lebesgue space on \mathbf{R}^n with norm $\|\cdot\|_p$; $\|\cdot\|$ and (\cdot, \cdot) denote the norm and scalar product in L^2 , respectively; $H^{m,s}$ denotes the weighted Sobolev space defined by $H^{m,s} = \{\psi \in \mathcal{S}' ; \|\psi\|_{m,s} \equiv \|\langle x \rangle^s (1 - \Delta)^{m/2} \psi\| < \infty\}$ for $m, s \in \mathbf{R}$, where \mathcal{S}' denotes the space of temperate distributions and $\langle x \rangle = (1 + |x|^2)^{1/2}$; $D_m = D(A^m) = \{\psi \in L^2 ; A^j \psi \in L^2 \text{ for all } j \text{ with } 1 \leq j \leq m\}$ denotes the domain of the m -th power of the self-adjoint dilation operator $A = (1/2i)(x \cdot \nabla + \nabla \cdot x)$, equipped with the graph norm, which is equivalent to $\|\psi\|_m \equiv \sum_{j=0}^m \|(x \cdot \nabla)^j \psi\|$ and to $\|\psi\| + \|(x \cdot \nabla)^m \psi\|$.

With these notations we state our main results. In this paper we prove

Theorem 1. Let $m \geq 1$ be an integer. Suppose that $\partial^\alpha V \in L^\infty$ for all α with $|\alpha| \leq 2m - 2$ and that $(x \cdot \nabla)^j V \in L^\infty$ for all j with $1 \leq j \leq m$. Then e^{-itH} is bounded from D_m to $H^{2m, -2m}$ for any $t \in \mathbb{R} \setminus \{0\}$ with the associated operator norm bounded by $C(1 + |t|^{-2m})$.

Combining Theorem 1 with the results in [27] (see also [20, 26, 29]), we obtain

Theorem 2. Let $m \geq 1$ be an integer. In addition to the assumptions in Theorem 1, if $x \cdot \nabla \partial^\alpha V \in L^\infty$ for all α with $|\alpha| \leq 2m - 2$ then e^{-itH} is bounded from $D_m + H^{0, 2m}$ to $H^{2m, -2m}$ for any $t \in \mathbb{R} \setminus \{0\}$ with the associated operator norm bounded by $C(1 + |t|^{-2m})$.

Remark 1. The condition that $|\partial^\alpha V(x)| \leq C_\alpha \langle x \rangle^{-|\alpha|}$ for all α may be the simplest one that ensures all the assumptions above. The following potentials illustrate other examples.

- (1) $V(x) = x^\alpha / |x|^{|\alpha|}$ with $\alpha \neq 0$ satisfies the assumptions with $m = 1$, while the first derivatives are not bounded at the origin. A general example in this direction is a smooth function on $\mathbb{R}^n \setminus \{0\}$ with homogeneity of degree zero. The asymptotic behavior of propagators for Schrödinger operators with such a class of potentials is studied by Herbst[17].
- (2) $V(x) = \sin \log(x^\alpha)$ with $\alpha \neq 0$ satisfies the assumptions with $m = 1$, while the first derivatives need not stay bounded along the axes.
- (3) $V(x) = \cos(|x|/(1 + |x|))$ satisfies the assumptions with $m = 1, 2$.
- (4) $V(x) = \exp(-|x|^\sigma)$ with $\sigma > 0$ satisfies the assumptions with $m \leq \sigma/2 + 1$. If σ is even, V satisfies all the assumptions.

Remark 2. Let V satisfy all the assumptions in Theorems 1 and 2, respectively. Then for any $t \neq 0$ the propagator e^{-itH} maps D_∞ and $D_\infty + H^{0, \infty}$ into C^∞ , respectively, where $D_\infty = \bigcap_{m \geq 0} D_m$ is the space of C^∞ -vectors for A and $H^{0, \infty} = \bigcap_{m \geq 0} H^{0, m}$.

Remark 3. $\phi(x) = |x|^{-n/2+\epsilon}(1 + |x|)^{-2\epsilon}$ with $\epsilon > 0$ satisfies $\phi \in D_\infty$ while $\phi \notin H^{0, \epsilon}$. By the preceding remark, $e^{-itH}\phi \in C^\infty$ for any $t \neq 0$, provided that V satisfies all the assumptions in Theorem 1. This example falls beyond the scope of the previous literature even when restricted to the case $V = 0$.

2. Proof of Theorem 1

The proof of Theorem 1 requires the following lemma.

Lemma. Let $m \geq 1$ be an integer. Then there exists a constant C such that for

all $\psi \in H^{2m, -2m}$

(1)

$$\|\psi\|_{2m-1, -2m+1} \leq C \left(\sum_{j=0}^{m-1} \|\Delta^j \psi\|_{0, -2j} \right)^{1/2} \|\Delta^m \psi\|_{0, -2m}^{1/2} + C \sum_{j=0}^{m-1} \|\Delta^j \psi\|_{0, -2j},$$

(2)

$$\sum_{j=0}^{2m} \|\psi\|_{j, -j} \leq C \sum_{k=0}^m \|\Delta^k \psi\|_{0, -2k},$$

(3)

$$\|\psi\|_{m, -m} \leq C \|\psi\|^{1/2} \left(\sum_{j=0}^m \|\Delta^j \psi\|_{0, -2j} \right)^{1/2}.$$

Proof. By density it suffices to prove the estimates for $\psi \in \mathcal{S}$. Let α satisfy $|\alpha| = 2m - 2$. By the relation $|\partial_k \partial^\alpha \psi|^2 = (1/2) \partial_k^2 |\partial^\alpha \psi|^2 - \text{Re}(\overline{\partial^\alpha \psi} \partial_k^2 \partial^\alpha \psi)$, integration by parts, and the Schwarz inequality, we obtain

$$\begin{aligned} \|\partial_k \partial^\alpha \psi\|_{0, -2m+1}^2 &= \frac{1}{2} (\langle x \rangle^{-4m+2}, \partial_k^2 |\partial^\alpha \psi|^2) - \text{Re}(\langle x \rangle^{-4m+2} \partial^\alpha \psi, \partial_k^2 \partial^\alpha \psi), \\ &\leq C \|\partial^\alpha \psi\|_{0, -2m+2}^2 + \|\partial^\alpha \psi\|_{0, -2m+2} \|\partial_k^2 \partial^\alpha \psi\|_{0, -2m}. \end{aligned}$$

This implies (1) since

$$(4) \quad \sum_{|\beta|=2l} \|\partial^\beta \psi\|_{0, -2l} \leq C \|\psi\|_{2l, -2l} \leq C \sum_{j=0}^l \|\Delta^j \psi\|_{0, -2j} \leq C \sum_{j=0}^l \|\Delta^j \psi\|_{0, -2j}$$

and the norm defined by $\|\psi\|_{k, s}^* = \|\psi\|_{0, s} + \sum_{|\alpha|=k} \|\partial^\alpha \psi\|_{0, s}$ is an equivalent norm in $H^{k, s}$ with $k \in \mathbb{N}$, $s \in \mathbb{R}$ [37]. Similarly, (2) follows from (1) and (4). The proof of (3) proceeds similarly since for $|\alpha| = m$

$$\begin{aligned} \|\partial^\alpha \psi\|_{0, -m}^2 &= (\partial^\alpha \psi, \langle x \rangle^{-2m} \partial^\alpha \psi) \\ &= (-1)^{|\alpha|} (\psi, \partial^\alpha (\langle x \rangle^{-2m} \partial^\alpha \psi)) \\ &\leq \|\psi\| \|\partial^\alpha (\langle x \rangle^{-2m} \partial^\alpha \psi)\| \leq C \|\psi\| \|\psi\|_{2m, -2m}. \quad \square \end{aligned}$$

Proof of Theorem 1. We first regularize V by a mollifier. Let $\rho \in C_0^\infty(\mathbb{R}^n)$ satisfy $\rho \geq 0$ and $\|\rho\|_1 = 1$ and let $\rho_\epsilon(x) = \epsilon^{-n} \rho(\epsilon^{-1}x)$ for $\epsilon > 0$. Let $V_\epsilon = V * \rho_\epsilon$ and let $u_\epsilon(t) = \exp(-it(H_0 + V_\epsilon))\phi$ for $\phi \in \mathcal{S}$. Then $u_\epsilon \in C^\infty(\mathbb{R}; \mathcal{S})$ since $\partial^\alpha V_\epsilon \in L^\infty$ for all α [18, 19, 28, 29, 32]. We prove that

$$(5.m) \quad \|\Delta^j u_\epsilon(t)\|_{0, -2j} \leq C(1 + |t|^{-2j}) \|\phi\|_j, \quad 0 \leq j \leq m, \quad t \in \mathbb{R} \setminus \{0\},$$

$$(6.m) \quad \|P^j u_\epsilon(t)\| \leq C(1 + |t|^j) \|\phi\|_j, \quad 0 \leq j \leq m, \quad t \in \mathbb{R} \setminus \{0\},$$

with constant C independent of ϵ , t and ϕ . We prove (5.m) and (6.m) by induction on m . The case $m = 0$ follows from the unitarity of the propagator $\exp(-it(H_0 + V_\epsilon))$. Let $m \geq 1$ and suppose that (5.m-1) and (6.m-1) hold. We proceed to the case m . By the commutation relation $[L, P] = 2L$, we have, omitting the time variable t ,

$$\begin{aligned} LP^m u_\epsilon &= (P+2)^m L u_\epsilon = (P+2)^m (V_\epsilon u_\epsilon) \\ &= \sum_{j=0}^m \binom{m}{j} 2^{m-j} P^j (V_\epsilon u_\epsilon) \\ &= V_\epsilon P^m u_\epsilon + \sum_{k=0}^{m-1} \sum_{j=0}^{m-k} \binom{m}{k} \binom{m-k}{j} 2^{m-j-k} (P^j V_\epsilon) P^k u_\epsilon. \end{aligned}$$

Taking the imaginary part of the scalar product in L^2 between $LP^m u_\epsilon$ with the last form and $P^m u_\epsilon$ and integrating by parts, we have

$$(7) \quad \frac{d}{dt} \|P^m u_\epsilon\|^2 = \sum_{k=0}^{m-1} \sum_{j=0}^{m-k} \binom{m}{k} \binom{m-k}{j} 2^{m-j-k+1} \operatorname{Im}((P^j V_\epsilon) P^k u_\epsilon, P^m u_\epsilon).$$

Noting that $(x \cdot \nabla) V * \rho_\epsilon = (x \cdot \nabla V) * \rho_\epsilon + V * \nabla \cdot (x \rho_\epsilon)$, we obtain

$$(8) \quad \sum_{j=0}^m \|P^j V_\epsilon\|_\infty \leq C \sum_{j=0}^m \|(x \cdot \nabla)^j V\|_\infty.$$

with constant C independent of ϵ . By (7), (8), and (6.m-1),

$$(9) \quad \left| \frac{d}{dt} (\|P^m u_\epsilon\|^2 + \delta) \right| \leq C \sum_{k=0}^{m-1} \|P^k u_\epsilon\| \|P^m u_\epsilon\| \\ \leq C(1 + |t|^{m-1}) \|\phi\|_{m-1} (\|P^m u_\epsilon\|^2 + \delta)^{1/2},$$

for any $\delta > 0$. Dividing both sides of (9) by $(\|P^m u_\epsilon\| + \delta)^{1/2}$, integrating in t , and letting $\delta \downarrow 0$, we obtain (6.m). We next prove (5.m) by making use of (5.m-1) and (6.m). By expanding the m -th power of P , $(2t)^m \partial_t^m$ is expressed as

$$\begin{aligned} (2t)^m \partial_t^m &= P^m - (x \cdot \nabla)^m - \sum_{k=0}^{m-1} a_{m,k} t^k \partial_t^k \\ &\quad - \sum_{j=1}^{m-1} \binom{m}{j} (x \cdot \nabla)^{m-j} \left((2t)^j \partial_t^j + \sum_{k=1}^{j-1} a_{j,k} t^k \partial_t^k \right) \end{aligned}$$

with constants $a_{j,k}$, $1 \leq j \leq m$ and $1 \leq k \leq m-1$, where we follow the convention $\sum_{k=1}^0(\dots) = 0$. Letting (10) act on u_ϵ , using the equation $i\partial_t u_\epsilon = (H_0 + V_\epsilon)u_\epsilon$, and estimating the result, we have

$$(11) \quad \begin{aligned} & (2|t|)^m \|(H_0 + V_\epsilon)^m u_\epsilon\|_{0,-2m} \\ & \leq \|P^m u_\epsilon\| + \|(x \cdot \nabla)^m u_\epsilon\|_{0,-2m} + C \sum_{k=1}^{m-1} |t|^k \|(H_0 + V_\epsilon)^k u_\epsilon\|_{0,-2m} \\ & \quad + C \sum_{j=1}^{m-j} \sum_{k=1}^j |t|^k \|(x \cdot \nabla)^{m-j} (H_0 + V_\epsilon)^k u_\epsilon\|_{0,-2m}. \end{aligned}$$

By expanding the j -th power of $H_0 + V_\epsilon$,

$$(12) \quad \begin{aligned} & (H_0 + V_\epsilon)^j \\ & = \sum_{k=0}^j \binom{j}{k} V_\epsilon^{j-k} H_0^k \\ & \quad + \sum_{k=1}^{j-1} \sum_{\substack{|\beta| \leq 2(j-k)-1 \\ |\alpha_1 + \dots + \alpha_k + \beta| = 2(j-k)}} C(j, k, \{\alpha_l\}, \beta) \left(\prod_{l=1}^k \partial^{\alpha_l} V_\epsilon \right) \partial^\beta \end{aligned}$$

and therefore by (2)

$$(13) \quad \begin{aligned} & \|((H_0 + V_\epsilon)^j - H_0^j) u_\epsilon\|_{0,-2j} \\ & \leq C \sum_{k=0}^{j-1} \|V_\epsilon\|_\infty^{j-k} \|H_0^k u_\epsilon\|_{0,-2k} \\ & \quad + C \sum_{k=1}^{j-1} \left(\sum_{|\alpha| \leq 2(j-k)} \|\partial^\alpha V_\epsilon\|_\infty \right)^k \sum_{|\beta| \leq 2(j-k)-1} \|\partial^\beta u_\epsilon\|_{0,-|\beta|} \\ & \leq C \left(1 + \sum_{|\alpha| \leq 2j-2} \|\partial^\alpha V\|_\infty^j \right) \sum_{k=0}^{j-1} \|\Delta^k u_\epsilon\|_{0,-2k}. \end{aligned}$$

By (13) and (5.m-1),

$$(14) \quad \begin{aligned} & \|(t\Delta)^m u_\epsilon\|_{0,-2m} \\ & \leq (2|t|)^m \|(H_0 + V_\epsilon)^m u_\epsilon\|_{0,-2m} + C(|t|^m + |t|^{-m+2}) \|\phi\|_{m-1}, \end{aligned}$$

$$(15) \quad \begin{aligned} \sum_{k=1}^{m-1} |t|^k \|(H_0 + V_\epsilon)^k u_\epsilon\|_{0,-2m} & \leq C \sum_{k=1}^{m-1} |t|^k \sum_{j=0}^k \|\Delta^j u_\epsilon\|_{0,-2j} \\ & \leq C(|t|^{m-1} + |t|^{-m+1}) \|\phi\|_{m-1}. \end{aligned}$$

Combining (6.m), (11), (14) and (15), we have

$$\begin{aligned}
(16) \quad & \| (t\Delta)^m u_\epsilon \|_{0, -2m} \\
& \leq C(|t|^m + |t|^{-m+1}) \|\phi\|_m + \| (x \cdot \nabla)^m u_\epsilon \|_{0, -2m} \\
& \quad + C \sum_{j=1}^{m-1} \sum_{k=1}^j |t|^k \| (x \cdot \nabla)^{m-j} (H_0 + V_\epsilon)^k u_\epsilon \|_{0, -2m}.
\end{aligned}$$

By (2), (3), (5.m - 1), the middle term of the RHS of (16) is estimated as

$$\begin{aligned}
(17) \quad & C \sum_{|\alpha| \leq m} \|\partial^\alpha u_\epsilon\|_{|\alpha| - 2m} \leq C \sum_{|\alpha| \leq m} \|\partial^\alpha u_\epsilon\|_{0, -|\alpha|} \\
& \leq C \sum_{j=0}^{m-1} \|u_\epsilon\|_{j, -j} + C \|u_\epsilon\|_{m, -m} \\
& \leq C \sum_{k=0}^{[m/2]} \|\Delta^k u_\epsilon\|_{0, -2k} + C(|t|^{-m} \|\phi\|)^{1/2} \left(\sum_{j=0}^m |t|^m \|\Delta^j u_\epsilon\|_{0, -2j} \right)^{1/2} \\
& \leq \frac{1}{4} \|(t\Delta)^m u_\epsilon\|_{0, -2m} + C(1 + |t|^m) \sum_{j=0}^{m-1} \|\Delta^j u_\epsilon\|_{0, -2j} + C|t|^{-m} \|\phi\| \\
& \leq \frac{1}{4} \|(t\Delta)^m u_\epsilon\|_{0, -2m} + C(|t|^m + |t|^{-m}) \|\phi\|_{m-1},
\end{aligned}$$

where $[m/2]$ is the largest integer less than or equal to $m/2$. By (1), (2), (12) and (5.m - 1), the last term of the RHS of (16) is estimated as

$$\begin{aligned}
(18) \quad & C \sum_{j=1}^{m-1} \sum_{k=1}^j \sum_{|\alpha| \leq m-j} |t|^k \|\partial^\alpha (H_0 + V_\epsilon)^k u_\epsilon\|_{0, |\alpha| - 2m} \\
& \leq C \sum_{j=1}^{m-1} \sum_{k=1}^j \sum_{|\alpha| \leq m-j} \sum_{|\beta| \leq |\alpha| + 2k} |t|^k \|\partial^\beta u_\epsilon\|_{0, |\alpha| - 2m} \\
& \leq C \sum_{j=1}^{m-1} \sum_{k=1}^j \sum_{|\alpha| \leq m-j} \sum_{|\beta| \leq |\alpha| + 2k} |t|^k \|\partial^\beta u_\epsilon\|_{0, -|\beta|} \\
& \leq C \sum_{j=1}^{m-1} \sum_{k=1}^j \sum_{l=0}^{m-j+2k} |t|^k \|u_\epsilon\|_{l, -l} \\
& \leq C \sum_{j=1}^{m-2} \sum_{l=0}^{m-1} (1 + |t|^j) \|u_\epsilon\|_{l, -l} + C|t|^{m-1} \|u_\epsilon\|_{2m-1, -2m+1}
\end{aligned}$$

$$\begin{aligned}
&\leq C(1 + |t|^{m-1}) \sum_{j=0}^{m-1} \|\Delta^j u_\epsilon\|_{0,-2j} \\
&\quad + C \left(\sum_{j=0}^{m-1} |t|^{m-2} \|\Delta^j u_\epsilon\|_{0,-2j} \right)^{1/2} \|(t\Delta)^m u_\epsilon\|_{0,-2m}^{1/2} \\
&\leq \frac{1}{4} \|(t\Delta)^m u_\epsilon\|_{0,-2m} + C(|t|^m + |t|^{-m}) \|\phi\|_{m-1}.
\end{aligned}$$

By (16), (17) and (18), we have $\|(t\Delta)^m u_\epsilon\|_{0,-2m} \leq C(|t|^m + |t|^{-m}) \|\phi\|_m$ and hence (5.m), as required. We note that (5.m) and (6.m) extends to $\phi \in D_m$ since \mathcal{S} is dense in D_m . By (5.m) we have

$$(19) \quad \|u_\epsilon(t)\|_{2m,-2m} \leq C(1 + |t|^{-2m}) \|\phi\|_m, \phi \in D_m, t \in \mathbf{R} \setminus \{0\},$$

where C is independent of ϵ , t and ϕ . Finally we consider (19) in the limit $\epsilon \downarrow 0$. For $\psi \in L^2$, $V_\epsilon \psi \rightarrow V\psi$ almost everywhere as $\epsilon \downarrow 0$ and $|(V_\epsilon - V)\psi| \leq 2\|V\|_\infty |\psi|$. This implies $\|(V_\epsilon - V)\psi\| \rightarrow 0$ as $\epsilon \downarrow 0$, which in turn implies

$$\begin{aligned}
&\|((H_0 + V_\epsilon + i)^{-1} - (H_0 + V + i)^{-1})\psi\| \\
&= \|(H_0 + V_\epsilon + i)^{-1}(V_\epsilon - V)(H_0 + V + i)^{-1}\psi\| \\
&\leq \|(V_\epsilon - V)(H_0 + V + i)^{-1}\psi\| \rightarrow 0 \quad \text{as } \epsilon \downarrow 0.
\end{aligned}$$

The Trotter-Kato theorem shows that $u_\epsilon(t) \rightarrow e^{-itH}\phi$ in L^2 locally uniformly on t -intervals as $\epsilon \downarrow 0$. Since the operator $\langle x \rangle^{-2m}(1 - \Delta)^m$ is closed in L^2 , we conclude from (19) that $e^{-itH}\phi \in H^{2m,-2m}$ for $t \in \mathbf{R} \setminus \{0\}$ and $\|e^{-itH}\phi\|_{2m,-2m} \leq C(1 + |t|^{-2m}) \|\phi\|_m$. \square

REFERENCES

1. M. Ben-Artzi, A. Devinatz, *Local smoothing and convergence properties of Schrödinger-type equations*, J. Funct. Anal. 101 (1991), 231–254.
2. P. Constantin, J. C. Saut, *Local smoothing properties of dispersive equations*, J. Amer. Math. Soc. 1 (1988), 413–439.
3. P. Constantin, J. C. Saut, *Local smoothing properties of Schrödinger equations*, Indiana Univ. Math. J. 38 (1989), 791–810.
4. J. M. Ghidaglia, J. C. Saut, *On the initial value problem for the Davey-Stewartson system*, Nonlinearity 3 (1990), 475–506.
5. J. Ginibre, N. Hayashi, *Almost global existence of small solutions to quadratic nonlinear Schrödinger equations in three space dimensions*, Math. Z. (to appear).
6. J. Ginibre, G. Velo, *Scattering theory in the energy space for a class of nonlinear Schrödinger equations*, J. Math. Pures Appl. 64 (1985), 363–401.
7. J. Ginibre, G. Velo, *Smoothing properties and retarded estimates for some dispersive evolution equations*, Commun. Math. Phys. 144 (1992), 163–188.
8. N. Hayashi, *Global existence of small solutions to quadratic nonlinear Schrödinger equations*, Comm. PDE 18 (1993), 1109–1124.

9. N. Hayashi, K. Kato, *Regularity in time of solutions to nonlinear Schrödinger equations*, J. Funct. Anal. (to appear).
10. N. Hayashi, K. Nakamitsu, M. Tsutsumi, *On solutions of the initial value problem for nonlinear Schrödinger equations in one space dimension*, Math. Z. 192 (1986), 637–650.
11. N. Hayashi, K. Nakamitsu, M. Tsutsumi, *On solutions of the initial value problem for nonlinear Schrödinger equations*, J. Funct. Anal. 71 (1987), 218–245.
12. N. Hayashi, T. Ozawa, *Smoothing effect for some Schrödinger equations*, J. Funct. Anal. 85 (1989), 307–348.
13. N. Hayashi, T. Ozawa, *On the derivative nonlinear Schrödinger equation*, Physica D 55 (1992), 14–36.
14. N. Hayashi, T. Ozawa, *Global, small radially symmetric solutions to nonlinear Schrödinger equations and a gauge transformation*, Preprint (1994).
15. N. Hayashi, S. Saitoh, *Analyticity and smoothing effect for the Schrödinger equation*, Ann. Inst. Henri Poincaré, Physique théorique 52 (1990), 163–173.
16. N. Hayashi, S. Saitoh, *Analyticity and global existence of small solutions to some nonlinear Schrödinger equations*, Commun. Math. Phys. 129 (1990), 27–42.
17. I. W. Herbst, *Spectral and scattering theory for Schrödinger operators with potentials independent of $|x|$* , Amer. J. Math. 113 (1991), 509–565.
18. M. J. Huang, *Commutators and invariant domains for Schrödinger propagators*, Preprint (1994).
19. W. Hunziker, *On the space-time behavior of Schrödinger wave functions*, J. Math. Phys. 7 (1966), 300–304.
20. A. Jensen, *Commutator methods and a smoothing property of the Schrödinger evolution group*, Math. Z. 191 (1986), 53–59.
21. T. Kato, *Wave operators and similarity for some non-selfadjoint operators*, Math. Ann. 162 (1966), 258–279.
22. T. Kato, *On nonlinear Schrödinger equations*, Ann. Inst. Henri Poincaré, Physique théorique 46 (1987), 113–129.
23. T. Kato, K. Yajima, *Some examples of smooth operators and the associated smoothing effect*, Rev. Math. Phys. 1 (1989), 481–496.
24. C. E. Kenig, G. Ponce, K. Vega, *Oscillatory integrals and regularity of dispersive equations*, Indiana Univ. Math. J. 40 (1991), 33–69.
25. C. E. Kenig, G. Ponce, K. Vega, *Small solutions to nonlinear Schrödinger equations*, Ann. Inst. Henri Poincaré, Analyse non linéaire 10 (1993), 255–288.
26. K. Nakamitsu, *Smoothing effects for Schrödinger evolution groups*, Tokyo Denki Univ. Kiyō 10 (1988), 49–52.
27. T. Ozawa, *Smoothing effect and dispersion of singularities for the Schrödinger evolution group*, Arch. Rational Mech. Anal. 110 (1990), 165–186.
28. T. Ozawa, *Invariant subspaces for the Schrödinger evolution group*, Ann. Inst. Henri Poincaré, Physique théorique 54 (1991), 43–57.
29. T. Ozawa, *Space-time behavior of propagators for Schrödinger evolution equations with Stark effect*, J. Funct. Anal. 92 (1991), 264–292.
30. T. Ozawa, Y. Tsutsumi, *Existence and smoothing effect of solutions for the Zakharov equations*, Publ. RIMS, Kyoto Univ. 28 (1992), 329–361.
31. G. Ponce, *Regularity of solutions to nonlinear dispersive equations*, J. Differential Equations 78 (1989), 122–135.
32. C. Radin, B. Simon, *Invariant domains for time-dependent Schrödinger equations*, J. Differential Equations 29 (1979), 289–296.
33. B. Simon, *Best constants in some operator smoothness estimates*, J. Funct. Anal. 107 (1992), 66–71.
34. P. Sjölin, *Regularity of solutions to the Schrödinger equations*, Duke Math. J. 55 (1987), 699–715.

35. P. Sjölin, *Local regularity of solutions to nonlinear Schrödinger equations*, Ark. Mat. 28 (1990), 145–157.
36. R. S. Strichartz, *Restrictions of Fourier transforms to quadratic surfaces and decay of solutions for wave equations*, Duke Math. J. 44 (1977), 705–714.
37. H. Triebel, *Space of distributions with weights. Multipliers in L_q -spaces with weights*, Math. Nachr. 78 (1977), 339–355.
38. L. Vega, *The Schrödinger equations: pointwise convergence to the initial data*, Proc. Amer. Math. Soc. 102 (1988), 874–878.
39. K. Watanabe, *Smooth perturbations of the selfadjoint operator $|\Delta|^{\alpha/2}$* , Tokyo J. Math. 14 (1991), 239–250.
40. K. Yajima, *Existence of solutions for Schrödinger evolution equations*, Commun. Math. Phys. 110 (1987), 415–426.
41. K. Yajima, *Schrödinger evolution equations with magnetic fields*, J. Analyse Math. 56 (1991), 29–76.
42. M. Yamazaki, *On the microlocal smoothing effect of dispersive partial differential equations. I. Second-order linear equations*, in "Algebraic Analysis," Academic Press, San Diego, CA, 1988.