



Title	The 28th Sapporo Symposium on Partial Differential Equations
Author(s)	Jimbo, Shuichi
Citation	Hokkaido University technical report series in mathematics, 77, 1
Issue Date	2003-01-01
DOI	10.14943/632
Doc URL	http://hdl.handle.net/2115/690 ; http://eprints3.math.sci.hokudai.ac.jp/0277/
Type	bulletin (article)
Note	The 28th Sapporo Symposium on Partial Differential Equations Organizers: T. Ozawa, Y. Giga, S. Jimbo, K. Tsutaya, Y. Tonegawa, G. Nakamura. Venues: Department of Mathematics, Faculty of Science, Hokkaido University July 23, 2003 (Wednesday) 9:30-10:30 Gregory SEREGIN (Steklov Institute/Keio Univ.) Interior regularity of L_3 inf- solutions to the Navier-Stokes equations 11:00-12:00 Takaaki NISHIDA (Kyoto Univ.) Heat convection problems and computer assisted proof 14:30-15:00 Akihiro SHIMOMURA (Gakushuin Univ.) Modified wave operators for Maxwell-Schrodinger equations 15:15-15:45 Hideaki SUNAGAWA (Osaka Univ.) Remarks on the large time asymptotics for nonlinear Klein-Gordon systems 16:00-16:30 Hirokazu NINOMIYA (Ryukoku Univ.) Curved traveling front of Allen-Cahn equations July 24, 2003 (Thursday) 9:30-10:30 Masahiro YAMAMOTO (Univ. Tokyo) Uniqueness in inverse scattering problems with a single incident wave 11:00-12:00 Shinya NISHIBATA (Tokyo Inst. Tech.) Asymptotic behavior of spherically symmetric solutions to the compressible Navier Stokes equation with external forces 14:30-15:00 Dening LI (West Virginia Univ.) Conical shock waves in supersonic flow 15:15-15:45 Yasushi TANIUCHI (Shinshu Univ.) Remarks on global solvability of 2-D Boussinesq equations with nondecaying initial data July 25, 2003 (Friday) 9:30-10:30 Ryuichi SUZUKI (Kokushikan Univ.) Blow-up of solutions of quasilinear parabolic equations with localized reactions 11:00-12:00 SAKAGUCHI (Ehime Univ.) Initial behavior of solutions of diffusion equations and symmetries of domains"
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Ninomiya.pdf



[Instructions for use](#)

The curved traveling front of the Allen-Cahn equation

Hirokazu Ninomiya (Ryukoku University)

and

Masaharu Taniguchi (Tokyo Institute of Technology)

In this talk we consider the Allen-Cahn equation:

$$u_t = \Delta u + f(u) \quad (x, y, t) \in \mathbf{R}^2 \times \mathbf{R}_+ \quad (1)$$

where f is of “bistable type”. The typical example of the nonlinear term f is

$$f(u) = u(1 - u)(u - a), \quad 0 < a < \frac{1}{2}. \quad (2)$$

The constant states 0 and 1 are stable under the diffusion-free system. By the assumption of a , the region of the state 1 is getting larger and larger and finally it covers the whole space. When the state 1 propagates, we can observe the characteristic profiles. In the one-dimensional space, one of the typical solutions is a *traveling wave* solution which never changes its shape without translation. Substituting $u(x, t) = \Phi(x - ct)$, we have

$$\Phi_{\xi\xi} - c\Phi_{\xi} + f(\Phi) = 0.$$

Actually for the nonlinearity (2), we have

$$\Phi(\xi) = \frac{1}{2} \left(1 - \tanh \frac{\xi}{2\sqrt{2}} \right), \quad c = \sqrt{2} \left(\frac{1}{2} - a \right).$$

As the appropriate singular limit, the interface between two states 1 and 0 becomes sharp and we can get the interface equation (see e.g. [4]):

$$V = H + k \quad (3)$$

where V is a normal velocity, H is the curvature, and k is a given constant. This equation is also observed in the filamentary vortex of the Ginzburg–Landau equation confined in a plane [3] and the BZ reaction [6].

The typical solutions of (3) are the circles and lines. In the case $k \neq 0$, unfortunately, some interfaces may possess some self-intersection points eventually, even if the initial interface has none. If the interface is represented by the graph $y = v(x, t)$, the equation (3) is reduced to

$$v_t = \frac{v_{xx}}{1 + v_x^2} + k\sqrt{1 + v_x^2} \quad x \in \mathbf{R}, t > 0, \quad (4)$$

Deckelnick et al in [3] proved the existence of the traveling curved front and studied the stability of the front under some restricted assumptions for u_0 . The authors relaxed the assumption for the initial data and classified all the traveling fronts in [7, 8]. They proved the following (see [7, Proposition 1.1, Theorem 1.2]).

Theorem 1 *Any traveling front of (3) with velocity $v(0, c)$ is one of the three, after appropriate translations,*

- (i) *lines $y = m_*x$, and $y = -m_*x$*
- (ii) *a traveling curved front $\Gamma_c(t)$ which possesses two asymptotes $y = \pm m_*x$,*
- (iii) *stationary circles with radius $1/|k|$ only in the case $c = 0$,*

where $m_* := \sqrt{c^2 - k^2}/k$. Moreover the explicit form of the traveling curved front $\Gamma_c(t) = \{y = \varphi(x) + ct\}$ with speed $c(\geq k)$ is given in

$$x(\theta; c) := \frac{\theta}{c} + \frac{k}{c\sqrt{c^2 - k^2}} \log \left| \frac{1 + \sqrt{\frac{c+k}{c-k}} \tan \frac{\theta}{2}}{1 - \sqrt{\frac{c+k}{c-k}} \tan \frac{\theta}{2}} \right|,$$

$$y(\theta; c) := -\frac{1}{c} \log \left(\frac{c \cos \theta - k}{c - k} \right),$$

for $\theta \in (-\arctan m_*, \arctan m_*)$.

The traveling curved front $\Gamma_c(t)$ is “V-shaped”, which connects two asymptotes. The existence of this traveling front is also reported in [2, 3] and in a liquid BZ reaction [6].

The asymptotic stability of the curved traveling front in (4) is discussed in [3, 8]. It is proved that the traveling curved front is asymptotically stable, if the initial perturbation is restricted to

$$BC_0^1 := \{v \in C^1(\mathbf{R}) \mid \sup_{-\infty < x < \infty} (|v(x)| + |v_x(x)|) < \infty, \lim_{|x| \rightarrow \infty} v(x) = 0\}.$$

and that if you take the perturbation space

$$BC^1 := \{v \in C^1(\mathbf{R}) \mid \sup_{-\infty < x < \infty} (|v(x)| + |v_x(x)|) < \infty\},$$

instead of BC_0^1 , the traveling curved front is not asymptotically stable (see [8, Theorem 1.1 and Theorem 4.1]).

By the above observation, we can expect that a “V-shaped” traveling wave solution of (1) exists. Actually we have the following theorem.

Theorem 2 *There exists a traveling wave solution $u(x, y, t) = U(x, y - ct)$ of (1) such that*

$$\lim_{R \rightarrow \infty} \sup_{(x, y) \in D_R} \left| U(x, y) - \Phi \left(\frac{k}{c}(y - m_*|x|) \right) \right| = 0$$

where

$$D_R := \{(x, y) \mid x^2 + y^2 \geq R^2\}.$$

Bonnet and Hamel [1] showed the existence of the “V-shaped” traveling wave solutions, if f is of the “ignition temperature” type (mono-stable type) instead of (2). Hamel and Monneau [5] shows the uniqueness of the traveling front of the corresponding singular limit problem.

References

- [1] A. BONNET AND F. HAMEL, Existence of nonplanar solutions of a simple model of premixed bunsen flames, *SIAM J. Math. Anal.*, **31**, (1999) 80–118.
- [2] P. K. BRAZHNIK, Exact solutions for the kinematic model of autowaves in two-dimensional excitable media, *Physica D*, **94**, (1996) 205–220.
- [3] K. Deckelnick, C. M. Elliott, and G. Richardson, Long time asymptotics for forced curvature flow with applications to the motion of a superconducting vortex, *Nonlinearity* **10** (1997), 655–678.
- [4] P. C. Fife, *Dynamics of Internal Layers and Diffusive Interfaces* (CBMS-NSF Reg. Conf. Ser. Appl. Math. 53), SIAM (1988).
- [5] F. HAMEL AND R. MONNEAU, Existence and uniqueness for a free boundary problem arising in combustion theory, *Interfaces and Free Boundaries*, **4**, (2002) 167–210.

- [6] V. Pérez-Muñuzuri, M. Gómez-Gesteira, A. P. Muñuzuri, V. A. Davydov, and V. Pérez-Villar, V-shaped stable nonspiral patterns, *Physical Review E* **51-2** (1995), 845–847.
- [7] H. Ninomiya and M. Taniguchi, Traveling curved fronts of a mean curvature flow with constant driving force, in "Free boundary problems: theory and applications, I" *GAKUTO Internat. Ser. Math. Sci. Appl.* **13** (2000), 206–221.
- [8] H. Ninomiya and M. Taniguchi, Stability of traveling curved fronts in a curvature flow with driving force, *Methods and Application of Analysis*, **8-2** (2001), 429–450.
- [9] M. H. Protter and H. F. Weinberger, *Maximum Principles in Differential Equations*, (1984) Springer-Verlag.