



Title	An Abel-Tauber theorem for Fourier sine transforms
Author(s)	Inoue, A.
Citation	Hokkaido University Preprint Series in Mathematics, 261, 1-6
Issue Date	1994-8-1
DOI	10.14943/83408
Doc URL	http://hdl.handle.net/2115/69012
Type	bulletin (article)
File Information	pre261.pdf



[Instructions for use](#)

**An Abel-Tauber theorem
for Fourier sine
transforms**

Akihiko Inoue

Series #261. August 1994

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 236: T. Mikami, Large deviations for the first exit time on small random perturbations of dynamical systems, 32 pages. 1994.
- # 237 T. Ozawa, Characterization of Trudinger's inequality, 7 pages. 1994.
- # 238 T. Hibi, Buchsbaum complexes with linear resolutions, 12 pages. 1994.
- # 239 Y. Giga, N. Mizoguchi, On time periodic solutions of the Dirichlet problem for degenerate parabolic equations of nondivergence type, 23 pages. 1994.
- # 240 C. Dohmen, Existence of Fast Decaying Solutions to a Haraux-Weissler Equation With a Prescribed Number of Zeroes, 12 pages. 1994.
- # 241 K. Sugano, Note on H-separable Frobenius extensions, 8 pages. 1994.
- # 242 J. Zhai, Some Estimates For The Blowing up Solutions of Semilinear Heat Equations, 11 pages. 1994.
- # 243 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and Smoothing Effect of the Schrödinger Evolution Group, 10 pages. 1994.
- # 244 D. Lehmann, T. Suwa, Residues of holomorphic vector fields relative to singular invariant subvarieties, 26 pages. 1994.
- # 245 H. Kubo, Slowly decaying solutions for semilinear wave equations in odd space dimensions, 30 pages. 1994.
- # 246 T. Nakazi, M. Yamada, (A_2) -Conditions and Carleson Inequalities, 27 pages. 1994.
- # 247 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and smoothing Effect of Solutions to the Benjamin-ono Equation, 17 pages. 1994.
- # 248 H. Kikuchi, Sheaf cohomology theory for measurable spaces, 12 pages. 1994.
- # 249 A. Inoue, Tauberian theorems for Fourier cosine transforms, 9 pages. 1994.
- # 250 S. Izumiya, G. T. Kossioris, Singularities for viscosity solutions of Hamilton-Jacobi equations, 23 pages. 1994.
- # 251 H. Kubo, K. Kubota, Asymptotic behaviors of radially symmetric solutions of $\square u = |u|^p$ for super critical values p in odd space dimensions, 51 pages. 1994.
- # 252 T. Mikami, Large Deviations and Central Limit Theorems for Eyraud-Farlie-Gumbel-Morgenstern Processes, 9 pages. 1994.
- # 253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.
- # 254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.
- # 255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.
- # 256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.
- # 257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.
- # 258 Y-G. Chen, Y. Giga, T. Hitaka and M. Honma, A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature, 18 pages. 1994.
- # 259 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 7 pages. 1994.
- # 260 T. Mikami, Copula fields and its applications, 14 pages. 1994.

An Abel-Tauber theorem for Fourier sine transforms

By Akihiko INOUE

Abstract. We prove an Abel-Tauberian theorem for Fourier sine transforms. It can be considered as the analogue of the Abel-Tauber theorem of Pitman in the boundary case. We apply it to Fourier sine series as well as to the tail behavior of a probability distribution.

1. Introduction and results

The aim of this paper is to prove an Abel-Tauber theorem for Fourier sine transforms. It characterizes, for example, the asymptotic behavior $f(t) \sim 1/t^2$ as $t \rightarrow \infty$ in terms of the Fourier sine transform of f , where f is a locally integrable, eventually non-increasing function on $[0, \infty)$ such that $\lim_{t \rightarrow \infty} f(t) = 0$. A similar result for Fourier sine series will be obtained as a corollary.

To state our results, we recall and introduce some notation. We denote by R_0 the whole class of slowly varying functions at infinity, that is, R_0 is the class of positive measurable l , defined on some neighborhood of infinity, satisfying

$$\forall \lambda > 0, \quad \lim_{x \rightarrow \infty} l(\lambda x)/l(x) = 1.$$

For $l \in R_0$, the class Π_l is the class of measurable g satisfying

$$\forall \lambda \geq 1, \quad \lim_{x \rightarrow \infty} \{g(\lambda x) - g(x)\}/l(x) = c \log \lambda$$

for some constant c called the l -index of g . It is useful to name the class of functions of which we define the Fourier sine transforms. The function $f : [0, \infty) \rightarrow \mathbb{R}$ belongs to $DL_{loc}^1[0, \infty)$ if it is locally integrable and eventually non-increasing on $[0, \infty)$, $\lim_{t \rightarrow \infty} f(t) = 0$. For $f \in DL_{loc}^1[0, \infty)$, we define its *Fourier sine transform* F_s by

$$(1.1) \quad F_s(\xi) = \int_0^{\infty-} f(t) \sin t\xi dt \quad (0 < \xi < \infty),$$

where we write $\int_0^{\infty-}$ to denote an improper integral obtained from \int_0^M by letting $M \uparrow \infty$. Since the improper integral on the right converges uniformly on each (a, ∞) with $a > 0$, F_s is a continuous function on $(0, \infty)$. See the proof of Theorem 6 of Titchmarsh [6].

Here is the main theorem of this paper:

1991 *Mathematics Subject Classification.* Primary 40E05; Secondary 42A32, 60E10.

THEOREM 1.1. Let $l \in R_0$ and $f \in DL_{loc}^1[0, \infty)$. Let F_s be the Fourier sine transform of f . Then

$$(1.2) \quad f(t) \sim t^{-2}l(t) \quad (t \rightarrow \infty)$$

if and only if

$$(1.3) \quad xF_s(1/x) \in \Pi_l \text{ in } x \text{ with } l\text{-index } 1.$$

The analogue for Fourier sine series is:

THEOREM 1.2. Let $l \in R_0$. Suppose that the real sequence $\{b_n\}$ is eventually non-increasing, and tends to 0 as $n \rightarrow \infty$. We set

$$(1.4) \quad g_s(\xi) = \sum_{n=1}^{\infty} b_n \sin n\xi \quad (0 < \xi < 2\pi).$$

Then

$$(1.5) \quad b_n \sim n^{-2}l(n) \quad (n \rightarrow \infty)$$

if and only if

$$(1.6) \quad xg_s(1/x) \in \Pi_l \text{ in } x \text{ with } l\text{-index } 1.$$

Now we recall the Abel-Tauber theorem of Pitman [4] which is closely related to Theorem 1.1. Let $l \in R_0$ and $0 < \alpha < 2$. Let $f \in DL_{loc}^1[0, \infty)$, and let F_s be the Fourier sine transform of f . Then, by Pitman [4],

$$(1.7) \quad f(t) \sim t^{-\alpha}l(t) \quad (t \rightarrow \infty)$$

if and only if

$$(1.8) \quad F_s(\xi) \sim \xi^{\alpha-1}l(1/\xi)\Gamma(1-\alpha)\cos(\pi\alpha/2) \quad (\xi \rightarrow 0+).$$

From this and Theorem 1.1, we see that the behavior (1.2) is a critical case; we need Π -variation to characterize (1.2) in terms of the Fourier sine transform of f . For the related Abel-Tauber theorems for integral transforms, we refer to Chapter 4 of Bingham, Goldie and Teugels [1].

In the proof of Theorem 1.1, we will find that it is enough to prove the theorem for $f \in DL_{loc}^1[0, \infty)$ finite and non-increasing on $[0, \infty)$. For such f , we have the inversion formula which represents f by the Fourier sine transform of f (see Theorem 7 of Titchmarsh [6]). However it is difficult to use to prove the Tauberian implication (1.3) \Rightarrow (1.2). The difficulty is in that the problem we have to deal with involves both Π -variation and improper integrals. The key to the proof is to reduce the problem to the analogous result of Inoue [2], [3] for Fourier cosine transforms.

The proofs of Theorems 1.1 and 1.2 will be given in section 2. In section 3, we apply Theorem 1.1 to the tail behavior of a probability distribution.

2. Proofs of Theorems 1.1 and 1.2

For $f \in DL_{loc}^1[0, \infty)$, we define its *Fourier cosine transform* F_c by

$$(2.1) \quad F_c(\xi) = \int_0^{\infty-} f(t) \cos t\xi dt \quad (0 < \xi < \infty).$$

We have the following Abel-Tauber theorem for Fourier cosine transforms:

THEOREM 2.1 (Pitman [4] and Inoue [2], [3]). Let $l \in R_0$ and $f \in DL_{loc}^1[0, \infty)$. Let F_c be the Fourier cosine transform of f . Then

$$(2.2) \quad f(t) \sim t^{-1}l(t) \quad (t \rightarrow \infty)$$

if and only if

$$(2.3) \quad F_c(1/\cdot) \in \Pi_l \text{ with } l\text{-index } 1.$$

The Abelian implication (2.2) \Rightarrow (2.3) is essentially Theorem 7 (iii) of Pitman [4], while the Tauberian assertion (2.3) \Rightarrow (2.2) is due to Inoue [2], [3].

Using Theorem 2.1, we shall prove Theorem 1.1.

PROOF OF THEOREM 1.1. Choose M so large that f is positive, finite and non-increasing on $[M, \infty)$. We set

$$g(t) = \begin{cases} f(M) & (0 \leq t < M), \\ f(t) & (M \leq t < \infty). \end{cases}$$

Let G_s be the Fourier sine transform of g . We set

$$(2.4) \quad a(x) = x^{-1} \sin x \quad (0 < x < \infty).$$

Then by the mean-value theorem,

$$(2.5) \quad |a(x) - a(y)| \leq \text{const.} |x - y| \quad (0 < x, y < 1),$$

so for any $\lambda > 1$,

$$\begin{aligned} & |\lambda x F_s(1/(\lambda x)) - x F_s(1/x) - \lambda x G_s(1/(\lambda x)) + x G_s(1/x)| / l(x) \\ &= \frac{1}{l(x)} \left| \int_0^M t \{f(t) - f(M)\} \{a(t/(\lambda x)) - a(t/x)\} dt \right| \\ &\leq \text{const.} \frac{(1 - \lambda^{-1})}{x l(x)} \int_0^M t^2 |f(t) - f(M)| dt \rightarrow 0 \quad (x \rightarrow \infty), \end{aligned}$$

whence (1.3) holds if and only if $x G_s(1/x) \in \Pi_l$ in x with l -index 1. Therefore we may assume that f is positive, finite and non-increasing on $[0, \infty)$.

First we assume (1.2). Then f is integrable over $[0, \infty)$, and so, by integration by parts,

$$(2.6) \quad F_s(\xi)/\xi = \int_0^{\infty-} h(t) \cos t \xi dt \quad (0 < \xi < \infty)$$

with

$$(2.7) \quad h(t) = \int_t^{\infty} f(s) ds \quad (0 \leq t < \infty).$$

Since (1.2) is equivalent to $h(t) \sim t^{-1}l(t)$ as $t \rightarrow \infty$, we immediately obtain (1.3) by the Abelian implication of Theorem 2.1.

Next, we prove that (1.3) implies (1.2). By the second mean-value theorem for integrals,

$$F_s(\xi) = \lim_{U \rightarrow \infty} \int_0^U f(t) \sin t\xi dt = \lim_{U \rightarrow \infty} f(0+) \int_0^\zeta \sin t\xi dt$$

for some $\zeta \in (0, U)$, and so $\xi F_s(\xi)$ is bounded on $(0, \infty)$. In particular, for any $x > 0$, $F_s(\xi)(1 - \cos x\xi)/\xi$ is integrable over $(0, \infty)$. Then by applying Theorem 38 of Titchmarsh [6] to the pair of $I_{[0, x]}$ and f , we obtain

$$\int_0^x f(t) dt = \frac{2}{\pi} \int_0^\infty F_s(\xi) \frac{1 - \cos x\xi}{\xi} d\xi \quad (0 < x < \infty).$$

By Theorem 3.7.4 of Bingham, Goldie and Teugels [1], (1.3) implies $|xF_s(1/x)| \in R_0$ in x , so $F_s(\xi)/\xi$ is integrable on $(0, 1)$ whence on $(0, \infty)$. Thus for any $x > 0$,

$$\int_0^x f(t) dt \leq \frac{4}{\pi} \int_0^\infty \frac{|F_s(\xi)|}{\xi} d\xi < \infty,$$

whence f is integrable over $(0, \infty)$. Again we arrive at (2.6) with (2.7), and similarly we obtain (1.2) from (1.3) by the Tauberian implication of Theorem 2.1. \square

Following the method of Soni and Soni [5], we shall prove Theorem 1.2 as a corollary of Theorem 1.1.

PROOF OF THEOREM 1.2. We set

$$f(t) = \begin{cases} 0 & (0 \leq t < 1/2), \\ b_n & (n - 1/2 \leq t < n + 1/2, \quad n = 1, 2, \dots). \end{cases}$$

Then f is in $DL_{loc}^1[0, \infty)$, and (1.5) is equivalent to

$$(2.8) \quad f(t) \sim t^{-2}l(t) \quad (t \rightarrow \infty).$$

Let F_s be the Fourier sine transform of f . Then by a simple calculation,

$$F_s(\xi) = a(\xi/2)g_s(\xi) \quad (0 < \xi < 2\pi),$$

where a is defined by (2.4). So we have for any $\lambda > 1$ and $x > 0$,

$$(2.9) \quad \begin{aligned} & \{\lambda x F_s(1/(\lambda x)) - x F_s(1/x)\} / l(x) \\ &= a(1/(2\lambda x)) \{\lambda x g_s(1/(\lambda x)) - x g_s(1/x)\} / l(x) \\ & \quad + x g_s(1/x) \{a(1/(2\lambda x)) - a(1/(2x))\} / l(x). \end{aligned}$$

By Theorem 3.7.4 of Bingham, Goldie and Teugels [1], (1.6) implies $|xg_s(1/x)| \in R_0$ in x , so by (2.5) the second term on the right of (2.9) tends to 0 as $x \rightarrow \infty$. Therefore

(1.6) implies $xF_s(1/x) \in \Pi_l$ in x with l -index 1, which, by the Tauberian implication of Theorem 1.1, implies (2.8) whence (1.5).

Conversely, we set $c(x) = 1/a(x)$ for $x > 0$. Then we also have

$$|c(x) - c(y)| \leq \text{const.}|x - y| \quad (0 < x, y < 1).$$

Arguing similarly, we obtain (1.6) from (1.5), which completes the proof. \square

3. Application to the tail behavior

In this section, we apply Theorem 1.1 to the tail behavior of a probability distribution. For the related results, we refer to Pitman [4] as well as pp. 336-337 of Bingham, Goldie and Teugels [1], and Inoue [3].

Let X be a real random variable defined on a probability space (Ω, \mathcal{F}, P) . The *tail-sum* of X is the function T defined by

$$T(x) = P(X \leq -x) + P(X > x) \quad (0 \leq x < \infty).$$

Let U be the real part of the characteristic function of X :

$$U(\xi) = E[\cos \xi X] \quad (\xi \in \mathbb{R}).$$

Then we have

$$\{1 - U(\xi)\}/\xi = \int_0^{\infty-} T(x) \sin \xi x dx \quad (0 < \xi < \infty).$$

Since T is finite and non-increasing on $[0, \infty)$, $\lim_{x \rightarrow \infty} T(x) = 0$, by Theorem 1.1 we immediately obtain

Theorem 3.1. *Let $l \in R_0$. Then $T(x) \sim x^{-2}l(x)$ as $x \rightarrow \infty$ if and only if $x^2\{1 - U(1/x)\} \in \Pi_l$ in x with l -index 1.*

REFERENCES

- [1] Bingham, N. H., Goldie, C. M. and J. L. Teugels, Regular variation, Cambridge University Press, Cambridge, 1987.
- [2] Inoue, A., Regularly varying correlation functions of solutions of delay Langevin equations, Preprint (1994).
- [3] Inoue, A., Tauberian theorems for Fourier cosine transforms, Preprint (1994).
- [4] Pitman, E. J. G., On the behaviour of the characteristic function of a probability distribution in the neighbourhood of the origin, J. Austral. Math. Soc. 8 (1968), 422-443.
- [5] Soni, K. and R. P. Soni, Slowly varying functions and asymptotic behaviour of a class of integral transforms III, J. Math. Anal. Appl. 49 (1975), 612-628.
- [6] Titchmarsh, E. C., Introduction to the Theory of Fourier Integrals, 2nd ed., Oxford University Press, London, 1948.

Department of Mathematics
Hokkaido University
Sapporo
060 Japan