



Title	Ginzburg landau equation and stable steady state solutions in a non-trivial domain
Author(s)	Jimbo, S.; Morita, Y.; Zhai, J.
Citation	Hokkaido University Preprint Series in Mathematics, 271, 1-17
Issue Date	1994-11-1
DOI	10.14943/83418
Doc URL	http://hdl.handle.net/2115/69022
Type	bulletin (article)
File Information	pre271.pdf



[Instructions for use](#)

**GINZBURG LANDAU EQUATION
AND STABLE STEADY
STATE SOLUTIONS IN A
NON-TRIVIAL DOMAIN**

S. Jimbo, Y. Morita and J. Zhai

Series #271. November 1994

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 245 H. Kubo, Slowly decaying solutions for semilinear wave equations in odd space dimensions, 30 pages. 1994.
- # 246 T. Nakazi, M. Yamada, (A_2) -Conditions and Carleson Inequalities, 27 pages. 1994.
- # 247 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and smoothing Effect of Solutions to the Benjamin-ono Equation, 17 pages. 1994.
- # 248 H. Kikuchi, Sheaf cohomology theory for measurable spaces, 12 pages. 1994.
- # 249 A. Inoue, Tauberian theorems for Fourier cosine transforms, 9 pages. 1994.
- # 250 S. Izumiya, G. T. Kossioris, Singularities for viscosity solutions of Hamilton-Jacobi equations, 23 pages. 1994.
- # 251 H. Kubo, K. Kubota, Asymptotic behaviors of radially symmetric solutions of $\square u = |u|^p$ for super critical values p in odd space dimensions, 51 pages. 1994.
- # 252 T. Mikami, Large Deviations and Central Limit Theorems for Eyraud-Farlie-Gumbel-Morgenstern Processes, 9 pages. 1994.
- # 253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.
- # 254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.
- # 255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.
- # 256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.
- # 257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.
- # 258 Y-G. Chen, Y. Giga, T. Hitaka and M. Honma, A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature, 18 pages. 1994.
- # 259 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 7 pages. 1994.
- # 260 T. Mikami, Copula fields and its applications, 14 pages. 1994.
- # 261 A. Inoue, An Abel-Tauber theorem for Fourier sine transforms, 6 pages. 1994.
- # 262 N. Kawazumi, Homology of hyperelliptic mapping class groups for surfaces, 13 pages. 1994.
- # 263 Y. Giga, M. E. Gurtin, A comparison theorem for crystalline evolution in the plane, 14 pages. 1994.
- # 264 J. Wierzbicki, On Commutativity of Diagrams of Type II_1 Factors, 26 pages. 1994.
- # 265 N. Hayashi, T. Ozawa, Schrödinger Equations with nonlinearity of integral type, 12 pages. 1994.
- # 266 T. Ozawa, On the resonance equations of long and short waves, 8 pages. 1994.
- # 267 T. Mikami, A sufficient condition for the uniqueness of solutions to a class of integro-differential equations, 9 pages. 1994.
- # 268 Y. Giga, Evolving curves with boundary conditions, 10 pages. 1994.
- # 269 A. Arai, Operator-theoretical analysis of representation of a supersymmetry algebra in Hilbert space, 12 pages. 1994.
- # 270 A. Arai, Gauge theory on a non-simply-connected domain and representations of canonical commutation relations, 18 pages. 1994.

GINZBURG LANDAU EQUATION AND STABLE STEADY STATE SOLUTIONS IN A NON-TRIVIAL DOMAIN

SHUICHI JIMBO*, YOSHIHISA MORITA** AND JIAN ZHAI*

*Department of Mathematics, Faculty of Science
Hokkaido University

**Department of Applied Mathematics and Informatics
Faculty of Science and Technology,
Ryukoku University

ABSTRACT. The Ginzburg Landau equation with a large parameter is studied in a bounded domain with the Neumann B.C. It is shown that if $n = 2$ or $n = 3$ and the domain is not simply connected, many kinds of non-constant stable equilibrium solutions exist.

§1. Introduction

In this paper we study the existence of non-constant stable equilibrium solutions of the following Ginzburg Landau equation,

$$(1.1) \quad \begin{cases} \frac{\partial \Phi}{\partial t} = \Delta \Phi + \lambda(1 - |\Phi|^2)\Phi & \text{in } (0, \infty) \times \Omega, \\ \frac{\partial \Phi}{\partial \nu} = 0 & \text{on } (0, \infty) \times \partial\Omega \quad (\Phi : \mathbb{C} - \text{valued}), \end{cases}$$

where $\Omega \subset \mathbb{R}^n$ is a bounded domain with C^3 boundary and $\lambda > 0$ is a parameter.

Ginzburg and Landau (1950) introduced a model of the superconductivity phenomena in the low temperature. Afterwards the important features of this phenomenon were analyzed through the GL equation. Moreover the successful idea of GL model has been used in many other fields of physics. The original GL equation is the system of equations of the wave function Φ of the electrons in a material and the vector potential A of the magnetic field made by the current of the electrons. The equation (1.1) is one of the simplified models and it is obtained by putting $A \equiv 0$ in the original equation and it is sometimes used in the case that the magnetic effect is expected to be very small and negligible inside the superconductor Ω . We are concerned with the occurrence of a permanent current of electrons without outer (magnetic) driving force. Mathematically speaking, it is the problem of existence of non-constant stable equilibrium solution of (1.1). In [15] it was proved that if Ω is convex, there is no non-constant stable solution to (1.1) for any $\lambda > 0$. This shows that the domain Ω should be more or less complicated so that (1.1) has non-constant stable steady state solutions. In [15], [16] it was proved that non-constant stable equilibria exist in a ring shaped domain (solid torus) for large $\lambda > 0$. In this

paper we deal with more general domains than that in [15], [16]. Precisely speaking, we will obtain the same result for the domains, which are not simply connected in the case $n = 2$ or $n = 3$ (cf. Fig. 1). From this result we see that the situation of solutions of the GL equation is very different from that of the (real valued) single reaction diffusion equation and the competition diffusion system (2 species) with Neumann B.C., because these equations do not have non-constant stable equilibria in an annulus (see also [7], [17], [18], [19], [20]).

Let us fix the meaning of the stability of an equilibrium solution. (1.1) defines a dynamical system in the function space $X = C^0(\bar{\Omega}; \mathbb{C})$. We consider the stability in the sense of Lyapunov, which is defined as follows.

Definition 1.1. An equilibrium solution Ψ of (1.1) is *stable* iff for any neighborhood $W (\ni \Psi)$ in X there exists a neighborhood of $W' (\ni \Psi)$ such that any solution $\Phi(t, \cdot)$ of (1.1) with $\Phi(0, \cdot) \in W'$ satisfies $\Phi(t, \cdot) \in W (\forall t \geq 0)$.

For other important problems (vortices, motion of vortices, magnetic quantization) of Ginzburg Landau equation with or without magnetic effect, see [2], [3], [4], [8], [9], [14], [25] and the references therein.

Acknowledgement

The authors would like to express their thanks to Prof. I. Takagi (Tohoku Univ.), Prof. M. Morimoto (Okayama Univ.) for valuable discussions and to Prof. Y. Giga (Hokkaido Univ.) for the information concerning the inequality in Prop. 4.1.

§2. Main results

In this section we present the main results. Let $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) be a bounded domain with C^3 boundary and consider the following equation,

$$(2.1) \quad \begin{cases} \Delta \Phi + \lambda(1 - |\Phi|^2)\Phi = 0 & \text{in } \Omega, \\ \frac{\partial \Phi}{\partial \nu} = 0 & \text{on } \partial\Omega \quad (\Phi : \mathbb{C} - \text{valued}), \end{cases}$$

where ν is the outward normal vector on $\partial\Omega$. We deal with the domains with the following condition.

(A) There exists a continuous map $\theta_0 : \bar{\Omega} \rightarrow S^1$ which is not homotopy equivalent to a constant value map.

Now we present the main theorem of this paper.

Theorem 2.1. Assume the condition (A). Then there exists a non-constant stable equilibrium solution Φ_λ to (2.1) for large $\lambda > 0$. Moreover $\Phi_\lambda(x) \neq 0$ in Ω and the map

$$(2.2) \quad \bar{\Omega} \ni x \rightarrow \Phi_\lambda(x)/|\Phi_\lambda(x)| \in S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$$

is homotopic to θ_0 for large $\lambda > 0$.

In the case of low dimensions, the condition (A) reduces to a more comprehensive property of Ω . Precisely speaking, (A) holds if $n = 2$ or 3 and Ω is not simply connected (cf. Proposition A.1 in Appendix). Hence the next result follows immediately from the above theorem.

Corollary 2.2. Assume that $n = 2$ or $n = 3$. If Ω is not simply connected, the same result as Theorem 2.1 holds.

Remark 2.3. In the case that Ω is a solid torus in \mathbb{R}^3 , the same result was obtained in [16].

The proofs of the above results will be given in the remaining sections of this paper.

§3. Preliminaries and S^1 -valued (harmonic) functions

In this section we prepare a formulation which we will use in the proof of the construction of non-constant solutions to (2.1). Let $\widehat{\Omega}$ be the universal covering space of Ω which is endowed with the canonical metric such that the projection $\iota_1 : \widehat{\Omega} \rightarrow \Omega$ is locally isometric. It is well known that \mathbb{R} is the universal cover of S^1 . Let $\iota_2 : \mathbb{R} \rightarrow S^1$ be the covering map. Remark also that the fundamental group $\pi_1(\Omega)$ acts as an isometric diffeomorphism on $\widehat{\Omega}$ (the covering transformation group). We denote its action as follows

$$\widehat{\Omega} \times \pi_1(\Omega) : (z, \gamma) \mapsto z \cdot \gamma \in \widehat{\Omega}.$$

We denote the generators of $\pi_1(\Omega)$ by β_1, \dots, β_m with the relations

$$\begin{aligned} \beta_{m(1,i)}^{p(1,i)} \cdot \beta_{m(2,i)}^{p(2,i)} \cdots \beta_{m(n(i),i)}^{p(n(i),i)} &= e \\ (1 \leq m(j,i) \leq m, p(j,i) = 1 \text{ or } -1, 1 \leq j \leq n(i), n(i) \in \mathbb{N}, 1 \leq i \leq k) \end{aligned}$$

where $e \in \pi_1(\Omega)$ is the unit. For any continuous map $\theta : \Omega \rightarrow S^1$, there exists a continuous map $F : \widehat{\Omega} \rightarrow \widehat{S^1} = \mathbb{R}$ such that the following diagram is commutative (cf. Proposition 5.3 in [13]).

$$\begin{array}{ccc} \widehat{\Omega} & \xrightarrow{F} & \widehat{S^1} = \mathbb{R} \\ \iota_1 \downarrow & & \downarrow \iota_2 \\ \Omega & \xrightarrow{\theta} & S^1 \end{array}$$

$$(3.1) \quad \theta(\iota_1(z)) = \iota_2(F(z)) \quad \text{for } \forall z \in \widehat{\Omega}.$$

F is unique up to an additive constant (integer $\times 2\pi$). Such F is called a lift of θ . On the other hand, what is the totality of \mathbb{R} -valued continuous functions in $\widehat{\Omega}$, which are obtained by lifting all the continuous maps on Ω into S^1 ? A standard argument in the theory of the covering spaces gives the following result.

Lemma 3.1. For a \mathbb{R} -valued continuous function F in $\widehat{\Omega}$, there exists a continuous map θ on Ω into S^1 satisfying (3.1) if and only if there exists a unique set of integers $(\ell_1, \dots, \ell_m) \in \mathbb{Z}^m$ with

$$p(1, i)\ell_{m(1, i)} + p(2, i)\ell_{m(2, i)} \cdots + p(n(i), i)\ell_{m(n(i), i)} = 0 \quad (1 \leq i \leq k)$$

such that

$$(3.2) \quad F(z \cdot \beta_i) = F(z) + 2\pi\ell_i \quad (\forall z \in \widehat{\Omega}, \quad 1 \leq i \leq m).$$

Here $(\ell_1, \dots, \ell_m) \in \mathbb{Z}^m$ depends only on the homotopy class of θ .

Remark 3.2. F satisfies the condition (3.2) for $(\ell_1, \dots, \ell_m) = (0, \dots, 0)$ if and only if there exists a \mathbb{R} -valued continuous function ρ in Ω such that $\rho(\iota_1(z)) = F(z)$ for $z \in \widehat{\Omega}$. This correspondence is one-to-one. Hence, F can be identified with a function on Ω . By the same consideration, for a C^1 class \mathbb{R} -valued function F on $\widehat{\Omega}$ which satisfies (3.2) for (ℓ_1, \dots, ℓ_m) , $\nabla_z F$ can be identified with a \mathbb{R}^n -valued function on Ω because $\nabla_z F(z \cdot \beta_j) = \nabla_z F(z)$ for $z \in \widehat{\Omega}$ and $1 \leq j \leq m$.

$$\begin{array}{ccc} \widehat{\Omega} & \xrightarrow{F} & \mathbb{R} \\ \iota_1 \downarrow & & \\ \Omega & & \end{array}$$

Using the above formulation, we will prove that there exists a S^1 -valued harmonic function in Ω (with the Neumann B. C.) in each homotopy class of continuous maps on $\overline{\Omega}$ into S^1 . This fact is known in more general framework in the theory of harmonic maps on manifolds (cf. [12]). However we give the proof below because the following argument is important in the construction of the solutions of (2.1) in §4. The equation is written as follows,

$$(3.3) \quad \operatorname{div}(\nabla\theta) = 0 \quad \text{in } \Omega, \quad \frac{\partial\theta}{\partial\nu} \equiv \langle \nabla\theta \cdot \nu \rangle = 0 \quad \text{on } \partial\Omega.$$

Here $\nabla\theta$ is well-defined as a \mathbb{R}^n -valued function in Ω , while θ itself is a S^1 -valued function. We suppose the solutions of (3.3) to be regular up to the boundary because of the elliptic regularity theory and hence we naturally regard solution θ as a function on $\overline{\Omega}$.

Lemma 3.3. There exists a solution to (3.3) in each homotopy class of continuous mappings on $\overline{\Omega}$ into S^1 . Moreover, the solution is unique up to uniform rotation of values in S^1 .

(Proof of Lemma 3.3) Given any continuous map θ_0 on $\overline{\Omega}$ into S^1 , we will make a solution of (3.3) with the same homotopy as θ_0 . We can assume, without loss of generality, that θ_0 is C^3 up to $\overline{\Omega}$, because we can mollify θ_0 (if necessary) in each local patch of $\overline{\Omega}$ without changing the homotopy. We “lift” the equation (3.3) to

one on the universal covering space $\widehat{\Omega}$ so that we can argue about solutions which are \mathbb{R} -valued functions. Using Lemma 3.1, $\widehat{\theta}_0$ satisfies (3.2) for a $(\ell_1, \dots, \ell_m) \in \mathbb{Z}^m$. The equation for the lift $\widehat{\theta} : \widehat{\Omega} \rightarrow \mathbb{R}$, is

$$(3.4) \quad \begin{cases} \operatorname{div}_z(\nabla_z \widehat{\theta}) = 0 & \text{in } \widehat{\Omega}, \quad \frac{\partial \widehat{\theta}}{\partial \nu_z} = 0 & \text{on } \partial \widehat{\Omega}, \\ \text{with } \xi(z \cdot \beta_i) = \xi(z) + 2\pi \ell_i & \text{for } \forall z \in \widehat{\Omega} \text{ and } 1 \leq i \leq m. \end{cases}$$

Thus the problem turns to seek for $\widehat{\theta}$ satisfying (3.4) and the condition (3.2) for the same (ℓ_1, \dots, ℓ_m) as $\widehat{\theta}_0$. By transforming the unknown variable $\widehat{\theta}$ to ξ by $\xi = \widehat{\theta} - \widehat{\theta}_0$, we get the equation

$$(3.5) \quad \begin{cases} \operatorname{div}_z(\nabla_z \xi) = -\operatorname{div}(\nabla \widehat{\theta}_0) & \text{in } \widehat{\Omega}, \quad \frac{\partial \xi}{\partial \nu_z} = -\frac{\partial \widehat{\theta}_0}{\partial \nu_z} & \text{on } \partial \widehat{\Omega}, \\ \text{with } \xi(z \cdot \beta_i) = \xi(z) & \text{for } \forall z \in \widehat{\Omega} \text{ and } 1 \leq i \leq m. \end{cases}$$

From Remark 3.2, ξ in (3.5) can be identified with a function in \mathbb{R} -valued function in Ω (also denoted by ξ) and hence the equation (3.5) reduces to the equation on Ω ,

$$(3.6) \quad \operatorname{div}(\nabla \xi) = -\operatorname{div}(\nabla \theta_0) \quad \text{in } \Omega, \quad \frac{\partial \xi}{\partial \nu} = -\frac{\partial \theta_0}{\partial \nu} \quad \text{on } \partial \Omega.$$

It is well-known that (3.6) has a unique equation up to an arbitrary additive real constant and thus we obtain a required solution with the same homotopy class as θ_0 to the original equation (3.3). \square

§4. Construction of equilibrium solutions

Before the construction of solutions, we prepare an auxiliary inequality (cf. (4.2)) due to S. Campanato [6], which gives an estimate of norm of the resolvent of the 2nd order elliptic operator in the Hölder space. We rely on this inequality in the proof of the existence of the solution and its asymptotic behavior. We state this inequality in the simplest version for our later use. Consider

$$(4.1) \quad -\Delta U + \lambda U = f \quad \text{in } \Omega, \quad \frac{\partial U}{\partial \nu} = 0 \quad \text{on } \partial \Omega,$$

where $\lambda > 0$ is a positive parameter. Let $\alpha \in (0, 1)$ be an arbitrarily fixed constant. It is well known that for any $f \in C^\alpha(\overline{\Omega})$ and $\lambda > 0$, there exists a unique solution $U_\lambda \in C^{2+\alpha}(\overline{\Omega})$. Moreover the following estimate holds for this solution U_λ .

Proposition 4.1 (Campanato [6]). There exists a constant $K > 0$ (independent of $\lambda > 0$) such that

$$(4.2) \quad \|U_\lambda\|_{C^\alpha(\overline{\Omega})} \leq \frac{K}{\lambda} \|f\|_{C^\alpha(\overline{\Omega})} \quad (\lambda > 0).$$

We will construct equilibrium solutions to the Ginzburg Landau equation in the following form

$$(4.3) \quad \Phi(x) = w(x)e^{i\theta(x)}$$

where

$$w : \Omega \longrightarrow (0, \infty), \quad \theta : \Omega \longrightarrow S^1 = \mathbb{R}/2\pi\mathbb{Z}.$$

From the equation (2.1) we easily derive the following equations for w and θ .

$$(4.4) \quad \begin{cases} \Delta w + (\lambda(1 - w^2) - |\nabla\theta|^2) w = 0 & \text{in } \Omega, \\ \frac{\partial w}{\partial \nu} = 0 & \text{on } \partial\Omega, \end{cases}$$

$$(4.5) \quad \operatorname{div}(w^2 \nabla\theta) = 0 \quad \text{in } \Omega, \quad \frac{\partial\theta}{\partial \nu} \equiv \langle \nabla\theta \cdot \nu \rangle = 0 \quad \text{on } \partial\Omega.$$

We remark again that $\nabla\theta$ is well-defined as a \mathbb{R}^n -valued function in Ω for a C^1 map $\Omega \longrightarrow S^1 = \mathbb{R}/2\pi\mathbb{Z}$. The purpose of this section is to construct a nontrivial solution (w, θ) to (4.4)-(4.5). We will prove the following result.

Proposition 4.2. There exists a $\lambda_* > 0$ such that the system of equations (4.4) and (4.5) has a pair of solutions $(w_\lambda, \theta_\lambda)$ such that θ_λ is homotopic to θ_0 (given in the assumption in Theorem 2.1) with

$$(4.6) \quad 1 - \frac{c}{\lambda} \leq w_\lambda \leq 1 \quad \text{in } \Omega,$$

$$(4.7) \quad \limsup_{\lambda \rightarrow \infty} \|\nabla\theta_\lambda\|_{L^\infty(\Omega)} < \infty, \quad \lim_{\lambda \rightarrow \infty} \|\Delta\theta_\lambda\|_{L^\infty(\Omega)} = 0,$$

$$(4.8) \quad \lim_{\lambda \rightarrow \infty} \|\lambda(1 - w_\lambda^2) - |\nabla\theta_\lambda|^2\|_{L^\infty(\Omega)} = 0.$$

The sketch of the proof is as follows. A solution is obtained by seeking for the fixed point of the map $\Psi : \theta \longmapsto w \longmapsto \tilde{\theta}$ where θ is homotopic to θ_0 , w is a positive solution of (4.4) for this θ and $\tilde{\theta}$ is a solution of (4.5) (for this w) which is homotopic to θ_0 . Then using several kinds of estimates, we set up a situation where the Schauder fixed point theorem works.

(Proof of Propostion 4.2)

(Step 1:) Without loss of generality, we can assume that $\theta_0 : \bar{\Omega} \longrightarrow S^1$ is C^3 . We fix arbitrary points $p \in \Omega$ and $q \in S^1$. From Proposition 3.1, there exists a unique harmonic map $\theta_* : \Omega \longrightarrow S^1$ with Neumann B.C. such that $\theta_*(p) = q$. Fix arbitrary points $\hat{p} \in \hat{\Omega}$ and $\hat{q} \in \hat{S}^1 = \mathbb{R}$ such that $\iota_1(\hat{p}) = p$ and $\iota_2(\hat{q}) = q$. Remark that any continuous map $\theta : \Omega \longrightarrow S^1$ with $\theta(p) = q$ is uniquely lifted to a map $\hat{\theta} : \hat{\Omega} \longrightarrow \mathbb{R}$ such that $\hat{\theta}(\hat{p}) = \hat{q}$ (cf. [13]). We introduce a certain class of continuous maps, in which we will find a solution of (4.5)

$$E = \{\theta \in C^{1+\alpha}(\bar{\Omega}; S^1) \mid \theta(p) = q, \theta \text{ is homotopic to } \theta_*, d(\theta, \theta_*) \leq 1\},$$

where

$$d(\theta_1, \theta_2) = \|\widehat{\theta}_1 - \widehat{\theta}_2\|_{C^{1+\alpha}(\overline{\Omega})},$$

for $\theta_1, \theta_2 \in C^{1+\alpha}(\overline{\Omega}; S^1)$ such that θ_i is homotopic to θ_* and $\theta_i(p) = q$ ($i = 1, 2$). Remark that $\widehat{\theta}_1 - \widehat{\theta}_2$ is well-defined as a \mathbb{R} -valued function in Ω (see Remark 3.2). Under this metric d , E is a complete metric space and it is isometrically diffeomorphic to the following set

$$\{\rho \in C^{1+\alpha}(\overline{\Omega}) \mid \rho(p) = 0, \|\rho\|_{C^{1+\alpha}(\overline{\Omega})} \leq 1\}.$$

(Step 2:) Applying the same arguments in [15] (making upper-lower solution pair), we have a $\lambda_0 > 0$ such that (4.4) has a unique positive solution $w = w(\lambda, \theta)$ for any $\lambda \geq \lambda_0$ and $\theta \in E$ and an estimate

$$(4.9) \quad 1 - c/\lambda \leq w(\lambda, \theta; x) \leq 1 \quad (x \in \Omega),$$

for a constant c which is independent of λ and $\theta \in E$ and this inequality yields

$$\sup_{\lambda \geq \lambda_0} \sup_{\theta \in E} \sup_{x \in \Omega} |\lambda(1 - w(\lambda, \theta; x))^2 w(\lambda, \theta; x) - |\nabla \theta|^2 w(\lambda, \theta; x)| < +\infty.$$

Applying the Schauder estimate to the equation (4.4), we obtain $\{w(\lambda, \theta)\}_{\lambda \geq \lambda_0, \theta \in E}$ is bounded in $C^{1+\alpha}(\overline{\Omega})$. To see a more detailed asymptotics of $w(\lambda, \theta)$. Define a function g by the relation $w(\lambda, \theta; x) = 1 - g(\lambda, \theta; x)/\lambda$ and derive the equation satisfied by g .

$$(4.10) \quad \begin{cases} (2 - \frac{1}{\lambda} \Delta)g + (w + 2)(w - 1)g - |\nabla \theta|^2 w = 0 & \text{in } \Omega, \\ \partial g / \partial \nu = 0 & \text{on } \partial \Omega. \end{cases}$$

Applying Propostion 3.1 to (4.5), we have

$$\begin{aligned} \|g\|_{C^\alpha(\overline{\Omega})} &\leq c_1 \|(w + 2)(w - 1)g - |\nabla \theta|^2 w\|_{C^\alpha(\overline{\Omega})} \\ &\leq c_1(\epsilon \|g\|_{C^\alpha(\overline{\Omega})} + c_2 \|g\|_{C^0(\overline{\Omega})}) + c_3, \end{aligned}$$

for $\lambda \geq \lambda_1(\epsilon)$. $\epsilon > 0$ can be taken arbitrarily small. By taking $\epsilon > 0$ such that $0 < c_1 \epsilon \leq 1/2$, we obtain that there exists $c_4 > 0$ such that

$$(4.11) \quad \|g(\lambda, \theta)\|_{C^\alpha(\overline{\Omega})} = \lambda \|w(\lambda, \theta) - 1\|_{C^\alpha(\overline{\Omega})} \leq c_4$$

for $\lambda \geq \lambda_1$ and $\theta \in E$ and consequently there exists also $c_5 > 0$ such that

$$\|\lambda w(\lambda, \theta)(1 - w(\lambda, \theta)^2) - |\nabla \theta|^2 w(\lambda, \theta)\|_{C^\alpha(\overline{\Omega})} \leq c_5 \quad (\lambda \geq \lambda_1, \theta \in E).$$

Applying again the Schauder estimate in (4.4), we get that $\|w(\lambda, \theta)\|_{C^{2+\alpha}(\overline{\Omega})}$ is bounded uniformly in $\theta \in E$ and $\lambda \geq \lambda_1$. From the compactness of the inclusion map $C^{2+\alpha}(\overline{\Omega}) \hookrightarrow C^2(\overline{\Omega})$, we obtain

$$(4.12) \quad \lim_{\lambda \rightarrow \infty} \sup_{\theta \in E} \|w(\lambda, \theta) - 1\|_{C^2(\overline{\Omega})} = 0.$$

(Step 3:) Next we consider (4.5) for $w = w(\lambda, \theta)$ obtained above. To seek for a solution θ of (4.5) which is homotopic to θ_0 and $\theta(p) = q$, we do as in §3. We consider the following lifted equation of (4.5) in $\widehat{\Omega}$, which is satisfied by $\widehat{\theta}$ (lift of θ).

$$(4.13) \quad \operatorname{div}_z(w(\iota_1(z))^2 \nabla_z \widehat{\theta}) = 0 \quad \text{in } \widehat{\Omega}, \quad \frac{\partial \widehat{\theta}}{\partial \nu_z} = 0 \quad \text{on } \partial \widehat{\Omega},$$

where the unknown function $\widehat{\theta}$ is a \mathbb{R} -valued function in $\widehat{\Omega}$. By changing the unknown function from $\widehat{\theta}$ to η' by $\eta' = \widehat{\theta} - \widehat{\theta}_0$, we obtain the equation

$$\operatorname{div}_z(w(\iota_1(z))^2 \nabla_z \widehat{\eta}) = -\operatorname{div}_z(w(\iota_1(z))^2 \nabla_z \widehat{\theta}_0) \quad \text{in } \widehat{\Omega}, \quad \frac{\partial \widehat{\eta}}{\partial \nu_z} = -\frac{\partial \widehat{\theta}_0}{\partial \nu_z} \quad \text{on } \partial \widehat{\Omega}.$$

Now remark that we are seeking for θ which is homotopic to θ_0 and hence, from Lemma 3.1 and Remark 3.2, $\eta' = \widehat{\theta} - \widehat{\theta}_0$ can be regarded as a \mathbb{R} -valued function in Ω . That is, a unique $\eta : \Omega \rightarrow \mathbb{R}$ corresponds to η' in such a way that $\eta'(z) = \eta(\iota_1(z))$ in $\widehat{\Omega}$. The equation for η is

$$(4.14) \quad \operatorname{div}(w^2 \nabla \eta) = -\operatorname{div}(w^2 \nabla \theta_0) \quad \text{in } \Omega, \quad \frac{\partial \eta}{\partial \nu} = -\frac{\partial \theta_0}{\partial \nu} \quad \text{on } \partial \Omega.$$

It is easy to see that (4.14) has the unique solution $\eta(x)$ such that $\eta(p) = \widehat{q} - \widehat{\theta}_0(\widehat{p})$. Thus we obtain the solution $\widehat{\theta}(z) = \widehat{\theta}_0 + \eta(\iota_1(z))$ of (4.13). Hence $\theta(x) = \iota_2(\widehat{\theta}(z))$ with $\iota_1(z) = x \in \Omega$ is a solution of (4.5) for $w(x) = w(\lambda, \theta)$ such that $\theta(p) = q$. Denote this solution $\widetilde{\theta}$ by $\Psi_\lambda(\theta)$.

(Step 4:) We will check the continuity of the map $\Psi_\lambda : E \rightarrow E$. First, the continuity of the map

$$E \ni \theta \rightarrow w(\lambda, \theta) \in C^{2+\alpha}(\overline{\Omega})$$

is verified by estimating the difference $w(\lambda, \theta_1) - w(\lambda, \theta_2)$ through

$$(4.15) \quad \begin{cases} (\Delta - 2\lambda - |\nabla \theta_2|^2)(w_2 - w_1) + \lambda(w_2 - w_1)(3 - (w_1^2 + w_1 w_2 + w_2^2)) \\ \quad = w_1(|\nabla \theta_2|^2 - |\nabla \theta_1|^2) \quad \text{in } \Omega, \\ (\partial/\partial \nu)(w_2 - w_1) = 0 \quad \text{on } \partial \Omega, \end{cases}$$

where $w_i(x) = w(\lambda, \theta_i; x)$ for $i = 1, 2$. By the aid of the Schauder estimate and (4.6), we see that $\|w_2 - w_1\|_{C^{2+\alpha}(\overline{\Omega})}$ is small if $d(\theta_1, \theta_2)$ is small for $\theta_1, \theta_2 \in E$. Here we used that the coefficient of the second term is bounded for large λ and hence it is absorbed in “ λ ” in the first term. Next we consider the difference of $\widetilde{\theta}_1 \equiv \Psi_\lambda(\theta_1)$ and $\widetilde{\theta}_2 \equiv \Psi_\lambda(\theta_2)$. Similarly as in the above argument, the difference $\xi = \widetilde{\theta}_1 - \widetilde{\theta}_2$ can be regarded as a \mathbb{R} -valued function in Ω , which satisfies

$$(4.16) \quad \begin{cases} \operatorname{div}(w_1^2 \nabla \xi) = \operatorname{div}((w_2^2 - w_1^2) \nabla \theta_2) \quad \text{in } \Omega, \\ \partial \xi / \partial \nu = 0 \quad \text{on } \partial \Omega \quad \text{with } \xi(p) = 0. \end{cases}$$

Applying again the Schauder estimate to (4.16) and obtain that $\|\xi\|_{C^{2+\alpha}(\bar{\Omega})}$ is small if $\|w_2 - w_1\|_{C^{1+\alpha}(\bar{\Omega})}$ is small. Consequently we complete the proof of the continuity of Ψ_λ in E . In the above argument it is easy to see that $\Psi_\lambda(E)$ is relatively compact in E for large $\lambda > 0$. Applying the Schauder fixed point theorem, we get a fixed point θ_λ in E and complete the proof of the existence of solution $(w_\lambda, \theta_\lambda)$ by putting $w_\lambda = w(\lambda, \theta_\lambda)$. The estimates (4.6), (4.7), (4.8) follow from (4.9), (4.12) and (4.5). \square

§5. Stability of $\Phi_\lambda = w_\lambda e^{i\theta_\lambda}$

In this section, we will discuss the stability of the equilibrium solution Φ_λ of (1.1), which has been constructed in the previous section. For the later discussion, we express Φ_λ in terms of real functions. Let u_λ and v_λ be the real and imaginary part of Φ_λ , i.e., $\Phi_\lambda(x) = u_\lambda(x) + v_\lambda(x)i$. Clearly (u_λ, v_λ) is a solution of the following system of equations,

$$(5.1) \quad \begin{cases} \Delta \begin{pmatrix} u \\ v \end{pmatrix} + \lambda(1 - u^2 - v^2) \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \text{in } \Omega, \\ \frac{\partial}{\partial \nu} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \text{on } \partial\Omega. \end{cases}$$

Let us consider the linearized eigenvalue problem of (5.1) at $\Phi_\lambda = (u_\lambda, v_\lambda)$.

$$(5.2) \quad \begin{cases} \Delta \begin{pmatrix} \phi \\ \psi \end{pmatrix} + \lambda M(u_\lambda, v_\lambda) \begin{pmatrix} \phi \\ \psi \end{pmatrix} + \mu \begin{pmatrix} \phi \\ \psi \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \text{in } \Omega, \\ \frac{\partial}{\partial \nu} \begin{pmatrix} \phi \\ \psi \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \text{on } \partial\Omega, \end{cases}$$

where

$$(5.3) \quad M(u, v) = \begin{pmatrix} 1 - 3u^2 - v^2 & -2uv \\ -2uv & 1 - u^2 - 3v^2 \end{pmatrix}.$$

It is easy to see that (5.2) is a selfadjoint eigenvalue problem with the real eigenvalues of finite multiplicity.

Definition 5.1. Let $\{\mu_k(\lambda)\}_{k=1}^\infty$ be the set of the eigenvalues of (5.2) which is arranged in increasing order (with counting multiplicity). It is known that these eigenvalues are real values with finite multiplicity and the corresponding eigenfunctions can be chosen as real valued.

For the solution (u_λ, v_λ) , $(-v_\lambda, u_\lambda)$ is an eigenfunction of (5.2) for an eigenvalue $\mu = 0$. This implies $\{\mu_k(\lambda)\}_{k=1}^\infty \ni 0$. Remark also that $\{e^{ic}\Phi_\lambda \mid c \in \mathbb{R}\}$ forms a one dimensional family of solutions of (2.1). Hence, to prove the stability (cf. Definition 1.1) of Φ_λ , it suffices to show that 0 is a simple eigenvalue and all other eigenvalues are positive. Such argument to prove the stability of an equilibrium point in a continuum of equilibria (of finite dimension) in a dynamical system, is

standard and can be found in several literature (cf. Chapter 5 in [11]). We can prove the following results.

Proposition 5.2.

$$(5.4) \quad \lim_{\lambda \rightarrow \infty} \mu_k(\lambda) = \mu_k \quad (k \geq 1),$$

where $\{\mu_k\}_{k=1}^{\infty}$ are the set of the eigenvalues arranged in increasing order (counting multiplicity) of the Laplacian with Neumann boundary condition, i.e.,

$$(5.5) \quad \begin{cases} \Delta \psi + \mu \psi = 0 & \text{in } \Omega, \\ \frac{\partial \psi}{\partial \nu} = 0 & \text{on } \partial \Omega. \end{cases}$$

Moreover, let $\{(\phi_{k,\lambda}, \psi_{k,\lambda})\}_{k=1}^{\infty}$ be an system of the corresponding eigenfunctions of (5.2) which is orthonormalized in $L^2(\Omega) \times L^2(\Omega)$. Then

$$(5.6) \quad \lim_{\lambda \rightarrow \infty} (\|\nabla(\phi_{k,\lambda} \cos \theta_\lambda + \psi_{k,\lambda} \sin \theta_\lambda)\|_{L^2(\Omega)}^2 + \lambda \|\phi_{k,\lambda} \cos \theta_\lambda + \psi_{k,\lambda} \sin \theta_\lambda\|_{L^2(\Omega)}^2) = 0 \quad (k \geq 1).$$

This yields the following result which implies the stability of Φ_λ for large $\lambda > 0$.

Corollary 5.3. There exists $\lambda_* > 0$ and $d > 0$ (d is independent of λ) such that

$$(5.7) \quad \mu_1(\lambda) \equiv 0, \quad \mu_2(\lambda) \geq d \quad \text{for } \lambda \geq \lambda_*.$$

(Proof of Corollary 5.3.) As was stated before, $\{\mu_k(\lambda)\}_{k=1}^{\infty} \ni 0, \forall \lambda > 0$ and hence from $\mu_1 = 0$ and $\mu_2 > 0$, Propostion 5.2 yields that $\mu_1(\lambda) \equiv 0$. The other part follows directly. \square

Proof of Proposition 5.2.

To see more detailed situation of the eigenvalues of (5.2), transform the unknown variable (ϕ, ψ) into $(\hat{\phi}, \hat{\psi})$ by

$$(5.8) \quad \begin{pmatrix} \phi(x) \\ \psi(x) \end{pmatrix} = \begin{pmatrix} \cos \theta_\lambda(x) & -\sin \theta_\lambda(x) \\ \sin \theta_\lambda(x) & \cos \theta_\lambda(x) \end{pmatrix} \begin{pmatrix} \hat{\phi}(x) \\ \hat{\psi}(x) \end{pmatrix}$$

and get a new eigenvalue problem (which is still selfadjoint with the same eigenvalues),

$$(5.9) \quad \begin{cases} \Delta \begin{pmatrix} \hat{\phi} \\ \hat{\psi} \end{pmatrix} + \begin{pmatrix} -\nabla \theta_\lambda \nabla \hat{\psi} \\ \nabla \theta_\lambda \nabla \hat{\phi} \end{pmatrix} + (\lambda(1 - w_\lambda^2) - |\nabla \theta_\lambda|^2) \begin{pmatrix} \hat{\phi} \\ \hat{\psi} \end{pmatrix} \\ + \Delta \theta_\lambda \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \hat{\phi} \\ \hat{\psi} \end{pmatrix} - 2\lambda w_\lambda^2 \begin{pmatrix} \hat{\phi} \\ 0 \end{pmatrix} + \mu \begin{pmatrix} \hat{\phi} \\ \hat{\psi} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \text{in } \Omega, \\ \begin{pmatrix} \partial \hat{\phi} / \partial \nu \\ \partial \hat{\psi} / \partial \nu \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \text{on } \partial \Omega. \end{cases}$$

For the simplicity of notation we use (ϕ, ψ) in place of $(\widehat{\phi}, \widehat{\psi})$ in the following arguments. To compare the eigenvalue problems (5.5) and (5.9), we use the variational characterization of the eigenvalues of a selfadjoint operator (cf. [23]). Let us define $\mathcal{E}_\lambda(\phi, \psi)$ and $\mathcal{E}_\infty(\psi)$ as follows,

$$(5.10) \quad \mathcal{E}_\lambda(\phi, \psi) = \int_{\Omega} \{ |\nabla\phi|^2 + |\nabla\psi|^2 + (\nabla\theta_\lambda \cdot \nabla\psi)\phi - (\nabla\theta_\lambda \cdot \nabla\phi)\psi \\ - (\lambda(1 - w_\lambda^2) - |\nabla\theta_\lambda|^2)(\phi^2 + \psi^2) + 2\lambda w_\lambda^2 \phi^2 \} dx,$$

$$(5.11) \quad \mathcal{E}_\infty(\psi) = \int_{\Omega} |\nabla\psi|^2 dx.$$

Then the eigenvalues $\{\mu_k(\lambda)\}_{k=1}^\infty$ of (5.9) and $\{\mu_k\}_{k=1}^\infty$ of (5.5) and the corresponding systems of the eigenfunctions $\{(\phi_{k,\lambda}, \psi_{k,\lambda})\}_{k=1}^\infty$ and $\{\psi_k\}_{k=1}^\infty$, are characterized inductively by the following series of minimizing problems (5.12) and (5.13), respectively. Put $\phi_{0,\lambda} = \psi_{0,\lambda} = \psi_0 = 0$ in Ω for a convenience of the definition.

$$(5.12) \quad \mu_k(\lambda) = \min\{\mathcal{E}_\lambda(\phi, \psi); (\phi, \psi) \in E_k(\lambda), \|\phi\|_{L^2}^2 + \|\psi\|_{L^2}^2 = 1\}$$

where

$$E_k(\lambda) = \{(\phi, \psi) \in H^1(\Omega) \times H^1(\Omega); \int_{\Omega} (\phi\phi_{\ell,\lambda} + \psi\psi_{\ell,\lambda})dx = 0, 0 \leq \ell \leq k-1\}.$$

Let $(\phi_{k,\lambda}, \psi_{k,\lambda})$ be a minimizer (which exists) of (5.12).

$$(5.13) \quad \mu_k = \min\{\mathcal{E}_\infty(\psi); \psi \in E_k(\infty), \|\psi\|_{L^2} = 1\},$$

where

$$E_k(\infty) = \{\psi \in H^1(\Omega); \int_{\Omega} \psi\psi_\ell dx = 0, 0 \leq \ell \leq k-1\}.$$

Let ψ_k be a minimizer (which exists) of (5.13).

We first prove that each eigenvalue $\mu_k(\lambda)$ is bounded from below when $\lambda \rightarrow \infty$. That is,

$$(5.14) \quad \liminf_{\lambda \rightarrow \infty} \mu_k(\lambda) > -\infty \quad \text{for } k \geq 1.$$

From Proposition 4.2-(4.7), $\|\nabla\theta_\lambda\|_{L^\infty(\Omega)}$ is bounded in λ , there exists a $c > 0$ (independent of $\lambda > 0$) such that

$$|\nabla\theta_\lambda \cdot \nabla\psi_{k,\lambda}\phi_{k,\lambda} - \nabla\theta_\lambda \cdot \nabla\phi_{k,\lambda}\psi_{k,\lambda}| \\ \leq \frac{1}{2}(|\nabla\phi_{k,\lambda}|^2 + |\nabla\psi_{k,\lambda}|^2) + c(\phi_{k,\lambda}^2 + \psi_{k,\lambda}^2).$$

and consequently,

$$(5.15) \quad \begin{aligned} \mu_k(\lambda) = \mathcal{E}_\lambda(\phi_{k,\lambda}, \psi_{k,\lambda}) &\geq \frac{1}{2} \int_{\Omega} (|\nabla \phi_{k,\lambda}|^2 + |\nabla \psi_{k,\lambda}|^2) dx \\ &\quad - c' \int_{\Omega} (\phi_{k,\lambda}^2 + \psi_{k,\lambda}^2) dx + 2\lambda \int_{\Omega} \phi_{k,\lambda}^2 dx \quad (k \geq 1). \end{aligned}$$

Hence, using $\|\phi_{k,\lambda}\|_{L^2(\Omega)}^2 + \|\psi_{k,\lambda}\|_{L^2(\Omega)}^2 = 1$, we obtain (5.14).

We can prove that $\lim_{\lambda \rightarrow \infty} \mu_k(\lambda) = \mu_k$ for $k = 1, 2, 3, \dots$ inductively. We are going to carry out this procedure only for $k = 1, 2$ because even these cases ($k = 1, 2$) are enough to deduce Corollary 5.3, which is the main purpose of this section (stability of Φ_λ) and also because it is easy to see how the further arguments for $k = 3, 4, \dots$ proceed, from these cases $k = 1, 2$. Similar arguments which will appear hereafter, are also found in [16].

CASE : $k = 1$

Claim 1.

$$(5.16) \quad \limsup_{\lambda \rightarrow \infty} \mu_1(\lambda) \leq \mu_1.$$

(Proof of Claim 1)

From (5.12)

$$\mu_1(\lambda) \leq \mathcal{E}_\lambda(0, \psi_1) = \int_{\Omega} (|\nabla \psi_1|^2 + (\lambda(1 - w_\lambda^2) - |\nabla \theta_\lambda|^2) \psi_1^2) dx.$$

Using Proposition 4.2-(4.8), we get (5.16).

From Claim 1 and the inequality (5.15) for $k = 1$, we have,

Claim 2.

$$(5.17) \quad \limsup_{\lambda \rightarrow \infty} \int_{\Omega} (|\nabla \phi_{1,\lambda}|^2 + |\nabla \psi_{1,\lambda}|^2) dx < \infty, \quad \limsup_{\lambda \rightarrow \infty} \lambda \int_{\Omega} \phi_{1,\lambda}^2 dx < \infty.$$

The next claim completes the case $k = 1$.

Claim 3.

$$(5.18) \quad \lim_{\lambda \rightarrow \infty} \mu_1(\lambda) = \mu_1, \quad \lim_{\lambda \rightarrow \infty} \int_{\Omega} (|\nabla \phi_{1,\lambda}|^2 + \lambda \phi_{1,\lambda}^2) dx = 0.$$

(Proof of claim 3)

Take an arbitrary positive sequence $\{\lambda_m\}_{m=1}^\infty$ which tends to ∞ as $m \rightarrow \infty$. From (5.14) and Claim 1, $|\mu_1(\lambda)|$ is bounded when $\lambda \rightarrow \infty$. By the aid of Claim 2, there exist a subsequence $\{\eta_m\}$ of $\{\lambda_m\}$ and $\psi'_1 \in H^1(\Omega)$ such that

$$(5.19) \quad \begin{cases} \lim_{m \rightarrow \infty} \mu_1(\eta_m) \equiv \exists \mu \quad (\leq \mu_1), \\ \lim_{m \rightarrow \infty} \psi_{1,\eta_m} = \psi'_1 \quad \text{weakly in } H^1(\Omega) \quad \text{and strongly in } L^2(\Omega). \end{cases}$$

From (5.10),

$$(5.20) \quad \mu_1(\lambda) = \mathcal{E}_\lambda(\phi_{1,\lambda}, \psi_{1,\lambda}) = \int_{\Omega} \{ |\nabla \phi_{1,\lambda}|^2 + |\nabla \psi_{1,\lambda}|^2 + (\nabla \theta_\lambda \cdot \nabla \psi_{1,\lambda}) \phi_{1,\lambda} - (\nabla \theta_\lambda \cdot \nabla \phi_{1,\lambda}) \psi_{1,\lambda} - (\lambda(1 - w_\lambda^2) - |\nabla \theta_\lambda|^2)(\phi_{1,\lambda}^2 + \psi_{1,\lambda}^2) + 2\lambda w_\lambda^2 \phi_{1,\lambda}^2 \} dx$$

From Propostion 4.2, (5.19) and the lower semicontinuity of the norm under the weak convergence in Hilbert space,

$$(5.21) \quad (\mu_1 \geq) \quad \mu = \lim_{m \rightarrow \infty} \mu_1(\eta_m) = \lim_{m \rightarrow \infty} \mathcal{E}_{\eta_m}(\phi_{1,\eta_m}, \psi_{1,\eta_m}) \\ = \limsup_{m \rightarrow \infty} \int_{\Omega} \{ |\nabla \phi_{1,\eta_m}|^2 + |\nabla \psi_{1,\eta_m}|^2 + 2\eta_m w_{\eta_m}^2 \phi_{1,\eta_m}^2 \} dx \\ = \liminf_{m \rightarrow \infty} \int_{\Omega} \{ |\nabla \phi_{1,\eta_m}|^2 + |\nabla \psi_{1,\eta_m}|^2 + 2\eta_m w_{\eta_m}^2 \phi_{1,\eta_m}^2 \} dx \\ \geq \limsup_{m \rightarrow \infty} \int_{\Omega} |\nabla \psi_{1,\eta_m}|^2 dx \geq \liminf_{m \rightarrow \infty} \int_{\Omega} |\nabla \psi_{1,\eta_m}|^2 dx \\ \geq \int_{\Omega} |\nabla \psi'_1|^2 dx = \mathcal{E}_\infty(\psi'_1).$$

From $\|\psi'_1\|_{L^2(\Omega)} = 1$ and (5.13), $\mathcal{E}_\infty(\psi) \geq \mu_1$. Therefore, from (5.21),

$$\mathcal{E}_\infty(\psi'_1) = \mu = \mu_1 = \lim_{m \rightarrow \infty} \mu_1(\eta_m)$$

follows and ψ_1 is an eigenfunction of (5.5) corresponding to μ_1 . Accordingly, we get

$$\lim_{m \rightarrow \infty} \eta_m \int_{\Omega} \phi_{1,\eta_m}^2 dx = 0, \quad \lim_{m \rightarrow \infty} \int_{\Omega} |\nabla \phi_{1,\eta_m}|^2 dx = 0.$$

The whole sequence $\{\lambda_m\}$ was arbitrary and hence these facts (just proved) concludes Claim 3 and consequently, (5.3) and (5.4) for $k = 1$.

CASE : $k = 2$

Claim 1.

$$(5.22) \quad \limsup_{\lambda \rightarrow \infty} \mu_2(\lambda) \leq \mu_2.$$

(Proof of Claim 1)

Take an arbitrary positive sequence $\{\lambda_m\}$ such that $\lim_{m \rightarrow \infty} \lambda_m = \infty$. By the argument in the case $k = 1$, there exists a subsequence $\{\eta_m\}$ and an eigenfunction ψ'_1 of (5.5) corresponding to μ_1 such that

$$(5.23) \quad \lim_{m \rightarrow \infty} \psi_{1,\eta_m} = \psi'_1 \quad \text{weakly in } H^1(\Omega) \quad \text{and strongly in } L^2(\Omega).$$

We can take $\psi \in L.h.[\psi_1, \psi_2] \equiv \{a\psi_1 + b\psi_2 \mid a, b \in \mathbb{R}\}$ such that

$$(5.24) \quad \|\psi\|_{L^2(\Omega)} = 1, \quad (\psi, \psi'_1)_{L^2(\Omega)} = 0.$$

Put

$$\begin{pmatrix} \phi'_\lambda \\ \psi'_\lambda \end{pmatrix} = \begin{pmatrix} 0 \\ \psi \end{pmatrix} - (\psi \cdot \psi_{1,\lambda})_{L^2} \begin{pmatrix} \phi_{1,\lambda} \\ \psi_{1,\lambda} \end{pmatrix}.$$

Remark that $\lim_{m \rightarrow \infty} (\psi, \psi_{1,\eta_m})_{L^2(\Omega)} = 0$ and $\mathcal{E}_\infty(\psi) \leq \mu_2$. From (5.12),

$$\mu_2(\eta_m) \leq \mathcal{E}_{\eta_m}(\phi'_{\eta_m}, \psi'_{\eta_m}) / (\|\phi'_{\eta_m}\|_{L^2}^2 + \|\psi'_{\eta_m}\|_{L^2}^2).$$

Using (5.23),(5.24) with (5.18), we have,

$$\mathcal{E}_{\eta_m}(\phi'_{\eta_m}, \psi'_{\eta_m}) \leq \int_{\Omega} |\nabla \psi|^2 dx + o(1) \leq \mu_2 + o(1) \quad \text{as } m \rightarrow \infty$$

$$\|\phi'_{\eta_m}\|_{L^2}^2 + \|\psi'_{\eta_m}\|_{L^2}^2 = \|\psi'_1\|_{L^2}^2 = 1 + o(1) \quad \text{as } m \rightarrow \infty.$$

Thus $\limsup_{m \rightarrow \infty} \mu_2(\eta_m) \leq \mu_2$ follows. Since the sequence $\{\lambda_m\}$ ($\uparrow \infty$) was arbitrary, (5.22) is proved.

From (5.15), Claim 1 and Proposition 4.2, we have,

Claim 2.

$$(5.25) \quad \limsup_{\lambda \rightarrow \infty} \int_{\Omega} (|\nabla \phi_{2,\lambda}|^2 + |\nabla \psi_{2,\lambda}|^2) dx < \infty, \quad \limsup_{\lambda \rightarrow \infty} \lambda \int_{\Omega} \phi_{2,\lambda}^2 dx < \infty.$$

The next claim completes the case $k = 2$.

Claim 3.

$$(5.26) \quad \lim_{\lambda \rightarrow \infty} \mu_2(\lambda) = \mu_2, \quad \lim_{\lambda \rightarrow \infty} \int_{\Omega} (|\nabla \phi_{2,\lambda}|^2 + \lambda \phi_{2,\lambda}^2) dx = 0.$$

(Proof of Claim 3)

Take an arbitrary positive sequence $\{\lambda_m\}$ such that $\lim_{m \rightarrow \infty} \lambda_m = \infty$. From Claim 1, 2 and the case $k = 1$, there exist a subsequence $\{\eta_m\} \subset \{\lambda_m\}$ and an eigenfunction ψ'_1 of (5.5) corresponding to μ_1 and $\psi'_2 \in H^1(\Omega)$ such that

$$(5.27) \quad \begin{cases} \lim_{m \rightarrow \infty} \mu_2(\eta_m) \equiv \exists \mu \quad (\leq \mu_2), \\ \lim_{m \rightarrow \infty} \psi_{1,\eta_m} = \psi'_1 \quad \text{weakly in } H^1(\Omega) \quad \text{and strongly in } L^2(\Omega), \\ \lim_{m \rightarrow \infty} \phi_{2,\eta_m} = 0 \quad \text{weakly in } H^1(\Omega) \quad \text{and strongly in } L^2(\Omega), \\ \lim_{m \rightarrow \infty} \psi_{2,\eta_m} = \psi'_2 \quad \text{weakly in } H^1(\Omega) \quad \text{and strongly in } L^2(\Omega). \end{cases}$$

Using Proposition 4.2 and (5.27), we get

$$\begin{aligned}
(5.28) \quad (\mu_2 \geq) \quad \mu &= \lim_{m \rightarrow \infty} \mu_2(\eta_m) = \lim_{m \rightarrow \infty} \mathcal{E}_{\eta_m}(\phi_{2,\eta_m}, \psi_{2,\eta_m}) \\
&= \limsup_{m \rightarrow \infty} \int_{\Omega} \{ |\nabla \phi_{2,\eta_m}|^2 + |\nabla \psi_{2,\eta_m}|^2 + 2\eta_m w_{\eta_m}^2 \phi_{1,\eta_m}^2 \} dx \\
&= \liminf_{m \rightarrow \infty} \int_{\Omega} \{ |\nabla \phi_{2,\eta_m}|^2 + |\nabla \psi_{2,\eta_m}|^2 + 2\eta_m w_{\eta_m}^2 \phi_{1,\eta_m}^2 \} dx \\
&\geq \limsup_{m \rightarrow \infty} \int_{\Omega} |\nabla \psi_{2,\eta_m}|^2 dx \geq \liminf_{m \rightarrow \infty} \int_{\Omega} |\nabla \psi_{2,\eta_m}|^2 dx \\
&\geq \int_{\Omega} |\nabla \psi'_2|^2 dx = \mathcal{E}_{\infty}(\psi'_2).
\end{aligned}$$

On the other hand, from

$$(\phi_{2,\eta_m}, \phi_{1,\eta_m})_{L^2} + (\psi_{2,\eta_m}, \psi_{1,\eta_m})_{L^2} = 0, \quad \|\phi_{2,\eta_m}\|_{L^2} + \|\psi_{2,\eta_m}\|_{L^2} = 1,$$

we obtain $(\psi'_2, \psi'_1)_{L^2(\Omega)} = 0$ and $\|\psi'_2\|_{L^2} = 1$ respectively. Hence we have $\mathcal{E}_{\infty}(\psi'_2) \geq \mu_2$. (5.28) and this inequality prove that

$$\mathcal{E}_{\infty}(\psi'_2) = \mu = \mu_2 = \lim_{m \rightarrow \infty} \mu_2(\eta_m)$$

and ψ'_2 is an eigenfunction of (5.5) corresponding to μ_2 .

$$(5.30) \quad \lim_{m \rightarrow \infty} \int_{\Omega} (|\nabla \phi_{2,\eta_m}|^2 + \eta_m \phi_{2,\eta_m}^2) dx = 0,$$

follows simultaneously. Since the whole sequence $\{\lambda_m\}_{m=1}^{\infty}$ was arbitrary, (5.26) is proved. Consequently the case $k = 2$ is completed.

The cases for $k \geq 3$ proceed similarly.

Appendix.

We will prove the topological result which we used in §2 to deduce Corollary 2.2 from Theorem 2.1. Concerning the following arguments, we are indebted to Professor M. Morimoto in Okayama university. We are very grateful to him. Let Ω be a domain in §2, that is, $\Omega \subset \mathbb{R}^n$ is a bounded domain with C^3 boundary. We have the following proposition.

Proposition A.1 ([21]). If $n = 2$ and $n = 3$, the condition (A) (in §2) holds if and only if Ω is not simply connected.

(Proof of Proposition A.1)

We omit the case $n = 2$ because it is easy. We prove the case $n = 3$. Put $X = \bar{\Omega}$. Let Ω be simply connected. Let $f : X \rightarrow S^1$ be an arbitrary continuous map. Because X is simply connected, there exists a continuous map $\hat{f} : X \rightarrow \mathbb{R}$ such that $f(x) = \iota(\hat{f}(x))$ ($x \in X$) where $\iota : \mathbb{R} \rightarrow S^1$ is a covering map (cf. Proposition

5.3 in [13]). Define $f_t(x) = \iota(tf(x))$ ($0 \leq t \leq 1$). From this, it is easy to see that f is homotopic to a constant map. Conversely, assume that any continuous map $X \rightarrow S^1$ is homotopic to a constant map. Then $H^1(X) = 0$ (1-dim cohomology vanishes) (cf. Theorem 7.1 in [13]). The coefficient ring of the homology groups and cohomology groups in this proof is \mathbb{Z} . Remark that X is a 3-dimensional orientable compact smooth manifold with C^3 boundary ∂X . Using the Poincaré duality theorem, we have $H_2(X, \partial X) \cong H^1(X) = 0$. From the universal coefficient theorem, $H^1(X) \cong \text{Hom}(H_1(X), \mathbb{Z}) = 0$ and hence $H_1(X)$ is a finite abelian group. On the other hand, ∂X is a finite disjoint union of orientable 2 dimensional compact manifolds and so $H_1(\partial X)$ is torsion free, i.e. $H_1(\partial X) = \mathbb{Z} \oplus \cdots \oplus \mathbb{Z}$. Hence, from the homology exact sequence,

$$\cdots \rightarrow H_2(X, \partial X) \rightarrow H_1(\partial X) \rightarrow H_1(X) \rightarrow \cdots$$

we have that $H_1(\partial X) = 0$. This implies that the euler number of each component of ∂X is 2 and hence it is diffeomorphic to S^2 . By deforming the components of ∂X (the spheres) which are located inside of X to points, we see that X is simply connected. \square

Remark A.2. For $n \geq 4$, it is known that there exists $\Omega \subset \mathbb{R}^n$ which does not satisfy the condition (A) while it is not simply connected.

References

- [1] H. Amann, On the existence of positive solutions of nonlinear elliptic boundary value problems, *Indiana Univ. Math. J.* **21** (1971), 125-146.
- [2] P. Bauman, N. N. Carlson and D. Phillips, On the zeros of solutions to Ginzburg Landau type systems, *SIAM J. Math. Anal.* **24** (1993), 1283-1293.
- [3] P. Bauman, C.N. Chen, D. Phillips and P. Sternberg, Vortex annihilation in nonlinear heat flow for Ginzburg Landau systems, preprint.
- [4] F. Bethuel, H. Brezis and F. Helein, *Ginzburg Landau Vortices*, Birkhäuser (1994).
- [5] K.J. Brown, P.C. Dunne and R.A. Gardner, A semilinear parabolic system arising in the theory of superconductivity, *J. Diff. Eqs.* **40** (1981), 232-252.
- [6] S. Campanato, Generation of analytic semigroups, in the Hölder topology, by elliptic operators of second order with Neumann boundary condition, *Estratto da Le Matematiche* **35** (1980), 61-72.
- [7] R.G. Casten and C.J. Holland, Instability results for reaction diffusion equations with Neumann boundary conditions, *J. Diff. Eqs.* **27** (1978), 266-273.
- [8] Q. Du, M. Gunzberger and J. Peterson, Analysis and approximation of the Ginzburg Landau model of superconductivity, *SIAM Review* **34** (1992), 54-81.
- [9] C. Elliot, H. Matano and Q. Tang, Vector Landau Ginzburg equation and superconductivity second order phase transition, to appear in *European J. Appl. Math.*

- [10] D. Gilbarg and N. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer, New York, 1977.
- [11] D. Henry, Geometric Theory of Semilinear Parabolic Equations, Springer-Verlag, New York 1981.
- [12] R. Hamilton, Harmonic Maps of Manifolds with Boundary, Lecture Notes in Mathematics 471, Springer.
- [13] S.T. Hu, Homotopy Theory, Academic press, London 1959.
- [14] A. Jaffe and C. Taubes, Vortices and Monopoles, Birkhauser 1980.
- [15] S. Jimbo and Y. Morita, Stability of non-constant steady state solutions to a Ginzburg Landau equation in higher space dimensions, Nonlinear Anal. TMA, 22 (1994), 753-770.
- [16] S. Jimbo and Y. Morita, Ginzburg Landau equation and stable solutions in a rotational domain, preprint.
- [17] S. Jimbo, A construction of the perturbed solutions of semilinear elliptic equation in the singularly perturbed domain, J. Fac. Sci. Univ. Tokyo 36 (1989), 163-189.
- [18] K. Kishimoto and H.F. Weinberger, The spatial homogeneity of stable equilibria of some reaction diffusion equations on convex domain, J. Diff. Eqs. 58 (1985), 15-21.
- [19] H. Matano, Asymptotic behavior and stability of solutions semilinear diffusion equations, Publ. RIMS, Kyoto Univ. 15 (1979), 401-455.
- [20] H. Matano and M. Mimura, Pattern formation in competition diffusion systems in nonconvex domains, Publ. RIMS, Kyoto Univ. 19 (1983), 1049-1079.
- [21] M. Morimoto, personal communication.
- [22] M.H. Protter and H.F. Weinberger, Maximum Principle in Differential Equations, Pentice-Hall, Englewood Cliffs, 1967.
- [23] M. Reed and B. Simon, Methods of Mathematical Physics, Academic Press, New York. Vol. 4, 1978.
- [24] D. Sattinger, Monotone methods in nonlinear elliptic and parabolic boundary value problems, Indiana Univ. Math. J. 21 (1972), 979-1000.
- [25] Y. Yang, Existence, regularity and asymptotic behavior of the solution to the Ginzburg Landau equations on \mathbb{R}^3 , Commun. Math. Phys. 123 (1989), 147-161.