



Title	Global existence of small amplitude solutions for the Klein-Gordon-Zakharov equations
Author(s)	Tsutaya, K.
Citation	Hokkaido University Preprint Series in Mathematics, 273, 1-11
Issue Date	1994-12-1
DOI	10.14943/83420
Doc URL	http://hdl.handle.net/2115/69024
Type	bulletin (article)
File Information	pre273.pdf



[Instructions for use](#)

**Global Existence of Small
Amplitude Solutions for the
Klein-Gordon-Zakharov Equations**

Kimitoshi Tsutaya

Series #273. December 1994

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 247 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and smoothing Effect of Solutions to the Benjamin-ono Equation, 17 pages. 1994.
- # 248 H. Kikuchi, Sheaf cohomology theory for measurable spaces, 12 pages. 1994.
- # 249 A. Inoue, Tauberian theorems for Fourier cosine transforms, 9 pages. 1994.
- # 250 S. Izumiya, G. T. Kossioris, Singularities for viscosity solutions of Hamilton-Jacobi equations, 23 pages. 1994.
- # 251 H. Kubo, K. Kubota, Asymptotic behaviors of radially symmetric solutions of $\Delta u = |u|^p$ for super critical values p in odd space dimensions, 51 pages. 1994.
- # 252 T. Mikami, Large Deviations and Central Limit Theorems for Eyraud-Farlie-Gumbel-Morgenstern Processes, 9 pages. 1994.
- # 253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.
- # 254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.
- # 255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.
- # 256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.
- # 257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.
- # 258 Y-G. Chen, Y. Giga, T. Hitaka and M. Honma, A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature, 18 pages. 1994.
- # 259 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 7 pages. 1994.
- # 260 T. Mikami, Copula fields and its applications, 14 pages. 1994.
- # 261 A. Inoue, An Abel-Tauber theorem for Fourier sine transforms, 6 pages. 1994.
- # 262 N. Kawazumi, Homology of hyperelliptic mapping class groups for surfaces, 13 pages. 1994.
- # 263 Y. Giga, M. E. Gurtin, A comparison theorem for crystalline evolution in the plane, 14 pages. 1994.
- # 264 J. Wierzbicki, On Commutativity of Diagrams of Type II_1 Factors, 26 pages. 1994.
- # 265 N. Hayashi, T. Ozawa, Schrödinger Equations with nonlinearity of integral type, 12 pages. 1994.
- # 266 T. Ozawa, On the resonance equations of long and short waves, 8 pages. 1994.
- # 267 T. Mikami, A sufficient condition for the uniqueness of solutions to a class of integro-differential equations, 9 pages. 1994.
- # 268 Y. Giga, Evolving curves with boundary conditions, 10 pages. 1994.
- # 269 A. Arai, Operator-theoretical analysis of representation of a supersymmetry algebra in Hilbert space, 12 pages. 1994.
- # 270 A. Arai, Gauge theory on a non-simply-connected domain and representations of canonical commutation relations, 18 pages. 1994.
- # 271 S. Jimbo, Y. Morita and J. Zhai, Ginzburg Landau equation and stable steady state solutions in a non-trivial domain, 17 pages. 1994.
- # 272 S. Izumiya, A. Takiyama, A time-like surface in Minkowski 3-space which contains light-like lines, 7 pages. 1994.

Global Existence of Small Amplitude Solutions for the Klein-Gordon-Zakharov Equations

Kimitoshi Tsutaya *

Department of Mathematics
Hokkaido University
Sapporo 060, Japan

Key words and phrases : Global existence, Klein-Gordon-Zakharov equations.

1 Introduction

We consider the Cauchy problem of the Klein-Gordon-Zakharov equations in three space dimensions :

$$\partial_t^2 u - \Delta u + u = -nu, \quad t > 0, \quad x \in \mathbf{R}^3, \quad (1.1)$$

$$\partial_t^2 n - \Delta n = \Delta |u|^2, \quad t > 0, \quad x \in \mathbf{R}^3, \quad (1.2)$$

$$u(0, x) = u_0(x), \quad \partial_t u(0, x) = u_1(x), \quad (1.3)$$

$$n(0, x) = n_0(x), \quad \partial_t n(0, x) = n_1(x),$$

where $\partial_t = \partial/\partial t$, and $u(t, x)$ and $n(t, x)$ are functions from $\mathbf{R}_+ \times \mathbf{R}^3$ to \mathbf{C}^3 and from $\mathbf{R}_+ \times \mathbf{R}^3$ to \mathbf{R} , respectively. The system (1.1)-(1.2) describes the propagation of strong turbulence of the Langmuir wave in a high frequency plasma (see [14]).

Many results have been obtained concerning the global existence of small amplitude solutions for the coupled systems of the Klein-Gordon and wave equations with quadratic nonlinearity (see, e.g., [1-4, 6, 7, 9-11]). Two methods are known to be applicable to solve those systems. One is to use the invariant Sobolev space with respect to the generators of the Lorentz group. This was developed by Klainerman [7]. He also introduced the notion of the null condition to prove the existence of global solutions for the wave equations with quadratic nonlinearity. We note that

*Partially supported by THE SUHARA MEMORIAL FOUNDATION.

the null condition technique is based on the Lorentz invariance of the equations. Recently, Bachelot [1] and Georgiev [3] improved the null condition technique to show the global existence result for the Dirac-Klein-Gordon equations and the Maxwell-Dirac equations, respectively, which are physically important (see also Georgiev [4]). Another method is based on the theory of normal forms introduced by Shatah [10], which is an extension of Poincaré's theory of normal forms for the ordinary differential equations to the partial differential equations. The idea of this method is to transform the original system with quadratic nonlinearity into a new system with cubic nonlinearity. See also [12] and its references. Recently, applying the argument of normal forms to (1.1)–(1.2), Ozawa, Tsutaya and Tsutsumi [9] have proved the existence of global solutions to (1.1)–(1.3) for small initial data. However, in [9] one needs the high regularity assumptions on the data to ensure the global existence. Moreover, the global solution n of (1.1)–(1.3) constructed by [9] must belong to the homogeneous Sobolev space \dot{H}^{-1} of negative index. In this paper we prove that there exist the global solutions of (1.1)–(1.3) for small data using the invariant Sobolev space but without applying the null condition technique. We improve the regularity requirements on the initial data. Our proof seems simpler and shorter than that of [9].

Before we state the main results in the present paper, we give several notations. We put $\partial_j = \partial/\partial x_j$ for $j = 1, 2, 3$. Let $\Gamma = (\Gamma_j; j = 1, \dots, 10)$ denote the generators of the Poincaré group $(\partial_t, \partial_1, \partial_2, \partial_3, L_1, L_2, L_3, \Omega_{12}, \Omega_{23}, \Omega_{13})$, where

$$\begin{aligned} L_j &= x_j \partial_t + t \partial_j, \quad j = 1, 2, 3, \\ \Omega_{ij} &= x_i \partial_j - x_j \partial_i, \quad 1 \leq i < j \leq 3, \end{aligned}$$

and we put

$$\partial = (\partial_t, \partial_1, \partial_2, \partial_3).$$

For a multi-index $\alpha = (\alpha_1, \alpha_2, \alpha_3)$, we put

$$\partial_x^\alpha = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}.$$

For a multi-index $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)$, we put

$$\partial^\alpha = \partial_t^{\alpha_1} \partial_1^{\alpha_2} \partial_2^{\alpha_3} \partial_3^{\alpha_4}.$$

For a multi-index $\alpha = (\alpha_1, \dots, \alpha_{10})$, we put

$$\Gamma^\alpha = \Gamma_1^{\alpha_1} \dots \Gamma_{10}^{\alpha_{10}}.$$

For $1 \leq p \leq \infty$, let L^p denote the standard L^p space on \mathbf{R}^3 . For $m \geq 0$ and $s \geq 0$, we define the weighted Sobolev space $H^{m,s}$ on \mathbf{R}^3 as follows:

$$H^{m,s} = \{v \in L^2; (1 + |x|^2)^{s/2} (1 - \Delta)^{m/2} v \in L^2\}$$

We put $H^m \equiv H^{m,0}$ for $m \geq 0$. Let $\omega = (1 - \Delta)^{1/2}$ and $\omega_0 = (-\Delta)^{1/2}$.

We have the following theorem concerning the global existence of solutions to (1.1)-(1.3) for small initial data.

Theorem 1.1 *Let $0 < \varepsilon < 1/6$ and $k \geq 4$. Assume that $u_0 \in H^{k+5, k+4}$, $u_1 \in H^{k+4, k+4}$, $n_0 \in H^{k+4, k+4}$ and $n_1 \in H^{k+3, k+4}$. Then, there exists a $\delta > 0$ such that if*

$$\|u_0\|_{H^{k+5, k+4}} + \|u_1\|_{H^{k+4, k+4}} + \|n_0\|_{H^{k+4, k+4}} + \|n_1\|_{H^{k+3, k+4}} \leq \delta, \quad (1.4)$$

then (1.1)-(1.3) has the unique global solutions (u, n) satisfying

$$u \in \bigcap_{j=0}^{k+4} C^j([0, \infty); H^{k+4-j}), \quad (1.5)$$

$$n \in \bigcap_{j=0}^{k+4} C^j([0, \infty); H^{k+4-j}), \quad (1.6)$$

$$\begin{aligned} & \sum_{|\alpha|=k+4} \sup_{t \geq 0} (1+t)^{-\varepsilon} \{ \|\partial_t \Gamma^\alpha u(t)\|_{L^2} + \|\omega \Gamma^\alpha u(t)\|_{L^2} \} \\ & + \sum_{|\alpha| \leq k+4} \sup_{t \geq 0} (1+t)^{-\varepsilon} \|\Gamma^\alpha u(t)\|_{L^2} + \sum_{|\alpha| \leq k+4} \sup_{t \geq 0} \|\Gamma^\alpha n(t)\|_{L^2} \\ & + \sum_{|\alpha| \leq k} \sup_{\substack{t \geq 0 \\ x \in \mathbf{R}^2}} \{ |(1+t+|x|)^{3/2-2\varepsilon} \Gamma^\alpha u(t, x)| \\ & \quad + |(1+t+|x|) \Gamma^\alpha n(t, x)| \} < \infty. \end{aligned} \quad (1.7)$$

Remark 1.1 (i) By (1.7), we see that the right hand sides of (1.1)-(1.2) are integrable in time and therefore we find that the solutions (u, n) of (1.1)-(1.3)

constructed by Theorem 1.1 asymptotically approach the free solutions as $t \rightarrow \infty$. See [9].

(ii) As compared to [9], we have brought down the regularity assumptions on the data significantly. Instead, we need some spatial decay on the data, which is inevitable as far as the method depends on the invariant Sobolev norms.

(iii) The estimate (1.7) is close to the optimal one as regards the space-time behavior of solutions.

The following corollary follows easily from the proof of Theorem 1.1.

Corollary 1.2 *In addition to all the assumptions in Theorem 1.1, if $u_0, u_1, n_0, n_1 \in \bigcap_{m \geq 1} H^m$, then the solutions (u, n) given by Theorem 1.1 satisfy*

$$u(t, x), n(t, x) \in C^\infty([0, \infty) \times \mathbb{R}^3).$$

We prove Theorem 1.1 by the contraction argument in the next section. The main tools in the proof are the decay estimate of the inhomogeneous linear Klein-Gordon equation due to Georgiev [4] and the Sobolev inequality in the Minkowski space by Klainerman [6, 8] and Hörmander [5]. We show the global existence result for (1.1)–(1.3) by using these two inequalities and adopting the weight function in the norm slightly different from [2, 3] and [1]. Finally, we remark that the various constants in the estimate are simply denoted by C .

2 Proof of Theorem 1.1.

In this section we describe the proof of Theorem 1.1. We solve (1.1)–(1.3) by the contraction mapping principle. For $k \in \mathbb{N}$, $a \in \mathbb{R}$, we define the norms

$$\begin{aligned} |u|_{k,a} &= \sup_{\substack{s \geq 0 \\ y \in \mathbb{R}^3}} (1 + s + |y|)^{-a} \sum_{|\alpha| \leq k} |\Gamma^\alpha u(s, y)|, \\ \|u\|_{k,a} &= \sup_{s \geq 0} (1 + s)^{-a} \|u(s)\|_k, \end{aligned}$$

$$\begin{aligned}\|u(t)\|_k &= \sum_{|\alpha| \leq k} \|\Gamma^\alpha u(t)\|, \\ \|u(t)\| &= \|u(t)\|_{L^2(\mathbb{R}^3)}.\end{aligned}$$

We put

$$\begin{aligned}X &= X(k, M) \\ &= \{(u(t), n(t)); u(t) \in \bigcap_{j=0}^{k+4} C^j([0, \infty); H^{k+4-j}), \\ &\quad n(t) \in \bigcap_{j=0}^{k+4} C^j([0, \infty); H^{k+4-j}), u(0, x) = u_0(x), \partial_t u(0, x) = u_1(x), \\ &\quad n(0, x) = n_0(x), \partial_t n(0, x) = n_1(x), \|(u, n)\|_X \leq M\},\end{aligned}$$

where

$$\|(u, n)\|_X = \|u\|_{k, -3/2+2\varepsilon} + \|u\|_{k+4, \varepsilon} + \|\partial_t u\|_{k+4, \varepsilon} + \|n\|_{k+4, 0} \quad (k \geq 4, 0 < \varepsilon < 1/6),$$

and M is a positive constant, which will be determined later. We introduce the metric $d((u, n), (v, m))$ into X as follows.

$$d((u, n), (v, m)) = \|(u - v, n - m)\|_X$$

for $(u, n), (v, m) \in X$. We note that X is complete with respect to the metric d .

Given $(u, n) \in X$, we define the maps

$$\chi_1 : (u, n) \rightarrow \psi = \chi_1(u, n),$$

$$\chi_2 : (u, n) \rightarrow \varphi = \chi_2(u, n) \quad \text{for } (u, n) \in X$$

by solving the Cauchy problem

$$\partial_t^2 \psi - \Delta \psi + \psi = -nu, \quad t > 0, \quad x \in \mathbb{R}^3, \quad (2.1)$$

$$\partial_t^2 \varphi - \Delta \varphi = \Delta |u|^2, \quad t > 0, \quad x \in \mathbb{R}^3, \quad (2.2)$$

$$\psi(0, x) = u_0(x), \quad \partial_t \psi(0, x) = u_1(x), \quad (2.3)$$

$$\varphi(0, x) = n_0(x), \quad \partial_t \varphi(0, x) = n_1(x).$$

We put $\chi(u, n) = (\chi_1(u, n), \chi_2(u, n))$. Let δ be a positive constant satisfying (1.4). We show that if δ is sufficiently small, then the map χ is a contraction from X into itself.

We first evaluate $\psi = \chi_1(u, n)$. To derive the L^∞ estimate of ψ , we use the following time decay estimate of solution of the linear Klein-Gordon equations due to Georgiev [4].

Lemma 2.1 *Let $u(t, x)$ be a solution of the inhomogeneous linear Klein-Gordon equation :*

$$\begin{cases} \partial_t^2 u - \Delta u + u = f, & t > 0, \quad x \in \mathbb{R}^3, \\ u(0, x) = u_0(x), \quad \partial_t u(0, x) = u_1(x), & x \in \mathbb{R}^3. \end{cases}$$

Then, we have

$$\begin{aligned} (1+t+|x|)^{3/2}|u(t, x)| &\leq C \sum_{j=0}^{\infty} \sum_{|\alpha| \leq 4} \sup_{s \in (0, t)} \varphi_j(s) \|(1+s+|y|)\Gamma^\alpha f(s, y)\|_{L^2(\mathbb{R}_y^3)} \\ &\quad + C \sum_{j=0}^{\infty} \sum_{|\alpha| \leq 5} \|(1+|y|)^{3/2} \varphi_j(|y|)\Gamma^\alpha u(0, y)\|_{L^2(\mathbb{R}_y^3)}, \end{aligned}$$

where $\{\varphi_j\}_{j=0}^{\infty}$ is a Littlewood-Paley partition of unity, i.e.,

$$\sum_{j=0}^{\infty} \varphi_j(s) = 1, \quad s \geq 0; \quad \varphi_j \in C_0^\infty(\mathbb{R}), \quad \varphi_j \geq 0 \quad \text{for all } j \geq 0;$$

$$\text{supp } \varphi_j = [2^{j-1}, 2^{j+1}] \quad \text{for } j \geq 1, \quad \text{supp } \varphi_0 \cap \overline{\mathbb{R}_+} = [0, 2].$$

We apply Γ^α to (2.1) and compute the equation of $\Gamma^\alpha \psi$ on the basis of the commutation relation

$$[\square + 1, \Gamma] = 0. \tag{2.4}$$

Then, it follows from Lemma 2.1 and a finite dependence domain argument that

$$\begin{aligned} &(1+t+|x|)^{3/2-2\epsilon} |\Gamma^\alpha \psi(t, x)| \\ &\leq C \sum_{j=0}^{\infty} \sum_{|\lambda| \leq 4} \sup_{s \in (0, t)} \varphi_j(s) \|(1+s+|y|)^{1-2\epsilon} \Gamma^{\alpha+\lambda}(nu)(s, y)\| \\ &\quad + C \sum_{j=0}^{\infty} \sum_{|\lambda| \leq 5} \|(1+|y|)^{3/2} \varphi_j(|y|)\Gamma^{\alpha+\lambda} \psi(0, y)\| \\ &\quad + C \sum_{j=0}^{\infty} \sum_{|\lambda| \leq 4} \|(1+|y|)^{5/2} \varphi_j(|y|)\Gamma^{\alpha+\lambda} \partial_t \psi(0, y)\| \end{aligned} \tag{2.5}$$

for $|\alpha| \leq k$ (see [3], p. 867). From (2.1), we easily see that

$$\begin{aligned} & \sum_{j=0}^{\infty} \sum_{|\lambda| \leq 5} \|(1+|y|)^{3/2} \varphi_j(|y|) \Gamma^{\alpha+\lambda} \psi(0, y)\| \\ & + \sum_{j=0}^{\infty} \sum_{|\lambda| \leq 4} \|(1+|y|)^{5/2} \varphi_j(|y|) \Gamma^{\alpha+\lambda} \partial_t \psi(0, y)\| \leq C\delta, \quad |\alpha| \leq k. \end{aligned} \quad (2.6)$$

On the other hand, by the Sobolev inequality in the Minkowski space \mathbf{R}^{3+1} due to Klainerman [6, 8] and Hörmander [5], we have

$$\begin{aligned} & \|(1+s+|y|)^{1-2\varepsilon} \Gamma^{\alpha+\lambda} (nu)(s, y)\| \\ & \leq C \sum_{|\beta|+|\gamma|=\alpha+\lambda} \|(1+s+|y|)^{1-2\varepsilon} \Gamma^{\beta} n \cdot \Gamma^{\gamma} u(s, y)\| \\ & \leq C(1+s)^{-\varepsilon} \|n\|_{[k/2]+2, -1} \|u\|_{k+4, \varepsilon} \\ & \quad + C(1+s)^{-1/2} \|n\|_{k+4, 0} \|u\|_{[k/2]+2, -3/2+2\varepsilon} \\ & \leq C(1+s)^{-\varepsilon} \|n\|_{[k/2]+5, 0} \|u\|_{k+4, \varepsilon} \\ & \quad + C(1+s)^{-1/2} \|n\|_{k+4, 0} \|u\|_{[k/2]+2, -3/2+2\varepsilon} \\ & \leq C(1+s)^{-\varepsilon} M^2, \quad |\alpha| \leq k, |\lambda| \leq 4. \end{aligned} \quad (2.7)$$

Since

$$\sum_{j=0}^{\infty} \sup_{s \in (0, t)} \varphi_j(s) (1+s)^{-\varepsilon} \leq C,$$

inequalities (2.5)-(2.7) yield

$$\sum_{|\alpha| \leq k} (1+t+|x|)^{3/2-2\varepsilon} |\Gamma^{\alpha} \psi(t, x)| \leq C\delta + CM^2. \quad (2.8)$$

We next show the energy estimate of ψ . Applying Γ^{α} to (2.1) for $|\alpha| \leq k+4$, we have

$$\begin{aligned} & \|\Gamma^{\alpha} \psi(t)\| + \|\partial \Gamma^{\alpha} \psi(t)\| \\ & \leq C \left(\|u_0\|_{H^{k+5, k+4}} + \|u_1\|_{H^{k+4, k+4}} + \|n_0\|_{H^{k+3, k+4}} + \|n_1\|_{H^{k+2, k+4}} \right. \\ & \quad \left. + \sum_{|\alpha| \leq k+4} \int_0^t \|\Gamma^{\alpha} (nu)(s)\| ds \right) \\ & \leq C\delta + C \sum_{|\beta|+|\gamma| \leq k+4} \int_0^t \|\Gamma^{\beta} n \cdot \Gamma^{\gamma} u(s)\| ds \end{aligned}$$

$$\begin{aligned}
&\leq C\delta + C \int_0^t \left\{ (1+s)^{-1+\varepsilon} \|n\|_{[k/2]+2,-1} \|u\|_{k+4,\varepsilon} \right. \\
&\quad \left. + (1+s)^{-3/2+2\varepsilon} \|n\|_{k+4,0} \|u\|_{[k/2]+2,-3/2+2\varepsilon} \right\} ds \\
&\leq C\delta + C(1+t)^\varepsilon M^2,
\end{aligned}$$

where we have used the Sobolev inequality in the Minkowski space \mathbf{R}^{3+1} as before.

Hence, using the fact

$$[\partial, \Gamma^\alpha] = \sum_{|\beta| \leq |\alpha|-1} C_\beta \Gamma^\beta,$$

we obtain

$$\begin{aligned}
&(1+s)^{-\varepsilon} (\|\psi(t)\|_{k+4} + \|\partial\psi(t)\|_{k+4}) \\
&\leq C\delta + CM^2.
\end{aligned} \tag{2.9}$$

We next evaluate $\varphi = \chi_2(u, n)$. To derive the L^2 estimate of φ , we write the solution of (2.2) as follows :

$$\varphi = \varphi_0 + \varphi_1,$$

where φ_0 is a solution of $\partial_t^2 \varphi_0 - \Delta \varphi_0 = 0$ with the data n_0 and n_1 , and φ_1 is a solution of

$$\partial_t^2 \varphi_1 - \Delta \varphi_1 = \Delta |u|^2 \tag{2.10}$$

with zero data. Then, using the $L^2 - L^2$ estimate by Klainerman [7], we have

$$\begin{aligned}
\|\Gamma^\alpha \varphi_0(t)\| &\leq \|\Gamma^\alpha \varphi_0(0)\| \\
&\leq C(\|n_0\|_{H^{k+4,k+4}} + \|n_1\|_{H^{k+3,k+4}}) \\
&\leq C\delta.
\end{aligned} \tag{2.11}$$

On the other hand, to derive the L^2 estimate of φ_1 , we rewrite (2.10) applied by Γ^α as the following integral equation :

$$\Gamma^\alpha \varphi_1(t) = \sum_{|\alpha| \leq k+4} \int_0^t \frac{\sin \omega_0(t-s)}{\omega_0} (\partial_s + \partial_x) \partial \Gamma^\alpha (|u|^2)(s) ds, \quad |\alpha| \leq k+4,$$

where we have used the facts

$$[\square, \Gamma] = 0, \quad [\Gamma, \partial] = \partial.$$

Then, by integrating by parts and the Hardy-Littlewood-Sobolev inequality, we have

$$\begin{aligned}
\|\Gamma^\alpha \varphi_1(t)\| &\leq C \sum_{|\alpha| \leq k+4} \|\partial \Gamma^\alpha(|u|^2)(0)\|_{L^{5/3}(\mathbb{R}^3)} \\
&\quad + C \sum_{|\alpha| \leq k+4} \int_0^t \|\partial \Gamma^\alpha(|u|^2)(s)\| ds \\
&\leq C \|\partial u(0)\|_{k+4} \sum_{|\beta| \leq [(k+5)/2]} \|\Gamma^\beta u(0)\|_{L^3(\mathbb{R}^3)} \\
&\quad + C \int_0^t (1+s)^{-3/2+3\varepsilon} \|u\|_{[(k+5)/2], -3/2+2\varepsilon} \|\partial u\|_{k+4, \varepsilon} ds \\
&\leq C\delta^2 + CM^2, \quad |\alpha| \leq k+4,
\end{aligned} \tag{2.12}$$

since $\varepsilon < 1/6$. Thus, from (2.11) and (2.12), we obtain

$$\|\varphi(t)\|_{k+4} \leq C\delta + C\delta^2 + CM^2. \tag{2.13}$$

Inequalities (2.8), (2.9) and (2.13) yield

$$\|(\psi, \varphi)\|_X \leq C\delta + CM^2, \tag{2.14}$$

where C does not depend on δ and t . If we choose $\delta > 0$ and $M > 0$ so that

$$M \leq \frac{1}{4C}, \quad \delta \leq \frac{M}{4C},$$

then we obtain by (2.12)

$$\|(\psi, \varphi)\|_X \leq \frac{1}{2}M.$$

In the same way, we can prove that

$$d(\chi(u, n), \chi(v, m)) \leq \frac{1}{2}d((u, n), (v, m)), \quad (u, n), (v, m) \in X,$$

if $\delta > 0$ and $M > 0$ are chosen as above. Thus, we see that if $\delta > 0$ is sufficiently small, then $\chi(u, n)$ is a contraction map from X into itself. Therefore, there exist unique solutions $(u, n) \in X$ of (1.1)–(1.3).

Finally, the uniqueness of the solutions satisfying (1.1)–(1.3) follows by a similar argument as above. For details, see [2].

Acknowledgements

I would like to thank Professor Yoshi Tsutsumi for his valuable comments. I am also grateful to Professor Tohru Ozawa who pointed out mistakes in the first version of this paper.

References

- [1] A. Bachelot, *Problème de Cauchy globale pour des systèmes de Dirac-Klein-Gordon*, Ann. Inst. Henri Poincaré, Physique théorique **48**, 387–422 (1988).
- [2] V. Georgiev, *Global solution of the system of wave and Klein-Gordon equations*, Math. Z. **203**, 683–698 (1990).
- [3] V. Georgiev, *Small amplitude solutions of the Maxwell-Dirac equations*, Indiana Univ. Math. J. **40**, 845–883 (1991).
- [4] V. Georgiev, *Decay estimates for the Klein-Gordon equations*, Commun. Part. Diff. Eqs. **17**, 1111–1139 (1992).
- [5] L. Hörmander, *On Sobolev spaces associated with some Lie algebras*, Report 4, 1985, Inst. Mittag-Leffler.
- [6] S. Klainerman, *Uniform decay estimates and the Lorentz invariance of the classical wave equations*, Comm. Pure Appl. Math. **38**, 321–332 (1985).
- [7] S. Klainerman, *The null condition and global existence to nonlinear wave equations*, Lect. in Appl. Math. **23**, 293–326 (1986).
- [8] S. Klainerman, *Remarks on the global Sobolev inequalities in the Minkowski space R^{n+1}* , Comm. Pure Appl. Math. **40**, 111–117 (1987).
- [9] T. Ozawa, K. Tsutaya and Y. Tsutsumi, *Normal Form and Global Solutions for the Klein-Gordon-Zakharov Equations*, to appear in Ann. Inst. Henri Poincaré, Analyse non linéaire.
- [10] J. Shatah, *Normal forms and quadratic nonlinear Klein-Gordon equations*, Comm. Pure Appl. Math. **38**, 685–696 (1985).

- [11] T. Sideris, *Decay estimates for the three-dimensional inhomogeneous Klein-Gordon equation and applications*, Commun. Part. Diff. Eqs. **14**, 1421–1455 (1989)
- [12] J.C.H. Simon and E. Taflin, *The Cauchy problem for non-linear Klein-Gordon equations*, Commun. Math. Phys. **152**, 433-478 (1993).
- [13] W.A. Strauss, *Nonlinear Wave Equations*, CBMS Regional Conference Series in Mathematics, no. 73, Amer. Math. Soc., Providence, RI, 1989.
- [14] S.G. Thornhill and D. ter Haar, *Langmuir turbulence and modulational instability*, Phys. Reports (Sect. C of Phys. Lett.) **43**, 43–99 (1978).