



Title	Lattices of closure operators
Author(s)	Higuchi,A.
Citation	Hokkaido University Preprint Series in Mathematics, 298, 1-6
Issue Date	1995-6-1
DOI	10.14943/83445
Doc URL	http://hdl.handle.net/2115/69049
Type	bulletin (article)
File Information	pre298.pdf



[Instructions for use](#)

Lattices of closure operators

Akira Higuchi

Series #298. June 1995

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 274 H. Kubo, On the critical decay and power for semilinear wave equations in odd space dimensions, 22 pages. 1994.
- # 275 N. Terai, T. Hibi, Alexander duality theorem and second Betti numbers of Stanley-Reisner rings, 2 pages. 1995.
- # 276 N. Terai, T. Hibi, Stanley-Reisner rings whose Betti numbers are independent of the base field, 12 pages. 1995.
- # 277 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with cyclic polytopes, 11 pages. 1995.
- # 278 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with stacked polytopes, 8 pages. 1995.
- # 279 N. Terai, T. Hibi, Finite free resolutions and 1-skeletons of simplicial $(d - 1)$ -spheres, 3 pages. 1995.
- # 280 N. Terai, T. Hibi, Monomial ideals and minimal non-faces of Cohen-Macaulay complexes, 6 pages. 1995.
- # 281 A. Arai, N. Tominaga, Analysis of a family of strongly commuting self-adjoint operators with applications to perturbed d'Alembertians and the external field problem in quantum field theory, 44 pages. 1995.
- # 282 T. Mikami, Asymptotic behavior of the first exit time of randomly perturbed dynamical systems with a repulsive equilibrium point, 29 pages. 1995.
- # 283 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 17 pages. 1995.
- # 284 T. Nakazi, M. Yamada, Riesz's Functions In Weighted Hardy And Bergman Spaces, 20 pages. 1995.
- # 285 K. Hidano, K. Tsutaya, Scattering theory for nonlinear wave equations in the invariant Sobolev space, 32 pages. 1995.
- # 286 A. Arai, Strong coupling limit of the zero-energy-state density of the Dirac-Weyl operator with a singular vector potential, 8 pages. 1995.
- # 287 T. Nakazi, Factorizations of outer functions and extremal problems, 15 pages. 1995.
- # 288 A. Kishimoto, The Rohlin property for automorphisms of UHF algebras, 15 pages. 1995.
- # 289 K. Goto, A. Yamaguchi and I. Tsuda, Nine-bit states cellular automata are capable of simulating the pattern dynamics of coupled map lattice, 24 pages. 1995.
- # 290 Y. Giga, Interior derivative blow-up for quasilinear parabolic equations, 16 pages. 1995.
- # 291 F. Hiroshima, Functional Integral Representation of a Model in QED, 48 pages. 1995.
- # 292 N. Kawazumi, A Generalization of the Morita-Mumford Classes to Extended Mapping Class Groups for Surfaces, 11 pages. 1995.
- # 293 P. Aviles and Y. Giga, The distance function and defect energy, 23 pages. 1995.
- # 294 S. Izumiya and A. Takiyama, A time-like surface in Minkowski 3-space which contains pseudocircles, 11 pages. 1995.
- # 295 S. Izumiya, Local classifications of multi-valued solutions of quasilinear first order partial differential equations, 12 pages. 1995.
- # 296 A. Kishimoto, A Rohlin property for one-parameter automorphism groups, 27 pages. 1995.
- # 297 F. Hiroshima, Diamagnetic Inequalities for Systems of Nonrelativistic Particles with a Quantized Field, 23 pages. 1995.

Lattices of closure operators

Akira Higuchi
Department of Mathematics
Faculty of Science
Hokkaido University
Sapporo 060, Japan

June 15, 1995

Abstract

The system of all the closure operators on a set V forms a lattice [3]. This lattice is isomorphic to the lattice of all the topped intersection structures on V . This paper describes basic properties of these lattices, and gives a method of listing up all the members of these lattices.

1 Closure operators

A map $C : \mathcal{P}(V) \rightarrow \mathcal{P}(V)$ is called a *closure operator on V* if it satisfies, for $A, B \subseteq V$,

- (i) $A \subseteq C(A)$,
- (ii) $A \subseteq B \Rightarrow C(A) \subseteq C(B)$,
- (iii) $C(C(A)) = C(A)$.

We denote by $\text{CO}(V)$ the set of all the closure operators.

Let C be a closure operator on V . A subset A of V is called *closed with respect to C* if it is a fixed point of C . We define

$$\mathcal{A}_C = \{A \subseteq V \mid C(A) = A\}.$$

Proposition 1 [1] Let C be a closure operator on V . Then for each $A \subseteq V$, $C(A) = \bigcap \{B \in \mathcal{A}_C \mid B \supseteq A\}$.

Consequently, for each family \mathcal{A} of subsets of V , there is at most one closure operator $C_{\mathcal{A}}$ whose set of closed sets is equal to \mathcal{A} .

2 Topped intersection structures

A family \mathcal{A} of subsets of V is called an *intersection structure on V* if $\bigcap \mathcal{S} \in \mathcal{A}$ whenever $\mathcal{S} \subseteq \mathcal{A}$, and \mathcal{A} is called *topped* if $V \in \mathcal{A}$. We denote by

$\mathbf{TIStr}(V)$ the set of all the topped intersection structures on V . Obviously the minimum element of $\mathbf{TIStr}(V)$ is $\{V\}$, and the maximum element is $\mathcal{P}(V)$.

Proposition 2 [2] The map $\beta : C \mapsto \mathcal{A}_C$ is a bijection of $\mathbf{CO}(V)$ onto $\mathbf{TIStr}(V)$, with the inverse map given by $\beta^{-1}(\mathcal{A}) = C_{\mathcal{A}}$.

Lemma 3 Let \mathcal{A} be a topped intersection structure on V , X a member of \mathcal{A} , and $\mathcal{B} = \mathcal{A} \setminus \{X\}$. Then,

$$X \neq \bigcap \{B \in \mathcal{B} \mid B \supseteq X\} \iff \mathcal{B} \in \mathbf{TIStr}(V).$$

Proof.

(\implies) Let $X \neq \bigcap \{B \in \mathcal{B} \mid B \supseteq X\}$, and let \mathcal{S} be a subset of \mathcal{B} . Since $\mathcal{S} \subseteq \mathcal{B} \subseteq \mathcal{A}$, $\bigcap \mathcal{S} \in \mathcal{A}$. We prove that $\bigcap \mathcal{S} \neq X$ by contradiction.

Suppose $\bigcap \mathcal{S} = X$. Because $S \supseteq X$ for each $S \in \mathcal{S}$, we have $\mathcal{S} \subseteq \{B \in \mathcal{B} \mid B \supseteq X\}$. Hence $\bigcap \mathcal{S} \supseteq \bigcap \{B \in \mathcal{B} \mid B \supseteq X\} \supseteq X$. This implies $\bigcap \{B \in \mathcal{B} \mid B \supseteq X\} = X$, a contradiction. As a result, $\bigcap \mathcal{S} \in \mathcal{A} \setminus \{X\} = \mathcal{B}$.

(\impliedby) Let $\mathcal{B} \in \mathbf{TIStr}(V)$. Since $\{B \in \mathcal{B} \mid B \supseteq X\} \subseteq \mathcal{B}$, we have $\bigcap \{B \in \mathcal{B} \mid B \supseteq X\} \in \mathcal{B}$. Because $X \notin \mathcal{B}$, it follows $X \neq \bigcap \{B \in \mathcal{B} \mid B \supseteq X\}$.

q.e.d.

Proposition 4 $\mathbf{TIStr}(V)$ is a topped intersection structure on $\mathcal{P}(V)$.

Proof. Obviously $\mathcal{P}(V) \in \mathbf{TIStr}(V)$. Take $\mathbf{Sub} \subseteq \mathbf{TIStr}(V)$. We prove that $\bigcap \mathbf{Sub} \in \mathbf{TIStr}(V)$. For each $\mathcal{A} \in \mathbf{Sub}$, $V \in \mathcal{A}$ because $\mathcal{A} \in \mathbf{TIStr}(V)$. Hence $V \in \bigcap \mathbf{Sub}$. Let $\mathcal{S} \subseteq \bigcap \mathbf{Sub}$. For each $\mathcal{A} \in \mathbf{Sub}$, $\bigcap \mathcal{S} \in \mathcal{A}$ since $\mathcal{S} \subseteq \bigcap \mathbf{Sub} \subseteq \mathcal{A}$ and $\mathcal{A} \in \mathbf{TIStr}(V)$. Hence $\bigcap \mathcal{S} \in \bigcap \mathbf{Sub}$. Therefore we obtain $\bigcap \mathbf{Sub} \in \mathbf{TIStr}(V)$.

q.e.d.

3 Lattices of closure operators

We define a binary relation \preceq on $\mathbf{CO}(V)$ as the pointwise order, and define \ll as the covering relation of the relation \preceq . i.e.,

$$\begin{aligned} C \preceq D &\iff C(A) \subseteq D(A) \text{ for all } A \subseteq V, \\ C \ll D &\iff C \prec D \text{ and } C \preceq E \prec D \text{ implies } C = E. \end{aligned}$$

Obviously the relation \preceq is a partial order on $\mathbf{CO}(V)$.

Proposition 5 The map $\beta : C \mapsto \mathcal{A}_C$ is an order-isomorphism from $(\mathbf{CO}(V), \preceq)$ to $(\mathbf{TIStr}(V), \supseteq)$.

Proof. Let $C \preceq D$. For $A \subseteq V$, if $D(A) = A$, then $A \subseteq C(A) \subseteq D(A) = A$, i.e., $C(A) = A$. Hence $\mathcal{A}_C \supseteq \mathcal{A}_D$. Conversely, let $\mathcal{A}_C \supseteq \mathcal{A}_D$. For each $A \subseteq V$, we have $\bigcap \{B \in \mathcal{A}_C \mid B \supseteq A\} \subseteq \bigcap \{B \in \mathcal{A}_D \mid B \supseteq A\}$, i.e., $C(A) = D(A)$. Hence $C \preceq D$.

q.e.d.

Lemma 6 $C, D \in \text{CO}(V)$ and $C \ll D$ implies $|\mathcal{A}_C \setminus \mathcal{A}_D| = 1$.

Proof. Let X be a maximal element of $\mathcal{A}_C \setminus \mathcal{A}_D$, and let $\mathcal{B} = \mathcal{A}_C \setminus \{X\}$. From $\mathcal{B} \subseteq \mathcal{A}_D$, it follows $\bigcap \{B \in \mathcal{B} \mid B \supseteq X\} \supseteq \bigcap \{B \in \mathcal{A}_D \mid B \supseteq X\} = D(X) \supseteq X$. Moreover, $X \notin \mathcal{A}_D$ implies $X \neq D(X)$. Hence $X \neq \bigcap \{B \in \mathcal{B} \mid B \supseteq X\}$. Using Lemma 3.2, we obtain $\mathcal{B} \in \text{TIStr}(V)$. Since $C \prec C_{\mathcal{B}} \preceq D$ and $C \ll D$, we have $C_{\mathcal{B}} = D$. Therefore $\mathcal{A}_D = \mathcal{B} = \mathcal{A}_C \setminus \{X\}$.

q.e.d.

Let \mathcal{A} be a topped intersection structure on V . Since $\inf \mathcal{B} = \bigcap \mathcal{B}$ and $\sup \mathcal{B} = C_{\mathcal{A}}(\bigcup \mathcal{B})$ where $\mathcal{B} \subseteq \mathcal{A}$, (\mathcal{A}, \subseteq) forms a complete lattice [2]. Moreover, $(\text{TIStr}(V), \subseteq)$ and $(\text{CO}(V), \preceq)$ are also complete lattices, because $\text{TIStr}(V)$ is a topped intersection structure on $\mathcal{P}(V)$.

Theorem 7 Let $C = C_0 \ll C_1 \ll C_2 \ll \dots \ll C_n = D$ be a chain of $(\text{CO}(V), \preceq)$. Then the length of the chain is $|\mathcal{A}_C \setminus \mathcal{A}_D|$.

Proof. Let $C = C_0 \ll C_1 \ll C_2 \ll \dots \ll C_n = D$. Lemma 6 shows that $|\mathcal{A}_{C_i} \setminus \mathcal{A}_{C_{i+1}}| = 1$ for each i . Hence $|\mathcal{A}_C \setminus \mathcal{A}_D| = n$, i.e., the length of the chain is $|\mathcal{A}_C \setminus \mathcal{A}_D|$.

q.e.d.

We define a binary relation $<$ of $\mathcal{P}(V)$ as a linear extension of \preceq , and for $\mathcal{A}, \mathcal{B} \subseteq \mathcal{P}(V)$, we define $\mathcal{A} \ll \mathcal{B}$ if and only if there exists $X \in \mathcal{P}(V)$ such that

- (i) $\mathcal{A} = \mathcal{B} \setminus \{X\}$,
- (ii) X is a maximum element of $(\mathcal{P}(V) \setminus \mathcal{A}, <)$,
- (iii) $X \neq \bigcap \{A \in \mathcal{A} \mid A \supseteq X\}$.

Proposition 8 Let V be a finite set, and \mathcal{A} be a family of subsets of V . Then

$$\mathcal{A} \in \text{TIStr}(V) \iff \mathcal{A} = \mathcal{P}(V) \text{ or } \mathcal{A} \ll \mathcal{B} \text{ for some } \mathcal{B} \in \text{TIStr}(V).$$

Proof.

(\implies) Let $\mathcal{A} \in \text{TIStr}(V)$, and $\mathcal{A} \neq \mathcal{P}(V)$. We prove that $\mathcal{A} \ll \mathcal{B}$ for some $\mathcal{B} \in \text{TIStr}(V)$. Let X be the minimum element of $(\mathcal{P}(V) \setminus \mathcal{A}, <)$, and let $\mathcal{B} = \mathcal{A} \cup \{X\}$. Because $X \neq V$, we have $V \in \mathcal{B}$. Take $\mathcal{S} \subseteq \mathcal{B}$. If $X \notin \mathcal{S}$, then $\mathcal{S} \subseteq \mathcal{A}$ implies $\bigcap \mathcal{S} \in \mathcal{A}$, hence $\bigcap \mathcal{S} \in \mathcal{B}$. Otherwise, $\bigcap \mathcal{S} \subseteq X$ and X is the minimum element of $\mathcal{P}(V) \setminus \mathcal{A}$, hence $\bigcap \mathcal{S} \in \mathcal{A} \cup \{X\} = \mathcal{B}$. Therefore $\mathcal{B} \in \text{TIStr}(V)$. Moreover, we have $X \neq \bigcap \{A \in \mathcal{A} \mid A \supseteq X\}$ using Lemma 3. Hence $\mathcal{A} \ll \mathcal{B}$.

(\impliedby) Let $\mathcal{A} \ll \mathcal{B}$, $\mathcal{B} \in \text{TIStr}(V)$. Because $\mathcal{A} = \mathcal{B} \setminus \{X\}$ and $X \neq \bigcap \{A \in \mathcal{A} \mid A \supseteq X\}$, we have $\mathcal{A} \in \text{TIStr}(V)$ using Lemma 3.

q.e.d.

Consequently, there exists a chain

$$\mathcal{A} = \mathcal{A}_0 \ll \mathcal{A}_1 \ll \dots \ll \mathcal{A}_n = \mathcal{P}(V)$$

if and only if $\mathcal{A} \in \text{TIStr}(V)$. Moreover, such a chain, if it exists, is unique.

Hence, the Hasse diagram of $(\mathbf{TIStr}(V), \ll)$ is a spanning tree of the diagram of $(\mathbf{TIStr}(V), \subset)$ (figure 1, 2). We can list up all the topped intersection structures (and all the closure operators) of V by traveling the tree (table 1).

$ V $	$ \mathbf{CO}(V) $
1	2
2	7
3	61
4	2480
5	1385552

Table 1: the number of all the closure operators on a finite set V

I would like to express my gratitude to Professor Toru Tsujishita for his useful advice.

References

- [1] John P. Cleave, A study of logics (Oxford University Press, 1991).
- [2] B. A. Davey and H. A. Priestley, Introduction to lattices and order (Cambridge University Press, Cambridge, 1990).
- [3] Oystein Ore, Combinations of closure relations, Annals of Mathematics, vol. 44 (1943) 514-533.




Figure 1: the diagram of $(\text{TIStr}(V), \subset)$ ($|V| = 3$)




Figure 2: the diagram of $(\text{TIStr}(V), \ll)$ ($|V| = 3$)