



Title	Slice maps and multipliers of invariant subspaces
Author(s)	Nakazi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 313, 1-11
Issue Date	1995-11-1
DOI	10.14943/83460
Doc URL	http://hdl.handle.net/2115/69064
Type	bulletin (article)
File Information	pre313.pdf



[Instructions for use](#)

**Slice Maps And Multipliers
Of Invariant Subspaces**

Takahiko Nakazi

Series #313. November 1995

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #289 K. Goto, A. Yamaguchi and I. Tsuda, Nine-bit states cellular automata are capable of simulating the pattern dynamics of coupled map lattice, 24 pages. 1995.
- #290 Y. Giga, Interior derivative blow-up for quasilinear parabolic equations, 16 pages. 1995.
- #291 F. Hiroshima, Functional Integral Representation of a Model in QED, 48 pages. 1995.
- #292 N. Kawazumi, A Generalization of the Morita-Mumford Classes to Extended Mapping Class Groups for Surfaces, 11 pages. 1995.
- #293 P. Aviles and Y. Giga, The distance function and defect energy, 23 pages. 1995.
- #294 S. Izumiya and A. Takiyama, A time-like surface in Minkowski 3-space which contains pseudocircles, 11 pages. 1995.
- #295 S. Izumiya, Local classifications of multi-valued solutions of quasilinear first order partial differential equations, 12 pages. 1995.
- #296 A. Kishimoto, A Rohlin property for one-parameter automorphism groups, 27 pages. 1995.
- #297 F. Hiroshima, Diamagnetic Inequalities for Systems of Nonrelativistic Particles with a Quantized Field, 23 pages. 1995.
- #298 A. Higuchi, Lattices of closure operators, 6 pages. 1995.
- #299 S. Izumiya and W-Z. Sun, Singularities of solution surfaces for quasilinear 1st order partial differential equations, 9 pages. 1995.
- #300 D. Lehmann, M. Soares and T. Suwa, On the index of a holomorphic vector field tangent to a singular variety, 14 pages. 1995.
- #301 J. Zhai, Harmonic maps and Ginzburg-Landau type elliptic system, 20 pages. 1995.
- #302 M.-H. Giga and Y. Giga, Geometric evolution by nonsmooth interfacial energy, 15 pages. 1995.
- #303 S. Jimbo and J. Zhai, Ginzburg-Landau equation with magnetic effect: non-simply-connected domains, 21 pages. 1995.
- #304 T. Ozawa, On the nonlinear Schrödinger equations of derivative type, 27 pages. 1995.
- #305 N.H. Bingham and A. Inoue, Jordan's theorem for fourier and hankel transforms, 30 pages. 1995.
- #306 T. Honda and T. Suwa, Residue formulas for singular foliations defined by meromorphic functions on surfaces, 19 pages. 1995.
- #307 J. Yoshizaki, On the structure of the singular set of a complex analytic foliation, 25 pages. 1995.
- #308 A. Arai, Representation of canonical commutation relations in a gauge theory, the Aharonov-Bohm effect, and Dirac Weyl operator, 17 pages. 1995.
- #309 A. Arai and M. Hirokawa, On the existence and uniqueness of ground states of the spin-boson Hamiltonian, 20 pages. 1995.
- #310 S. Izumiya and T. Sano, Generic affine differential geometry of plane curves, 8 pages. 1995.
- #311 N. Kawazumi, On the stable cohomology algebra of extended mapping class groups for surfaces, 13 pages. 1995.
- #312 H.M.Ito and T. Mikami, Poissonian asymptotics of a randomly perturbed dynamical system: Flip-flop of the Stochastic Disk Dynamo, 20 pages. 1995.

Slice Maps And Multipliers Of Invariant Subspaces

by

Takahiko Nakazi*

Department of Mathematics
Hokkaido University
Sapporo 060, Japan

AMS Subject Classification(1991) : Primary 47 A 15, 32 A 35

Key Words And Phrases : Hardy space, several variables, invariant subspace,
slice map

*This research was partially supported by Grant-in-Aid for Scientific Research,
Ministry of Education

Abstract. Let $\overline{D^2}$ be the closed bidisc and T^2 be its distinguished boundary. For $(\alpha, \beta) \in \overline{D^2}$, let $\Phi_{\alpha\beta}$ be a slice map, that is, $(\Phi_{\alpha\beta}f)(\lambda) = f(\alpha\lambda, \beta\lambda)$ for $\lambda \in D$ and $f \in H^2(D^2)$. Then $\ker\Phi_{\alpha\beta}$ is an invariant subspace, and it is not difficult to describe $\ker\Phi_{\alpha\beta}$ and $\mathcal{M}(\ker\Phi_{\alpha\beta}) = \{\phi \in L^\infty(T^2) : \phi\ker\Phi_{\alpha\beta} \subset H^2(D^2)\}$. In this paper, we study the set $\mathcal{M}(M)$ of all multipliers for an invariant subspace M such that the common zero set of M contains that of $\ker\Phi_{\alpha\beta}$.

§1 Introduction

Let D^2 be the open unit disc in \mathbb{C}^2 and T^2 be its distinguished boundary. Normalized Lebesgue measure on T^2 is denoted by dm . For $1 \leq p < \infty$, $H^p(D^2)$ is the Hardy space and $L^p(T^2)$ is the Lebesgue space on T^2 . Let $N(D^2)$ denote the Nevanlinna class. Each f in $N(D^2)$ has radial limits f^* defined on T^2 a.e. Moreover, there is a singular measure $d\sigma_f$ on T^2 determined by f such that the least harmonic majorant $u(\log |f|)$ of $\log |f|$ is given by $u(\log |f|)(\zeta) = P_\zeta(\log |f^*| + d\sigma_f)$ where P_ζ denotes Poisson integration and $\zeta = (z, w) \in D^2$. Put $N_*(D^2) = \{f \in N(D^2) ; d\sigma_f \leq 0\}$, then $H^p(D^2) \subset N_*(D^2) \subset N(D^2)$ and $H^p(D^2) = N_*(D^2) \cap L^p(T^2) \subset N(D^2) \cap L^p(T^2)$. These facts are shown in [6, Theorem 3.3.5].

A closed subspace M of $H^2(D^2)$ is said to be invariant if $zM \subset M$ and $wM \subset M$. For an invariant subspace M of $H^2(D^2)$, set

$$\mathcal{M}(M) = \{\phi \in L^\infty(T^2) ; \phi M \subseteq H^2(D^2)\}.$$

If $M = qH^2(D^2)$ for some inner function q , it is trivial to see $\mathcal{M}(M) = \bar{q}H^\infty(D^2)$. In the case of one variable, an arbitrary invariant subspace M has the form $qH^2(D)$ for some inner function q by the famous Beurling theorem [1]. Hence $\mathcal{M}(M) = \bar{q}H^\infty(D)$. Hence the map $M \rightarrow \mathcal{M}(M)$ is one-to-one. However this result for invariant subspaces of $H^2(D^2)$ is not true. The author [4] studied the relation between M and $\mathcal{M}(M)$. To study $\mathcal{M}(M)$, D. Douglas and K. Yan [2] introduced the common zero set $\mathcal{Z}(M)$ and the singular measure $\mathcal{Z}_\partial(M)$, that is,

$$\mathcal{Z}(M) = \{\zeta \in D^2 ; f(\zeta) = 0 \text{ for } f \in M\}$$

and

$$\mathcal{Z}_\partial(M) = \inf\{-d\sigma_f ; f \in M, f \neq 0\}.$$

They showed that if the real 2-dimensional Hausdorff measure of $\mathcal{Z}(M)$ is zero and $\mathcal{Z}_\partial(M) = 0$, then $\mathcal{M}(M) = H^\infty(D^2)$. In this paper, we are interested in invariant subspaces M of $H^2(D^2)$ such that the real 2-dimensional Hausdorff measure of $\mathcal{Z}(M)$ is positive and $\mathcal{Z}_\partial(M) = 0$.

Fix $(\alpha, \beta) \in \overline{D^2}$. For f in $H^p(D^2)$,

$$(\Phi_{\alpha\beta}^p f)(\lambda) = f(\alpha\lambda, \beta\lambda) \quad (\lambda \in D).$$

$\Phi_{\alpha\beta}^p$ is called a slice map. $\Phi_{\alpha\beta}^2$ maps $H^2(D^2)$ into $L_a^2(D)$, where $L_a^2(D)$ is the Bergman space (cf. [p53]). In this paper, we study the kernel $\ker \Phi_{\alpha\beta}^p$ and the range $\text{ran} \Phi_{\alpha\beta}^p$ for $p = 2, \infty$. $\ker \Phi_{\alpha\beta}^2$ is an invariant subspace of $H^2(D^2)$ and the closure of $\text{ran} \Phi_{\alpha\beta}^2$ is an invariant subspace of $L_a^2(D)$. Put

$$\mathcal{D}_{\alpha\beta} = \{(\alpha\lambda, \beta\lambda) \in D^2 ; \lambda \in D\},$$

then $\mathcal{Z}(\ker\Phi_{\alpha\beta}^2) = \mathcal{D}_{\alpha\beta}$ if $(\alpha, \beta) \in T^2 \cup T \times D \cup D \times T$. The 2-dimensional Hausdorff measure of $\mathcal{Z}(\ker\Phi_{\alpha\beta}^2)$ is positive and $\mathcal{Z}_{\beta}(\ker\Phi_{\alpha\beta}^2) = 0$. In this paper, we show $\mathcal{M}(M) = H^\infty(D^2)$ when $\mathcal{Z}(M) = \mathcal{D}_{\alpha\beta}$ for some $(\alpha, \beta) \in T^2$ and $\mathcal{Z}_{\beta}(M) = 0$ and M satisfies some additional natural condition. The main result in this paper is Theorem 4 in Section 3. Theorem 1 of [2] has a lot of corollaries on the rigidity of invariant subspaces. Theorem 3 in this paper has similarly such corollaries. Hence our results can be seen as the generalizations of results of R.G.Douglas and K.Yan.

For f in $N(D^2)$, $f(\zeta) = \sum_{j=0}^{\infty} F_j(\zeta)$ is a homogeneous expansion of f and F_j is a polynomial which is homogeneous of degree j . The smallest $j = j(f)$ such that F_j is not the zero-polynomial is called the order of the zero which f has at $\zeta = (0, 0)$. For $p \in D^2$, the order of the zero of f at p is simply the order of the zero of $f(p + \zeta)$ at $\zeta = (0, 0)$. We will write $f_p(\zeta) = f(p + \zeta)$.

§2. Slice maps

In this section, we study the slice map $\Phi_{\alpha\beta} = \Phi_{\alpha\beta}^p$ for $(\alpha, \beta) \in \overline{D^2}$.

Proposition 1. Let $(\alpha, \beta) \in \overline{D^2}$.

- (1) $\Phi_{\alpha\beta}^2$ is a contractive map from $H^2(D^2)$ to $L_a^2(D)$.
- (2) If $(\alpha, \beta) \in D^2$, then $\text{ran}\Phi_{\alpha\beta}^2$ is a subset of analytic functions on \bar{D} .
- (3) If $(\alpha, \beta) \in T^2$, then $\Phi_{\alpha\beta}^2$ is an onto map from $H^2(D^2)$ to $L_a^2(D)$ with $\|\Phi_{\alpha\beta}^2\| = 1$.
- (4) If $(\alpha, \beta) \in T \times D \cup D \times T$, then $\Phi_{\alpha\beta}^2$ is an onto map from $H^2(D^2)$ to $H^2(D)$ with $\|\Phi_{\alpha\beta}^2\| \leq (1 - |\beta|^2)^{-1}$.

Proof. (1) For $f \in H^2(D^2)$, let $f(z, w) = \sum_{j=0}^{\infty} F_j(z, w)$ be a homogeneous expansion of f . Then $F_j(z, w) = \sum_{\ell=0}^j a_\ell z^{j-\ell} w^\ell$ and $\int |F_j|^2 dm = \sum_{\ell=0}^j |a_\ell|^2$. Moreover

$$\int |f|^2 dm = \sum_{j=0}^{\infty} \int |F_j|^2 dm = \sum_{j=0}^{\infty} \sum_{\ell=0}^j |a_\ell|^2 < \infty.$$

$(\Phi_{\alpha\beta} f)(\lambda) = \sum_{j=0}^{\infty} F_j(\alpha, \beta) \lambda^j$ and

$$|F_j(\alpha, \beta)|^2 \leq \left(\sum_{\ell=0}^j |a_\ell|^2 \right) \left(\sum_{\ell=0}^j |\beta|^{2\ell} \right) \leq (j+1) \left(\sum_{\ell=0}^j |a_\ell|^2 \right).$$

Hence

$$\int_0^1 \int_0^{2\pi} |\Phi_{\alpha\beta} f|^2 (re^{i\theta}) r d\theta dr / \pi$$

$$\begin{aligned}
&= \int_0^1 \sum_{j=0}^{\infty} |F_j(\alpha, \beta)|^2 r^{2j+1} 2dr = \sum_{j=0}^{\infty} |F_j(\alpha, \beta)|^2 \frac{1}{j+1} \\
&\leq \sum_{j=0}^{\infty} \sum_{\ell=0}^j |a_{\ell}|^2 = \int |f|^2 dm.
\end{aligned}$$

Thus $\Phi_{\alpha\beta}f \in L_a^2(D)$ and $\|\Phi_{\alpha\beta}\| \leq 1$.

(2) is clear. (3). For $g \in L_a^2(D)$ with $g(\lambda) = \sum_{j=0}^{\infty} b_j \lambda^j$, put $f(z, w) = \sum_{j=0}^{\infty} \frac{b_j}{j+1} (\bar{\beta}w)^{j-\ell} (\bar{\alpha}z)^{\ell}$. Then $f \in H^2(D^2)$ and $(\Phi_{\alpha\beta}f)(\lambda) = g(\lambda)$. This and (1) imply (3). (4). We may assume $(\alpha, \beta) \in T \times D$. Then

$$|F_j(\alpha, \beta)|^2 \leq (1 - |\beta|^2)^{-1} \sum_{\ell=0}^j |a_{\ell}|^2$$

and hence

$$\begin{aligned}
&\int_0^{2\pi} |\Phi_{\alpha\beta}f|^2 (re^{i\theta}) d\theta / 2\pi \\
&\leq \sum_{j=0}^{\infty} \frac{1}{1 - |\beta|^2} \sum_{\ell=0}^j |a_{\ell}|^2 \leq \frac{1}{1 - |\beta|^2} \int |f|^2 dm.
\end{aligned}$$

For $g \in H^2(D)$ with $g(\lambda) = \sum_{j=0}^{\infty} b_j \lambda^j$, put $f(z, w) = \sum_{j=0}^{\infty} b_j (\bar{\alpha}z)^j$. Then $f \in H^2(D^2)$ and $(\Phi_{\alpha\beta}f)(\lambda) = g(\lambda)$. This implies (4).

(3) of Proposition 1 is essentially known (see [6, p53]). Now we study the slice map $\Phi_{\alpha\beta}^{\infty}$ on $H^{\infty}(D^2)$. Let L be the norm closed linear span of $\overline{H^{\infty}(D^2)} \cap H^{\infty}(D^2)$ in $L^{\infty}(T^2)$. Then $L \neq L^{\infty}(T^2)$ (see[5]).

Proposition 2. Let $(\alpha, \beta) \in \overline{D^2}$.

- (1) $\Phi_{\alpha\beta}^{\infty}$ is a contractive homomorphism from $H^{\infty}(D^2)$ to $H^{\infty}(D)$.
- (2) If $(\alpha, \beta) \in T^2 \cup T \times D \cup D \times T$, then $\Phi_{\alpha\beta}^{\infty}$ is a contractive homomorphism from $H^{\infty}(D^2)$ onto $H^{\infty}(D)$.
- (3) If $(\alpha, \beta) \in T^2$, there exists a contractive $*$ -homomorphism $\tilde{\Phi}_{\alpha\beta}^{\infty}$ from L onto $L^{\infty}(T)$ such that $\tilde{\Phi}_{\alpha\beta}^{\infty} | H^{\infty}(T^2) = \Phi_{\alpha\beta}^{\infty} | H^{\infty}(T^2)$.

Proof. (1) is clear. (2). If $g \in H^{\infty}(D)$ with $g(\lambda) = \sum_{j=0}^{\infty} b_j \lambda^j$ and $|\alpha| = 1$, then $f(z, w) = \sum_{j=0}^{\infty} b_j (\bar{\alpha}z)^j \in H^{\infty}(D^2)$ and $(\Phi_{\alpha\beta}f)(\lambda) = g(\lambda)$. This and (1) imply (2).

- (3) For $f_j, \bar{g}_j \in H^{\infty}(D^2)$ and $j = 1, \dots, n$, put

$$\{\tilde{\Phi}_{\alpha\beta}^{\infty}(\sum_{j=1}^n f_j \bar{g}_j)\}(\lambda) = \sum_{j=1}^n f_j(\alpha\lambda, \beta\lambda) \overline{g_j(\alpha\lambda, \beta\lambda)}$$

for $\lambda \in D$, then $\tilde{\Phi}_{\alpha\beta}^{\infty}(\sum_{j=1}^n f_j \bar{g}_j)$ can be seen as an element in $L^{\infty}(T)$ by its radial limits. Hence for a.e. $\lambda \in T$

$$\begin{aligned}
& | \Phi_{\alpha\beta}(\sum_{j=1}^n f_j \bar{g}_j)(\lambda) | \\
& \leq \text{ess sup}_{\lambda \in T} | \sum_{j=1}^n f_j(\alpha\lambda, \beta\lambda) \overline{g_j(\alpha\lambda, \beta\lambda)} | \\
& \leq \text{ess sup}_{(z,w) \in T^2} | \sum_{j=1}^n f_j(\alpha z, \beta w) \overline{g_j(\alpha z, \beta w)} | = \| \sum_{j=1}^n f_j \bar{g}_j \|_{\infty}
\end{aligned}$$

because $(\alpha, \beta) \in T^2$. Then $\tilde{\Phi}_{\alpha\beta}$ is the extension of $\Phi_{\alpha\beta}$ from $H^\infty(D^2)$ to L , then $\tilde{\Phi}_{\alpha\beta}$ is a contractive *-homomorphism from L to $L^\infty(T)$. If $U(\lambda) = \sum_{j=1}^n F_j(\lambda) \bar{G}_j(\lambda)$ a.e. on T where $F_j, G_j \in H^\infty(D)$, then $u(z, w) = \sum_{j=1}^n F_j(\bar{\alpha}z) \bar{G}_j(\beta w)$ belongs to L and $(\tilde{\Phi}_{\alpha\beta}u)(\lambda) = U(\lambda)$ a.e. on T . Since arbitrary function U in $L^\infty(T)$ can be approximated by such functions, $\tilde{\Phi}_{\alpha\beta}$ is onto.

The following lemma will be used in the proofs in the following proposition and the main theorem. We can prove it by an approximation method as in [4] but we prove it using Proposition 2.

Lemma. If $\phi \in L^\infty(T^2)$, $(\alpha, \beta) \in T^2$ and $\phi(z, w)(\beta z - \alpha w) \in H^\infty(D^2)$, then $\phi \in H^\infty(D^2)$.

Proof. Note that $\beta z - \alpha w \in \ker \Phi_{\alpha\beta}$. If $\phi(\beta z - \alpha w) = g$ for some $g \in H^\infty(D^2)$, then g belongs to $\ker \Phi_{\alpha\beta}$. In fact, $\hat{\phi}(\beta z - \alpha w)^\wedge = \hat{g}$ on $\text{Spec } L^\infty(T^2)$ which is the maximal ideal space of $L^\infty(T^2)$ and $(\beta z - \alpha w)^\wedge = 0$ on $\text{hull}(\ker \tilde{\Phi}_{\alpha\beta})$. Hence $\hat{g} = 0$ on $\text{hull}(\ker \tilde{\Phi}_{\alpha\beta}) \cap \text{Spec } L^\infty(T^2)$. Since L is a commutative C^* -algebra, every element of $\text{Spec } L$ extends to an element of $\text{Spec } L^\infty(T^2)$. Therefore $\hat{g} = 0$ on $\text{hull}(\ker \tilde{\Phi}_{\alpha\beta})$. Thus $g \in (\ker \tilde{\Phi}_{\alpha\beta}) \cap H^\infty(D^2) = \ker \Phi_{\alpha\beta}$. Hence if $g = \sum_{j=0}^\infty G_j$ and $G_j(z, w) = \sum_{\ell=0}^j b_\ell z^{j-\ell} w^\ell$, then

$$G_j(z, w) = z^j \sum_{\ell=0}^j b_\ell (\bar{z}w)^\ell = k \prod_{\ell=1}^j (w - k_\ell z)$$

where $k \in \mathbb{C}$ and $k_\ell \in \mathbb{C}$ for $1 \leq \ell \leq j$ and $G_j(\alpha\lambda, \beta\lambda) \equiv 0$ for $\lambda \in D$ because $g \in \ker \Phi_{\alpha\beta}^2$. Thus $G_j(z, w) = m(\beta z - \alpha w) \prod_{\ell=2}^j (w - m_\ell z)$ where $m \in \mathbb{C}$ and $m_\ell \in \mathbb{C}$ for $2 \leq \ell \leq j$ and hence $g/(\beta z - \alpha w)$ is analytic on D^2 . Since $d\sigma_{\beta z - \alpha w} = 0$, $g/(\beta z - \alpha w) \in N_*(D^2) \cap L^\infty(T^2) = H^\infty(D^2)$ and hence ϕ belongs to $H^\infty(D^2)$.

Proposition 3. Let $(\alpha, \beta) \in \overline{D^2}$.

- (1) For any $r \in (0, 1]$, $\ker \Phi_{\alpha\beta}^2 = \ker \Phi_{r\alpha, r\beta}^2$.
- (2) $\ker \Phi_{\alpha\beta}^2$ is an invariant subspace of $H^2(D^2)$,

$$\mathcal{Z}(\ker \Phi_{\alpha\beta}^2) = \mathcal{D}_{\alpha\beta} \text{ and } \mathcal{Z}_\partial(\ker \Phi_{\alpha\beta}^2) = 0.$$

For any $p \in \mathcal{D}_{\alpha\beta}$, $\beta z - \alpha w \in \ker \Phi_{\alpha\beta}^2$ has a zero of order 1 at p .

(3) If $(\alpha, \beta) \in T^2$, then $(\beta z - \alpha w)H^2(D^2)$ is dense in $\ker \Phi_{\alpha\beta}^2$ but $\ker \Phi_{\alpha\beta}^2 \neq (\beta z - \alpha w)H^2(D^2)$. If $(\alpha, \beta) \in T \times D \cup D \times T$, then $\ker \Phi_{\alpha\beta}^2 = (\beta z - \alpha w)H^2(D^2)$.

(4) If $(\alpha, \beta) \in T^2$, then $\mathcal{M}(\ker \Phi_{\alpha\beta}^2) = H^\infty(D^2)$ and if $(\alpha, \beta) \in T \times D \cup D \times T$, then $\mathcal{M}(\ker \Phi_{\alpha\beta}^2) = (\beta z - \alpha w)^{-1}H^\infty(D^2)$.

(5) If $(\alpha, 0) \in \bar{D} \times D$ and $\alpha \neq 0$, then $\ker \Phi_{\alpha 0}^2 = wH^2(D^2)$ and hence $\mathcal{M}(\ker \Phi_{\alpha 0}^2) = w^{-1}H^\infty(D^2)$.

(6) Let M be an invariant subspace of $H^2(D^2)$ with $\ker \Phi_{\alpha\beta}^2 \subsetneq M$, $\mathcal{M}(M) = H^\infty(D^2)$. If $(\alpha, \beta) \in T^2$, then $\mathcal{Z}(M) = \{(\alpha a_j, \beta a_j) \in D^2; \sum_{j=1}^\infty (1 - |a_j|) \times [-\log(1 - |a_j|)]^{1-\varepsilon} < \infty \text{ for all } \varepsilon > 0\}$. If $(\alpha, \beta) \in T \times D \cup D \times T$, then $\mathcal{Z}(M) = \{(\alpha a_j, \beta a_j) \in D^2; \sum_{j=1}^\infty (1 - |a_j|) < \infty\}$. If $(\alpha, 0) \in \bar{D} \times D$ and $\alpha \neq 0$, then $M = qH^2(D) \oplus wH^2(D^2)$ where q is a one variable inner function with $q = q(z)$ and hence $\mathcal{Z}(M) = \{(s, 0) \in D^2; q(s) = 0 \text{ and } s \in D\}$.

Proof. (1) and (2) are clear. (3). Let $(\alpha, \beta) \in T^2$. If $f \in \ker \Phi_{\alpha\beta}$, $f = \sum_{j=0}^\infty F_j$ and $F_j(z, w) = \sum_{\ell=0}^j a_\ell z^{j-\ell} w^\ell$, then $F_j(z, w) = c(\beta z - \alpha w) \prod_{\ell=2}^j (w - c_\ell z)$ and hence f can be approximated by the functions in $(\beta z - \alpha w)H^2(D^2)$. This implies that $(\beta z - \alpha w)H^2(D^2)$ is dense in $\ker \Phi_{\alpha\beta}^2$. Suppose $\ker \Phi_{\alpha\beta} = (\beta z - \alpha w)H^2(D^2)$, then the multiplication operator by $\beta z - \alpha w$ is a left invertible operator from $H^2(D^2)$ to $\ker \Phi_{\alpha\beta}$. Hence there exists a positive constant ε such that

$$\int_{T^2} |g|^2 |\beta z - \alpha w|^2 dm \geq \varepsilon \int_{T^2} |g|^2 dm$$

for all $g \in H^\infty(D^2)$ and so

$$\int_{T^2} u |\beta z - \alpha w|^2 dm \geq \varepsilon \int_{T^2} u dm$$

for all nonnegative continuous functions u on T^2 . Thus $|\beta z - \alpha w|^2 \geq \varepsilon > 0$ a.e. on T^2 and this contradiction implies that $\ker \Phi_{\alpha\beta} \neq (\beta z - \alpha w)H^2(D^2)$. Let $(\alpha, \beta) \in T \times D \cup D \times T$. Since $\beta z - \alpha w$ is invertible in L^∞ , $\ker \Phi_{\alpha\beta} = (\beta z - \alpha w)H^2(D^2)$ because $(\beta z - \alpha w)H^2(D^2)$ is dense in $\ker \Phi_{\alpha\beta}$.

(4) Let $(\alpha, \beta) \in T^2$. If $\phi \in \mathcal{M}(\ker \Phi_{\alpha\beta})$, then $\phi(\beta z - \alpha w) = g$ for some $g \in H^\infty(D^2)$. By Lemma, ϕ belongs to $H^\infty(D^2)$ and hence $\mathcal{M}(\ker \Phi_{\alpha\beta}) = H^\infty(D^2)$. Let $(\alpha, \beta) \in T \times D \cup D \times T$ and $\phi \in \mathcal{M}(\ker \Phi_{\alpha\beta})$. By (3), $\ker \Phi_{\alpha\beta} = (\beta z - \alpha w)H^2(D^2)$ and hence

$$\phi(\beta z - \alpha w)H^2(D^2) \subset H^2(D^2).$$

This implies that $\phi(\beta z - \alpha w) \in H^\infty(D^2)$ and so $\mathcal{M}(\ker \Phi_{\alpha\beta}) = (\beta z - \alpha w)^{-1}H^\infty(D^2)$. (5) is easy to see. (6) If $(\alpha, \beta) \in T^2$ and $\ker \Phi_{\alpha\beta} \subsetneq M$, then by (3) of Proposition 1 and [3, Corollary 3.6], $\mathcal{Z}([\Phi_{\alpha\beta} M]_2) = \{a_j \in D : \sum_{j=1}^\infty (1 - |a_j|) [-\log(1 - |a_j|)]^{1-\varepsilon} < \infty \text{ for all } \varepsilon > 0\}$. This implies the first part. For $(\alpha, \beta) \in T \times D \cup D \times T$, we can show similarly by (4) of Proposition 1.

§3. Multipliers

By (4) of Proposition 3, we know the set of all multipliers $\mathcal{M}(M)$ of an invariant subspace such that $\ker\Phi_{\alpha\beta}^2 \subseteq M \subseteq H^2(D^2)$ when $(\alpha, \beta) \in T^2 \cup T \times D \cup D \times T$ or $(\alpha, 0) \in D \times D \setminus (0, 0)$. When $(\alpha, \beta) \in D \times D$ and $|\alpha| = |\beta| > 0$, there exists $(\alpha_0, \beta_0) \in T \times T$ such that $\alpha = r\alpha_0$ and $\beta = r\beta_0$ for some $r \in (0, 1)$. When $(\alpha, \beta) \in D \times D$ and $0 \leq |\alpha| < |\beta|$, there exists $(\alpha_0, \beta_0) \in D \times T$ such that $\alpha = r\alpha_0$ and $\beta = r\beta_0$ for some $r \in (0, 1)$. (1) of Proposition 3 implies $\ker\Phi_{\alpha\beta}^2 = \ker\Phi_{\alpha_0\beta_0}^2$. Hence for arbitrary $(\alpha, \beta) \in \bar{D} \times \bar{D} \setminus (0, 0)$, we can describe $\mathcal{M}(M)$ by Proposition 3. In this section, we study $\mathcal{M}(M)$ without such a condition. In this section, for example, we study $\mathcal{M}(M)$ of an invariant subspace such that $M \subseteq \ker\Phi_{\alpha\beta}^2$. In fact, we study such a problem more generally, that is, when the 2-dimensional Hausdorff measure of $\mathcal{Z}(M) \cap \mathcal{D}_{\alpha\beta}^c$ is zero. For $\Lambda \subset T^2 \cup T \times D \cup D \times T$, put

$$\mathcal{D}_\Lambda = \{\cup \mathcal{D}_{\alpha\beta} ; (\alpha, \beta) \in \Lambda\} \setminus \{(0, 0)\}.$$

Note that if $\mathcal{Z}(M) \supseteq \mathcal{D}_\Lambda$ and Λ is an infinite set such that $\mathcal{D}_{\alpha\beta} \cap \mathcal{D}_{\gamma\delta} = \{(0, 0)\}$ when $(\alpha, \beta) \neq (\gamma, \delta)$, then $M = \{0\}$.

Theorem 4. Let Λ be a finite set of T^2 . If M is an invariant subspace of $H^2(D^2)$ which satisfies the following (1) ~ (3), then $\mathcal{M}(M) = H^\infty(D^2)$.

(1) For any $p \in \mathcal{Z}(M) \cap \mathcal{D}_\Lambda$, there exists a function f in M such that f has a zero of order 1 at p .

(2) The 2-dimensional Hausdorff measure of $\mathcal{Z}(M) \cap \mathcal{D}_\Lambda^c$ is zero.

(3) $\mathcal{Z}_\beta(M) = 0$.

Proof. Suppose $\phi \in \mathcal{M}(M)$. Fix $p \in \mathcal{Z}(M) \cap \mathcal{D}_\Lambda$. By (1), let f be a function in M such that f has a zero of order 1 at p . Let $(\alpha, \beta) \in \Lambda$ with $p \in \mathcal{D}_{\alpha\beta}$. By definition of $\mathcal{M}(M)$, $\phi f = g$ for some $g \in H^2(D^2)$. Put $k(z, w) = \beta z - \alpha w$, then $k_p(\zeta) = k(\zeta + p) = k(\zeta)$ and $k_p(\zeta)\phi_p(\zeta)f_p(\zeta) = k(\zeta)g_p(\zeta)$. Suppose $f_p(\zeta) = \sum_{j=0}^{\infty} F_j(\zeta)$ is a homogeneous expansion of f_p . Since $1 = s(f_p)$, $F_1(0, w) = cw$ for $c \neq 0$. By the Weierstrass preparation theorem (cf [6, Theorem 1.2.1]), there exists a polydisc Δ in \mathbb{C}^2 , centered at $(0, 0)$, such that

$$f_p(z, w) = W(z, w)h(z, w)$$

for $(z, w) \in \Delta$ where h is analytic in Δ , h has no zero in Δ , $W(z, w) = w + b_0(z)$ and b_0 is analytic in Δ with $b_0(0) = 0$. Since $f_p(\alpha\lambda, \beta\lambda) \equiv 0$ on D , $\beta\lambda + b_0(\alpha\lambda) = 0$ if $(\alpha\lambda, \beta\lambda) \in \Delta$ and hence $b_0(\alpha\lambda) = -\beta\lambda$. Thus $b_0(z) = -\frac{\beta}{\alpha}z$ and $W(z, w) = -\frac{1}{\alpha}(\beta z - \alpha w)$. Therefore $k_p\phi_p = k g_p/f_p$ is analytic in Δ and so $k\phi$ is analytic in $\Delta + p$, in a sense of D. Douglas and K. Yan [2]. Therefore

$$\prod_{(\alpha, \beta) \in \Lambda} (\beta z - \alpha w)\phi(z, w)$$

is analytic in a neighborhood of $\mathcal{Z}(M) \cap \mathcal{D}_\Lambda$.

If $p \notin \mathcal{Z}(M)$, then there exists a function k in M such that k has no zero in some polydisc Δ_p , centered at p . As in the proof above, $\phi(z, w)$ is analytic in Δ_p and hence ϕ is analytic in $D^2 \setminus \mathcal{Z}(M)$. Thus $\Pi(\beta z - \alpha w)\phi(z, w)$ is analytic in $D^2 \setminus \mathcal{Z}(M) \cap \mathcal{D}_\Lambda^c$. By (2), $\mathcal{Z}(M) \cap \mathcal{D}_\Lambda^c$ is a removable singularity for analytic functions and hence $\psi(z, w) = \Pi(\beta z - \alpha w)\phi(z, w)$ is analytic in D^2 . By the proof of [2, Theorem 1], $\psi \in N(D^2) \cap L^\infty(T^2)$ and $d\sigma_\psi \leq -\mathcal{Z}_\theta(M)$ because $d\sigma_\phi = d\sigma_\psi$. By (3), $d\sigma_\psi = 0$ and hence $\psi \in H^\infty(D^2)$. By Lemma, ϕ belongs to $H^\infty(D^2)$ and hence $\mathcal{M}(M) = H^\infty(D^2)$.

Theorem 5. Let Λ be a finite set of $T \times D \cup D \times T$. If M is an invariant subspace of $H^2(D^2)$ which $\mathcal{Z}(M) \supseteq \mathcal{D}_\Lambda$ and satisfies the following (1) ~ (3), then

$$\mathcal{M}(M) = \prod_{(\alpha, \beta) \in \Lambda} (\beta z - \alpha w)^{-1} H^\infty(D^2).$$

(1) For any $p \in \mathcal{Z}(M)$, there exists a function f in M such that f has a zero of order 1 at p .

(2) The 2-dimensional Hausdorff measure of $\mathcal{Z}(M) \cap \mathcal{D}_\Lambda^c$ is zero.

(3) $\mathcal{Z}_\theta(M) = 0$.

Proof. By the proof of Theorem 4, if $\phi \in \mathcal{M}(M)$, then $\Pi(\beta z - \alpha w)\phi(z, w) \in H^\infty(D^2)$ where (α, β) ranges over Λ . Hence $\phi \in \Pi(\beta z - \alpha w)^{-1} H^\infty(D^2)$. Conversely if $\phi \in \Pi(\beta z - \alpha w)^{-1} H^\infty(D^2)$ and $f \in M$, then $f = 0$ on \mathcal{D}_Λ , hence by the Weierstrass preparation theorem, $\Pi(\beta z - \alpha w)^{-1} f(z, w)$ is analytic in D^2 and ϕ belongs to $\mathcal{M}(M)$.

§4. Two general cases and remarks

Let a and b be two functions in $H^\infty(D)$ with $\|a\|_\infty \leq 1$ and $\|b\|_\infty \leq 1$. For f in $H^p(D^2)$,

$$(\Phi_{ab}^p f)(\lambda) = f(a(\lambda), b(\lambda)) \quad (\lambda \in D).$$

If $a(\lambda) = \alpha\lambda$ and $b(\lambda) = \beta\lambda$, then Φ_{ab}^p was called a slice map $\Phi_{\alpha\beta}^p$ in the previous sections. For arbitrary pair a and b , we know only very trivial results. It is easy to see that Φ_{ab}^∞ maps $H^\infty(D^2)$ into $H^\infty(D)$. If $\|a\|_\infty < 1$ and $\|b\|_\infty < 1$, then Φ_{ab}^p maps $H^p(D^2)$ into $H^\infty(D)$. In general, $\ker \Phi_{ab}^2$ is still an invariant subspace of $H^2(D^2)$, and

$$\mathcal{Z}(\ker \Phi_{ab}^2) = \mathcal{D}_{ab} = \{(a(\lambda), b(\lambda)) \in D^2; \lambda \in D\}.$$

The function $b(z) - a(w)$ may not belong to $\ker \Phi_{ab}^2$. If $a(\lambda) = \alpha\lambda$ and $b(\lambda) = \beta\lambda$, then $(boa)(\lambda) = (aob)(\lambda)$ for $\lambda \in D$ and hence $b(z) - a(w)$ belongs to $\ker \Phi_{ab}^2$. If $a(\lambda) = \lambda$ and $b(\lambda)$ is an inner function, then $(boa)(\lambda) = (aob)(\lambda)$ for $\lambda \in D$ and hence $b(z) - a(w) =$

$b(z) - w$ belongs to $\ker \Phi_{ab}^2$. In this case, $\mathcal{Z}_\partial(\ker \Phi_{ab}^2) = 0$. For any $p \in \mathcal{D}_{ab}$, $b(z) - w$ has a zero of order 1 at $p \in \mathcal{D}_{ab}$. If $\phi \in L^\infty(T^2)$ and $(b(z) - w)\phi(z, w) \in H^\infty(D^2)$, then $\phi \in H^\infty(D^2)$. This can be shown as in [4, Proposition 3 and Theorem 7]. This implies (4) of Proposition 3. The proof of the following theorem is almost parallel to that of Theorem 4.

Theorem 6. Let $a(\lambda) = \lambda$ and $b(\lambda)$ be an inner function. If M is an invariant subspace of $H^2(D^2)$ which satisfies the following (1) ~ (3), then $\mathcal{M}(M) = H^\infty(D^2)$.

- (1) For any $p \in \mathcal{Z}(M) \cap \mathcal{D}_{ab}$, there exists a function f in M such that f has a zero of order 1 at $p \in \mathcal{Z}(M) \cap \mathcal{D}_{ab}$.
- (2) The 2-dimensional Hausdorff measure of $\mathcal{Z}(M) \cap \mathcal{D}_{ab}^c$ is zero.
- (3) $\mathcal{Z}_\partial(M) = 0$.

If $a(\lambda) = \lambda$ and $b(\lambda) = cq(\lambda)$ where c is a constant with $|c| < 1$ and q is an inner function, we can show a version of Theorem 5 as Theorem 6 which is that of Theorem 4.

Let D^n be the open unit polydisc in \mathbb{C}^n and T^n be its distinguished boundary. Fix $\alpha = (\alpha_1, \dots, \alpha_n) \in \overline{D^n}$. For f in $H^p(D^n)$

$$(\Phi_\alpha^p f)(\lambda) = f(\alpha_1 \lambda, \dots, \alpha_n \lambda) \quad (\lambda \in D).$$

(1), (2) and (3) of Proposition 1 can be proved for arbitrary n . If $\alpha_j \in T$ for some j with $1 \leq j \leq n$ and $\alpha_i \in D$ for all i with $1 \leq i \leq n$ and $i \neq j$, we can show that Φ_α^2 is an onto map from $H^2(D^n)$ to $H^2(D)$ with $\|\Phi_\alpha^2\| \leq \prod_{i \neq j} (1 - |\alpha_j|^2)^{-1}$. This is a generalization of (4) of Proposition 1. Similarly we can generalize Proposition 2. If $\phi \in L^\infty(T^n)$ and $(\alpha_i z_j - \alpha_j z_i)\phi(z_1, \dots, z_n) \in H^\infty(D^n)$ where $1 \leq i \neq j \leq n$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in T^n$, then $\phi \in H^\infty(D^n)$. This also can be shown as in [4, Proposition 3 and Theorem 7]. $\ker \Phi_\alpha^2$ is an invariant subspace and a generalization of (1) and (2) of Proposition 3 is true. Suppose $n > 2$. If M is an invariant subspace of $H^2(D^n)$, $\mathcal{Z}(M) = \mathcal{D}_\alpha = \{(\alpha_1 \lambda, \dots, \alpha_n \lambda); \lambda \in D\}$ for $\alpha \in T^n$ and $\mathcal{Z}_\partial(M) = 0$, then $\mathcal{M}(M) = H^\infty(D^n)$. For it is a result of R.G. Douglas and K. Yan [2, Theorem 1] because the real $2n - 2$ dimensional Hausdorff measure of $\mathcal{Z}(M)$ is zero.

Remark. (i) As in Theorem 1 of [2], Theorem 4 can be stated as the following: If M is an invariant subspace of $H^2(D^2)$ which satisfies (1) and (2), then $\phi \in \mathcal{M}(M)$ if and only if $\phi \in N(D^2) \cap L^\infty(T^2)$ and $d\sigma_\phi \leq \mathcal{Z}_\partial(M)$. (ii) By Lemma 7 in [2] and Theorem 4, if M and N are quasi-similar invariant subspaces of $H^2(D^2)$ and M satisfies (1) ~ (3) in Theorem 4, then $M \subseteq N$. This is a generalization of Theorem 2 in [2]. Similarly we can generalize Corollaries 9 and 12. (iii) Let M, N be invariant subspaces of $H^2(D^2)$ satisfying (a) the 2-dimensional Hausdorff measures of $\mathcal{Z}(M) \cap \mathcal{D}_\Lambda^c$ and $\mathcal{Z}(N) \cap \mathcal{D}_\Lambda^c$ are zero. (b) $\mathcal{Z}_\partial(M) = \mathcal{Z}_\partial(N)$. (c) M and N satisfy the condition (1) in Theorem 4 about $\mathcal{Z}(M) \cap \mathcal{D}_\Lambda$ and $\mathcal{Z}(N) \cap \mathcal{D}_\Lambda$. If M and N are quasi-similar, then $M = N$.

References

1. A.Beurling, On two problems concerning linear transformations in Hilbert space, Acta Math. 81(1949) 239-255.
2. R.G.Douglas and K.Yan, On the rigidity of Hardy submodules, Integral Eq Op. Th. 13(1990), 350-363.
3. C.Horowitz, Zeros of functions in the Bergman spaces, Duke Math. J. 41(1974).
4. T.Nakazi, Multipliers of invariant subspaces in the bidisc, Proc. Edinburgh Math. Soc. 37(1994), 193-199.
5. M.Range, A small boundary for H^∞ on the polydisc, Proc. Amer. Math. Soc 32(1972), 253-255.
6. W.Rudin, Function Theory in Polydisks, Benjamin, New York (1969).