



Title	Asymptotic stability of planar rarefaction wave for scalar viscous conservation law
Author(s)	Ito, K.
Citation	Hokkaido University Preprint Series in Mathematics, 384, 1-8
Issue Date	1997-6-1
DOI	10.14943/83530
Doc URL	<a href="http://hdl.handle.net/2115/69134">http://hdl.handle.net/2115/69134</a>
Type	bulletin (article)
File Information	pre384.pdf



[Instructions for use](#)

**Asymptotic Stability of  
Planar Rarefaction Wave for  
Scalar Viscous Conservation Law**

**Kazuo Ito**

Series #384. June 1997

HOKKAIDO UNIVERSITY  
PREPRINT SERIES IN MATHEMATICS

- #359 I. Tsuda and K. Tadaki, A logic-based dynamical theory for a genesis of biological threshold, 49 pages. 1996.
- #360 I. Tsuda and A. Yamaguchi, Singular-continuous nowhere-differentiable attractors in neural systems, 40 pages. 1996.
- #361 M. Nakamura and T. Ozawa, Low energy scattering for nonlinear Schrödinger equations in fractional order Sobolev spaces, 17 pages. 1996.
- #362 I. Nakamura, Hilbert schemes and simple singularities  $E_6$ ,  $E_7$  and  $E_8$ , 21 pages. 1996.
- #363 T. Mikami, Equivalent conditions on the central limit theorem for a sequence of probability measures on  $R$ , 7 pages. 1996.
- #364 S. Izumiya and T. Sano, Generic affine differential geometry of space curves, 23 pages. 1996.
- #365 T. Tsukada, Stability of reticular optical caustics, 12 pages. 1996.
- #366 A. Arai and M. Hirokawa, On the existence and uniqueness of ground states of a generalized spin-boson model, 40 pages. 1996.
- #367 A. Arai, A class of representations of the  $*$ -algebra of the canonical commutation relations over a Hilbert space and instability of embedded eigenvalues in quantum field models, 12 pages. 1996.
- #368 K. Ito, BV-solutions of a hyperbolic-elliptic system for a radiating gas, 33 pages. 1997.
- #369 M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, 20 pages. 1997.
- #370 N.H. Bingham and A. Inoue, An Abel-Tauber theorem for Hankel transforms, 8 pages. 1997.
- #371 T. Nakazi and H. Sawada, The commutator ideal in Toeplitz algebras for uniform algebras and the analytic structure, 9 pages. 1997.
- #372 M.-H. Giga and Y. Giga, Stability for evolving graphs by nonlocal weighted curvature, 70 pages. 1997.
- #373 T. Nakazi, Brown-Halmos type theorems of weighted Toeplitz operators, 14 pages. 1997.
- #374 J. Inoue and S.-E. Takahashi, On characterizations of the image of Gelfand transform of commutative Banach algebras, 30 pages. 1997.
- #375 L. Solomon and H. Terao, The double coxeter arrangement, 21 pages. 1997.
- #376 G. Ishikawa and T. Morimoto, Solution surfaces of Monge-Ampère equations, 15 pages. 1997.
- #377 G. Ishikawa, A relative transversality theorem and its applications, 16 pages. 1997.
- #378 J. Inoue and T. Nakazi, On the zeroes of solutions of an extremal problem in  $H^1$ , 14 pages. 1997.
- #379 Y. Giga and K. Ito, On pinching of curves moved by surface diffusion, 12 pages. 1997.
- #380 F. Hiroshima, Weak coupling limit removing an ultraviolet cut-off for a Hamiltonian of particles interacting with a scalar field, 39 pages. 1997.
- #381 Y. Giga, S. Matsui and Y. Shimizu, On estimates in Hardy spaces for the Stokes flow in a half space, 13 pages. 1997.
- #382 A. Arai, A new estimate for the ground state energy of Schrödinger operators, 12 pages. 1997.
- #383 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the Sobolev space of critical order, 24 pages. 1997.

# Asymptotic Stability of Planar Rarefaction Wave for Scalar Viscous Conservation Law

KAZUO ITO

Department of Mathematics  
Faculty of Science  
Hokkaido University  
Sapporo 060, Japan

## Abstract

We study asymptotic stability of the planar rarefaction wave in one or two space dimensional scalar viscous conservation law for nonsmooth initial data.

**AMS subject classification:** 35L65, 35L67, 76L05.

**Keywords:** planar rarefaction wave, asymptotic stability, viscous conservation law.

## 1 Introduction

We study the initial value problem of the scalar viscous conservation law in one or two space dimensions:

$$\begin{cases} u_t + \sum_{i=1}^N (f_i(u))_{x_i} = \sum_{i=1}^N u_{x_i x_i}, \\ u(0, x) = u_0(x). \end{cases} \quad (1)$$

Here  $N = 1$  or  $2$  and  $u = u(t, x)$  is unknown real-valued function of time  $t \geq 0$  and position  $x = x_1 \in R$  or  $x = (x_1, x_2) \in R^2$ . We assume that each  $f_i$  is smooth and  $f_1$  is convex, i.e., there is an  $\alpha > 0$  such that  $f_1''(\lambda) \geq \alpha$  for  $\lambda \in R$ . We also assume that initial data  $u_0(x)$  satisfy

$$u_0(x) \rightarrow u_{\pm} \quad \text{as } x_1 \rightarrow \infty \text{ for any } x_2 \in R,$$

where  $u_-$  and  $u_+$  are given constants with  $u_- < u_+$ . Our purpose in this paper is to study asymptotic behavior of solutions to (1) as  $t \rightarrow \infty$  when initial data  $u_0(x)$  are not necessarily smooth.

Asymptotic behavior of solutions to (1) as  $t \rightarrow \infty$  is related to the planar rarefaction wave  $r(t, x_1)$ , which is defined by the unique weak solution of the Riemann problem of the form

$$\begin{cases} r_t + (f(r))_{x_1} = 0, & t > 0, x_1 \in R, \\ r(0, x_1) = r_0(x_1), & x_1 \in R, \end{cases} \quad (2)$$

where

$$r_0(x_1) = \begin{cases} u_-, & x_1 < 0, \\ u_+, & x_1 > 0. \end{cases} \quad (3)$$

In fact there are several works on asymptotic stability of the planar rarefaction wave in (1), see [4, 1, 8, 5, 7] and references cited there. Especially Xin [8] first studied the case  $N = 2$  and showed that the planar rarefaction wave  $r(t, x_1)$  was stable in  $L^\infty(R^2)$  when  $u_0(x)$  were close to smooth functions  $u_0^r(x_1)$  in  $H^2(R^2)$ ; here  $u_0^r(x_1)$  have the properties such as

$$\lim_{x_1 \rightarrow \pm} u_0^r(x_1) = u_{\pm}, \quad \frac{du_0^r}{dx_1}(x_1) > 0, \quad \left| \frac{d^2 u_0^r}{dx_1^2}(x_1) \right| \leq k_0 \frac{du_0^r}{dx_1}(x_1),$$

where  $k_0$  is a positive constant. Under the same assumption as [8], the author [5] gave the convergence rate of solutions to (1) toward  $r(t, x_1)$  when  $|u_+ - u_-|$  was small and  $u_0 - u_0^r \in L^1(\mathbb{R}^2)$ . Recently Nishikawa and Nishihara [7] obtained the same result as [5] without smallness conditions.

Our aim in this paper is to give the same result as [7] for a class containing nonsmooth initial data, which properly includes the class treated in [7]. In the next section we give our main result. After stating some preliminaries in Section 3, we prove main result in the last section. The ordering principle plays an essential role in the proof.

## 2 Main result

To begin with, we define smooth rarefaction wave  $w(t, x_1)$ , which is first introduced by Nishikawa and Nishihara [7], by the unique solution of the following initial value problem:

$$\begin{cases} w_t + (f_1(w))_{x_1} = w_{x_1 x_1} + \frac{f'''(w)}{f''(w)} w_{x_1}^2, & t > -1, x_1 \in \mathbb{R}, \\ w(-1, x_1) = r_0(x_1), & x_1 \in \mathbb{R}. \end{cases} \quad (4)$$

Put  $w_0(x_1) \equiv w(0, x_1)$ . The smooth rarefaction wave  $w(t, x_1)$  has the following properties.

**Lemma 1** [2], [6] *The smooth rarefaction wave  $w(t, x_1)$  satisfies the following properties on  $[0, \infty) \times \mathbb{R}$ : for any  $p \in [1, \infty]$ , there are constants  $c_d > 0$ ,  $C_d > 0$  (depending on  $d = |u_+ - u_-|$ ), and  $C > 0$  such that*

$$\begin{aligned} u_- < w(t, x_1) < u_+, \quad |w(t, x_1) - u_{\pm}| \leq C_d e^{-c_d x_1^2}, \\ w_{x_1}(t, x_1) > 0, \\ \|w_{x_1}(t)\|_{L^p(\mathbb{R})} \leq C d^{1/p} (1+t)^{-(1-1/p)}, \\ \|(w-r)(t)\|_{L^p(\mathbb{R})} \leq C \begin{cases} (1+t)^{-(1-1/p)/2}, & p \in (1, \infty], \\ \log(2+dt^{1/2}), & p = 1. \end{cases} \end{aligned}$$

Let us introduce the viscous rarefaction wave  $U(t, x_1)$  by the unique solution of the following initial value problem:

$$\begin{cases} U_t + (f_1(U))_{x_1} = U_{x_1 x_1}, & t > 0, x_1 \in \mathbb{R}, \\ U(0, x_1) = w_0(x_1), & x_1 \in \mathbb{R}. \end{cases} \quad (5)$$

$U(t, x_1)$  has the following properties.

**Lemma 2** [8], [5]  *$U(t, x_1)$  is smooth and  $U_{x_1}(t, x_1) > 0$  in  $[0, \infty) \times \mathbb{R}$ . Furthermore,*

$$\begin{aligned} U - w &\in C([0, \infty); H^2(\mathbb{R})), \\ \|U(t) - r(t)\|_{L^2(\mathbb{R})} &= O(t^{-1/4} \log t) \quad \text{as } t \rightarrow \infty. \end{aligned}$$

We define a series of sets of functions:

$$\begin{aligned} E_0(\mathbb{R}^N) &= \{v_0 \in (L^2 \cap L^1)(\mathbb{R}^N); \exists \varphi_0, \exists \psi_0 \in (H^2 \cap L^1)(\mathbb{R}^N) \text{ s.t.} \\ &\quad \varphi_0(x) \leq v_0(x) \leq \psi_0(x) \text{ a.e.}\}, \\ Y^0(\mathbb{R}^N) &= \{v \in C([0, \infty); (L^2 \cap L^1)(\mathbb{R}^N)) \cap L^\infty((0, \infty) \times \mathbb{R}^N); \\ &\quad Dv \in L^2(0, T; L^2(\mathbb{R}^N)) (\forall T > 0)\}, \\ Y^2(\mathbb{R}^N) &= \{v \in C([0, \infty); (H^2 \cap L^1)(\mathbb{R}^N)) \cap L^\infty((0, \infty) \times \mathbb{R}^N); \\ &\quad Dv \in L^2(0, T; H^2(\mathbb{R}^N)), \\ &\quad t^{1/2} Dv \in L^\infty(0, T; L^1(\mathbb{R}^N)) (\forall T > 0)\} \end{aligned}$$

where  $D = \partial_{x_1}$  for  $N = 1$  and  $D = (\partial_{x_1}, \partial_{x_2})$  for  $N = 2$ ,

$$D_0(R^N) = \{w_0\} + E_0(R^N),$$

$$X(R) = \{w\} + Y^0(R), \quad X(R^2) = \{U\} + Y^0(R^2).$$

**Definition** Let  $u_0 \in D_0(R^N)$ . We call  $u(t, x)$  a global *mild solution* of (1), if  $u(t, x)$  satisfies the following conditions:  $u \in X(R^N)$  and

$$u(t) = S(t)u_0 - \sum_{i=1}^N \int_0^t \partial_{x_i} S(t-\tau) f_i(u)(\tau) d\tau \quad \text{for } t \geq 0.$$

Here

$$(S(t)u_0)(x) = \int_{R^N} G(t, x-x') u_0(x') dx'$$

and  $G(t, x) = (4\pi t)^{-N/2} \exp(-|x|^2/(4t))$  is the Gauss kernel.

Then our main result is stated as follows.

**Theorem 3** (*Asymptotic stability of rarefaction wave*). Let  $N = 1$  or  $2$ . For any  $u_0 \in D_0(R^N)$ , there exists a unique global mild solution  $u \in X(R^N)$  of (1) satisfying

$$\sup_{x_2 \in R} \|u(t, \cdot, x_2) - r(t, \cdot)\|_{L^2(R)} = O(t^{-1/4} \log t) \quad (6)$$

as  $t \rightarrow \infty$ .

*Remark.* When  $N = 1$ , it follows from Lemma 1 that

$$D_0(R) = \{r_0\} + E_0(R).$$

This shows that  $r(t, x_1)$  is asymptotically stable for  $E_0(R)$ -perturbations to  $r_0(x_1)$ .

### 3 Preliminaries

In this section we state some preliminaries needed in the proof of Theorem 3.

First of all, with the aid of Theorems 4, 5, 6 and 7 in [7], we define the mapping

$$\begin{aligned} F: (H^2 \cap L^1)(R^N) &\rightarrow Y^2(R^N), \\ \varphi_0 &\mapsto \varphi = F(\varphi_0) \end{aligned}$$

by utilizing the unique solution of the following initial value problem:

$$\begin{cases} \varphi_t + (f_1(\varphi + w) - f_1(w))_{x_1} = \varphi_{x_1 x_1} - \frac{f'''(w)}{f''(w)} w_{x_1}^2, & t > 0, x_1 \in R, \\ \varphi(0, x_1) = \varphi_0(x_1), & x_1 \in R \end{cases}$$

for  $N = 1$  and

$$\begin{cases} \varphi_t + (f_1(\varphi + U) - f_1(U))_{x_1} + (f_2(\varphi + U))_{x_2} = \sum_{i=1}^2 \varphi_{x_i x_i}, & t > 0, x \in R^2, \\ \varphi(0, x) = \varphi_0(x), & x \in R^2 \end{cases}$$

for  $N = 2$ .

From now on we will often use the following notations: Let  $j(x_1)$ ,  $x_1 \in R$ , be a nonnegative smooth function with  $\int_R j(x_1) dx_1 = 1$ , whose support is in  $[-1, 1]$ . For  $k = 1, 2, \dots$ , set  $j_k(x_1) = kj(kx_1)$  and

$$h^k(x_1) = \int_R j_k(x_1 - x_1') h(x_1') dx_1'$$

for functions  $h(x_1)$ . This is called the smoothing of  $h$ . We denote by  $C$  universal positive constants, whose values are different in each occasion.

The mapping  $\varphi = F(\varphi_0)$  has the following properties.

**Theorem 4** [7] (i) Let  $N = 1$ . Then,

$$\begin{aligned} & \|\varphi(t)\|_{H^m(R)}^2 + \int_0^t \int_R w_{x_1} \varphi^2 dx_1 d\tau \\ & + \int_0^t \|\varphi_{x_1}(\tau)\|_{H^m(R)}^2 d\tau \leq C(\|\varphi_0\|_{H^m(R)}^2 + 1), \quad t \geq 0, \end{aligned}$$

for  $m = 0, 1, 2$ , and

$$\|\varphi(t)\|_{L^2(R)} = O(t^{-1/4} \log t) \quad \text{as } t \rightarrow \infty.$$

(ii) Let  $N = 2$ . Then,

$$\begin{aligned} & \|\varphi(t)\|_{H^m(R^2)}^2 + \int_0^t \int_{R^2} U_{x_1} \varphi^2 dx d\tau \\ & + \int_0^t \|D\varphi(\tau)\|_{H^m(R^2)}^2 d\tau \leq C\|\varphi_0\|_{H^m(R^2)}^2, \quad t \geq 0, \end{aligned}$$

for  $m = 0, 1, 2$ , and

$$\sup_{x_2 \in R} \|\varphi(t, \cdot, x_2)\|_{L^2(R)} = O(t^{-1/4} \log t) \quad \text{as } t \rightarrow \infty.$$

**Lemma 5** Let  $N = 1$  or  $2$ .

(i) ( $L^1$ -contraction). Let  $v_0, V_0 \in (H^2 \cap L^1)(R^N)$ . Then,

$$\|F(v_0)(t) - F(V_0)(t)\|_{L^1(R^N)} \leq \|v_0 - V_0\|_{L^1(R^N)}, \quad t \geq 0. \quad (7)$$

(ii) (Ordering principle). In (i), if  $v_0(x) \leq V_0(x)$  a.e., then

$$F(v_0)(t, x) \leq F(V_0)(t, x) \quad \text{a.e.} \quad (8)$$

*Proof.* We only show the proof for the case  $N = 1$ , since the case  $N = 2$  can be treated in a similar way.

(i): For  $v = F(v_0)$  and  $V = F(V_0)$ , set  $z = v - V$ . Then  $z$  satisfies

$$\begin{cases} z_t + (f_1(v+w) - f_1(V+w))_{x_1} = z_{x_1 x_1}, \\ z(0, x_1) = z_0(x_1) \equiv (v_0 - V_0)(x_1). \end{cases} \quad (9)$$

Let

$$\chi(x_1) = \begin{cases} -1, & x_1 < 0, \\ 0, & x_1 = 0, \\ 1, & x_1 > 0. \end{cases}$$

Multiplying (9) by  $\chi^k(z)$  ( $\chi^k$  is the smoothing of  $\chi$ ) and integrating it over  $(0, t) \times R$ , we have

$$\begin{aligned} & \int_R \int_0^{z(t, x_1)} \chi^k(\lambda) d\lambda dx_1 - \int_R \int_0^{z_0(x_1)} \chi^k(\lambda) d\lambda dx_1 \\ & + \int_0^t \int_R \chi^k(z) (f_1(v+w) - f_1(V+w))_{x_1} dx_1 d\tau \\ & = - \int_0^t \int_R (\chi^k)'(z) (z_{x_1})^2 dx_1 d\tau \leq 0. \end{aligned} \quad (10)$$

The third term in the left hand side (denoting it by  $I(t)$ ) is estimated as follows by using the boundedness of  $v, V$  and  $w$ :

$$\begin{aligned} I(t) &= - \int_0^t \int_R (\chi^k)'(z) z_{x_1} z \frac{f_1(v+w) - f_1(V+w)}{z} dx_1 d\tau \\ &= - \int_0^t \int_R \partial_{x_1} \int_0^z (\chi^k)'(\lambda) \lambda d\lambda \int_0^1 f_1'(V+w+\theta z) d\theta dx_1 d\tau \\ &= \int_0^t \int_R \int_0^z (\chi^k)'(\lambda) \lambda d\lambda \int_0^1 f_1''(V+w+\theta z) (V+w+\theta z)_{x_1} d\theta dx_1 d\tau \\ &\leq \frac{C}{k} \int_0^t (\|v_{x_1}\|_{L^1(R)} + \|V_{x_1}\|_{L^1(R)} + \|w_{x_1}\|_{L^1(R)}) d\tau. \end{aligned}$$

Noting  $v, V \in Y^2(R)$  and using Lemma 1, we find that the last term is integrable for any  $t \geq 0$ . Letting  $k \rightarrow \infty$  in (10), we get (7).

(ii): It is easy to see

$$\int_R z(t, x_1) dx_1 = \int_R z_0(x_1) dx_1.$$

Adding the above and (7), we have

$$\int_R (|z| + z)(t, x_1) dx_1 \leq \int_R (|z_0| + z_0)(x_1) dx_1 = 0,$$

since  $z_0(x_1) \leq 0$  a.e.. Nonnegativeness of  $(|z| + z)(t, x_1)$  implies  $(|z| + z)(t, x_1) = 0$  a.e., which gives  $z(t, x_1) \leq 0$  a.e.. This completes the proof.  $\square$

## 4 Proof of the main theorem

*Proof of Theorem 3.* We only show the proof for  $N = 2$ , since the case  $N = 1$  can be proved in a similar way.

Let  $u_0(x) = w_0(x) + v_0(x)$ , where  $v_0 \in E_0(R^2)$ . Put  $v^k \equiv F(v_0^k)$ , where  $v_0^k$  is the smoothing of  $v_0$ . We begin by showing the following lemma.

**Lemma 6** *There are a subsequence of  $\{v^k\}_{k=1}^\infty$  (denoting it by  $\{y^k\}_{k=1}^\infty$ ) and a  $v \in Y^0(R^2)$  such that the following convergences hold:*

$$\begin{aligned} y^k &\rightarrow v && \text{in } C([0, T]; L^1(R^2)), \\ y^k &\rightarrow v && \text{in } L^\infty(0, T; L^2(R^2)) \text{ weakly}^*, \\ Dy^k &\rightarrow Dv && \text{in } L^2(0, T; L^2(R^2)) \text{ weakly}, \\ y^k &\rightarrow v && \text{a.e. } (0, T) \times R^2 \end{aligned}$$

as  $k \rightarrow \infty$  for any  $T > 0$ .

*Proof of Lemma 6.* Throughout this proof, we denote  $Q_n = C([0, n]; L^1(R^2))$  for  $n = 1, 2, \dots$ . It follows from Lemma 5 (i) that

$$\|v^k(t) - v^l(t)\|_{L^1(R^2)} \leq \|v_0^k - v_0^l\|_{L^1(R^2)}, \quad t \geq 0, \quad (11)$$

for  $k, l = 1, 2, \dots$ , which shows that in particular  $\{v^k\}_{k=1}^\infty$  is a Cauchy sequence in  $Q_1$ . Then, there are a subsequence  $\{v^{k,1}\}_{k=1}^\infty$  of  $\{v^k\}_{k=1}^\infty$  and a  $q^1 \in Q_1$  such that

$$v^{k,1} \rightarrow q^1 \quad \text{in } Q_1 \text{ as } k \rightarrow \infty.$$

Inductively, it follows from (11) that  $\{v^{k,n}\}_{k=1}^\infty$  is a Cauchy sequence in  $Q_{n+1}$ . Then, there are a subsequence  $\{v^{k,n+1}\}_{k=1}^\infty$  of  $\{v^{k,n}\}_{k=1}^\infty$  and a  $q^{n+1} \in Q_{n+1}$  such that

$$v^{k,n+1} \rightarrow q^{n+1} \quad \text{in } Q_{n+1} \text{ as } k \rightarrow \infty$$

and

$$q^{n+1} = q^n \quad \text{a.e. } [0, n] \times R^2.$$

Define

$$\begin{aligned} y^k &= v^{k,k}, \\ v(t, x) &= q^n(t, x) \quad \text{if } (t, x) \in [0, n] \times R^2. \end{aligned}$$

Then, we obtain

$$y^k \rightarrow v \quad \text{in } C([0, T]; L^1(R^2)) \text{ as } k \rightarrow \infty$$

for any  $T > 0$ . We can also choose a subsequence of  $\{y^k\}_{k=1}^\infty$  (denoting it again by  $\{y^k\}_{k=1}^\infty$ ) such as

$$y^k \rightarrow v \quad \text{as } k \rightarrow \infty \text{ a.e. } (0, \infty) \times R^2.$$



On the other hand, it follows from Theorem 4 that

$$\begin{aligned} & \|y^k(t)\|_{L^2(R^2)}^2 + \int_0^t \int_{R^2} U_{x_1}(y^k)^2 dx d\tau \\ & + \int_0^t \|Dy^k(\tau)\|_{L^2(R^2)}^2 d\tau \leq C \|y^k(0)\|_{L^2(R^2)}^2 \leq \|v_0\|_{L^2(R^2)}^2, \end{aligned} \quad (12)$$

which shows the boundednesses of  $\{y^k\}_{k=1}^\infty$  in  $L^\infty(0, T; L^2(R^2))$  and of  $\{Dy^k\}_{k=1}^\infty$  in  $L^2(0, T; L^2(R^2))$  for any  $T > 0$ . Then, there is a subsequence of  $\{y^k\}_{k=1}^\infty$  (denoting it again by  $\{y^k\}_{k=1}^\infty$ ) such that

$$\begin{aligned} y^k & \rightarrow v & \text{in } L^\infty(0, T; L^2(R^2)) \text{ weakly}^*, \\ Dy^k & \rightarrow Dv & \text{in } L^2(0, T; L^2(R^2)) \text{ weakly} \end{aligned}$$

as  $k \rightarrow \infty$  for any  $T > 0$ .

We show  $v \in L^\infty((0, \infty) \times R^2)$ . Note

$$\varphi_0^k(x) \leq y^k(0, x) \leq \psi_0^k(x) \quad \text{a.e.,}$$

since  $v_0 \in E_0(R^2)$ . It then follows from Lemma 5 (ii) that

$$\varphi^k(t, x) \leq y^k(t, x) \leq \psi^k(t, x) \quad \text{a.e.,} \quad (13)$$

where  $\varphi^k = F(\varphi_0^k)$  and  $\psi^k = F(\psi_0^k)$ . Furthermore, we can show by using the above procedure that

$$\varphi^k \rightarrow \varphi, \quad \psi^k \rightarrow \psi \quad \text{as } k \rightarrow \infty \text{ a.e. } (0, \infty) \times R^2,$$

where  $\varphi = F(\varphi_0)$  and  $\psi = F(\psi_0)$ . With this fact in mind, letting  $k \rightarrow \infty$  in (13), we obtain

$$\varphi(t, x) \leq v(t, x) \leq \psi(t, x) \quad \text{a.e..} \quad (14)$$

Since  $\varphi$  and  $\psi \in L^\infty((0, \infty) \times R^2)$ , we thus prove  $v \in L^\infty((0, \infty) \times R^2)$ .

It remains to prove  $v \in C([0, \infty); L^2(R^2))$ . For this purpose, we first check that

$$\lim_{k \rightarrow \infty} \sup_{t \in [0, T]} \|(f_i(y^k + U) - f_i(v + U))(t)\|_{L^1(R^2)} = 0 \quad (15)$$

for any  $T > 0$  and  $i = 1, 2$ . In fact,

$$\begin{aligned} & \sup_{t \in [0, T]} \|(f_i(y^k + U) - f_i(v + U))(t)\|_{L^1(R^2)} \\ & = \sup_{t \in [0, T]} \left\| \int_0^1 f_i'(v + U + \theta(y^k - v)) d\theta (y^k - v) \right\|_{L^1(R^2)}(t) \\ & \leq C \sup_{t \in [0, T]} \|(y^k - v)(t)\|_{L^1(R^2)} \rightarrow 0 \quad \text{as } k \rightarrow \infty, \end{aligned}$$

where we have used Lemma 2, (13), Theorem 4 and (14).

Note  $y^k$  satisfies

$$\begin{aligned} y^k(t) & = S(t)y(0)^k - \int_0^t \partial_{x_1} S(t-\tau)(f_1(y^k + U) - f_1(U))(\tau) d\tau \\ & \quad - \int_0^t \partial_{x_2} S(t-\tau)f_2(y^k + U)(\tau) d\tau. \end{aligned}$$

Letting  $k \rightarrow \infty$  in  $C([0, T]; L^1(R))$  for any fixed  $T > 0$  with the aid of (15), we obtain

$$\begin{aligned} v(t) & = S(t)v_0 - \int_0^t \partial_{x_1} S(t-\tau)(f_1(v + U) - f_1(U))(\tau) d\tau \\ & \quad - \int_0^t \partial_{x_2} S(t-\tau)f_2(v + U)(\tau) d\tau. \end{aligned} \quad (16)$$

Let  $t_0 \geq 0$  be any fixed time. We show that the last term in the right hand side of (16) is continuous at  $t_0$  in  $L^2(\mathbb{R}^2)$ :

$$\begin{aligned} & \left\| \int_0^t \partial_{x_2} S(t-\tau) f_2(v+U)(\tau) d\tau - \int_0^{t_0} \partial_{x_2} S(t_0-\tau) f_2(v+U)(\tau) d\tau \right\|_{L^2(\mathbb{R}^2)} \\ & \leq \left\| \int_{t_0}^t \partial_{x_2} S(t-\tau) f_2(v+U)(\tau) d\tau \right\|_{L^2(\mathbb{R}^2)} \\ & \quad + \left\| \int_0^{t_0} \partial_{x_2} (S(t-\tau) - S(t_0-\tau)) f_2(v+U)(\tau) d\tau \right\|_{L^2(\mathbb{R}^2)} \\ & \leq C \left| \int_{t_0}^t \|v_{x_2}(\tau)\|_{L^2(\mathbb{R}^2)} d\tau \right| + \int_0^{t_0} \|(S(t-\tau) - S(t_0-\tau))(f_2'(v+U)v_{x_2})(\tau)\|_{L^2(\mathbb{R}^2)} d\tau. \end{aligned}$$

Then the Lebesgue dominated convergence theorem implies that the last integral tends to 0 as  $t \rightarrow t_0$ . The continuity of the second term in the right hand side of (16) in  $L^2(\mathbb{R}^2)$  can be shown in a similar way. Thus it is proved that  $v \in C([0, \infty); L^2(\mathbb{R}^2))$ , and hence  $v \in Y^0(\mathbb{R}^2)$ .

This completes the proof of Lemma 6.  $\square$

We show  $u(t, x) = U(t, x_1) + v(t, x)$  is the desired mild solution of (1). Adding (16) and

$$U(t) = S(t)w_0 - \int_0^t \partial_{x_1} S(t-\tau) f_1(U)(\tau) d\tau,$$

we get

$$u(t) = S(t)u_0 - \sum_{i=1}^2 \int_0^t \partial_{x_i} S(t-\tau) f_i(u)(\tau) d\tau,$$

which shows that  $u(t, x)$  is the desired global mild solution of (1).

We show (6). Utilizing (14), Theorem 4 and Lemma 2, we have

$$\begin{aligned} & \sup_{x_2 \in \mathbb{R}} \|u(t, \cdot, x_2) - \tau(t, \cdot)\|_{L^2(\mathbb{R})} \\ & \leq \sup_{x_2 \in \mathbb{R}} \|u(t, \cdot, x_2) - U(t, \cdot)\|_{L^2(\mathbb{R})} + \|U(t) - \tau(t)\|_{L^2(\mathbb{R})} \\ & \leq \sup_{x_2 \in \mathbb{R}} \max(\|\varphi(t, \cdot, x_2)\|_{L^2(\mathbb{R})}, \|\psi(t, \cdot, x_2)\|_{L^2(\mathbb{R})}) + \|U(t) - \tau(t)\|_{L^2(\mathbb{R})} \\ & \leq O(t^{-1/4} \log t) \quad \text{as } t \rightarrow \infty, \end{aligned}$$

which shows (6).

It remains to prove uniqueness of mild solutions of (1). Let  $u$  and  $\bar{u}$  be two mild solutions of (1) with initial data  $u(0) = \bar{u}(0) = u_0$ . Then,

$$(u - \bar{u})(t) = - \sum_{i=1}^N \int_0^t \partial_{x_i} S(t-\tau) (f_i(u) - f_i(\bar{u}))(\tau) d\tau.$$

It follows from the boundednesses of  $u$  and  $\bar{u}$  that

$$\begin{aligned} \|(u - \bar{u})(t)\|_{L^1(\mathbb{R}^N)} & \leq C \sum_{i=1}^N \int_0^t \partial_{x_i} S(t-\tau) \|(f_i(u) - f_i(\bar{u}))(\tau)\|_{L^1(\mathbb{R}^2)} d\tau \\ & \leq C \int_0^t (t-\tau)^{-1/2} \|(u - \bar{u})(\tau)\|_{L^1(\mathbb{R}^2)} d\tau. \end{aligned}$$

Applying Lemma 7.1.1 of [3], we get  $\|(u - \bar{u})(t)\|_{L^1(\mathbb{R}^2)} = 0$ ,  $t \geq 0$ .

The proof of Theorem 3 is complete.  $\square$

## References

- [1] E. Harabetian, *Rarefactions and large time behavior for parabolic equations and monotone schemes*, Commun. Math. Phys. 114, 527–536(1988).
- [2] Y. Hattori and K. Nishihara, *A note on the stability of the rarefaction wave of the Burgers equation*, Jpn. J. Indust. Appl. Math. 8, 85–96(1991).
- [3] D. Henry, *Geometric theory of semilinear parabolic equations*, Lecture Notes in Math. 840, Springer-Verlag, Berlin, Heidelberg, 1981.
- [4] A. M. Il'in and O. A. Oleinik, *Asymptotic behavior of solutions of the Cauchy problem for certain quasilinear equations for large time* (Russian), Math. Sob. 51, 191–216(1960).
- [5] K. Ito, *Asymptotic decay toward the planar rarefaction waves of solutions for viscous conservation laws in several space dimensions*, Math. Models Methods Appl. Sci. 6, 315–338(1996).
- [6] S. Kawashima and Y. Tanaka, *Stability of rarefaction waves for a model equation of a radiating gas*, in preparation.
- [7] M. Nishikawa and K. Nishihara, *Asymptotics toward the planar rarefaction wave for viscous conservation law in two space dimensions*, preprint.
- [8] Z. P. Xin, *Asymptotic stability of planar rarefaction waves for viscous conservation laws in several space dimensions*, Trans. Amer. Math. Soc. 319, 805–820(1990).