



Title	A remark on the quantale structure of multisets
Author(s)	Arase, T.
Citation	Hokkaido University Preprint Series in Mathematics, 393, 1-10
Issue Date	1997-10-1
DOI	10.14943/83539
Doc URL	http://hdl.handle.net/2115/69143
Type	bulletin (article)
File Information	pre393.pdf



[Instructions for use](#)

**A remark on the quantale structure
of multisets**

Tadamune Arase

Series #393. October 1997

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #368 K. Ito, BV-solutions of a hyperbolic-elliptic system for a radiating gas, 33 pages. 1997.
- #369 M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, 20 pages. 1997.
- #370 N.H. Bingham and A. Inoue, An Abel-Tauber theorem for Hankel transforms, 8 pages. 1997.
- #371 T. Nakazi and H. Sawada, The commutator ideal in Toeplitz algebras for uniform algebras and the analytic structure, 9 pages. 1997.
- #372 M.-H. Giga and Y. Giga, Stability for evolving graphs by nonlocal weighted curvature, 70 pages. 1997.
- #373 T. Nakazi, Brown-Halmos type theorems of weighted Toeplitz operators, 14 pages. 1997.
- #374 J. Inoue and S.-E. Takahashi, On characterizations of the image of Gelfand transform of commutative Banach algebras, 30 pages. 1997.
- #375 L. Solomon and H. Terao, The double coxeter arrangement, 21 pages. 1997.
- #376 G. Ishikawa and T. Morimoto, Solution surfaces of Monge-Ampère equations, 15 pages. 1997.
- #377 G. Ishikawa, A relative transversality theorem and its applications, 16 pages. 1997.
- #378 J. Inoue and T. Nakazi, On the zeroes of solutions of an extremal problem in H^1 , 14 pages. 1997.
- #379 Y. Giga and K. Ito, On pinching of curves moved by surface diffusion, 12 pages. 1997.
- #380 F. Hiroshima, Weak coupling limit removing an ultraviolet cut-off for a Hamiltonian of particles interacting with a scalar field, 39 pages. 1997.
- #381 Y. Giga, S. Matsui and Y. Shimizu, On estimates in Hardy spaces for the Stokes flow in a half space, 13 pages. 1997.
- #382 A. Arai, A new estimate for the ground state energy of Schrödinger operators, 12 pages. 1997.
- #383 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the Sobolev space of critical order, 24 pages. 1997.
- #384 K. Ito, Asymptotic stability of planar rarefaction wave for scalar viscous conservation law, 8 pages. 1997.
- #385 A. Arai, Representation-theoretic aspects of two-dimensional quantum systems in singular vector potentials: canonical commutation relations, quantum algebras, and reduction to lattice quantum systems, 32 pages. 1997.
- #386 P. Aviles and Y. Giga, On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields, 21 pages. 1997.
- #387 T. Nakazi and T. Yamamoto, Norms of some singular integral operators and their inverse operators, 28 pages. 1997.
- #388 M.-H. Giga and Y. Giga, Remarks on convergence of evolving graphs by nonlocal curvature, 18 pages. 1997.
- #389 T. Tsukada, Reticular Lagrangian singularities, 41 pages. 1997.
- #390 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the homogeneous Sobolev space, 12 pages. 1997.
- #391 Y. Giga, M. Ohnuma and M.-H. Sato, On strong maximum principle and large time behaviour of generalized mean curvature flow with the Neumann boundary condition, 24 pages. 1997.
- #392 T. Tsujishita and H. Watanabe, Monoidal closedness of the category of simulations, 24 pages. 1997.

A remark on the quantale structure of multisets

Tadamune Arase
Department of Mathematics
Hokkaido University

July 28, 1997

abstract

The set $\mathcal{P}^*(I)$ of finite multisets on a set I has a natural quantale structure. In this note we study the quantic nuclei of this quantale, and show that they are parametrized by those subsets of the infinite multisets $\mathcal{P}^{**}(I)$ whose elements are mutually uncomparable. We also study the quantic nuclei of another natural quantale structure on $\mathcal{P}^*(I)$ with respect to the opposite order.

1991 Mathematics Subject Classification: 18B35, 06F99

1 Introduction

1.1 Quantale and quantic nucleus

We recall briefly basic notions of quantale and fix notations. We use standard notations and terminologies on lattices [2].

A quantale is a complete lattice Q endowed with an associative binary operation \otimes satisfying

$$\begin{aligned}\bigvee S \otimes q &= \bigvee \{s \otimes q \mid s \in S\}, \\ q \otimes \bigvee S &= \bigvee \{q \otimes s \mid s \in S\}.\end{aligned}$$

for $S \subseteq Q$ and $q \in Q$. In this note, we consider only commutative quantales, i.e, we assume \otimes is commutative. A quantale has another binary operation " \rightarrow " defined by

$$a \rightarrow b = \bigvee \{q \mid q \otimes a \leq b\}.$$

A quantic nucleus $j : Q \rightarrow Q$ is a closure operator satisfying

$$j(a \otimes b) \leq ja \otimes jb$$

for $a, b \in Q$. Its fixed point set

$$S_j = \{q \in Q | jq = q\}$$

is meet-closed and satisfies

$$a \rightarrow s \in S_j$$

for all $a \in Q$ and $s \in S_j$.

Conversely any subset $S \subseteq Q$ satisfying these two conditions is the fixed point set of the quantic nucleus j_S defined by

$$j_S(q) = \bigwedge \{s \in S | q \leq s\},$$

[3.Proposition 3.2.1].

1.2 The quantales of multisets

In this note we study the quantale $\mathcal{P}^*(I)$ of finite multisets on a set I defined as follows.

Let N be the set of natural numbers and put $N^* = N \cup \{0, \infty\}$. We regard N^* as an ordered set with $0 < 1 < 2 < \dots < \infty$.

Let I be a finite set. The set $\mathcal{P}^*(I)$ is defined by

$$\mathcal{P}^*(I) = \{(n_i)_{i \in I} | n_i \in N^* \setminus \{\infty\}\} \cup \{\infty\}.$$

This set has a quantale structure defined by

$$\begin{cases} (n \otimes m)_i = n_i + m_i, & \text{if } n, m \neq \top, \\ n \otimes m = \top, & \text{otherwise,} \end{cases}$$

and

$$n \leq m \iff m = \top \text{ or } n_i \leq m_i \text{ for all } i,$$

where $\top = \infty$. This quantale is called the quantale of finite multisubsets of I .

For $S \subseteq \mathcal{P}^*(I)$

$$(\bigvee S)_i = \sup\{n_i | n \in S\}$$

if $\sup\{n_i | n \in S\}$ is finite for all $i \in I$ and

$$\bigvee S = \top$$

if $\sup\{n_i | n \in S\} = \infty$ for some i .

The meet of S is given by

$$(\bigwedge S)_i = \inf\{n_i | n \in S\}.$$

1.3 The quantale of infinite multisets on I

The product $(N^*)^I$ has a commutative quantale structure defined by

$$n \leq m \iff n_i \leq m_i \quad \text{for all } i \in I$$

and

$$(n \otimes m)_i = n_i + m_i \quad \text{for } i \in I$$

where

$$n + \infty = \infty$$

for all $n \in N^*$. The top is the element \top defined by $(\top)_i = \infty$ for all i . This quantale will be called *the quantale of infinite multisets on I* , and will be denoted by $\mathcal{P}^{**}(I)$.

Note that the inclusion $i : \mathcal{P}^*(I) \hookrightarrow \mathcal{P}^{**}(I)$ is a monoid homomorphism but does not preserve the join. For example, suppose $I = \{a, b\}$ and define $s_k (k = 1, 2, \dots)$ by

$$(s_k)_a = k, \quad (s_k)_b = 0.$$

Then $\bigvee \{s_k\} = \top$ in $\mathcal{P}^*(I)$, but

$$(\bigvee \{s_k\})_a = \infty, \quad (\bigvee \{s_k\})_b = 0$$

in $\mathcal{P}^{**}(I)$.

2 Quantic nuclei on $\mathcal{P}^*(I)$

Since $n \rightarrow m = \bigvee \{x \mid x \otimes n \leq m\}$ for $n, m \in \mathcal{P}^*(I)$, we have

$$n \rightarrow m = \begin{cases} \top, & \text{if } m = \top, \\ 0, & \text{if } m \neq \top \text{ and } m \leq n, \\ m \ominus n, & \text{otherwise,} \end{cases}$$

where $(m \ominus n)_i = [m_i - n_i]_+$, where $[n]_+ = n$ if $n \geq 0$ and 0 if $n < 0$. From this, we can show the following proposition.

Proposition 1 *Let S be a subset of $\mathcal{P}^*(I)$. Then S is the set of the fixed points of a quantic nucleus of $\mathcal{P}^*(I)$ if and only if S contains both \top and the downset $\downarrow s := \{u \mid u \leq s\}$ of every $s \in S \setminus \{\top\}$, namely,*

$$S \supseteq \{\top\} \cup \bigcup_{s \in S \setminus \{\top\}} \downarrow s. \quad (1)$$

Proof. For $s \in S$

$$\{q \rightarrow s | q \in \mathcal{P}^*(I)\} = \begin{cases} \downarrow s, & \text{if } s \neq \top, \\ \{\top\}, & \text{if } s = \top. \end{cases}$$

In fact $u = (s \ominus u) \rightarrow s$ if $u \leq s \neq \top$, and $q \rightarrow \top = \top$. From this, for the subset $S \subseteq \mathcal{P}^*(I)$ which contains \top ,

$$\bigcup_{s \in S} \{q \rightarrow s | q \in \mathcal{P}^*(I)\} = \{\top\} \cup \bigcup_{s \in S \setminus \{\top\}} \downarrow s. \quad (2)$$

Let S be the fixed point set of a quantic nucleus. Then S is meet-closed and

$$S \supseteq \bigcup_{s \in S} \{q \rightarrow s | q \in \mathcal{P}^*(I)\}.$$

The meet-closedness of S implies S contains $\top = \bigwedge \emptyset$, and hence the equation (2) holds. Therefore, S satisfies the condition (1).

Conversely, let S be the set which satisfies the condition (1). Since S contains \top , the equation (2) holds, whence

$$S \supseteq \bigcup_{s \in S} \{q \rightarrow s | q \in \mathcal{P}^*(I)\}.$$

Since the inverse inclusion of condition (1) holds for any $S \subseteq \mathcal{P}^*(I)$, we obtain

$$S = \bigcup_{s \in S} \{q \rightarrow s | q \in \mathcal{P}^*(I)\}.$$

This implies S is meet closed. Therefore, S is the fixed point set of a quantic nucleus of $\mathcal{P}^*(I)$. ■

Corollary 2 Let S_j be the fixed point set of a quantic nucleus j of $\mathcal{P}^*(I)$. Then

$$j(n) = \begin{cases} n, & n \in S_j, \\ \top, & \text{otherwise.} \end{cases}$$

Proof. Recall

$$j(n) = \bigwedge \{s \in S_j | n \leq s\}.$$

But $n \leq s$ ($s \in S_j$) implies $n \in S_j$. Hence if $n \notin S_j$ then

$$j(n) = \bigwedge \emptyset = \top. \quad \blacksquare$$

Let $\mathcal{D}(\mathcal{P}^{**}(I))$ be the set of those subsets of the quantale $\mathcal{P}^{**}(I)$ whose elements are mutually uncomparable, i.e., $M \in \mathcal{D}(\mathcal{P}^{**}(I))$ if and only if no two elements $u, v \in M$ satisfies either $u \leq v$ or $v \leq u$. To prove the set $M \subseteq \mathcal{D}(\mathcal{P}^{**}(I))$ is finite, we show next lemma.

Lemma 3 *There is no infinite sequence $x_1, x_2, \dots \in (N^*)^n$ satisfying*

$$i < j \implies x_i \not\leq x_j, \quad \text{for all } i, j \in N^*.$$

Proof. Suppose $n = 1$. Because $x \not\leq y$ if and only if $x > y$ for $n = 1$, there is no infinite sequence $x_1, x_2, \dots \in N^*$ satisfying

$$i < j \implies x_i > x_j, \quad \text{for all } i, j \in N^*.$$

Now we assume there is no infinite sequence $x_1, x_2, \dots \in (N^*)^n$ satisfying

$$i < j \implies x_i \not\leq x_j, \quad \text{for all } i, j \in N^*$$

for $n \leq k$. Suppose there is an infinite sequence $x_1, x_2, \dots \in (N^*)^{k+1}$ satisfying

$$i < j \implies x_i \not\leq x_j, \quad \text{for all } i, j \in N^*.$$

Write $x_j \in (N^*)^{k+1}$ as

$$x_j = (x'_j, p_j), \text{ with } x'_j \in (N^*)^k \text{ and } p_j \in N^*.$$

Put $I = \{p_j | j = 1, 2, \dots\}$.

If I is finite, then there is a $p \in I$ with $J = \{j | p_j = p\}$ is infinite. Let $J = \{j_1 < j_2 < \dots\}$ and put $y_i = x'_{j_i}$. Then the infinite sequence $y_1, y_2, \dots \in (N^*)^k$ satisfies

$$i < j \implies x_i \not\leq x_j, \quad \text{for all } i, j \in N^*.$$

This contradicts the assumption.

If I is infinite, then there is an infinite sequence

$$j_1 < j_2 < \dots$$

with

$$p_{j_1} < p_{j_2} < \dots$$

Put $y_1 = x'_{j_1}$, then the infinite sequence $y_1, y_2, \dots \in (N^*)^k$ satisfies

$$i < j \implies x_i \not\leq x_j, \quad \text{for all } i, j \in N^*.$$

This contradicts the assumption. ■

Proposition 4 *The set $M \subseteq \mathcal{D}(\mathcal{P}^{**}(I))$ is finite.*

Proof. Suppose M is infinite. Then there is an infinite sequence $x_1, x_2, \dots \in M$, which satisfies

$$i < j \implies x_i \not\leq x_j, \quad \text{for all } i, j \in N^*$$

since all elements of M are mutually incomparable. This contradicts the conclusion of Lemma 3. Hence M must be finite. ■

Because for $M \in \mathcal{D}(\mathcal{P}^{**}(I))$

$$S_M = (\{\top\} \cup \bigcup_{m \in M} \downarrow m) \cap \mathcal{P}^*(I)$$

satisfies the condition of the Proposition 1, we can define the quantic nucleus j_M by

$$j_M(n) = \begin{cases} n, & \text{if } n \in S_M, \\ \top, & \text{otherwise.} \end{cases}$$

Theorem 5 *let $\mathcal{N}(\mathcal{P}^*(I))$ be the set of all the quantic nucleus of $\mathcal{P}^*(I)$. Then the map*

$$\pi : \mathcal{D}(\mathcal{P}^{**}(I)) \longrightarrow \mathcal{N}(\mathcal{P}^*(I)),$$

*which sends $M \in \mathcal{D}(\mathcal{P}^{**}(I))$ to the quantic nucleus j_M is a bijection.*

Proof. For the subset $S \subseteq \mathcal{P}^*(I)$ we define $\bar{S} \subseteq \mathcal{P}^{**}(I)$ as the set of

$$\bar{S} = \{\bigvee i(A) \mid A \subseteq S \setminus \{\top\}, A \text{ is directed}\}$$

where $i : \mathcal{P}^*(I) \longrightarrow \mathcal{P}^{**}(I)$ is the inclusion map (cf §1.3). Let

$$\mathcal{M}(S) = \{m \mid m \text{ is the maximal element of } S\}.$$

Let S_j be the fixed point set of a quantic nucleus j . From the above definition,

$$S_{\mathcal{M}(S_j)} = (\{\top\} \cup \bigcup_{m \in \mathcal{M}(S_j)} \downarrow m) \cap \mathcal{P}^*(I) = \{\top\} \cup \bigcup_{s \in S_j \setminus \{\top\}} \downarrow s = S_j.$$

This implies

$$\pi(\mathcal{M}(\bar{S}_j)) = j,$$

hence, π is surjection.

On the other hand,

$$\mathcal{M}(\bar{S}_M) = M,$$

In fact, we have $M \subseteq \mathcal{M}(\bar{S}_M)$ since $m = \bigvee (\downarrow m \cup \mathcal{P}^*(I))$.

We show now $M \supseteq \mathcal{M}(\bar{S}_M)$. For any $n \in \bar{S}_M$ there is a directed set A such that

$$n = \bigvee i(A)$$

where $i : \mathcal{P}^*(I) \rightarrow \mathcal{P}^{**}(I)$ is the inclusion map. We show that there is an upperbound of A in M . Suppose that for any $m \in M$ there is an $a_m \in A$ such that $a_m \not\leq m$. Since M is finite and A is directed, there is $a \in A$ such that for all $m \in M$, $a_m \leq a$. Because every $m \in M$ is maximal element of \bar{S}_M , we have there is a $m' \in M$ such that $a < m'$. From these $a_m \leq m'$, and this is contradiction. Therefore, for all $a \in A$ there is $m \in M$ such that m is comparable with a , and so $a \leq m$.

This implies π is injection. In fact, $S_M = S'_M$ implies $\mathcal{M}(\bar{S}_M) = \mathcal{M}(\bar{S}'_M)$, so $M = M'$. \blacksquare

3 The opposite order

The set $\mathcal{P}^*(I)$ has another quantale structure with the same binary product \otimes and with the opposite order \preceq defined by

$$n \preceq m \iff n = \perp \text{ or } n_i \geq m_i \text{ for all } i$$

where we write ∞ by \perp now. This type of quantales arises as the ideal quantales of the principal ideal domains [1]. The binary operator \rightarrow is the same as before but can be written now as

$$n \rightarrow m = \begin{cases} \perp, & \text{if } m = \perp, n \neq \perp, \\ 0, & \text{if } m = n = \perp \text{ or } n \preceq m \neq \perp, \\ m \ominus n, & \text{otherwise.} \end{cases}$$

This implies as before the following proposition.

Proposition 6 *Let S be a subset of $\mathcal{P}^*(I)$. Then S is the fixed point set of a quantic nucleus of $\mathcal{P}^*(I)$ if and only if S contains the all upsets $\uparrow s := \{u \mid u \succeq s\}$ of $s \in S \setminus \{\perp\}$, namely,*

$$S \supseteq \bigcup_{s \in S \setminus \{\perp\}} \uparrow s,$$

and S is meet-closed.

Proof. Because

$$\{q \rightarrow s \mid q \in \mathcal{P}^*(I)\} = \begin{cases} \uparrow s, & \text{if } s \neq \perp, \\ \{0, \perp\}, & \text{if } s = \perp, \end{cases}$$

we have

$$\bigcup_{s \in S} \{q \rightarrow s \mid q \in \mathcal{P}^*(I)\} = \begin{cases} \bigcup_{s \in S \setminus \{\perp\}} \uparrow s, & \text{if } \perp \notin S, \\ \{\perp\} \cup \bigcup_{s \in S \setminus \{\perp\}} \uparrow s, & \text{if } \perp \in S. \end{cases}$$

From this,

$$S \supseteq \bigcup_{s \in S \setminus \{\perp\}} \uparrow s \iff S \supseteq \bigcup_{s \in S} \{q \rightarrow s \mid q \in \mathcal{P}^*(I)\}. \quad (3)$$

Therefore,

$$S \supseteq \bigcup_{s \in S \setminus \{\perp\}} \uparrow s \text{ and } S \text{ is meet-closed,}$$

if and only if

$$S \supseteq \bigcup_{s \in S} \{q \rightarrow s \mid q \in \mathcal{P}^*(I)\} \text{ and } S \text{ is meet-closed,}$$

that is, S is the set of the fixed points of a quantic nucleus of $\mathcal{P}^*(I)$. ■

Note that the set S of the fixed points of a quantic nucleus is written by

$$S = \begin{cases} \bigcup_{s \in S \setminus \{\perp\}} \uparrow s, & \text{if } \perp \notin S, \\ \{\perp\} \cup \bigcup_{s \in S \setminus \{\perp\}} \uparrow s, & \text{if } \perp \in S. \end{cases}$$

since the inverse inclusion of the right hand side of (3), i.e.,

$$S \supseteq \bigcup_{s \in S} \{q \rightarrow s \mid q \in \mathcal{P}^*(I)\},$$

holds for any $S \subseteq \mathcal{P}^*(I)$.

There are two types of sets which satisfy the conditions of the proposition 3. The one is

$$S_m^1 = (\{\perp\} \cup \uparrow m) \cap \mathcal{P}^*(I),$$

for $m \in \mathcal{P}^{**}(I)$. We can define the quantic nucleus j_m^1 with the fixed point set S_m^1 by

$$j_m^1(n) = \begin{cases} n \vee m, & \text{if } n \neq \perp, \\ \perp, & \text{if } n = \perp, \end{cases}$$

for $n \in \mathcal{P}^*(I)$. The other one is

$$S_m^2 = \uparrow m,$$

for $m \in \mathcal{P}^*(I)$. We can define the quantic nucleus j_m^2 with the fixed point set S_m^2 by

$$j_m^2(n) = n \vee m,$$

for $n \in \mathcal{P}^*(I)$.

Lemma 7 Define the maps

$$\xi_1 : \mathcal{P}^{**}(I) \longrightarrow \mathcal{N}(\mathcal{P}^*(I)), \quad \xi_2 : \mathcal{P}^*(I) \longrightarrow \mathcal{N}(\mathcal{P}^*(I))$$

by

$$\xi_1(m) = j_m^1, \quad \xi_2(m) = j_m^2.$$

Then these maps are injection.

Proof. Let $m \neq m'$, then

$$S_m^1 \neq S_{m'}^1, \quad \text{and} \quad S_m^2 \neq S_{m'}^2.$$

Theorem 8 Then the map

$$\eta : \mathcal{P}^{**}(I) \amalg (\mathcal{P}^*(I) \setminus \{\perp\}) \longrightarrow \mathcal{N}(\mathcal{P}^*(I))$$

defined by

$$\eta = \xi_1 \amalg \xi_2$$

is a bijection.

Proof. Let S_j be the fixed point set of a quantic nucleus j , and $m = \bigwedge_{s \in S_j} (S_j \setminus \{\perp\})$.

If $\perp \notin S_j$, $m \in \mathcal{P}^*(I)$. Then

$$S_j = \bigcup_{s \in S_j \setminus \{\perp\}} \uparrow s = \uparrow m = S_m^2.$$

If $\perp \in S_j$, $m \in \mathcal{P}^{**}(I)$. Then

$$S_j = \{\perp\} \cup \bigcup_{s \in S_j \setminus \{\perp\}} \uparrow s = (\{\perp\} \cup \uparrow m) \cap \mathcal{P}^*(I) = S_m^1.$$

Hence, η is a surjection. Since ξ_1, ξ_2 are injection, η is a injection too. ■

References

- [1] T.Arase, The ideal quntale of a principal ideal domains and its quantic nuclei, Master's Thesis, Hokkaido university, 1997.
- [2] B.A.Davey and H.A.Priestley, Introduction to Lattices and Order, (Cambridge University Press, Cambridge, 1990)
- [3] K.I.Rosenthal, Quantaes and their applications, (Longman, Edinburgh, 1990).
- [4] K.I.Rosenthal, The theory of quantaloids, (Longman, Edinburgh, 1996).