



Title	Global existence of classical solutions to systems of wave equations with critical nonlinearity in three space dimensions
Author(s)	Yokoyama, K.
Citation	Hokkaido University Preprint Series in Mathematics, 405, 1-25
Issue Date	1998-2-1
DOI	10.14943/83551
Doc URL	http://hdl.handle.net/2115/69155
Type	bulletin (article)
File Information	pre405.pdf



[Instructions for use](#)

Global Existence of Classical Solutions
to Systems of Wave Equations
with Critical Nonlinearity
in Three Space Dimensions

Kazuyoshi Yokoyama

Series #405. February 1998

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #380 F. Hiroshima, Weak coupling limit removing an ultraviolet cut-off for a Hamiltonian of particles interacting with a scalar field, 39 pages. 1997.
- #381 Y. Giga, S. Matsui and Y. Shimizu, On estimates in Hardy spaces for the Stokes flow in a half space, 13 pages. 1997.
- #382 A. Arai, A new estimate for the ground state energy of Schrödinger operators, 12 pages. 1997.
- #383 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the Sobolev space of critical order, 24 pages. 1997.
- #384 K. Ito, Asymptotic stability of planar rarefaction wave for scalar viscous conservation law, 8 pages. 1997.
- #385 A. Arai, Representation-theoretic aspects of two-dimensional quantum systems in singular vector potentials: canonical commutation relations, quantum algebras, and reduction to lattice quantum systems, 32 pages. 1997.
- #386 P. Aviles and Y. Giga, On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields, 21 pages. 1997.
- #387 T. Nakazi and T. Yamamoto, Norms of some singular integral operators and their inverse operators, 28 pages. 1997.
- #388 M.-H. Giga and Y. Giga, Remarks on convergence of evolving graphs by nonlocal curvature, 18 pages. 1997.
- #389 T. Tsukada, Reticular Lagrangian singularities, 41 pages. 1997.
- #390 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the homogeneous Sobolev space, 12 pages. 1997.
- #391 Y. Giga, M. Ohnuma and M.-H. Sato, On strong maximum principle and large time behaviour of generalized mean curvature flow with the Neumann boundary condition, 24 pages. 1997.
- #392 T. Tsujishita and H. Watanabe, Monoidal closedness of the category of simulations, 24 pages. 1997.
- #393 T. Arase, A remark on the quantale structure of multisets, 10 pages. 1997.
- #394 N. H. Bingham and A. Inoue, Extension of the Drasin-Shea-Jordan theorem, 16 pages. 1997.
- #395 N. H. Bingham and A. Inoue, Ratio Mercerian theorems with applications to Hankel and Fourier transforms, 30 pages. 1997.
- #396 Y. Nishiura and D. Ueyama, A skeleton structure of self-replicating dynamics, 27 pages. 1997.
- #397 K. Hirata and K. Sugano, On semisimple extensions of serial rings, 6 pages. 1997.
- #398 D. Pei, Singularities of $\mathbb{R}P^2$ -valued Gauss maps of surfaces in Minkowski 3-space, 15 pages. 1997.
- #399 T. Mikami, Markov marginal problems and their applications to Markov optimal control, 28 pages. 1997.
- #400 M. Tsujii, A simple proof for monotonicity of entropy in the quadratic family, 8 pages. 1998.
- #401 M. Nakamura and T. Ozawa, Global solutions in the critical Sobolev space for the wave equations with nonlinearity of exponential growth, 9 pages. 1998.
- #402 D. Lehmann and T. Suwa, Generalization of variations and Baum-Bott residues for holomorphic foliations on singular varieties, 19 pages. 1998.
- #403 T. Nakazi and K. Okubo, ρ -contraction and 2×2 matrix, 6 pages. 1998.
- #404 Y. Kohsaka, Free boundary problem for quasilinear parabolic equation with fixed angle of contact to a boundary, 28 pages. 1998.

Global Existence of Classical Solutions to Systems of Wave Equations with Critical Nonlinearity in Three Space Dimensions

Kazuyoshi Yokoyama

Department of Mathematics, Hokkaido University,
Sapporo 060, Japan

Abstract

We discuss the existence of a global small solution to the Cauchy problem for a system of quasilinear wave equations in three space dimensions, when its nonlinear term have a critical exponent. Global existence is established on the null condition which is extended to the condition for systems of wave equations with different propagation speeds.

1 Introduction.

We consider the Cauchy problem for a system of quasilinear wave equations

$$\square_i u^i = F^i(\partial u, \partial^2 u) \quad \text{in } [0, \infty) \times \mathbf{R}^3, \quad (1.1)$$

$$u^i(0, \cdot) = \varepsilon f^i, \quad \partial_i u^i(0, \cdot) = \varepsilon g^i \quad \text{in } \mathbf{R}^3, \quad i = 1, \dots, m, \quad (1.2)$$

where $\square_i = \partial_0^2 - c_i^2 \sum_{j=1}^3 \partial_j^2$, $c_i > 0$, $\partial_\alpha = \partial/\partial x^\alpha$, $t = x^0$, $x = (x^1, x^2, x^3)$ and $u^i = u^i(t, x)$ are real-valued unknown functions. We denote by ∂ the space-time derivatives, i.e.

$$\partial u = (\partial_\alpha u^i)_{\alpha,i}, \partial^2 u = (\partial_\alpha \partial_\beta u^i)_{\alpha,\beta,i},$$

where α, β range over $0, 1, 2, 3$ and i over $1, \dots, m$. f^i and g^i belong to $C_0^\infty(\mathbf{R}^3)$ and ε is a positive small parameter. We also assume that each $F^i(\partial u, \partial^2 u)$ takes the form

$$F^i(\partial u, \partial^2 u) = \sum_{j=1}^m \sum_{\alpha,\beta=0}^3 C_{\alpha\beta}^{ij}(\partial u) \partial_\alpha \partial_\beta u^j + D^i(\partial u), \quad (1.3)$$

$$C_{\alpha\beta}^{ij}(\partial u) = C_{\beta\alpha}^{ij}(\partial u), \quad (1.4)$$

$$C_{\alpha\beta}^{ij}(0) = 0, D^i(0) = \partial D^i / \partial(\partial_\alpha u^j) = 0, \quad (1.5)$$

where $C_{\alpha\beta}^{ij}, D^i$ are C^∞ -functions near $\partial u = 0$.

We give a condition of global existence for the initial value problem (1.1)-(1.2) in three space dimensions. Small solutions exist globally when $F^i(\partial u, \partial^2 u)$ do not have quadratic parts, but in general we cannot expect global solutions when they have quadratic parts even if ε is small. However, Klainerman [7] introduced the null condition for the quadratic parts of single wave equations (or systems of wave equations with the same propagation speeds) and proved a global existence theorem on that condition. We extend the Klainerman's null condition to the case where the propagation speeds are different. In the two spatial dimensions, it was shown in [1] that the null condition for the systems with different propagation speeds could be derived by applying John-Shatah observation provided $D^i(\partial u) = 0$. We can prove the following null condition by applying similar argument to our case. That is,

$$\sum_{\alpha,\beta,\gamma=0}^3 C_{\alpha\beta\gamma}^{iii} X_\alpha^i X_\beta^i X_\gamma^i = 0, \sum_{\alpha,\beta=0}^3 D_{\alpha\beta}^{iii} X_\alpha^i X_\beta^i = 0 \quad (i = 1, \dots, m)$$

for all real vector $X^i = (X_0^i, X_1^i, X_2^i, X_3^i)$ satisfying

$$(X_0^i)^2 - c_i^2 \sum_{j=1}^3 (X_j^i)^2 = 0. \quad (1.6)$$

Here we have set

$$C_{\alpha\beta\gamma}^{ijk} = \frac{\partial C_{\alpha\beta}^{ij}}{\partial(\partial_\gamma u^k)}(0), D_{\alpha\beta}^{ijk} = \frac{\partial^2 D^i}{\partial(\partial_\alpha u^j) \partial(\partial_\beta u^k)}(0).$$

The aim of this paper is to prove a global existence theorem on the null condition (1.6).

Theorem 1.1 *Let c_i to be different from each other. Assume that the nonlinear terms $F^i(\partial u, \partial^2 u)$ given by (1.3)-(1.5) satisfy the null condition (1.6) and*

$$C_{\alpha\beta}^{ij}(\partial u) = C_{\alpha\beta}^{ji}(\partial u). \quad (1.7)$$

Then there exists a positive constant ε_0 such that the initial value problem (1.1)-(1.2) has a unique C^∞ -solution in $[0, \infty) \times \mathbf{R}^3$ for ε with $0 \leq \varepsilon < \varepsilon_0$.

In the two spatial dimensions, Hoshiga and Kubo have proved in [2] a corresponding global existence theorem. They obtained necessary estimates with the use of only the angular derivatives and the scaling operator to avoid the difficulty coming from the difference of speeds. We improve their method to prove our theorem. Comparing the two spatial dimensions, the weighted L^∞ -estimates involve ‘log t ’ (Proposition 3.1), and it is shown that the null condition is used to take away ‘log t ’ from our estimates (section 3.2). Further, by virtue of the null condition, we can prove L^2 -boundedness of the derivatives of the solution (section 4), which leads to the global existence.

2 Notation.

Set

$$\partial_\alpha = \partial / \partial x^\alpha, \quad x^0 = t,$$

$$\partial = (\partial_0, \partial_1, \partial_2, \partial_3),$$

$$\nabla = (\partial_1, \partial_2, \partial_3),$$

$$r = |x| \quad \text{for } x \in \mathbf{R}^3.$$

We denote by $\Gamma = (\Gamma_0, \dots, \Gamma_7)$ the collection of differential operators ∂, Ω, S where

$$\Omega = x \wedge \nabla, \quad (2.1)$$

$$\Gamma_7 = S = t\partial_t + r\partial_r, \quad (2.2)$$

$$\partial_r = \frac{x}{r} \cdot \nabla. \quad (2.3)$$

Then we find that the bracket $[\Gamma_\alpha, \Gamma_\beta]$ of any Γ_α and Γ_β is written by another Γ_γ . Moreover, we have

$$[\Gamma_\alpha, \square_i] = 0 \text{ for } 0 \leq \alpha \leq 6 \text{ and } [\Gamma_7, \square_i] = -2\square_i. \quad (2.4)$$

We also note that

$$\nabla = \frac{x}{r} \partial_r - \frac{x}{r^2} \wedge \Omega. \quad (2.5)$$

For $a = (a_1, \dots, a_k)$ ($a_i \in \{0, \dots, 7\}, 1 \leq i \leq k$) we define

$$\Gamma^a = \Gamma_{a_1} \cdots \Gamma_{a_k} \text{ and } |a| = k. \quad (2.6)$$

Let $u = {}^t(u^1, \dots, u^m)$ be a vector and set

$$w_i(t, r) = (1+r)(1+|c_i t - r|) \quad (i = 1, \dots, m). \quad (2.7)$$

Then we define

$$[\partial u]_{k,t} = \sum_{|a| \leq k} \sum_{i=1}^m \sum_{\alpha=0}^3 \sup_{0 \leq s \leq t} \sup_{x \in \mathbf{R}^3} |w_i(s, |x|) \Gamma^a \partial_\alpha u^i(s, x)|, \quad (2.8)$$

$$\|\partial u(t)\|_k = \sum_{|a| \leq k} \sum_{i=1}^m \sum_{\alpha=0}^3 \|\Gamma^a \partial_\alpha u^i(s, \cdot)\|_{L^2(\mathbf{R}^3)}. \quad (2.9)$$

$$\|\partial u\|_{k,t} = \sum_{|a| \leq k} \sum_{i=1}^m \sum_{\alpha=0}^3 \sup_{0 \leq s \leq t} \|\Gamma^a \partial_\alpha u^i(s, \cdot)\|_{L^2(\mathbf{R}^3)}. \quad (2.10)$$

3 Weighted L^∞ -estimates.

3.1 Estimates for solutions of scalar wave equations.

Let $v = v(t, x)$ be the smooth solution of the Cauchy problem

$$\partial_t^2 v - c_0^2 \Delta v = F \quad \text{in } [0, T] \times \mathbf{R}^3, \quad (3.1)$$

$$v(0, \cdot) = \partial_t v(0, \cdot) = 0 \quad \text{in } \mathbf{R}^3, \quad (3.2)$$

where $F \in C^\infty([0, T] \times \mathbf{R}^3)$ and $F(t, \cdot) \in C_0^\infty(\mathbf{R}^3)$ for each t . In this subsection we present the decay estimates for v that we will use later on.

Proposition 3.1 *Let v be the solution of (3.1)-(3.2). For $1 \leq \mu$, $0 < \nu$ and $0 \leq c$, we set*

$$z_{\mu,\nu}(s, \lambda) = (1 + |cs - \lambda|)^\mu (1 + s + \lambda)^\nu,$$

$$\Phi_\theta(t) = \begin{cases} \log(2 + t) & (\theta = 0) \\ 1 & (\theta > 0), \end{cases} \quad (3.3)$$

$$M_{\mu,\nu;k}(F) = \sum_{|a| \leq k} \sup_{0 \leq s \leq t} \sup_{y \in \mathbb{R}^3} |y| z_{\mu,\nu}(s, |y|) |\Gamma^a F(s, y)|. \quad (3.4)$$

Then we have

$$(i) \quad |v(t, x)| \leq C(1 + t + |x|)^{-1} \Phi_{\mu-1}(t) \Phi_{\nu-1}(t) M_{\mu,\nu;0}(F) \quad (3.5)$$

for $1 \leq \mu, 1 \leq \nu$,

$$(ii) \quad |\partial v(t, x)| \leq C(1 + |x|)^{-1} (1 + |c_0 t - |x||)^{-\nu} \Phi_{\mu-1}(t) M_{\mu,\nu;1}(F) \quad (3.6)$$

for $1 \leq \mu, 0 < \nu, c \neq c_0$, and

$$(iii) \quad |\partial v(t, x)| \leq C(1 + |x|)^{-1} \{ (1 + |c_0 t - |x||)^{-\nu} \Phi_{\mu-1}(t) \\ + (1 + |c_0 t - |x||)^{-\mu} \Phi_{\nu-1}(t) \} M_{\mu,\nu;1}(F) \quad (3.7)$$

for $1 \leq \mu, 1 \leq \nu, c = c_0$.

Proof. By a change of coordinates, the proof can be reduced to the case where $c_0 = 1$. So we let $c_0 = 1$. Set $|x| = r$. In appendix in [3], F. John showed that the solution of (3.1)-(3.2) could be expressed in the form

$$v(t, x) = (4\pi r)^{-1} \int_0^t ds \int_{|r-t+s|}^{r+t-s} \lambda d\lambda \int_0^{2\pi} F(s, \lambda \Theta) d\varphi, \quad (3.8)$$

where

$$\Theta = \Theta(s, \lambda, \varphi) = R(\sin \psi \cos \varphi, \sin \psi \sin \varphi, \cos \psi),$$

$$R \text{ is an orthogonal transformation with } R(0, 0, r) = x, \quad (3.9)$$

$$\cos \psi = (2r\lambda)^{-1} (r^2 + \lambda^2 - (t-s)^2), \quad \sin \psi = (1 - \cos^2 \psi)^{1/2}.$$

Thus, v can be written as

$$v(t, x) = (4\pi r)^{-1} \int_D \lambda d\lambda ds \int_0^{2\pi} F(s, \lambda \Theta) d\varphi, \quad (3.10)$$

where

$$\begin{aligned} D &= \{(s, \lambda) \mid 0 < s < t, \lambda_1 < \lambda < \lambda_2\}, \\ \lambda_1 &= |r - t + s|, \lambda_2 = r + t - s. \end{aligned} \tag{3.11}$$

We first prove (3.5). By (3.10),

$$|v(t, x)| \leq CI_0 M_{\mu, \nu; 0}(F)$$

where

$$I_0 = r^{-1} \int_D z_{\mu, \nu}(s, \lambda)^{-1} d\lambda ds.$$

We obtain (3.5) from

$$I_0 \leq C(1 + t + r)^{-1} \Phi_{\mu-1}(t) \Phi_{\nu-1}(t), \tag{3.12}$$

which we will prove now. In order to prove (3.12), four cases

1. $r \leq 1$,
2. $1 \leq r$ and $(2 + c)t \leq r$,
3. $1 \leq r$ and $2^{-1}t \leq r \leq (2 + c)t$,
4. $1 \leq r$ and $r \leq 2^{-1}t$

are considered separately.

In the first case, we have

$$\lambda_2 - \lambda_1 \leq 2r (\leq 2),$$

and hence

$$\begin{aligned} I_0 &\leq Cr^{-1} \int_0^t ds \int_{\lambda_1}^{\lambda_2} z_{\mu, \nu}(s, \lambda)^{-1} d\lambda \\ &\leq C \int_0^t (1 + |cs - \lambda_2|)^{-\mu} (1 + s + \lambda_2)^{-\nu} ds \\ &\leq C(1 + t + r)^{-\nu} \Phi_{\mu-1}(t). \end{aligned}$$

Therefore, (3.12) is proved for the case 1.

In the second case, the inequalities

$$\lambda - cs \geq \min\{r - t, r - ct\} \geq (2 + c)^{-1}(1 + \min\{1, c\})r \quad (3.13)$$

hold for $(s, \lambda) \in D$. Hence it follows from

$$z_{\mu, \nu}(s, \lambda)^{-1} \leq C(1 + r)^{-\mu - \nu} \quad (3.14)$$

that

$$\begin{aligned} I_0 &\leq Cr^{-1} \int_D (1 + r)^{-\mu - \nu} d\lambda ds \\ &\leq C(1 + t + r)^{1 - \mu - \nu}. \end{aligned}$$

Consequently, we have (3.12) for the case 2.

In the third and the fourth case, we introduce the new variables of integration

$$\begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ -c & 1 \end{pmatrix} \begin{pmatrix} s \\ \lambda \end{pmatrix}. \quad (3.15)$$

Set

$$\alpha_0 = 2^{-1}\{(1 - c)\alpha + (1 + c)(r - t)\}. \quad (3.16)$$

If $|t - r| < \alpha < t + r$, then $-\alpha < \alpha_0 < \alpha$. Hence,

$$\begin{aligned} &\int_D z_{\mu, \nu}(s, \lambda)^{-1} d\lambda ds \\ &\leq \int_D (1 + |cs - \lambda|)^{-\mu} (1 + s + \lambda)^{-\nu} d\lambda ds \\ &\leq \int_{|r-t|}^{r+t} (1 + \alpha)^{-\nu} d\alpha \int_{\alpha_0}^{\alpha} (1 + |\beta|)^{-\mu} d\beta \\ &\leq \int_{|r-t|}^{r+t} (1 + \alpha)^{-\nu} \Phi_{\mu-1}(\alpha) d\alpha. \end{aligned}$$

Therefore in the third case, it follows that

$$\begin{aligned} I_0 &\leq C(1 + r)^{-1} \Phi_{\mu-1}(t) \int_{|r-t|}^{r+t} (1 + \alpha)^{-\nu} d\alpha \\ &\leq C(1 + t + r)^{-1} \Phi_{\mu-1}(t) \Phi_{\nu-1}(t), \end{aligned}$$

and in the fourth case, it follows that

$$\begin{aligned} I_0 &\leq Cr^{-1}(1 + t - r)^{-\nu} \Phi_{\mu-1}(t) \int_{t-r}^{t+r} d\alpha \\ &\leq C(1 + t + r)^{-\nu} \Phi_{\mu-1}(t). \end{aligned}$$

Hence we have proved (3.12) in all cases.

We next prove (3.6) and (3.7). In order to prove (3.6) and (3.7), we give representation formulae for $\partial_\alpha v$.

Let $(s, \lambda) \in D$ and $0 \leq \varphi \leq 2\pi$. By (2.5),

$$(\nabla F)(s, \lambda\Theta) = \Theta(\partial_r F)(s, \lambda\Theta) - \lambda^{-1}\Theta \wedge (\Omega F)(s, \lambda\Theta). \quad (3.17)$$

Since $\partial_\lambda \Theta \cdot \Theta = 0$, it follows that

$$\begin{aligned} & (\partial_r F)(s, \lambda\Theta) \\ &= \partial_\lambda \{F(s, \lambda\Theta)\} - \lambda \partial_\lambda \Theta \cdot (\nabla F)(s, \lambda\Theta) \\ &= \partial_\lambda \{F(s, \lambda\Theta)\} - \lambda \partial_\lambda \Theta \cdot \{\Theta(\partial_r F)(s, \lambda\Theta) - \lambda^{-1}\Theta \wedge (\Omega F)(s, \lambda\Theta)\} \\ &= \partial_\lambda \{F(s, \lambda\Theta)\} + \partial_\lambda \Theta \cdot (\Theta \wedge (\Omega F)(s, \lambda\Theta)). \end{aligned} \quad (3.18)$$

In a similar manner,

$$(\partial_t F)(s, \lambda\Theta) = \partial_s \{F(s, \lambda\Theta)\} + \partial_s \Theta \cdot (\Theta \wedge (\Omega F)(s, \lambda\Theta)). \quad (3.19)$$

If we substitute (3.18) into (3.17), we have

$$\begin{aligned} (\nabla F)(s, \lambda\Theta) &= \Theta \partial_\lambda \{F(s, \lambda\Theta)\} + \Theta \{\partial_\lambda \Theta \cdot (\Theta \wedge (\Omega F)(s, \lambda\Theta))\} \\ &\quad - \lambda^{-1}\Theta \wedge (\Omega F)(s, \lambda\Theta). \end{aligned} \quad (3.20)$$

We use these expressions in

$$\partial_\alpha v(t, x) = (4\pi r)^{-1} \int_D \lambda d\lambda ds \int_0^{2\pi} (\partial_\alpha F)(s, \lambda\Theta) d\varphi,$$

which are obtained from (3.10). We split the domain of integration D into D_1 and D_2 , where

$$\begin{aligned} D_1 &= \{(s, \lambda) \in D \mid \lambda_1 < \lambda < \lambda_1 + \delta \text{ or } \lambda_2 - \delta < \lambda < \lambda_2\}, \\ D_2 &= D \setminus D_1, \\ \delta &= \min\{1, r\}. \end{aligned} \quad (3.21)$$

Using (3.19),(3.20) and integrating by parts on the domain D_2 , we obtain the following representation formulae.

$$\begin{aligned}
& 4\pi r \partial_t v(t, x) \\
&= \int_{D_1} \lambda d\lambda ds \int_0^{2\pi} (\partial_t F)(s, \lambda\Theta) d\varphi \\
&\quad + \int_{\partial D_2} n_s d\sigma \int_0^{2\pi} \lambda F(s, \lambda\Theta) d\varphi \\
&\quad + \int_{D_2} \lambda d\lambda ds \int_0^{2\pi} \partial_s \Theta \cdot (\Theta \wedge (\Omega F)(s, \lambda\Theta)) d\varphi
\end{aligned} \tag{3.22}$$

$$\begin{aligned}
& 4\pi r \nabla v(t, x) \\
&= \int_{D_1} \lambda d\lambda ds \int_0^{2\pi} (\nabla F)(s, \lambda\Theta) d\varphi \\
&\quad + \int_{\partial D_2} n_\lambda d\sigma \int_0^{2\pi} \lambda \Theta F(s, \lambda\Theta) d\varphi \\
&\quad - \int_{D_2} d\lambda ds \int_0^{2\pi} \{\Theta F(s, \lambda\Theta) + \Theta \wedge (\Omega F)(s, \lambda\Theta)\} d\varphi \\
&\quad + \int_{D_2} \lambda d\lambda ds \int_0^{2\pi} \{-\partial_\lambda \Theta F(s, \lambda\Theta) \\
&\quad\quad + \Theta \{\partial_\lambda \Theta \cdot (\Theta \wedge (\Omega F)(s, \lambda\Theta))\}\} d\varphi
\end{aligned} \tag{3.23}$$

Here, (n_s, n_λ) is the unit outer normal vector field on ∂D_2 , and $d\sigma$ is the line element on ∂D_2 .

Note that $D_2 = \emptyset$ if $r < 1$. Therefore it follows from (3.22) and (3.23) that

$$|\partial v(t, x)| \leq C \sum_{i=1}^4 I_i M_{\mu, \nu; 1} \tag{3.24}$$

where

$$\begin{aligned}
I_1 &= r^{-1} \int_{D_1} z_{\mu, \nu}(s, \lambda)^{-1} d\lambda ds, \\
I_2 &= (1+r)^{-1} \int_{\partial D_2} z_{\mu, \nu}(s, \lambda)^{-1} d\sigma, \\
I_3 &= (1+r)^{-1} \int_{D_2} (1+\lambda)^{-1} z_{\mu, \nu}(s, \lambda)^{-1} d\lambda ds, \\
I_4 &= (1+r)^{-1} \int_{D_2} \sup_{0 \leq \varphi \leq 2\pi} |\partial \Theta| z_{\mu, \nu}(s, \lambda)^{-1} d\lambda ds.
\end{aligned}$$

By the definition of the domain D_1 , we can easily see

$$I_1 \leq C I_2. \tag{3.25}$$

Concerning I_2 , I_3 and I_4 , we will prove

$$I_2 \leq C(1+r)^{-1}(1+|t-r|)^{-\nu}\Phi_{\mu-1}(t) \quad (3.26)$$

for $1 \leq \mu, 0 < \nu, c \neq 1$,

$$I_2 \leq C(1+r)^{-1}\{(1+|t-r|)^{-\nu}\Phi_{\mu-1}(t) + (1+|t-r|)^{-\mu}\Phi_{\nu-1}(t)\} \quad (3.27)$$

for $1 \leq \mu, 1 \leq \nu, c = 1$,

$$I_3 \leq C(1+r)^{-1}(1+|t-r|)^{-\nu}\Phi_{\mu-1}(t) \quad (3.28)$$

for $1 \leq \mu, 0 < \nu$, and

$$I_4 \leq C(1+r)^{-1}(1+|t-r|)^{-\nu}\Phi_{\mu-1}(t) \quad (3.29)$$

for $1 \leq \mu, 0 < \nu, c \neq 1$ or $1 \leq \mu, 1/2 < \nu, c = 1$. If we have proved (3.26)-(3.29), then (3.6) and (3.7) are obtained through (3.24), (3.25) and these estimates.

(a) Proof of (3.26) and (3.27).

If $r \geq (2+c)t$, (3.26) and (3.27) are easily obtained from (3.13). So we let $r \leq (2+c)t$.

By the definition of I_2 , we have

$$(1+r)I_2 \leq C(I_2' + I_2'' + I_2'''),$$

where

$$\begin{aligned} I_2' &= \int_0^t z_{\mu,\nu}(s, \lambda_1)^{-1} ds, \\ I_2'' &= \int_0^t z_{\mu,\nu}(s, \lambda_2)^{-1} ds, \\ I_2''' &= \int_{|t-r|}^{t+r} z_{\mu,\nu}(0, \lambda)^{-1} d\lambda. \end{aligned}$$

If $c \neq 1$, it follows that

$$\begin{aligned} I_2' &\leq \int_0^t (1+|cs - \lambda_1|)^{-\mu}(1+s+\lambda_1)^{-\nu} ds \\ &\leq C(1+|t-r|)^{-\nu}\Phi_{\mu-1}(t), \end{aligned}$$

because $s + \lambda_1 \geq |t - r|$. However, if $c = 1$,

$$\begin{aligned}
I_2' &\leq \int_0^t (1 + |s - \lambda_1|)^{-\mu} (1 + s + \lambda_1)^{-\nu} ds \\
&= \int_0^{(t-r)_+} (1 + |s - \lambda_1|)^{-\mu} (1 + t - r)^{-\nu} ds \\
&\quad + \int_{(t-r)_+}^t (1 + |t - r|)^{-\mu} (1 + s + \lambda_1)^{-\nu} ds \\
&\leq C(1 + |t - r|)^{-\nu} \Phi_{\mu-1}(t) + C(1 + |t - r|)^{-\mu} \Phi_{\nu-1}(t).
\end{aligned}$$

As for I_2'' and I_2''' , we obtain, straightforwardly,

$$I_2'' \leq C(1 + t + r)^{-\nu} \Phi_{\mu-1}(t),$$

$$I_2''' \leq C(1 + |t - r|)^{1-\mu-\nu}$$

for $1 \leq \mu$ and $0 < \nu$. Thus we have proved (3.26) and (3.27).

(b) Proof of (3.28).

In case $r \geq (2 + c)t$, (3.28) results from (3.13).

Let $r \leq (2 + c)t$. If $c = 0$, by the change of variables (3.15),

$$\begin{aligned}
(1 + r)I_3 &\leq \int_{|t-r|}^{t+r} (1 + \alpha)^{-\nu} d\alpha \int_{\alpha_0}^{\alpha} (1 + \beta)^{-1-\mu} d\beta \\
&\leq C \int_{|t-r|}^{t+r} (1 + |t - r|)^{-\nu} (1 + \alpha_0)^{-\mu} d\alpha \\
&\leq C(1 + |t - r|)^{-\nu} \Phi_{\mu-1}(t).
\end{aligned}$$

If $c > 0$, set

$$\begin{aligned}
\Lambda_0 &= \{(s, \lambda) | 0 \leq \lambda \leq 2^{-1}cs\}, \\
\Lambda_1 &= \{(s, \lambda) | 0 \leq 2^{-1}cs \leq \lambda\}.
\end{aligned} \tag{3.30}$$

Since

$$(s, \lambda) \in \Lambda_0 \Rightarrow (1 + \lambda)^{-1} z_{\mu, \nu}(s, \lambda)^{-1} \leq C(1 + \lambda)^{-1} (1 + s + \lambda)^{-\mu-\nu}, \tag{3.31}$$

$$(s, \lambda) \in \Lambda_1 \Rightarrow (1 + \lambda)^{-1} z_{\mu, \nu}(s, \lambda)^{-1} \leq C(1 + |cs - \lambda|)^{-\mu} (1 + s + \lambda)^{-1-\nu}, \tag{3.32}$$

the inequality

$$(1 + \lambda)^{-1} z_{\mu, \nu}(s, \lambda)^{-1} \leq C \{(1 + \lambda)^{-\mu} + (1 + |cs - \lambda|)^{-\mu}\} (1 + s + \lambda)^{-1 - \nu} \quad (3.33)$$

holds. Hence, adapting (3.15) for each term of (3.33), we obtain (3.28).

(c) Proof of (3.29).

We first need estimates of $|\partial\Theta|$. By the definition (3.9) of Θ , we have

$$\begin{aligned} |\partial_\lambda \Theta| &= |\lambda^2 + \bar{\lambda}_1 \lambda_2| \lambda^{-1} \{(\lambda^2 - \lambda_1^2)(\lambda_2^2 - \lambda^2)\}^{-1/2}, \\ |\partial_s \Theta| &= (\bar{\lambda}_1 + \lambda_2) \{(\lambda^2 - \lambda_1^2)(\lambda_2^2 - \lambda^2)\}^{-1/2}, \end{aligned}$$

where $\bar{\lambda}_1 = t - s - r$. Noting

$$\bar{\lambda}_1 = \begin{cases} -\lambda_1 & \text{for } (t - r)_+ < s < t \\ \lambda_1 & \text{for } 0 < s < t - r, \end{cases}$$

it follows from

$$\begin{aligned} \lambda^2 + \bar{\lambda}_1 \lambda_2 &= \lambda(\lambda + \bar{\lambda}_1) + \bar{\lambda}_1(\lambda_2 - \lambda), \\ \bar{\lambda}_1 + \lambda_2 &= (\lambda + \bar{\lambda}_1) + (\lambda_2 - \lambda) \end{aligned}$$

that

$$(t - r)_+ < s < t \Rightarrow |\partial\Theta| \leq (\lambda^2 - \lambda_1^2)^{-1/2} + (\lambda_2^2 - \lambda^2)^{-1/2}, \quad (3.34)$$

$$0 < s < t - r \Rightarrow |\partial\Theta| \leq (\lambda^2 - \lambda_1^2)^{-1/2} + \{(\lambda - \lambda_1)(\lambda_2 - \lambda)\}^{-1/2}. \quad (3.35)$$

As before, the case $r \geq (2 + c)t$ is easy. So we consider $r \leq (2 + c)t$. Set

$$\begin{aligned} D_2^{(1)} &= \{(s, \lambda) \in D_2 \mid (t - r)_+ < s < t\}, \\ D_2^{(2)} &= \{(s, \lambda) \in D_2 \mid 0 < s < t - r\}. \end{aligned} \quad (3.36)$$

Then from (3.34) and (3.35),

$$\begin{aligned} (s, \lambda) \in D_2^{(1)} &\Rightarrow |\partial\Theta| \leq C \{(1 + \lambda - \lambda_1)^{-1/2} + (1 + \lambda_2 - \lambda)^{-1/2}\} (1 + \lambda)^{-1/2}, \\ (s, \lambda) \in D_2^{(2)} &\Rightarrow |\partial\Theta| \leq C \{(1 + \lambda)^{-1/2} + (1 + \lambda_2 - \lambda)^{-1/2}\} (1 + \lambda - \lambda_1)^{-1/2}. \end{aligned}$$

Hence, noting (3.31) and (3.32), we obtain the following estimates:

$$(s, \lambda) \in D_2^{(i)} \Rightarrow |\partial\Theta|z_{\mu,\nu}(s, \lambda)^{-1} \leq C\{p_{\mu,\nu}^{(i)}(s, \lambda) + q_{\mu,\nu}^{(i)}(s, \lambda)\}, \quad i = 1, 2 \quad (3.37)$$

where

$$p_{\mu,\nu}^{(i)}(s, \lambda) = \xi_{\mu,\nu}^{(i),1}(s, \lambda) + \xi_{\mu,\nu}^{(i),2}(s, \lambda), \quad i = 1, 2, \quad (3.38)$$

$$q_{\mu,\nu}^{(i)}(s, \lambda) = \eta_{\mu,\nu}^{(i),1}(s, \lambda) + \eta_{\mu,\nu}^{(i),2}(s, \lambda), \quad i = 1, 2, \quad (3.39)$$

$$\xi_{\mu,\nu}^{(1),1}(s, \lambda) = (1 + \lambda - \lambda_1)^{-1/2}(1 + \lambda)^{-1/2-\mu}(1 + s + \lambda)^{-\nu},$$

$$\xi_{\mu,\nu}^{(1),2}(s, \lambda) = (1 + \lambda_2 - \lambda)^{-1/2}(1 + \lambda)^{-1/2-\mu}(1 + s + \lambda)^{-\nu},$$

$$\eta_{\mu,\nu}^{(1),1}(s, \lambda) = (1 + \lambda - \lambda_1)^{-1/2}(1 + |cs - \lambda|)^{-\mu}(1 + s + \lambda)^{-1/2-\nu},$$

$$\eta_{\mu,\nu}^{(1),2}(s, \lambda) = (1 + \lambda_2 - \lambda)^{-1/2}(1 + |cs - \lambda|)^{-\mu}(1 + s + \lambda)^{-1/2-\nu},$$

$$\xi_{\mu,\nu}^{(2),1}(s, \lambda) = (1 + \lambda)^{-1/2-\mu}(1 + \lambda - \lambda_1)^{-1/2}(1 + s + \lambda)^{-\nu},$$

$$\xi_{\mu,\nu}^{(2),2}(s, \lambda) = (1 + \lambda_2 - \lambda)^{-1/2}(1 + \lambda - \lambda_1)^{-1/2}(1 + \lambda)^{-\mu}(1 + s + \lambda)^{-\nu},$$

$$\eta_{\mu,\nu}^{(2),1}(s, \lambda) = \eta_{\mu,\nu}^{(1),1}(s, \lambda),$$

$$\eta_{\mu,\nu}^{(2),2}(s, \lambda) = (1 + \lambda_2 - \lambda)^{-1/2}(1 + \lambda - \lambda_1)^{-1/2}(1 + |cs - \lambda|)^{-\mu}(1 + s + \lambda)^{-\nu}.$$

We change the variables of integration by (3.15). Here, we let $c = 0$ to adapt (3.15) for $p_{\mu,\nu}^{(1)}(s, \lambda)$ and $p_{\mu,\nu}^{(2)}(s, \lambda)$. Then we can prove

$$\int_{D_2^{(i)}} \xi_{\mu,\nu}^{(i),j}(s, \lambda) d\lambda ds \leq C(1 + |t - r|)^{-\nu} \Phi_{\mu-1}(t) \quad (3.40)$$

for $1 \leq \mu, 0 < \nu$,

$$\int_{D_2^{(i)}} \eta_{\mu,\nu}^{(i),j}(s, \lambda) d\lambda ds \leq C(1 + |t - r|)^{-\nu} \Phi_{\mu-1}(t) \quad (3.41)$$

for $1 \leq \mu, 0 < \nu, (i, j) \neq (1, 1)$, and

$$\int_{D_2^{(1)}} \eta_{\mu,\nu}^{(1),1}(s, \lambda) d\lambda ds \leq C(1 + |t - r|)^{-\nu} \Phi_{\mu-1}(t) \quad (3.42)$$

for $1 \leq \mu, 0 < \nu, c \neq 1$ or $1 \leq \mu, 1/2 < \nu, c = 1$. We show the estimate of the integral of $\xi_{\mu,\nu}^{(1),1}(s, \lambda)$ and $\eta_{\mu,\nu}^{(1),1}(s, \lambda)$ here. The others are easy to treat.

We first prove

$$\begin{aligned} & \int_{\alpha_0}^{\alpha} (1 + \beta - \alpha_0)^{-1/2} (1 + |\beta|)^{-\tau} d\beta \\ & \leq C \{(1 + |\alpha_0|)^{1/2-\tau} + \chi(\alpha_0)(1 + |\alpha_0|)^{-1/2} \Phi_{\tau-1}(\alpha)\}, \end{aligned} \quad (3.43)$$

for $\tau \geq 1$ and $|t - r| < \alpha < t + r$, where χ is the characteristic function of the interval $(-\infty, 0)$. Let $(a, b) \subset (\alpha_0, \alpha)$ to be the interval where $1 + \beta - \alpha_0 < 1 + |\beta|$. If $\alpha_0 \geq 0$, then $(a, b) = (\alpha_0, \alpha)$, and if $\alpha_0 < 0$, then $(a, b) = (\alpha_0, \alpha_0/2)$. Integrating by parts,

$$\begin{aligned} & \int_a^b (1 + \beta - \alpha_0)^{-1/2} (1 + |\beta|)^{-\tau} d\beta \\ & = \int_a^b \partial_{\beta} \{2(1 + \beta - \alpha_0)^{1/2}\} (1 + |\beta|)^{-\tau} d\beta \\ & \leq 2(1 + |b|)^{1/2-\tau} + 2\tau \int_a^b (1 + |\beta|)^{-1/2-\tau} d\beta \\ & \leq C \{(1 + |a|)^{1/2-\tau} + (1 + |b|)^{1/2-\tau}\} \\ & \leq C(1 + |\alpha_0|)^{1/2-\tau}. \end{aligned}$$

Hence,

$$\begin{aligned} & \int_{\alpha_0}^{\alpha} (1 + \beta - \alpha_0)^{-1/2} (1 + |\beta|)^{-\tau} d\beta \\ & = \int_a^b + \int_b^{\alpha} \\ & \leq C(1 + |\alpha_0|)^{1/2-\tau} + C\chi(\alpha_0) \int_b^{\alpha} (1 + |\alpha_0|)^{-1/2} (1 + |\beta|)^{-\tau} d\beta, \end{aligned}$$

and we obtain (3.43).

Since $\lambda - \lambda_1 = 2(1 + c)^{-1}(\beta - \alpha_0)$ in $D_2^{(1)}$, it follows from (3.43) that

$$\begin{aligned} & \int_{D_2^{(1)}} \xi_{\mu, \nu}^{(1), 1}(s, \lambda) d\lambda ds \\ & \leq C \int_{|t-r|}^{t+r} (1 + \alpha)^{-\nu} d\alpha \int_{\alpha_0}^{\alpha} (1 + \beta - \alpha_0)^{-1/2} (1 + \beta)^{-1/2-\mu} d\beta \\ & \leq C \int_{|t-r|}^{t+r} (1 + \alpha)^{-\nu} (1 + \alpha_0)^{-\mu} d\alpha \\ & \leq C(1 + |t - r|)^{-\nu} \Phi_{\mu-1}(t), \end{aligned}$$

which prove (3.40) for $i = j = 1$. Concerning $\eta_{\mu,\nu}^{(1),1}(s, \lambda)$, we have

$$\begin{aligned} & \int_{D_2^{(1)}} \eta_{\mu,\nu}^{(1),1}(s, \lambda) d\lambda ds \\ & \leq C \int_{|t-r|}^{t+r} (1+\alpha)^{-1/2-\nu} d\alpha \int_{\alpha_0}^{\alpha} (1+\beta-\alpha_0)^{-1/2} (1+|\beta|)^{-\mu} d\beta \\ & \leq C \int_{|t-r|}^{t+r} (1+\alpha)^{-1/2-\nu} (1+|\alpha_0|)^{-1/2} d\alpha \Phi_{\mu-1}(t). \end{aligned} \quad (3.44)$$

If $c = 1$, we note that $\alpha_0 = r - t$. If $c \neq 1$, let $h(\alpha)$ be a primitive of $(1+|\alpha_0|)^{-1/2}$. Integrating by parts, it follows that

$$\begin{aligned} & \int_{|t-r|}^{t+r} (1+\alpha)^{-1/2-\nu} (1+|\alpha_0|)^{-1/2} d\alpha \\ & \leq (1+\alpha)^{-1/2-\nu} |h(\alpha)| \Big|_{\alpha=|t-r|}^{t+r} + (1/2+\nu) \int_{|t-r|}^{t+r} (1+\alpha)^{-3/2} |h(\alpha)| d\alpha \\ & \leq C(1+|t-r|)^{-\nu}, \end{aligned}$$

since $|h(\alpha)| \leq C(1+\alpha)^{1/2}$. Hence we obtain (3.42).

3.2 An estimate for the quasilinear system with the null condition.

Proposition 3.2 *Let $u = (u^1, \dots, u^m)$ be the smooth solution of*

$$\square_i u^i = F^i(\partial u, \partial^2 u) \quad \text{in } [0, T) \times \mathbf{R}^3, \quad (3.45)$$

$$u^i(0, \cdot) = \varepsilon f^i, \quad \partial_t u^i(0, \cdot) = \varepsilon g^i \quad \text{in } \mathbf{R}^3, \quad (3.46)$$

$$i = 1, \dots, m.$$

Assume that F^i satisfy (1.6) and c_i are different from each other. If $\varepsilon < 1$ and $[\partial u]_{[(N+7)/2], t} < 1$, then

$$[\partial u]_{N, t} \leq C_N(\varepsilon + \|\partial u\|_{N+9, t}^6). \quad (3.47)$$

Proof. Let u_0^i be the solutions of the homogeneous equations

$$\square_i u_0^i = 0 \quad \text{in } [0, \infty) \times \mathbf{R}^3, \quad (3.48)$$

$$u_0^i(0, \cdot) = \varepsilon f^i, \quad \partial_t u_0^i(0, \cdot) = \varepsilon g^i \quad \text{in } \mathbf{R}^3, \quad (3.49)$$

$$i = 1, \dots, m.$$

From (2.4), each $\Gamma^a u_0^i$ satisfies (3.48). Hence by the proof of Theorem 1 in [6] we have

$$\begin{aligned} |\Gamma^a u_0^i(t, x)| &\leq C_N \varepsilon (1+r)^{-1} (1 + |c_i t - r|)^{-1} \\ &\leq C_N \varepsilon (1+t+r)^{-1}, \end{aligned} \quad (3.50)$$

$$|\partial \Gamma^a u_0^i(t, x)| \leq C_N \varepsilon (1+r)^{-1} (1 + |c_i t - r|)^{-1}, \quad (3.51)$$

for $|a| \leq N$.

Set

$$u_1 = u - u_0. \quad (3.52)$$

Then, each $\Gamma^a u_1^i$ satisfies the equation of the form

$$\square_i \Gamma^a u_1^i = \sum_{b \leq a} C_{ab} \Gamma^b F^i(\partial u, \partial^2 u) \quad \text{in } [0, T) \times \mathbf{R}^3, \quad (3.53)$$

$$\Gamma^a u_1^i(0, \cdot) = \partial_i \Gamma^a u_1^i(0, \cdot) = 0 \quad \text{in } \mathbf{R}^3. \quad (3.54)$$

We apply Proposition 3.1 to a solution of (3.53)-(3.54) by replacing the weight $z_{\mu, \nu}(s, \lambda)$ with

$$z(s, \lambda) = \min_{1 \leq i \leq m+1} z_{1,1}^{(i)}(s, \lambda),$$

where

$$z_{\mu, \nu}^{(i)}(s, \lambda) = (1 + |c_i s - \lambda|)^\mu (1 + s + \lambda)^\nu, \quad c_{m+1} = 0. \quad (3.55)$$

Then it follows from (3.5)-(3.7) that

$$|\Gamma^a u_1^i(t, x)| \leq C_N (1+t+r)^{-1} \{\log(2+t)\}^2 M_N(F^i), \quad (3.56)$$

$$|\partial \Gamma^a u_1^i(t, x)| \leq C_N (1+r)^{-1} (1 + |c_i t - r|)^{-1} \log(2+t) M_{N+1}(F^i), \quad (3.57)$$

for $|a| \leq N$, where

$$M_k(F^i) = \sum_{|a| \leq k} \sup_{0 \leq s \leq t} \sup_{y \in \mathbf{R}^3} |y| z(s, |y|) |\Gamma^a F^i(\partial u, \partial^2 u)(s, y)|.$$

In order to estimate $M_k(F^i)$, we use the Sobolev inequality (Lemma 4.2 in [8]):

$$|y| |f(y)| \leq C \left\{ \sum_{|a| \leq 2} \|\Omega^a f\|_{L^2(\mathbf{R}^3)} + \sum_{|a| \leq 1} \|\partial_r \Omega^a f\|_{L^2(\mathbf{R}^3)} \right\}. \quad (3.58)$$

If $|b| + |c| \leq k$ and $0 \leq s \leq t$, we have

$$|y|z(s, |y|)|\partial\Gamma^b u^j(s, y)||\partial\Gamma^c u^l(s, y)| \leq C_k[\partial u]_{[k/2], t} \|\partial u\|_{k+2, t},$$

since $z(s, |y|) \leq Cw_i(s, |y|)$ ($i = 1, \dots, m$). Therefore,

$$M_k(F^i) \leq C_k[\partial u]_{[(k+1)/2], t} \|\partial u\|_{k+3, t}, \quad (3.59)$$

provided $|\partial u|, |\partial^2 u| < 1$. Therefore, it follows from (3.50), (3.51), (3.56), (3.57) and (3.59) that

$$|\Gamma^a u^i(t, x)| \leq C_N(1+t+r)^{-1} \{\log(2+t)\}^2 (\varepsilon + [\partial u]_{[(N+1)/2], t} \|\partial u\|_{N+3, t}), \quad (3.60)$$

$$\begin{aligned} |\partial\Gamma^a u^i(t, x)| &\leq C_N(1+r)^{-1}(1+|c_i t - r|)^{-1} \log(2+t) \cdot \\ &\cdot (\varepsilon + [\partial u]_{[(N+2)/2], t} \|\partial u\|_{N+4, t}) \end{aligned} \quad (3.61)$$

for $|a| \leq N$.

Next, we estimate the nonlinear terms by making use of (3.60), (3.61). We separate F^i into three parts:

$$F^i(\partial u, \partial^2 u) = N^i(\partial u, \partial^2 u) + R^i(\partial u, \partial^2 u) + G^i(\partial u, \partial^2 u), \quad (3.62)$$

where

$$N^i(\partial u, \partial^2 u) = \sum_{0 \leq \alpha, \beta, \gamma \leq 3} C_{\alpha\beta\gamma}^{iii} \partial_\gamma u^i \partial_\alpha \partial_\beta u^i + \sum_{0 \leq \alpha, \beta \leq 3} D_{\alpha\beta}^{iii} \partial_\alpha u^i \partial_\beta u^i, \quad (3.63)$$

$$R^i(\partial u, \partial^2 u) = \sum_{(j,k) \neq (i,i)} \left(\sum_{0 \leq \alpha, \beta, \gamma \leq 3} C_{\alpha\beta\gamma}^{ijk} \partial_\gamma u^k \partial_\alpha \partial_\beta u^j + \sum_{0 \leq \alpha, \beta \leq 3} D_{\alpha\beta}^{ijk} \partial_\alpha u^j \partial_\beta u^k \right), \quad (3.64)$$

and $G^i(\partial u, \partial^2 u)$ are higher order terms. Moreover, by the null condition (1.6), $N^i(\partial u, \partial^2 u)$ can be expressed in the form

$$\begin{aligned} N^i(\partial u, \partial^2 u) &= \sum_{0 \leq \alpha \leq 3} T_\alpha^i Q^i(u^i, \partial_\alpha u^i) + \sum_{0 \leq \alpha, \beta, \gamma \leq 3} T_{\alpha\beta\gamma}^i Q_{\alpha\beta}(u^i, \partial_\gamma u^i) \\ &+ \sum_{0 \leq \alpha \leq 3} \bar{T}_\alpha^i \partial_\alpha u^i \square_i u^i + T^i Q^i(u^i, u^i), \end{aligned} \quad (3.65)$$

where

$$Q^i(u, v) = \partial_i u \partial_t v - c_i^2 \nabla u \cdot \nabla v, \quad (3.66)$$

$$Q_{\alpha\beta}(u, v) = \partial_\alpha u \partial_\beta v - \partial_\beta u \partial_\alpha v. \quad (3.67)$$

These forms gain good decay near $c_i t - r = 0$. Indeed, the following estimates hold for $|c_i t - r| < c_i t/2$:

$$\begin{aligned} |Q^i(u, v)| &\leq C|c_i t - r|(1+t+r)^{-1}|\partial u||\partial v| \\ &\quad + C(1+t+r)^{-1}(|\Gamma u||\partial v| + |\partial u||\Gamma v|), \end{aligned} \quad (3.68)$$

$$|Q_{\alpha\beta}(u, v)| \leq C(1+t+r)^{-1}(|\partial u||\Gamma v| + |\Gamma u||\partial v|), \quad (3.69)$$

$$\begin{aligned} |\square_i u| &\leq C|c_i t - r|(1+t+r)^{-1}|\partial^2 u| \\ &\quad + C(1+t+r)^{-1}(|\partial u| + |\partial \Gamma u|). \end{aligned} \quad (3.70)$$

Let us give a proof of (3.68)-(3.70). Following [2], we define

$$S_i^\pm = \partial_t \pm c_i \partial_r. \quad (3.71)$$

Noting

$$\begin{aligned} S_i^+ u &= t^{-1}(c_i t - r)\partial_r u + t^{-1} S u, \\ |r^{-1} \Omega u| &\leq C|\nabla u|, \end{aligned}$$

(3.68) follows from the identity

$$\begin{aligned} Q^i(u, v) &= 2^{-1}(S_i^+ u S_i^- v + S_i^- u S_i^+ v) \\ &\quad - c_i^2 r^{-2} (\hat{x} \wedge \Omega u) \cdot (\hat{x} \wedge \Omega v), \hat{x} = x/r, \end{aligned}$$

which is derived from (2.5). If we rewrite $Q_{\alpha\beta}(u, v)$ by using (2.5) and

$$\partial_t u = -rt^{-1}\partial_r u + t^{-1} S u,$$

we can prove (3.69). Finally, (3.70) is the consequence of

$$\square_i u = S_i^+ S_i^- u - c_i^2 (2r^{-1}\partial_r u + r^{-2}\Omega u \cdot \Omega u).$$

Hence, it follows from (3.65), (3.68)-(3.70) that

$$\begin{aligned} |\Gamma^a N^i(\partial u, \partial^2 u)| &\leq C|c_i t - r|(1+t+r)^{-1} \sum_{|b|+|c|\leq|a|+1} |\partial \Gamma^b u^i| |\partial \Gamma^c u^i| \\ &\quad + C(1+t+r)^{-1} \sum_{|b|+|c|\leq|a|+2} |\Gamma^b u^i| |\partial \Gamma^c u^i| \end{aligned} \quad (3.72)$$

for $|c_i t - r| < c_i t/2$. Here we have used Lemma 1.2 in [7]. Therefore, if $|c_i t - r| < c_i t/2$, the estimates (3.60), (3.61) and (3.72) yield

$$\begin{aligned} |\Gamma^a N^i(\partial u, \partial^2 u)| &\leq C_N(1+t+r)^{-3}(1+|c_i t - r|)^{-1} \{\log(2+t)\}^3 \cdot \\ &\quad \cdot (\varepsilon + [\partial u]_{[(N+4)/2], t} \|\partial u\|_{N+6, t}^2) \end{aligned} \quad (3.73)$$

for $|a| \leq N$, since $\varepsilon < 1$ and $[\partial u]_{[(N+4)/2], t} < 1$. In case $|c_i t - r| \geq c_i t/2$, we find from (3.61) that

$$\begin{aligned} |\Gamma^a N^i(\partial u, \partial^2 u)| &\leq C_N \sum_{|b|+|c|\leq N+1} |\partial \Gamma^b u^i| |\partial \Gamma^c u^i| \\ &\leq C_N(1+r)^{-2}(1+t+r)^{-2} \{\log(2+t)\}^2 \cdot \\ &\quad \cdot (\varepsilon + [\partial u]_{[(N+4)/2], t} \|\partial u\|_{N+6, t}^2). \end{aligned} \quad (3.74)$$

Similarly, by (3.61)

$$\begin{aligned} |\Gamma^a R^i(\partial u, \partial^2 u)| &\leq C_N \sum_{(j,k) \neq (i,i)} \sum_{|b|+|c|\leq N+1} |\partial \Gamma^b u^j| |\partial \Gamma^c u^k| \\ &\leq C_N \{(1+r)^{-2}(1+t+r)^{-2} \\ &\quad + \sum_{j \neq i} (1+t+r)^{-2}(1+|c_j t - r|)^{-2}\} \{\log(2+t)\}^2 \cdot \\ &\quad \cdot (\varepsilon + [\partial u]_{[(N+4)/2], t} \|\partial u\|_{N+6, t}^2), \end{aligned} \quad (3.75)$$

$$\begin{aligned} |\Gamma^a G^i(\partial u, \partial^2 u)| &\leq C_N \sum_{1 \leq j, k, l \leq m} \sum_{|b|+|c|+|d|\leq N+1} |\partial \Gamma^b u^j| |\partial \Gamma^c u^k| |\partial \Gamma^d u^l| \\ &\leq C_N \{(1+r)^{-3}(1+t+r)^{-3} \\ &\quad + \sum_{1 \leq j \leq m} (1+t+r)^{-3}(1+|c_j t - r|)^{-3}\} \{\log(2+t)\}^3 \cdot \\ &\quad \cdot (\varepsilon + [\partial u]_{[(N+4)/2], t} \|\partial u\|_{N+6, t}^3), \end{aligned} \quad (3.76)$$

for $|a| \leq N$.

Therefore, it follows from (3.73)-(3.76) that

$$\begin{aligned}
& |\Gamma^a F^i(\partial u, \partial^2 u)| \\
& \leq C_N \{(1+t+r)^{-1} z_{1+\gamma, 1+\kappa}^{(i)}(t, r)^{-1} \\
& \quad + (1+t+r)^{-1} \sum_{j \neq i} z_{2, 1-\rho}^{(j)}(t, r)^{-1} \\
& \quad + (1+r)^{-1} z_{1+\gamma, 1+\kappa}^{(m+1)}(t, r)^{-1}\} (\varepsilon + [\partial u]_{[(N+4)/2], t} \|\partial u\|_{N+6, t}^3)
\end{aligned} \tag{3.77}$$

for $0 < \gamma < 1, 0 < \kappa < 1 - \gamma, 0 < \rho < 1$ and $|a| \leq N$. (3.77) give estimates of the right-hand side of the equations (3.53), and hence by Proposition 3.1 and (3.51),

$$|\partial \Gamma^a u^i(t, x)| \leq C_N (1+r)^{-1} (1 + |c_i t - r|)^{-1+\rho} (\varepsilon + [\partial u]_{[(N+5)/2], t} \|\partial u\|_{N+7, t}^3). \tag{3.78}$$

Using (3.78), we estimate $\Gamma^a R^i(\partial u, \partial^2 u)$ again. Then

$$\begin{aligned}
|\Gamma^a R^i(\partial u, \partial^2 u)| & \leq C_N \{(1+r)^{-2} (1+t+r)^{-2+2\rho} \\
& \quad + \sum_{j \neq i} (1+t+r)^{-2} (1 + |c_j t - r|)^{-2+2\rho}\} \cdot \\
& \quad \cdot (\varepsilon + [\partial u]_{[(N+6)/2], t} \|\partial u\|_{N+8, t}^6),
\end{aligned} \tag{3.79}$$

for $|a| \leq N$. We take $\rho < 1/2$ and replace the estimate (3.75) with (3.79). Then we have

$$\begin{aligned}
& |\Gamma^a F^i(\partial u, \partial^2 u)| \\
& \leq C_N \{(1+t+r)^{-1} z_{1+\gamma, 1+\kappa}^{(i)}(t, r)^{-1} \\
& \quad + (1+t+r)^{-1} \sum_{j \neq i} z_{1+\gamma, 1}^{(j)}(t, r)^{-1} \\
& \quad + (1+r)^{-1} z_{1+\gamma, 1+\kappa}^{(m+1)}(t, r)^{-1}\} (\varepsilon + [\partial u]_{[(N+6)/2], t} \|\partial u\|_{N+8, t}^6)
\end{aligned} \tag{3.80}$$

for some $0 < \gamma, \kappa < 1$. Then, applying Proposition 3.1 again, we have

$$|\partial \Gamma^a u^i(t, x)| \leq C_N (1+r)^{-1} (1 + |c_i t - r|)^{-1} (\varepsilon + [\partial u]_{[(N+7)/2], t} \|\partial u\|_{N+9, t}^6). \tag{3.81}$$

Consequently we have finished the proof.

4 Energy estimates.

Proposition 4.1 *Let u be the solution of (3.45)-(3.46). Assume that F^i satisfy (1.6) and c_i are different from each other. Assume moreover that $C_{\alpha\beta}^{ij}(\partial u)$ satisfy (1.7) and $[\partial u]_{[(N+10)/2],t} \leq \omega_N$, where $\omega_N \leq 1$ is a small number depending on N and given functions. Then,*

$$\|\partial u\|_{N,t} \leq C_N \varepsilon. \quad (4.1)$$

The proof of Theorem 1.1 is now easy. Uniqueness is proved by the application of the method in John [4], Appendix. Since the Cauchy problem (1.1)-(1.2) can be converted to a Cauchy problem for a symmetric hyperbolic system, we know the local existence of the smooth solution to (1.1)-(1.2).

Let $N \geq 10$. Fix $C_N \gg 1$ in (3.47) and (4.1), and set $\varepsilon_0 = \omega_N/2C_N$. According to Kato [5] and Majda [10] Chapter 2, we can take $T > 0$ so that there exists a solution to (1.1)-(1.2) in $[0, T] \times \mathbf{R}^3$ with $[\partial u]_{[(N+10)/2],T} \leq 2C_N \varepsilon$. Suppose that $T_* < \infty$ is the maximal of such T . Then, for arbitrary $t < T_*$, $[\partial u]_{[(N+10)/2],t} \leq \omega_N$. Therefore by Proposition 3.2 and Proposition 4.1,

$$\begin{aligned} [\partial u]_{N,t} &\leq C_N \varepsilon + C_N^6 \varepsilon^6 \\ &\leq C_N \varepsilon + C_N \varepsilon 2^{-5} \\ &\leq (3/2)C_N \varepsilon. \end{aligned}$$

This contradicts the maximality of T_* , and hence $T_* = \infty$. This proves the Theorem 1.1.

Proof of Proposition 4.1. If $v = (v^1, \dots, v^m)$ satisfies

$$\sum_{0 \leq \alpha, \beta \leq 3} \sum_{1 \leq j \leq m} a_{\alpha\beta}^{ij} \partial_\alpha \partial_\beta v^j = b^i, \quad i = 1, \dots, m \quad (4.2)$$

with

$$a_{\alpha\beta}^{ij} = a_{\beta\alpha}^{ij} = a_{\alpha\beta}^{ji}, \quad (4.3)$$

then we have the energy identity

$$\begin{aligned} & \sum_{0 \leq \alpha, \beta \leq 3} \sum_{1 \leq i, j \leq m} \{ \partial_\alpha (a_{\alpha\beta}^{ij} \partial_0 v^i \partial_\beta v^j) - \partial_\alpha a_{\alpha\beta}^{ij} \partial_0 v^i \partial_\beta^j - 2^{-1} \partial_0 (a_{\alpha\beta}^{ij} \partial_\alpha^i \partial_\beta^j) + 2^{-1} \partial_0 a_{\alpha\beta}^{ij} \partial_\alpha v^i \partial_\beta v^j \} \\ & = \sum_{1 \leq i \leq m} b^i \partial_0 v^i, \end{aligned} \quad (4.4)$$

by multiplying both side of (4.2) by $\partial_0 v^i$. Integrating (4.4) on $[0, t] \times \mathbf{R}^3$, we obtain

$$\begin{aligned} & 2^{-1} \int_{\mathbf{R}^3} \langle \partial v(t), \partial v(t) \rangle dx - 2^{-1} \int_{\mathbf{R}^3} \langle \partial v(0), \partial v(0) \rangle dx \\ & = \int_0^t ds \int_{\mathbf{R}^3} \left\{ \sum_{0 \leq \alpha, \beta \leq 3} \sum_{1 \leq i, j \leq m} (\partial_\alpha a_{\alpha\beta}^{ij} \partial_0 v^i \partial_\beta v^j - 2^{-1} \partial_0 a_{\alpha\beta}^{ij} \partial_\alpha v^i \partial_\beta v^j) + \sum_{1 \leq i \leq m} b^i \partial_0 v^i \right\} dx, \end{aligned} \quad (4.5)$$

where

$$\langle \partial v, \partial w \rangle = \sum_{1 \leq i, j \leq m} (a_{00}^{ij} \partial_0 v^i \partial_0 w^j - \sum_{1 \leq k, l \leq 3} a_{kl}^{ij} \partial_k v^i \partial_l w^j). \quad (4.6)$$

We first set

$$a_{\alpha\beta}^{ij} = \eta_{\alpha\beta}^i \delta^{ij} - C_{\alpha\beta}^{ij}(\partial u), \quad (4.7)$$

$$b^i = \sum_{1 \leq j \leq m} \sum_{0 \leq \alpha, \beta \leq 3} \{ a_{\alpha\beta}^{ij} \partial_\alpha \partial_\beta \Gamma^a u^j - \Gamma^a (a_{\alpha\beta}^{ij} \partial_\alpha \partial_\beta u^j) \} + \Gamma^a D^i(\partial u), \quad (4.8)$$

where

$$(\eta_{\alpha\beta}^i)_{0 \leq \alpha, \beta \leq 3} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -c_i^2 & 0 & 0 \\ 0 & 0 & -c_i^2 & 0 \\ 0 & 0 & 0 & -c_i^2 \end{pmatrix}. \quad (4.9)$$

Then each $a_{\alpha\beta}^{ij}$ satisfies (4.3), and $\Gamma^a u$ is a solution of (4.2). Therefore, it follows from (4.5) that

$$\begin{aligned} & 2^{-1} \int_{\mathbf{R}^3} \langle \partial \Gamma^a u(t), \partial \Gamma^a u(t) \rangle dx - 2^{-1} \int_{\mathbf{R}^3} \langle \partial \Gamma^a u(0), \partial \Gamma^a u(0) \rangle dx \\ & = \int_0^t ds \int_{\mathbf{R}^3} \left\{ \sum_{0 \leq \alpha, \beta \leq 3} \sum_{1 \leq i, j \leq m} (\partial_\alpha a_{\alpha\beta}^{ij} \partial_0 \Gamma^a u^i \partial_\beta \Gamma^a u^j - 2^{-1} \partial_0 a_{\alpha\beta}^{ij} \partial_\alpha \Gamma^a u^i \partial_\beta \Gamma^a u^j) \right. \\ & \quad \left. + \sum_{1 \leq i \leq m} b^i \partial_0 \Gamma^a u^i \right\} dx. \end{aligned}$$

Noting

$$\begin{aligned} & \sum_{1 \leq j \leq m} \sum_{0 \leq \alpha, \beta \leq 3} \{a_{\alpha\beta}^{ij} \partial_\alpha \partial_\beta \Gamma^a u^j - \Gamma^a (a_{\alpha\beta}^{ij} \partial_\alpha \partial_\beta u^j)\} \\ &= [\square_i, \Gamma^a] u^i - \sum_{1 \leq j, k \leq m} \sum_{0 \leq \alpha, \beta \leq 3} \{C_{\alpha\beta}^{ij} (\partial u) \partial_\alpha \partial_\beta \Gamma^a u^j - \Gamma^a (C_{\alpha\beta}^{ij} (\partial u) \partial_\alpha \partial_\beta \Gamma^a u^j)\}, \end{aligned}$$

we obtain

$$\begin{aligned} & 2^{-1} \int_{\mathbf{R}^3} \langle \partial \Gamma^a u(t), \partial \Gamma^a u(t) \rangle dx - 2^{-1} \int_{\mathbf{R}^3} \langle \partial \Gamma^a u(0), \partial \Gamma^a u(0) \rangle dx \\ & \leq C_N \sum_{\substack{|b|+|c| \leq N+1 \\ |b|, |c| \neq 0}} \sum_{1 \leq i, j, k \leq m} \int_0^t ds \int_{\mathbf{R}^3} |\Gamma^a \partial u^i| |\Gamma^b \partial u^j| |\Gamma^c \partial u^k| dx \\ & \leq C_N [\partial u]_{[(N+1)/2], t} \int_0^t (1+s)^{-1} \|\partial u(s)\|_N^2 ds \end{aligned}$$

for $|a| \leq N$. Since

$$\sum_{|a| \leq N} \int_{\mathbf{R}^3} \langle \partial \Gamma^a u(t), \partial \Gamma^a u(t) \rangle dx \geq C_N \|\partial u(t)\|_N^2$$

if $[\partial u]_{0, t}$ is small, it follows that

$$\|\partial u(t)\|_N^2 \leq C_N \{\varepsilon^2 + [\partial u]_{[(N+1)/2], t} \int_0^t (1+s)^{-1} \|\partial u(s)\|_N^2 ds\}.$$

Hence, by Gronwall's lemma,

$$\|\partial u(t)\|_N^2 \leq C_N \varepsilon^2 (1+t)^{C_N [\partial u]_{[(N+1)/2], t}}. \quad (4.10)$$

Next, we set

$$a_{\alpha\beta}^{ij} = \eta_{\alpha\beta}^i \delta^{ij}, \quad (4.11)$$

$$b^i = [\square_i, \Gamma^a] u^i + \Gamma^a F^i(\partial u, \partial^2 u). \quad (4.12)$$

Then by (4.5), we have

$$\|\partial u(t)\|_N^2 \leq C_N (\varepsilon^2 + \sum_{|a|, |b| \leq N} \sum_{1 \leq i \leq m} \int_0^t ds \int_{\mathbf{R}^3} |\Gamma^b F^i(\partial u, \partial^2 u)| |\partial_0 \Gamma^a u^i| dx). \quad (4.13)$$

Using (3.80) and (3.81), we have

$$|\Gamma^b F^i(\partial u, \partial^2 u)| |\partial_0 \Gamma^a u^i|$$

$$\begin{aligned}
&\leq C_N \{(1+s+r)^{-1} z_{1+\gamma,1+\kappa}^{(i)}(s,r)^{-1} \\
&\quad + (1+s+r)^{-1} \sum_{j \neq i} z_{1+\gamma,1}^{(j)}(s,r)^{-1} \\
&\quad + (1+r)^{-1} z_{1+\gamma,1+\kappa}^{(m+1)}(s,r)^{-1}\} (1+r)^{-1} (1+|c_i s - r|)^{-1} \cdot \\
&\quad \cdot (\varepsilon^2 + [\partial u]_{[(N+7)/2],s} \|\partial u\|_{N+9,s}^{12}) \\
&\leq C_N \{(1+s+r)^{-3-\kappa} (1+|c_i s - r|)^{-2-\gamma} \\
&\quad + (1+s+r)^{-4} \sum_{j \neq i} (1+|c_j s - r|)^{-1-\gamma} \\
&\quad + (1+r)^{-3-\gamma} (1+s+r)^{-2-\kappa}\} (\varepsilon^2 + [\partial u]_{[(N+7)/2],s} \|\partial u\|_{N+9,s}^{12}) \\
&\leq C_N (1+s)^{-1-\kappa} \sum_{j=1}^{m+1} (1+|c_j s - r|)^{-1-\gamma} r^{-2} (\varepsilon^2 + [\partial u]_{[(N+7)/2],s} \|\partial u\|_{N+9,s}^{12}). \quad (4.14)
\end{aligned}$$

Moreover, by (4.10) and (4.14) it follows that

$$\begin{aligned}
&|\Gamma^b F^i(\partial u, \partial^2 u)| |\partial_0 \Gamma^a u^i| \\
&\leq C_N (1+s)^{-1-\kappa} \sum_{j=1}^{m+1} (1+|c_j s - r|)^{-1-\gamma} r^{-2} (\varepsilon^2 + [\partial u]_{[(N+7)/2],s} \varepsilon^{12} (1+s)^{C_N [\partial u]_{[(N+10)/2],s}}) \\
&\leq C_N \varepsilon^2 (1 + [\partial u]_{[(N+7)/2],s}) (1+s)^{-1-\kappa + C_N [\partial u]_{[(N+10)/2],s}} \sum_{j=1}^{m+1} (1+|c_j s - r|)^{-1-\gamma} r^{-2}. \quad (4.15)
\end{aligned}$$

Therefore, if $[\partial u]_{[(N+10)/2],t} \leq 2^{-1} C_N^{-1} \kappa$, we obtain (4.1) from (4.13) and (4.15).

Acknowledgement.

I would like to thank Rentaro Agemi for his valuable advice. I would also like to express my gratitude to Kôji Kubota for helpful comments.

References

- [1] R. Agemi, K. Yokoyama, *The null condition and global existence to the system of wave equations with different speeds*, to appear in Series on Adv. in Math. for Appli. Sci., World Scientific.

- [2] A. Hoshiga, H. Kubo, *Global small amplitude solutions of nonlinear hyperbolic systems with a critical exponent under the null condition*, preprint.
- [3] F. John, *Lower bounds for the life span of solutions of nonlinear wave equations in three space dimensions*, Comm. Pure Appl. Math. **36**, 1983, 1-35.
- [4] F. John, *Nonlinear wave equations, Formation of singularities*, Pitcher lectures in the Math. Sci., Lehigh Univ., American Math. Soc., 1990.
- [5] T. Kato, *The Cauchy problem for quasi-linear symmetric hyperbolic systems*, Arch. Rational Mech. Anal. **58**, 1975, 181-205.
- [6] S. Klainerman, *Weighted L^∞ and L^1 estimates for solutions to the classical wave equation in three space dimensions*, Comm. Pure Appl. Math. **37**, 1984, 269-288.
- [7] S. Klainerman, *The null condition and global existence to nonlinear wave equations*, Lectures in Appl. Math. **23**, 1986, Amer. Math. Soc., 293-326.
- [8] S. Klainerman, T. C. Sideris, *On almost global existence for nonrelativistic wave equations in 3D*, Comm. Pure Appl. Math. **49**, 1996, 307-321.
- [9] M. Kovalyov, *Resonance-type behaviour in a system of nonlinear wave equations*, J. Differential Equations **77**, 1989, 73-83.
- [10] A. Majda, *Compressible fluid flow and systems of conservation laws in several space variables*, Springer.