Title	Existence of a forward self-similar stagnation flow of the Navier-Stokes equations
Author(s)	Ishikawa, M.; Matsui, S.
Citation	Hokkaido University Preprint Series in Mathematics, 418, 1-7
Issue Date	1998-7-1
DOI	10.14943/83564
Doc URL	http://hdl.handle.net/2115/69168
Type	bulletin (article)
File Information	pre418.pdf

[Instructions for use](#)

**Existence of a forward
self-similar stagnation flow
of the Navier-Stokes equations**

M. Ishikawa and S. Matsui

Series #418. July 1998

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

- #393 T. Arase, A remark on the quantale structure of multisets, 10 pages. 1997.
- #394 N. H. Bingham and A. Inoue, Extension of the Drasin-Shea-Jordan theorem, 16 pages. 1997.
- #395 N. H. Bingham and A. Inoue, Ratio Mercerian theorems with applications to Hankel and Fourier transforms, 30 pages. 1997.
- #396 Y. Nishiura and D. Ueyama, A skeleton structure of self-replicating dynamics, 27 pages. 1997.
- #397 K. Hirata and K. Sugano, On semisimple extensions of serial rings, 6 pages. 1997.
- #398 D. Pei, Singularities of $\mathbb{R}P^2$ -valued Gauss maps of surfaces in Minkowski 3-space, 15 pages. 1997.
- #399 T. Mikami, Markov marginal problems and their applications to Markov optimal control, 28 pages. 1997.
- #400 M. Tsujii, A simple proof for monotonicity of entropy in the quadratic family, 8 pages. 1998.
- #401 M. Nakamura and T. Ozawa, Global solutions in the critical Sobolev space for the wave equations with nonlinearity of exponential growth, 9 pages. 1998.
- #402 D. Lehmann and T. Suwa, Generalization of variations and Baum-Bott residues for holomorphic foliations on singular varieties, 19 pages. 1998.
- #403 T. Nakazi and K. Okubo, ρ -contraction and 2×2 matrix, 6 pages. 1998.
- #404 Y. Kohsaka, Free boundary problem for quasilinear parabolic equation with fixed angle of contact to a boundary, 28 pages. 1998.
- #405 K. Yokoyama, Global existence of classical solutions to systems of wave equations with critical nonlinearity in three space dimensions, 25 pages. 1998.
- #406 F. Hiroshima, Ground states and spectrum of quantum electrodynamics of non-relativistic particles, 58 pages. 1998.
- #407 N. Kawazumi and T. Uemura, Riemann-Hurwitz formula for Morita-Mumford classes and surface symmetries, 9 pages. 1998.
- #408 T. nakazi and K. Okubo, Generalized Numerical Radius And Unitary ρ -Dilation, 12 pages. 1998.
- #409 Y. Giga and K. Ito, Loss of convexity of simple closed curves moved by surface diffusion, 16 pages. 1998.
- #410 Y. Giga, K. Inui and S. Matsui, On the Cauchy problem for the Navier-Stokes equations with nondecaying initial data, 34 pages. 1998.
- #411 S. Izumiya, D. Pei and T. Sano, The lightcone gauss map and the lightcone developable of a spacelike curve in Minkowski 3-space, 16 pages. 1998.
- #412 M. Tsujii, Absolutely continuous invariant measures for piecewise real-analytic expanding maps on the plane, 18 pages. 1998.
- #413 J.-P. Brasselet, D. Lehmann, J. Seade and T. Suwa, Milnor classes of local complete intersections, 40 pages. 1998.
- #414 T. Suwa, Dual class of a subvariety, 19 pages. 1998.
- #415 M. Tsujii, Piecewise expanding maps on the plane with singular ergodic properties, 9 pages. 1998.
- #416 T. Yoshida, Categorical aspects of generating functions(I): Exponential formulas and Krull-Schmidt categories, 44 pages. 1998.
- #417 K. Yamaguchi, G_2 -Geometry of overdetermined systems of second order, 22 pages. 1998.

Existence of a forward self-similar stagnation flow of the Navier-Stokes equations

Dedicated to Professor Rentaro Agemi on his sixtieth birthday

Masayoshi ISHIKAWA and Shin'ya MATSUI*

Abstract

In this paper we construct an exact forward self-similar solution, representing a stagnation flow, for the Navier-Stokes equations by solving a third order ordinary differential equations.

1 Introduction

The solution of the form

$$u(x, t) = \frac{\sqrt{\nu}}{\sqrt{2(T-t)}} \cdot U\left(\frac{x}{\sqrt{2\nu(T-t)}}\right), p(x, t) = \frac{\nu}{\sqrt{2(T-t)}} \cdot P\left(\frac{x}{\sqrt{2\nu(T-t)}}\right)$$

is called (Leray's) backward self-similar solutions of the Navier-Stokes equations:

$$(NS) \quad \begin{cases} u_t - \nu \Delta u + (u, \nabla)u + p = 0, \\ \operatorname{div} u = 0 \end{cases}$$

for $x \in \mathbb{R}^3$ and $0 < t < T$, where $U = U(X) \in \mathbb{R}^3$ and $P = P(X)$ with $X \in \mathbb{R}^3$. In 1934 Leray [5] proposed to study backward self-similar solutions to construct blowing up solutions. His original question whether backward self-similar solutions with finite energy could be blow up, was answered negatively by [6] and [9] recently. They showed that backward self-similar solutions with the global energy estimates (in [6]) or the local energy estimates (in [9]) must be zero.

*Partly supported by the Grant-In-Aid for Scientific Research No. 10304010, the Japan Ministry of Education, Science and Culture.

In the interesting paper [8], Okamoto obtained so many exact solutions of backward and forward self-similar solutions (see, also [7]). Here, forward self-similar solutions are solutions in the form:

$$u(x, t) = \frac{\sqrt{\nu}}{\sqrt{2(T+t)}} \cdot U\left(\frac{x}{\sqrt{2\nu(T+t)}}\right), p(x, t) = \frac{\nu}{\sqrt{2(T+t)}} \cdot P\left(\frac{x}{\sqrt{2\nu(T+t)}}\right),$$

where $U = (U_1(X), U_2(X), U_3(X))$ and $P = P(X)$ satisfy the following equations:

$$(L) \quad \begin{cases} \Delta U - \nabla P = -U - (X, \nabla)U + (U, \nabla)U, \\ \operatorname{div} U = 0 \end{cases}$$

for $X \in \mathbb{R}^3$. It is known that forward self-similar solutions with finite energy do not exist ([3]). However, there is a systematic way based on global existence theorem, to construct forward self-similar solutions (with finite energy) as studied in [1, 3, 4]. Their solutions are not explicitly written by elementary functions or solutions of ordinary differential equations. In the section 12 in [8] he defined a forward self-similar stagnation flow of (NS) in the following manner. In (L), putting

$$U_1 = xf'(y), \quad U_2 = -f(y), \quad U_3 = 0$$

with $X = (x, y, z)$, then $f(y)$ satisfies

$$(L_1) \quad \begin{cases} -2f'(y) + (f'(y))^2 - yf''(y) - f(y)f''(y) - f'''(y) = -2k + k^2 \text{ for } y > 0, \\ f(0) = f'(0) = 0, \quad f'(\infty) = k, \end{cases}$$

where k is any number. The pressure P is determined by $P = -f'(y) - yf(y) - f^2(y)/2 + (-k + k^2/2)(x^2 + y^2)$. He, however, did not show existence of solutions to (L_1) . The purpose of the present paper is to construct solutions of (L_1) for $k = 1$ and to show results of numerical computation to (L_1) for $k = 1$. For this purpose we reduce (L_1) to a second order ordinary differential equations. Setting

$$v(y) = f'(y) - k, \tag{1}$$

we obtain

$$(L_2) \quad \begin{cases} v''(y) + ((1+k)y + w(y))v'(y) + 2(1-k)v(y) = v^2(y) \text{ for } y > 0, \\ v(0) = -k, \quad v(\infty) = 0, \quad w(y) = \int_0^y v(s)ds. \end{cases}$$

If $1 - k > 0$, Cauchy problem of linearized equation of (L_2) might have oscillate solutions. So, it seems to be impossible to construct solutions of (L_2) because of the condition $v(\infty) = 0$. On the other hand, according to our numerical

computations, linearized problems have so many blow up solution in the case of $1 - k < 0$. That is why we solve the equations (L_2) for $k = 1$.

To construct solutions, we employ the shooting method which is closely related to [2] although the equations discussed there are different. We consider the Cauchy problem:

$$(E_\alpha) \quad \begin{cases} v''(y) + (2y + w(y))v'(y) = v^2(y) & \text{for } y > 0, \\ v(0) = -1, \quad v'(0) = \alpha, \quad w(y) = \int_0^y v(s)ds. \end{cases}$$

We will find a suitable constant $\alpha = \alpha^*$ such that a solution of (E_{α^*}) satisfies $v(\infty) = 0$.

We are now in position to state our main theorem.

Theorem 1 *There exists the global solution $v(y)$ of (E_α) for some $\alpha > 0$, which tends to zero as $y \rightarrow \infty$. Thus, if $k = 1$, the equations (L_1) has a solution.*

In the section 2 we prove Theorem 1. After the proof we show a result of numerical experiment for (E_α) .

Finally, we are grateful to Professor Yoshikazu Giga for useful discussions and encouragement.

2 Existence of the exact solution

It is well known that the unique solution of (E_α) exists on $[0, T_\alpha)$ for some $T_\alpha > 0$ and any $\alpha \in \mathbb{R}$, where T_α may not be finite. Here and hereafter, we assume $\alpha \geq 0$ and denote the solution of (E_α) by $v_\alpha(y)$.

Lemma 1 *Let $v_\alpha(y)$ be the solution of (E_α) on $[0, T_\alpha)$ for $\alpha \geq 0$. Then we have*

$$v'_\alpha(y) \geq \alpha e^{-A_\alpha(y)} \quad \text{for } y \in [0, T_\alpha), \quad (2)$$

where $A_\alpha(y) = y^2 + \int_0^y w_\alpha(s)ds$ and $w_\alpha(y) = \int_0^y v_\alpha(s)ds$.

Proof. Since $v''_\alpha + (2y + w_\alpha) \cdot v'_\alpha = v_\alpha^2 \geq 0$, we have

$$\left(e^{A_\alpha(y)} \cdot v'_\alpha(y) \right)' \geq 0$$

for $y \in [0, T_\alpha)$. This implies (2).

Next to this, we define

$$J \equiv \{ \alpha \in [0, \infty); \text{ There exists } Y_\alpha \in (0, T_\alpha) \text{ such that } v_\alpha(Y_\alpha) = 0. \}$$

Here, if $\alpha > 0$, then (2) implies that Y_α is unique as far as it exists.

Lemma 2 *The solution $v_0(y)$ of (E_0) has no zeros. That is, $0 \notin J$.*

Proof. We first note that

$$v_0(y) > -1 \text{ for any } y \in (0, T_0). \quad (3)$$

Indeed, suppose that there exists some $\bar{y} \in (0, T_0)$ such that $v_0(\bar{y}) \leq -1$. Since $v_0(0) = -1$ and (2), we obtain $v_0(y) = -1$ for any $y \in [0, \bar{y}]$. However, the constant is not the solution of (E_0) . Thus we get (3).

We now prove our lemma. Suppose that $0 \in J$, that is, there exists $Y_0 \in (0, T_0)$ such that $v_0(y) < 0$ for $y \in [0, Y_0)$ and $v_0(Y_0) = 0$. Then we have $-1 < v_0(y) < 0$ for any $y \in (0, Y_0)$ by (3). Thus, we also obtain $w_0 \equiv \int_0^y v_0(s) ds < -y$ for $y \in (0, Y_0)$. These inequalities and (2) imply

$$v_0''(y) + yv_0'(y) = (-v_0(y)) \cdot (-v_0(y)) + (-w_0(y) - y) \cdot v_0'(y) < -v_0(y)$$

for $y \in (0, Y_0)$ using (E_0) . So we get $\{e^{y^2/2}v_0'(y)\}' + e^{y^2/2}v_0(y) < 0$ for $y \in (0, Y_0)$. Multiplying $e^{-y^2/2}$ to this inequality and integrating on $[0, Y_0]$, we have

$$\int_0^{Y_0} e^{-y^2/2} \cdot \{e^{y^2/2}v_0'(y)\}' dy + \int_0^{Y_0} v_0(y) dy < 0. \quad (4)$$

By the integration by parts, the first term in the left-hand side of (4) equals

$$\begin{aligned} v_0'(Y_0) + \int_0^{Y_0} yv_0'(y) dy \\ = v_0'(Y_0) + Y_0 \cdot v_0(Y_0) - \int_0^{Y_0} v_0(y) dy = v_0'(Y_0) - \int_0^{Y_0} v_0(y) dy, \end{aligned}$$

since $v_0'(0) = v_0(Y_0) = 0$. Hence, (4) yields $v_0'(Y_0) < 0$, which contradicts (2). We have proved $0 \notin J$.

Lemma 3 *The set J is open in $[0, \infty)$.*

Proof. Let $\alpha \in J$. We note that $\alpha > 0$ by Lemma 2. From the definition of J , there exists unique Y_α such that $v_\alpha(Y_\alpha) = 0$. By (2) there exists some positive δ such that $0 < v_\alpha(Y_\alpha + \delta) < \infty$. Thus, the continuity of $v_\alpha(y)$ with respect to y and α implies that there exists positive ε such that

$$v_\beta(Y_\alpha + \delta) > 0 \text{ for any } \beta \in (\alpha - \varepsilon, \alpha + \varepsilon).$$

Therefore, we have $v_\beta(Y_\beta) = 0$ for some $Y_\beta > 0$, as $v_\beta(0) = -1$ and (2) hold. This means that $\beta \in J$. This proves our lemma.

Lemma 4 *If $\alpha > 2/\pi$, then $\alpha \in J$. That is, the set J is not empty.*

Proof. Suppose that there exists $\beta > 2/\sqrt{\pi}$ with $\beta \notin J$. By (2) we have $v_\beta(y) < 0$ for any $y \geq 0$ and $w_\beta(y) = \int_0^y v_\beta(s)ds < 0$. Thus, the equation (E_β) and (2) imply

$$v_\beta''(y) + 2yv_\beta'(y) = -w_\beta(y) \cdot v_\beta'(y) + v_\beta(y)^2 > 0.$$

So we obtain

$$(e^{y^2} v_\beta'(y))' = e^{y^2} \cdot (v_\beta''(y) + 2yv_\beta'(y)) > 0.$$

Integrating this inequality twice on $[0, y]$, we have

$$v_\beta(y) > v_\beta(0) + v_\beta'(0) \cdot \int_0^y e^{-s^2} ds = -1 + \beta \cdot \int_0^y e^{-s^2} ds.$$

Letting $y \rightarrow \infty$ at the last inequality, then

$$\lim_{y \rightarrow \infty} v_\beta(y) \geq -1 + \beta \cdot \frac{\sqrt{\pi}}{2}$$

holds. From our assumption $\beta \notin J$ it follows that $\lim_{y \rightarrow \infty} v_\beta(y) \leq 0$. Thus we have $\beta \leq 2/\sqrt{\pi}$. This is contradiction. Thus we get our lemma.

The next lemma proves Theorem 1.

Lemma 5 *Put $\alpha^* = \inf J$. Then the solution v_{α^*} exists globally and satisfies $\lim_{y \rightarrow \infty} v_{\alpha^*}(y) = 0$.*

Proof. First, we note that $\alpha^* \notin J$ by Lemma 4. Thus, since the solution of (E_{α^*}) satisfies $v_{\alpha^*}(y) < 0$ for $y \in [0, T_{\alpha^*})$ by (2), then $T_{\alpha^*} = \infty$.

Let a sequence $\alpha_j \in J$ satisfy $\alpha_j \rightarrow \alpha^*$ as $j \rightarrow \infty$. From the definition of J , there exists $Y_{\alpha_j} > 0$ such that $v_{\alpha_j}(Y_{\alpha_j}) = 0$ for any j . Then we have

$$Y_{\alpha_j} \rightarrow \infty \text{ as } j \rightarrow \infty. \quad (5)$$

Indeed, suppose that the sequence $\{Y_{\alpha_j}\}$ is bounded. Then, for some subsequence $\{\alpha_{j(k)}\}$ and $Y_* \in [0, \infty)$ we have $Y_{\alpha_{j(k)}} \rightarrow Y_*$ as $k \rightarrow \infty$. Thus we get

$$0 = v_{\alpha_{j(k)}}(Y_{\alpha_{j(k)}}) \rightarrow v_{\alpha^*}(Y_{\alpha^*}) \text{ as } k \rightarrow \infty.$$

This means $v_{\alpha^*}(Y_{\alpha^*}) = 0$, that is $\alpha^* = \inf J \in J$. However, this contradicts Lemma 4. Hence we have (5).

Finally, we shall show $\lim_{y \rightarrow \infty} v_{\alpha^*}(y) = 0$, using (5). If there exists some $\varepsilon > 0$ such that $\lim_{y \rightarrow \infty} v_{\alpha^*}(y) = -\varepsilon < 0$. Then by (5) we obtain

$$v_{\alpha_j}(Y_{\alpha_j}) < -\frac{\varepsilon}{2} \text{ for any } j \geq N$$

for some natural number N . On the other hand, by the definition of Y_{α_j} we have $v_{\alpha_j}(Y_{\alpha_j}) = 0$. This is a contradiction. The proof is now complete.

The following figure is the result of our numerical experiment for the following equation

$$\begin{cases} -2f'(y) + (f'(y))^2 - yf''(y) - f(y)f''(y) - f'''(y) = -1 \text{ for } y > 0, \\ f(0) = f'(0) = 0, \quad f''(0) = 0.533, \end{cases}$$

which is equivalent to (E_α) with $\alpha = 0.533$, using the Runge-Kutta methods. In this figure each arrows indicate the vector fields $(xf'(y), -f(y))$ at each point. Here, the horizontal line is the x axis and the vertical line is the y axis.

References

- [1] Canone, M., Ondelettes, paraproducts et Navier-Stokes, Diderot editeur, Arta et Sciences, Paris, 1995.
- [2] Giga, Y., On elliptic equations related to self-similar solution for heat equations, Hiroshima Math. J., vol. 16 (1986), 539-552.
- [3] Giga, Y. and Miyakawa, T, Navier-Stokes flow in $\mathbb{R}^3$ with measures as initial vorticity and Morray spaces, Comm. in P.D.E., 14(1989), 577-618.
- [4] Kozono, H. and Yamazaki, M., Semilinear heat equations and the Navier-Stokes equation with distributions in new function spaces as initial data, Comm in P.D.E., 19(1994), 959-1014.
- [5] Leray, J., Sur le mouvement d'un liquide visquex emplissant l'espace, Acta Math. 63 (1934), 193-248.
- [6] Nečas, J., Růžička, M, and Šverák, V., On Leray's self-similar solution of the Navier-Stokes equations, Acta Math., 176 (1996), 283-394.
- [7] Okamoto, H., Axisymmetric stagnation flow converging obliquely on a cylinder, J. of the Phy. Soc. of Japan, vol. 64, No. 8 (1995), 2714-2717.
- [8] Okamoto, H., Exact solution of the Navier-Stokes equations via Leray's scheme, J.J.I.A.M., vol. 14, No.2 (1997), 169-197.
- [9] Tsai, T-P., On Leray's self-similar solutions of the Navier-Stokes equations satisfying local energy estimates, pre-print.

Masayoshi Ishikawa

Government & Public Corporation Information Systems Division,
Hitachi Ltd.
Shinsuna Plaza 6-27, Shinsuna 1 Chome, Koto ku, Tokyo
136, Japan
E mail; isshin@po.kumagaya.or.jp

Shin'ya Matsui

Hokkaido Information University
59-2, Nishi-Nopporo, Ebetsu
069-8585, Japan
E Mail; matsui@do-johodai.ac.jp