



Title	On the asymptotic behavior of the prediction error of a stationary process
Author(s)	Inoue, A.; Kasahara, Y.
Citation	Hokkaido University Preprint Series in Mathematics, 439, 1-12
Issue Date	1998-12-1
DOI	10.14943/83585
Doc URL	http://hdl.handle.net/2115/69189
Type	bulletin (article)
File Information	pre439.pdf



[Instructions for use](#)

**On the asymptotic behavior
of the prediction error
of a stationary process**

A. Inoue and Y. Kasahara

Series #439. December 1998

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #414 T. Suwa, Dual class of a subvariety, 19 pages. 1998.
- #415 M. Tsujii, Piecewise expanding maps on the plane with singular ergodic properties, 9 pages. 1998.
- #416 T. Yoshida, Categorical aspects of generating functions(I): Exponential formulas and Krull-Schmidt categories, 44 pages. 1998.
- #417 K. Yamaguchi, G_2 -Geometry of overdetermined systems of second order, 22 pages. 1998.
- #418 M. Ishikawa and S. Matsui, Existence of a forward self-similar stagnation flow of the Navier-Stokes equations, 8 pages. 1998.
- #419 S. Izumiya, H. Katsumi and T. Yamasaki, The rectifying developable and the spherical Darboux image of a space curve, 16 pages. 1998.
- #420 R. Kobayashi and Y. Giga, Equations with singular diffusivity, 45 pages. 1998.
- #421 D. Pei and T. Sano, The focal developable and the binormal indicatrix of a nonlightlike curve in Minkowski 3-space, 14 pages. 1998.
- #422 R. Kobayashi, J. A. Warren and W. C. Carter, Modeling grain boundaries using a phase field technique, 12 pages. 1998.
- #423 T. Tsukada, Reticular Legendrian Singularities, 28 pages. 1998.
- #424 A. N. Kirillov and M. Shimozone, A generalization of the Kostka-Foulkes polynomials, 37 pages. 1998.
- #425 M. Nakamura, Strichartz estimates for wave equations in the homogeneous Besov space, 17 pages. 1998.
- #426 A. Arai, On the essential spectra of quantum field Hamiltonians, 18 pages. 1998.
- #427 T. Sano, Bifurcations of affine invariants for one parameter family of generic convex plane curves, 11 pages. 1998.
- #428 F. Hiroshima, Ground states of a model in quantum electrodynamics, 48 pages. 1998.
- #429 F. Hiroshima, Uniqueness of the ground state of a model in quantum electrodynamics: A functional integral approach, 32 pages. 1998.
- #430 J. F. Van Diejen and A. N. Kirillov, Formulas for q -spherical functions using inverse scattering theory of reflectionless Jacobi operators, 33 pages. 1998.
- #431 G. Ishikawa, Determinacy, transversality and Lagrange stability, 13 pages. 1998.
- #432 T. Yoshida, Categorical aspects of generating functions(II): Operations on categories and functors, 65 pages. 1998.
- #433 K. Ito, Loss of convexity of compact hypersurfaces moved by surface diffusion, 20 pages. 1998.
- #434 Y. Shimizu, L^∞ -estimate of first-order space derivatives of Stokes flow in a half space, 22 pages. 1998.
- #435 T. Uemura, Morita-Mumford classes on finite cyclic subgroups of the mapping class group of closed surfaces, 14 pages. 1998.
- #436 A. Inoue and H. Kikuchi, Abel-Tauber theorems for Hankel and Fourier transforms and a problem of Boas, 20 pages. 1998.
- #437 T. Nakazi and T. Osawa, Finite rank intermediate Hankel operators on the Bergman space, 16 pages. 1998.
- #438 R. Yoneda, Compact Toeplitz operators on Bergman spaces, 14 pages. 1998.

**On the asymptotic behavior of the prediction error
of a stationary process**

Akihiko INOUE (Sapporo) and Yukio KASAHARA (Sapporo)

Abstract. We present an example of stationary process with long-time memory for which we can calculate explicitly the prediction error from the finite part of past. The long-time behavior of the prediction error is discussed.

1. INTRODUCTION AND RESULTS

Let $X = (X(t), t \in \mathbb{R})$ be a real, centered, weakly stationary process defined on a probability space (Ω, F, P) . For $T \geq 0$, we denote by $P_{[-T,0]}$ the orthogonal projection operator of $L^2(\Omega, F, P)$ to the subspace spanned by $\{X(u) : -T \leq u \leq 0\}$. Similarly, we write $P_{(-\infty,0]}$ for the orthogonal projection operator to the subspace spanned by $\{X(u) : -\infty < u \leq 0\}$. For $T > 0$ and $t > 0$, we define $Q(T, t)$ and $Q(\infty, t)$ by

$$\begin{aligned} Q(T, t) &:= E[\{X(t) - P_{[-2T,0]}X(t)\}^2], \\ Q(\infty, t) &:= E[\{X(t) - P_{(-\infty,0]}X(t)\}^2]. \end{aligned}$$

The reason why we put $2T$ rather than T in the above is that we follow the notation of Dym and McKean [3].

We are concerned with the asymptotic behavior of $Q(T, t) - Q(\infty, t)$ as $T \rightarrow \infty$. We are especially interested in the case where the stationary process X has the so called "long-time memory". The main difficulty of this problem comes from that of the calculation of $Q(T, t)$. In this paper, we present an example of X with long-time memory for which we can calculate $Q(T, t)$ explicitly. The authors know no other example of such X .

We write R for the autocovariance function of X : $R(t) = E[X(t)X(0)]$ for $t \in \mathbb{R}$. Let μ be the spectral measure of X : $R(t) = \int_{-\infty}^{\infty} e^{it\gamma} \mu(d\gamma)$. For $\frac{1}{2} < \alpha < 1$, we define the constants $c = c(\alpha)$ and $d = d(\alpha)$ by

$$c := \left\{ \pi \frac{\Gamma(1+\alpha)}{\Gamma(1-\alpha)} \right\}^{\frac{1}{2\alpha}}, \quad d := \frac{2^{3-2\alpha} \sin(\alpha\pi)}{\Gamma(\alpha - \frac{1}{2})^2}. \quad (1.1)$$

As usual, we write K_ν for the modified Bessel function (cf. Watson [5, 3.7]).

Theorem 1. Let $T > 0$, $t > 0$ and $\frac{1}{2} < \alpha < 1$. Set $T_1 := T + t$. Let X be a real, centered, weakly stationary process with spectral measure μ on \mathbb{R} of the form

$$\mu(d\gamma) = \frac{\sin(\alpha\pi)}{\pi} \cdot \frac{|\gamma|^{1-2\alpha}}{1+\gamma^2} d\gamma. \quad (1.2)$$

Then $Q(T, t) = d\{Q_1(T, t) + Q_2(T, t)\}$ with

$$Q_1(T, t) := \int_T^{T_1} \left\{ \int_s^{T_1} u^{1-\alpha} (T_1^2 - u^2)^{\alpha-\frac{3}{2}} K_{-\alpha}(u) du \right\}^2 \frac{T_1^2}{sK_{-\alpha}(s)^2} ds,$$

$$Q_2(T, t) := \int_T^{T_1} \left\{ \int_s^{T_1} u^{2-\alpha} (T_1^2 - u^2)^{\alpha-\frac{3}{2}} K_{1-\alpha}(u) du \right\}^2 \frac{ds}{sK_{1-\alpha}(s)^2}.$$

For the stationary process X in the theorem above, we can show that

$$R(t) \sim \frac{1}{\Gamma(2\alpha-1)} t^{-(2-2\alpha)} \quad (t \rightarrow \infty) \quad (1.3)$$

(see §4). In other words, X has long-time memory (see Beran [1]).

Theorem 2. Let α , t and X be as in Theorem 1. Then

$$Q(T, t) - Q(\infty, t) \sim \sin(\alpha\pi) \left\{ \frac{1}{\Gamma(\alpha-\frac{1}{2})} \int_0^t s^{\alpha-\frac{1}{2}} e^{s-t} ds \right\}^2 \frac{1}{T} \quad (T \rightarrow \infty).$$

Let X be as in Theorem 1 and let f be the spectral density of X :

$$f(\gamma) = \frac{\sin(\alpha\pi)}{\pi} \cdot \frac{|\gamma|^{1-2\alpha}}{1+\gamma^2} \quad (\gamma \in \mathbb{R}). \quad (1.4)$$

We write h for the outer function of X :

$$h(\zeta) := \exp \left\{ \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1+\gamma\zeta}{\gamma-\zeta} \cdot \frac{\log f(\gamma)}{1+\gamma^2} d\gamma \right\} \quad (\Im z > 0). \quad (1.5)$$

The canonical representation kernel F of X is defined by $F := (2\pi)^{-1/2} \hat{h}$, where \hat{h} is the Fourier transform of $h(\cdot) := \text{l.i.m.}_{\eta \downarrow 0} h(\cdot + i\eta) \in L^2(\mathbb{R})$. We have the following relation

$$h(\zeta) = (2\pi)^{-\frac{1}{2}} \int_0^{\infty} e^{i\zeta t} F(t) dt \quad (\Im z > 0). \quad (1.6)$$

Corollary. Let $\frac{1}{2} < \alpha < 1$ and $t > 0$. Let X be as in Theorem 1 with autocovariance function R and canonical representation kernel F . Then

$$Q(\frac{1}{2}T, t) - Q(\infty, t) \sim \left(\int_0^t F(s) ds \right)^2 \times \int_T^{\infty} \left\{ \frac{R(u)}{\int_{-u}^u R(s) ds} \right\}^2 du \quad (T \rightarrow \infty). \quad (1.7)$$

Recently, the authors obtained similar results to the corollary above for the stationary processes with autocovariance function R of the form

$$R(t) = \int_0^\infty e^{-|t|\lambda} \sigma(d\lambda) \quad (t \in \mathbb{R}),$$

where σ is a finite Borel measure on $(0, \infty)$. In them, we assumed that

$$R(t) \sim t^{-p} \ell(t) \quad (t \rightarrow \infty)$$

with $0 < p < \infty$ and ℓ slowly varying. In this case, the index p may be larger than or equal to 1 but the asymptotic relation (1.7) still holds as it is. The proofs of them are quite different from that of the present paper. The first author also obtained some relevant results for the partial autocorrelation coefficients of stationary time series. The details will appear elsewhere.

2. PROOF OF THEOREM 1

In the proof below, we apply the theory of “strings” of Krein as described in [3]. The key is to use Rule 6.9.4 in [3, p. 268].

Step 1. Recall $\frac{1}{2} < \alpha < 1$ and c from (1.1). We write m for the function

$$m(x) := \frac{c^2}{\alpha(1-\alpha)} x^{(1-\alpha)/\alpha} \quad (0 \leq x < \infty).$$

The purpose of this step is to obtain the functions A, B, C, D, K and the measure $d\Delta$ associated with the string specified by m ; see [3, Ch. 5] for background.

For $z \in \mathbb{C}$, we consider the following differential equation

$$\begin{cases} \frac{\partial^2}{\partial x^2} A(x, z) = -z^2 A(x, z) m'(x) & (0 < x < \infty), \\ A(0+, z) = 1, \quad \frac{\partial}{\partial x} A(0+, z) = 0. \end{cases}$$

The solution to the above is given by

$$A(x, z) := \pi \left\{ \frac{\alpha}{\sin(\alpha\pi)} \right\}^{\frac{1}{2}} z^\alpha x^{\frac{1}{2}} J_{-\alpha}(2czx^{\frac{1}{2\alpha}}) \quad (0 < x < \infty),$$

where J_ν is the Bessel function of the first kind ([5, 3.1]).

We set, as in [3, p. 172],

$$C(x, z) := A(x, z) \int_0^x \{A(y, z)\}^{-2} dy \quad (0 < x < \infty, \Im z \neq 0).$$

Since

$$\frac{\pi}{2 \sin(\alpha\pi)} \cdot \frac{d}{dx} \left\{ \frac{J_\alpha(x)}{J_{-\alpha}(x)} \right\} = \frac{1}{x J_{-\alpha}(x)^2}$$

(cf. [5, 5.11(1)]), we have

$$\frac{\alpha\pi}{\sin(\alpha\pi)} \cdot \frac{d}{dx} \left\{ \frac{J_\alpha(2cx x^{\frac{1}{2\alpha}})}{J_{-\alpha}(2cx x^{\frac{1}{2\alpha}})} \right\} = \frac{1}{x J_{-\alpha}(2cx x^{\frac{1}{2\alpha}})^2},$$

and so

$$C(x, z) = \left\{ \frac{\alpha}{\sin(\alpha\pi)} \right\}^{\frac{1}{2}} z^{-\alpha} x^{\frac{1}{2}} J_\alpha(2cx x^{\frac{1}{2\alpha}}) \quad (0 < x < \infty, \Im z \neq 0),$$

where we have used the following asymptotic representation (cf. [5, 3.12]):

$$J_\nu(x) \sim \left(\frac{2}{x} \right)^\nu \frac{1}{\Gamma(\nu+1)} \quad (x \rightarrow 0+). \quad (2.1)$$

As usual, we write $H_\nu^{(1)}$ for the Bessel function of the third kind ([5, 3.6]). Let the function D be as in [3, §5.4]. By [3, p. 175] and the asymptotic representations

$$J_\nu(z) = (\pi z/2)^{-\frac{1}{2}} [\cos \{z - \frac{1}{4}\pi(1+2\nu)\} + O(z^{-1})] \quad (|z| \rightarrow \infty), \quad (2.2)$$

$$H_\nu^{(1)}(z) \sim (\pi z/2)^{-\frac{1}{2}} \exp[i\{z - \frac{1}{4}\pi(1+2\nu)\}] \quad (|z| \rightarrow \infty) \quad (2.3)$$

(see [5, 7.2]), we have, for $\Im z \neq 0$,

$$\begin{aligned} D(0, z) &= \lim_{x \rightarrow \infty} \frac{C(x, z)}{A(x, z)} = \pi^{-1} z^{-2\alpha} \lim_{x \rightarrow \infty} \frac{J_\alpha(2cx x^{\frac{1}{2\alpha}})}{J_{-\alpha}(2cx x^{\frac{1}{2\alpha}})} \\ &= \pi^{-1} z^{-2\alpha} \lim_{x \rightarrow \infty} \left\{ e^{i\alpha\pi} - i \sin(\alpha\pi) \frac{H_{-\alpha}^{(1)}(2cx x^{\frac{1}{2\alpha}})}{J_{-\alpha}(2cx x^{\frac{1}{2\alpha}})} \right\} \\ &= \pi^{-1} z^{-2\alpha} e^{i\alpha\pi \cdot \text{sgn}(\Im z)}. \end{aligned}$$

Therefore, by [3, p. 175] and [5, 3.7(2)], for $0 < x < \infty$ and $\Im z \neq 0$,

$$\begin{aligned} D(x, z) &= D(0, z)A(x, z) - C(x, z) \\ &= \left\{ \frac{\alpha}{\sin(\alpha\pi)} \right\}^{\frac{1}{2}} z^{-\alpha} x^{\frac{1}{2}} \left\{ e^{i\alpha\pi \cdot \text{sgn}(\Im z)} J_{-\alpha}(2cx x^{\frac{1}{2\alpha}}) - J_\alpha(2cx x^{\frac{1}{2\alpha}}) \right\} \end{aligned}$$

or, by [5, 3.6(2)],

$$D(x, z) = \begin{cases} i\{\alpha \sin(\alpha\pi)\}^{\frac{1}{2}} z^{-\alpha} x^{\frac{1}{2}} H_{-\alpha}^{(1)}(2cx x^{\frac{1}{2\alpha}}), \\ -i\{\alpha \sin(\alpha\pi)\}^{\frac{1}{2}} z^{-\alpha} x^{\frac{1}{2}} H_{-\alpha}^{(2)}(2cx x^{\frac{1}{2\alpha}}). \end{cases}$$

In particular, by [5, 3.7(8)],

$$D(x, i) = \frac{2}{\pi} \{\alpha \sin(\alpha\pi)\}^{\frac{1}{2}} x^{\frac{1}{2}} K_{-\alpha}(2cx x^{\frac{1}{2\alpha}}) \quad (0 < x < \infty). \quad (2.4)$$

Recall μ from (1.2). We write $d\Delta$ for the measure $(1 + \gamma^2)d\mu(\gamma)$ on \mathbb{R} :

$$d\Delta(\gamma) := \frac{\sin(\alpha\pi)}{\pi} |\gamma|^{1-2\alpha} d\gamma.$$

By simple calculation,

$$\int_0^\infty \frac{\gamma^{1-2\alpha}}{\gamma^2 - z^2} d\gamma = \frac{\pi}{2 \sin(\alpha\pi)} z^{-2\alpha} \exp\{i\alpha\pi \cdot \operatorname{sgn}(\Im z)\} \quad (\Im z \neq 0),$$

whence

$$D(0, z) = \frac{1}{\pi} \int_{-\infty}^\infty \frac{d\Delta(\gamma)}{\gamma^2 - z^2} \quad (\Im z \neq 0).$$

See [3, §5.5] for the implication of this equality.

As in Rule 6.9.4 in [3, p. 268], we set

$$K(x) := -\frac{D(x, i)}{(\partial D / \partial x)(x, i)} \quad (x > 0).$$

By (2.4) and [5, 3.71(6)],

$$\frac{\partial D}{\partial x}(x, i) = -\frac{2c}{\alpha\pi} \{\alpha \sin(\alpha\pi)\}^{\frac{1}{2}} x^{\frac{1-\alpha}{2\alpha}} K_{1-\alpha}(2cx^{\frac{1}{2\alpha}})$$

whence

$$K(x) = c^{-1} \alpha x^{1-\frac{1}{2\alpha}} \frac{K_{-\alpha}(2cx^{\frac{1}{2\alpha}})}{K_{1-\alpha}(2cx^{\frac{1}{2\alpha}})} \quad (0 < x < \infty).$$

Let B as in [3, §5.7]. Then, for $0 < x < \infty$ and $\gamma \in \mathbb{R}$,

$$B(x, \gamma) = -\frac{1}{\gamma} \frac{\partial}{\partial x} A(x, \gamma) = c\pi \{\alpha \sin(\alpha\pi)\}^{-\frac{1}{2}} \gamma^\alpha x^{\frac{1-\alpha}{2\alpha}} J_{1-\alpha}(2c\gamma x^{\frac{1}{2\alpha}}).$$

Step 2. Recall $T > 0$, $t > 0$ and $T_1 = T + t$. By [3, §6.10] and Rule 6.9.4 in [3, p. 268] applied to the string specified by m , we obtain

$$Q(T, t) = Q_{\text{even}}(T, t) + Q_{\text{odd}}(T, t),$$

where

$$Q_{\text{even}}(T, t) = \pi \int_x^\infty \left[\frac{2}{\pi} \int_0^\infty \cos(\gamma T_1) \{A(y, \gamma) - \gamma K(y)B(y, \gamma)\} \frac{d\Delta(\gamma)}{1 + \gamma^2} \right]^2 \frac{dy}{K(y)^2},$$

$$Q_{\text{odd}}(T, t) = \pi \int_x^\infty \left[\frac{2}{\pi} \int_0^\infty \sin(\gamma T_1) \{\gamma A(y, \gamma) + K(y)B(y, \gamma)\} \frac{d\Delta(\gamma)}{1 + \gamma^2} \right]^2 dm(y)$$

with $T = \int_0^x \{m'(y)\}^{1/2} dy$, or $x = \{T/(2c)\}^{2\alpha}$. By change of variables $s = 2cy^{\frac{1}{2\alpha}}$,

$$Q_{\text{even}}(T, t) = \frac{2 \sin(\alpha\pi)}{\pi} \int_T^\infty \{I_1(s) - I_2(s)\}^2 \frac{ds}{sK_{-\alpha}(s)^2},$$

$$Q_{\text{odd}}(T, t) = \frac{2 \sin(\alpha\pi)}{\pi} \int_T^\infty \{I_3(s) - I_4(s)\}^2 \frac{ds}{sK_{1-\alpha}(s)^2},$$

where

$$I_1(s) := sK_{1-\alpha}(s) \int_0^\infty \cos(\gamma T_1) J_{-\alpha}(s\gamma) \frac{\gamma^{1-\alpha}}{1+\gamma^2} d\gamma,$$

$$I_2(s) := sK_{-\alpha}(s) \int_0^\infty \cos(\gamma T_1) J_{1-\alpha}(s\gamma) \frac{\gamma^{2-\alpha}}{1+\gamma^2} d\gamma,$$

$$I_3(s) := sK_{1-\alpha}(s) \int_0^\infty \sin(\gamma T_1) J_{-\alpha}(s\gamma) \frac{\gamma^{2-\alpha}}{1+\gamma^2} d\gamma,$$

$$I_4(s) := sK_{-\alpha}(s) \int_0^\infty \sin(\gamma T_1) J_{1-\alpha}(s\gamma) \frac{\gamma^{1-\alpha}}{1+\gamma^2} d\gamma.$$

Step 3. Recall the constant d from (1.1). In this step, we show that $Q_{\text{even}}(T, t) = d \cdot Q_1(T, t)$. We first note that $I_1 - I_2$ is continuous on $(0, \infty)$ by the asymptotic representations (2.1) and (2.2).

By [4, p. 68], (13.19),

$$\int_0^\infty \frac{x^{\nu+1}}{1+x^2} J_\nu(ax) \cos(xy) d\gamma = \cosh(y) K_\nu(a) \quad (y < a, -1 < \nu < \frac{3}{2}),$$

whence $I_1(s) - I_2(s) = 0$ for $s > T_1$ and also for $s = T_1$ by continuity. The calculation of $I_1(s) - I_2(s)$ for $T < s < T_1$ is more tricky. By

$$\frac{d}{dx} \{x^\alpha J_{-\alpha}(x)\} = -x^\alpha J_{1-\alpha}(x), \quad \frac{d}{dx} \{x^{1-\alpha} J_{1-\alpha}(x)\} = x^{1-\alpha} J_{-\alpha}(x),$$

$$\frac{d}{dx} \{x^\alpha K_{-\alpha}(x)\} = -x^\alpha K_{1-\alpha}(x), \quad \frac{d}{dx} \{x^{1-\alpha} K_{1-\alpha}(x)\} = -x^{1-\alpha} K_{-\alpha}(x)$$

(see [5, 3.2 and 3.71]), we have

$$\frac{\partial}{\partial s} \{I_1(s) - I_2(s)\} = -sK_{-\alpha}(s) \int_0^{\infty-} \cos(\gamma T_1) \gamma^{1-\alpha} J_{-\alpha}(s\gamma) d\gamma \quad (T < s < T_1).$$

In fact, by (2.2) and the second integral mean-value theorem ([6, §4.14]), the improper integral on the right converges uniformly in s on each compact subset of (T, T_1) , whence we may interchange derivative and integral. By [4, p. 67], (13.13),

$$\frac{\partial}{\partial s} \{I_1(s) - I_2(s)\} = -\frac{2^{1-\alpha} \pi^{\frac{1}{2}} T_1}{\Gamma(\alpha - \frac{1}{2})} s^{1-\alpha} (T_1^2 - s^2)^{\alpha - \frac{3}{2}} K_{-\alpha}(s)$$

for $T < s < T_1$, and so

$$I_1(s) - I_2(s) = \frac{2^{1-\alpha}\pi^{\frac{1}{2}}T_1}{\Gamma(\alpha - \frac{1}{2})} \int_s^{T_1} u^{1-\alpha}(T_1^2 - u^2)^{\alpha-\frac{3}{2}} K_{-\alpha}(u) du \quad (T < s < T_1).$$

Thus $Q_{\text{even}}(T, t) = d \cdot Q_1(T, t)$.

Step 4. Finally, we show that $Q_{\text{odd}}(T, t) = d \cdot Q_2(T, t)$. The proof is quite analogous to that for Q_{even} in Step 3. First, $I_3 - I_4$ is continuous on $(0, \infty)$. Next, since

$$\int_0^\infty \frac{x^\nu}{1+x^2} J_\nu(ax) \sin(xy) d\gamma = \sinh(y) K_\nu(a) \quad (y < a, -1 < \nu < \frac{5}{2})$$

(see [4, p. 166], (13.20)), $I_3 - I_4$ vanishes on $[T_1, \infty)$. Finally, by [4, p. 164], (13.9), for $T < s < T_1$,

$$\begin{aligned} \frac{\partial}{\partial s} \{I_3(s) - I_4(s)\} &= -s K_{1-\alpha}(s) \int_0^{\infty-} \sin(\gamma T_1) \gamma^{1-\alpha} J_{1-\alpha}(s\gamma) d\gamma \\ &= -\frac{2^{1-\alpha}\pi^{\frac{1}{2}}}{\Gamma(\alpha - \frac{1}{2})} s^{2-\alpha} (T_1^2 - s^2)^{\alpha-\frac{3}{2}} K_{1-\alpha}(s), \end{aligned}$$

whence

$$I_3(s) - I_4(s) = \frac{2^{1-\alpha}\pi^{\frac{1}{2}}}{\Gamma(\alpha - \frac{1}{2})} \int_s^{T_1} u^{2-\alpha} (T_1^2 - u^2)^{\alpha-\frac{3}{2}} K_{1-\alpha}(u) du \quad (T < s < T_1).$$

Thus $Q_{\text{odd}}(T, t) = d \cdot Q_2(T, t)$. This completes the proof. \square

3. PROOF OF THEOREM 2

Step 1. First we consider $Q_1(T, t)$. By change of variables $s' = s - T$, $u' = u - T$, we obtain

$$\begin{aligned} Q_1(T, t) &= 2^{2\alpha-3} \int_0^t ds \frac{(T+t)^2}{\left\{ (T+s)^{\frac{1}{2}} K_{-\alpha}(T+s) \right\}^2} \\ &\times \left[\int_s^t (T+u)^{\frac{1}{2}} K_{-\alpha}(T+u) (T+u)^{\frac{1}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} (t-u)^{\alpha-\frac{3}{2}} du \right]^2. \end{aligned}$$

By the asymptotic representation

$$x^{\frac{1}{2}} K_\nu(x) = (\pi/2)^{\frac{1}{2}} e^{-x} \left\{ 1 + \frac{1}{8}(4\nu^2 - 1)x^{-1} + O(x^{-2}) \right\} \quad (x \rightarrow \infty) \quad (3.1)$$

(see [5, 7.23]), we have, as $T \rightarrow \infty$,

$$(T+u)^{\frac{1}{2}} K_{-\alpha}(T+u) = (\pi/2)^{\frac{1}{2}} e^{-T-u} \left\{ 1 + \frac{1}{8}(4\alpha^2 - 1)T^{-1} + O(T^{-2}) \right\}.$$

This, together with

$$(1+x)^p = 1 + px + O(x^2) \quad (x \rightarrow 0+), \quad (3.2)$$

gives

$$\begin{aligned}
& (T+u)^{\frac{1}{2}} K_{-\alpha}(T+u)(T+u)^{\frac{1}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} \\
&= (\pi/2)^{\frac{1}{2}} e^{-T-u} T^{-1} \\
&\quad \times \left[1 + \left\{ \frac{1}{8}(4\alpha^2 - 1) + \left(\frac{1}{2} - \alpha\right)u + \frac{1}{2}\left(\alpha - \frac{3}{2}\right)(t+u) \right\} T^{-1} + O(T^{-2}) \right] \\
&= (\pi/2)^{\frac{1}{2}} e^{-T-u} T^{-1} \\
&\quad \times \left[1 + \frac{1}{2} \left\{ (\alpha^2 - \frac{1}{4}) - 2t + (\alpha + \frac{1}{2})(t-u) \right\} T^{-1} + O(T^{-2}) \right].
\end{aligned}$$

Hence, as $T \rightarrow \infty$,

$$\begin{aligned}
& \left[\int_s^t (T+u)^{\frac{1}{2}} K_{-\alpha}(T+u)(T+u)^{\frac{1}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} (t-u)^{\alpha-\frac{3}{2}} du \right]^2 \\
&= \frac{\pi e^{-2(t+T)}}{2T^2} \left[\left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right)^2 + \left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right) \right. \\
&\quad \left. \times \left(\int_0^{t-s} \left\{ (\alpha^2 - \frac{1}{4}) - 2t + (\alpha + \frac{1}{2})u \right\} u^{\alpha-\frac{3}{2}} e^u du \right) T^{-1} + O(T^{-2}) \right].
\end{aligned}$$

Similarly,

$$\frac{(T+t)^2}{(T+s)K_{-\alpha}(T+s)^2} = \frac{2T^2 e^{2(T+s)}}{\pi} \left[1 + \left\{ 2t - (\alpha^2 - \frac{1}{4}) \right\} T^{-1} + O(T^{-2}) \right].$$

Combining,

$$\begin{aligned}
& \frac{(T+t)^2}{(T+s)K_{-\alpha}(T+s)^2} \\
&\quad \times \left[\int_s^t (T+u)^{\frac{1}{2}} K_{-\alpha}(T+u)(T+u)^{\frac{1}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} (t-u)^{\alpha-\frac{3}{2}} du \right]^2 \\
&= e^{-2(t-s)} \left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right)^2 \\
&\quad + (\alpha + \frac{1}{2}) e^{-2(t-s)} \left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right) \left(\int_0^{t-s} u^{\alpha-\frac{1}{2}} e^u du \right) T^{-1} + O(T^{-2}).
\end{aligned}$$

Thus

$$Q_1(T, t) = 2^{2\alpha-3} J_0(t) + 2^{2\alpha-3} (\alpha + \frac{1}{2}) J_1(t) T^{-1} + O(T^{-2}) \quad (T \rightarrow \infty) \quad (3.3)$$

with

$$\begin{aligned}
J_0(t) &:= \int_0^t e^{-2s} \left(\int_0^s u^{\alpha-\frac{3}{2}} e^u du \right)^2 ds, \\
J_1(t) &:= \int_0^t e^{-2s} \left(\int_0^s u^{\alpha-\frac{3}{2}} e^u du \right) \left(\int_0^s u^{\alpha-\frac{1}{2}} e^u du \right) ds.
\end{aligned}$$

Step 2. Next we consider $Q_2(T, t)$. By change of variables $s' = s - T$, $u' = u - T$, we obtain

$$Q_2(T, t) = 2^{2\alpha-3} \int_0^t ds \frac{1}{\left\{ (T+s)^{\frac{1}{2}} K_{1-\alpha}(T+s) \right\}^2} \\ \times \left[\int_s^t (T+u)^{\frac{1}{2}} K_{1-\alpha}(T+u) (T+u)^{\frac{3}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} (t-u)^{\alpha-\frac{3}{2}} du \right]^2.$$

By (3.1), we have, as $T \rightarrow \infty$,

$$(T+u)^{\frac{1}{2}} K_{1-\alpha}(T+u) = (\pi/2)^{\frac{1}{2}} e^{-T-u} \left[1 + \frac{1}{8} \{4(1-\alpha)^2 - 1\} T^{-1} + O(T^{-2}) \right].$$

Therefore,

$$\begin{aligned} & (T+u)^{\frac{1}{2}} K_{1-\alpha}(T+u) (T+u)^{\frac{3}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} \\ &= (\pi/2)^{\frac{1}{2}} e^{-T-u} \\ & \quad \times \left[1 + \frac{1}{2} \left\{ (1-\alpha)^2 - \frac{1}{4} + (\alpha - \frac{3}{2})(t-u) \right\} T^{-1} + O(T^{-2}) \right]. \end{aligned}$$

Hence, as $T \rightarrow \infty$,

$$\begin{aligned} & \left[\int_s^t (T+u)^{\frac{1}{2}} K_{1-\alpha}(T+u) (T+u)^{\frac{3}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} (t-u)^{\alpha-\frac{3}{2}} du \right]^2 \\ &= \frac{\pi e^{-2(t+T)}}{2} \left[\left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right)^2 + \left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right) \right. \\ & \quad \left. \times \left(\int_0^{t-s} \left\{ (1-\alpha)^2 - \frac{1}{4} + (\alpha - \frac{3}{2})u \right\} u^{\alpha-\frac{3}{2}} e^u du \right) T^{-1} + O(T^{-2}) \right]. \end{aligned}$$

Similarly,

$$\frac{1}{(T+s)K_{1-\alpha}(T+s)^2} = \frac{2e^{2(T+s)}}{\pi} \left[1 - \left\{ (1-\alpha)^2 - \frac{1}{4} \right\} T^{-1} + O(T^{-2}) \right].$$

Combining,

$$\begin{aligned} & \frac{1}{(T+s)K_{1-\alpha}(T+s)^2} \\ & \times \left[\int_s^t (T+u)^{\frac{1}{2}} K_{1-\alpha}(T+u) (T+u)^{\frac{3}{2}-\alpha} \left\{ T + \frac{1}{2}(t+u) \right\}^{\alpha-\frac{3}{2}} (t-u)^{\alpha-\frac{3}{2}} du \right]^2 \\ &= e^{-2(t-s)} \left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right)^2 \\ & \quad + (\alpha - \frac{3}{2}) e^{-2(t-s)} \left(\int_0^{t-s} u^{\alpha-\frac{3}{2}} e^u du \right) \left(\int_0^{t-s} u^{\alpha-\frac{1}{2}} e^u du \right) T^{-1} + O(T^{-2}). \end{aligned}$$

Thus

$$Q_2(T, t) = 2^{2\alpha-3} J_0(t) + 2^{2\alpha-3} (\alpha - \frac{3}{2}) J_1(t) T^{-1} + O(T^{-2}) \quad (T \rightarrow \infty). \quad (3.4)$$

Step 3. Since $Q(T, t) \downarrow Q(\infty, t)$ as $T \rightarrow \infty$, we obtain, by (3.3) and (3.4),

$$Q(\infty, t) = \frac{2 \sin(\alpha\pi)}{\Gamma(\alpha - \frac{1}{2})^2} \int_0^t e^{-2s} \left(\int_0^s u^{\alpha-\frac{3}{2}} e^u du \right)^2 ds \quad (3.5)$$

and

$$Q(T, t) - Q(\infty, t) = \frac{(2\alpha - 1) \sin(\alpha\pi)}{\Gamma(\alpha - \frac{1}{2})^2} J_1(t) T^{-1} + O(T^{-2}). \quad (T \rightarrow \infty).$$

It remains to show that

$$(2\alpha - 1) J_1(t) = \left(\int_0^t s^{\alpha-\frac{1}{2}} e^{s-t} ds \right)^2. \quad (3.6)$$

Integrating by parts,

$$(\alpha - \frac{1}{2}) \int_0^s u^{\alpha-\frac{3}{2}} e^u du = s^{\alpha-\frac{1}{2}} e^s - \int_0^s u^{\alpha-\frac{1}{2}} e^u du,$$

whence the left-hand side of (3.6) is equal to $2\{J_2(t) - J_3(t)\}$, where

$$J_2(t) := \int_0^t s^{\alpha-\frac{1}{2}} e^s \left(\int_0^s u^{\alpha-\frac{1}{2}} e^u du \right) e^{-2s} ds,$$

$$J_3(t) := \int_0^t e^{-2s} \left(\int_0^s u^{\alpha-\frac{1}{2}} e^u du \right)^2 ds.$$

Again, integrating by parts,

$$J_2(t) = \left(\int_0^t s^{\alpha-\frac{1}{2}} e^{s-t} ds \right)^2 + 2J_3(t) - J_2(t).$$

Thus (3.6) follows. This completes the proof. \square

4. PROOF OF COROLLARY

First, we prove (1.3). We set

$$\ell(x) := \frac{x^2}{1+x^2}, \quad k(x) := x^{2\alpha-3} \cos(1/x) \quad (0 < x < \infty).$$

Then, for $t > 0$,

$$R(t) = \frac{2 \sin(\alpha\pi)}{\pi} \int_0^\infty \frac{\gamma^{1-2\alpha} \cos(\gamma t)}{1+\gamma^2} d\gamma = \frac{2 \sin(\alpha\pi)}{\pi} \int_0^\infty k(x) \ell(xt) dx.$$

Choose $\delta > 0$ so that $\delta < \min(2 - 2\alpha, 2\alpha - 1)$. Then the improper integrals

$$\int_{0+}^1 x^{-\delta} k(x) dx, \quad \int_1^{\infty-} x^{-\delta} k(x) dx$$

exist. Therefore, by the Bojanic-Karamata theorem (cf. Bingham, Goldie and Teugels [2, Th. 4.1.5]),

$$\int_0^{\infty} k(x)\ell(xt)dx \rightarrow \int_{0+}^{\infty-} k(x)dx \quad (t \rightarrow \infty).$$

Since

$$\int_{0+}^{\infty-} k(x)dx = \int_0^{\infty-} x^{1-2\alpha} \cos x dx = \frac{\pi}{2 \sin(\alpha\pi)\Gamma(2\alpha-1)},$$

(1.3) follows.

By (1.3),

$$\frac{R(u)}{\int_{-u}^u R(s)ds} \sim (\alpha - \frac{1}{2})u^{-1} \quad (u \rightarrow \infty),$$

whence

$$\int_T^{\infty} \left\{ \frac{R(u)}{\int_{-u}^u R(s)ds} \right\}^2 du \sim (\alpha - \frac{1}{2})^2 T^{-1} \quad (T \rightarrow \infty). \quad (4.1)$$

Recall f and h from (1.4) and (1.5). By applying Exercises 2.3.4 and 2.7.2 of [3] to the rational functions $1/(1-i\zeta)$ and $-i\zeta/(1-i\zeta)^2$, we obtain

$$\begin{aligned} \frac{1}{1-i\zeta} &= \exp \left\{ \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1+\gamma\zeta}{\gamma-\zeta} \cdot \frac{\log(1+\gamma^2)^{-1}}{1+\gamma^2} d\gamma \right\} \quad (\Im z > 0), \\ -i\zeta &= \exp \left\{ \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1+\gamma\zeta}{\gamma-\zeta} \cdot \frac{\log(\gamma^2)}{1+\gamma^2} d\gamma \right\} \quad (\Im z > 0) \end{aligned}$$

(note that $1/(1-i\zeta)$ and $-i\zeta$ are both positive on the upper imaginary axis). Therefore

$$h(\zeta) = \left\{ \frac{\sin(\alpha\pi)}{\pi} \right\}^{\frac{1}{2}} \frac{(-i\zeta)^{\frac{1}{2}-\alpha}}{1-i\zeta} \quad (\Im z > 0). \quad (4.2)$$

We set

$$G(t) := \frac{\{2 \sin(\alpha\pi)\}^{\frac{1}{2}}}{\Gamma(\alpha - \frac{1}{2})} \int_0^t e^{s-t} s^{\alpha-\frac{3}{2}} ds \quad (0 < t < \infty).$$

Then, by (4.2) and (1.6),

$$\begin{aligned} (2\pi)^{-\frac{1}{2}} \int_0^{\infty} e^{i\zeta t} G(t) dt &= \left\{ \frac{\sin(\alpha\pi)}{\pi} \right\}^{\frac{1}{2}} \frac{(-i\zeta)^{\frac{1}{2}-\alpha}}{1-i\zeta} \\ &= h(\zeta) = (2\pi)^{-\frac{1}{2}} \int_0^{\infty} e^{i\zeta t} F(t) dt, \end{aligned}$$

whence $F = G$. Therefore,

$$\int_0^t F(s)ds = \frac{\{2 \sin(\alpha\pi)\}^{\frac{1}{2}}}{(\alpha - \frac{1}{2})\Gamma(\alpha - \frac{1}{2})} \int_0^t e^{s-t} s^{\alpha-\frac{1}{2}} du. \quad (4.3)$$

This, together with Theorem 2 and (4.1), gives the corollary. \square

REFERENCES

- [1] Beran, J. (1994). *Statistics for long-memory processes*. Chapman & Hall, New York.
- [2] Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1989). *Regular variation*. 2nd edn, Cambridge University Press; 1st edn 1987.
- [3] Dym, H. and McKean, H.P. (1976). *Gaussian processes, function theory, and the inverse spectral problem*. Academic Press, New York.
- [4] Oberhettinger, F. (1990). *Tables of Fourier transforms and Fourier transforms of distributions*. Springer-Verlag, New York.
- [5] Watson, G.N. (1944). *A treatise on the theory of Bessel functions*. 2nd edn, Cambridge University Press.
- [6] Whittaker, E.T. and Watson, G.N. (1927). *A course of modern analysis*. 4th edn, Cambridge University Press.

Akihiko INOUE *Department of Mathematics, Hokkaido University, Sapporo 060, Japan.*

E-mail: `inoue@math.sci.hokudai.ac.jp`

Yukio KASAHARA *Department of Mathematics, Hokkaido University, Sapporo 060, Japan.*

E-mail: `y-kasa@math.sci.hokudai.ac.jp`