



Title	Some special bounded homomorphisms of a uniform algebra
Author(s)	Nakazi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 443, 1-10
Issue Date	1999-2-1
DOI	10.14943/83589
Doc URL	http://hdl.handle.net/2115/69193
Type	bulletin (article)
File Information	pre443.pdf



[Instructions for use](#)

**Some Special Bounded
Homomorphisms Of A Uniform**

Takahiko Nakazi

Series #443. February 1999

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #418 M. Ishikawa and S. Matsui, Existence of a forward self-similar stagnation flow of the Navier-Stokes equations, 8 pages. 1998.
- #419 S. Izumiya, H. Katsumi and T. Yamasaki, The rectifying developable and the spherical Darboux image of a space curve, 16 pages. 1998.
- #420 R. Kobayashi and Y. Giga, Equations with singular diffusivity, 45 pages. 1998.
- #421 D. Pei and T. Sano, The focal developable and the binormal indicatrix of a nonlightlike curve in Minkowski 3-space, 14 pages. 1998.
- #422 R. Kobayashi, J. A. Warren and W. C. Carter, Modeling grain boundaries using a phase field technique, 12 pages. 1998.
- #423 T. Tsukada, Reticular Legendrian Singularities, 28 pages. 1998.
- #424 A. N. Kirillov and M. Shimozono, A generalization of the Kostka-Foulkes polynomials, 37 pages. 1998.
- #425 M. Nakamura, Strichartz estimates for wave equations in the homogeneous Besov space, 17 pages. 1998.
- #426 A. Arai, On the essential spectra of quantum field Hamiltonians, 18 pages. 1998.
- #427 T. Sano, Bifurcations of affine invariants for one parameter family of generic convex plane curves, 11 pages. 1998.
- #428 F. Hiroshima, Ground states of a model in quantum electrodynamics, 48 pages. 1998.
- #429 F. Hiroshima, Uniqueness of the ground state of a model in quantum electrodynamics: A functional integral approach, 32 pages. 1998.
- #430 J. F. Van Diejen and A. N. Kirillov, Formulas for q -spherical functions using inverse scattering theory of reflectionless Jacobi operators, 33 pages. 1998.
- #431 G. Ishikawa, Determinacy, transversality and Lagrange stability, 13 pages. 1998.
- #432 T. Yoshida, Categorical aspects of generating functions(II): Operations on categories and functors, 65 pages. 1998.
- #433 K. Ito, Loss of convexity of compact hypersurfaces moved by surface diffusion, 20 pages. 1998.
- #434 Y. Shimizu, L^∞ -estimate of first-order space derivatives of Stokes flow in a half space, 22 pages. 1998.
- #435 T. Uemura, Morita-Mumford classes on finite cyclic subgroups of the mapping class group of closed surfaces, 14 pages. 1998.
- #436 A. Inoue and H. Kikuchi, Abel-Tauber theorems for Hankel and Fourier transforms and a problem of Boas, 20 pages. 1998.
- #437 T. Nakazi and T. Osawa, Finite rank intermediate Hankel operators on the Bergman space, 16 pages. 1998.
- #438 R. Yoneda, Compact Toeplitz operators on Bergman spaces, 14 pages. 1998.
- #439 A. Inoue and Y. Kasahara, On the asymptotic behavior of the prediction error of a stationary process, 12 pages. 1998.
- #440 A. Arai, M. Hirokawa and F. Hiroshima, On the absence of eigenvectors of Hamiltonians in a class of massless quantum field models without infrared cutoff, 26 pages. 1998.
- #441 A. N. Kirillov, A. Schilling and M. Shimozono, A bijection between Littlewood-Richardson tableaux and rigged configurations, 66 pages. 1999.
- #442 M. Tsujii, Absolutely continuous invariant measures for expanding piecewise linear maps, 28 pages. 1999.

Some Special Bounded Homomorphisms Of A Uniform Algebra

Takahiko Nakazi

ABSTRACT. Let $L(H)$ be the algebra of all bounded linear operators on a Hilbert space H and let A be a uniform algebra. In this paper we study the following questions. When is a unital bounded homomorphism Φ of A in $L(H)$ completely bounded? When is the norm $\|\Phi\|$ of Φ equal to the completely bounded norm $\|\Phi\|_{cb}$? In some special cases we answer this question. Suppose Φ is ρ -contractive ($0 < \rho < \infty$) where Φ is contractive if $\rho = 1$. We show that if A is a Dirichlet algebra or $\dim A/\ker \Phi = 2$ then Φ has a ρ -dilation. If Φ is a ρ -contractive homomorphism then $\|\Phi\| = \max(1, \rho)$ and if it has a ρ -dilation then $\|\Phi\|_{cb} = \max(1, \rho)$. Moreover we give a new example of a hypo-Dirichlet algebra in which a unital contractive homomorphism has a contractive dilation.

1. Introduction

Let X be a compact Hausdorff space, let $C(X)$ be the algebra of complex-valued continuous functions on X , and let A be a uniform algebra on X . Let H be a complex Hilbert space and $L(H)$ the algebra of all bounded linear operators on H . $I = I_H$ is the identity operator in H . An algebra homomorphism $f \rightarrow \Phi(f)$ of A in $L(H)$, which satisfies

$$\Phi(1) = I \text{ and } \|\Phi(f)\| \leq \gamma \|f\|_\infty$$

for some positive constant $\gamma \geq 1$, is called a unital bounded homomorphism of A . If $\gamma = 1$, it is called a unital contractive homomorphism.

For a subspace B of A , let $M_n(B)$ denote the set of $n \times n$ matrices with entries from B . For a map $\phi : B \rightarrow L(H)$, we obtain maps $\phi_n : M_n(B) \rightarrow M_n(L(H))$ via the formula

$$\phi_n((a_{ij})) = (\phi(a_{ij})).$$

If ϕ is a bounded map, then each ϕ_n will be bounded, and when $\sup_n \|\phi_n\|$ is finite, we call ϕ a completely bounded map of B in $L(H)$. We write

$$\|\phi\|_{cb} = \sup_n \|\phi_n\|.$$

The following problem is natural and important.

©0000 (copyright holder)

1991 *Mathematics Subject Classification*. Primary 47A20, 46J25.

The author was supported in part by Grant-in-Aid for Scientific Research, Ministry of Education.

PROBLEM . Suppose Φ is a unital bounded homomorphism of A .

- I. When is Φ completely bounded ?
- II. When is Φ completely bounded and $\|\Phi\| = \|\Phi\|_{cb}$?

A unital contractive homomorphism $v \rightarrow \tilde{\Phi}(v)$ of $C(X)$ on a Hilbert space K is called a contractive dilation of the unital contractive homomorphism $f \rightarrow \Phi(f)$ of A on H if H is a Hilbert subspace of K and

$$\Phi(f) = P\tilde{\Phi}(f)|_H \quad (f \in A)$$

where P is the orthogonal projection of K onto H . If Φ has a contractive dilation then Φ is completely contractive and hence $\|\Phi\| = \|\Phi\|_{cb}$. If Φ is completely contractive then Φ has a contractive dilation. This is well known (see [18, Corollary 6.7]).

If A is a uniform algebra and the uniform closure of $A + \bar{A}$, that is, $[A + \bar{A}]$ has finite codimension n in $C(X)$ then A is called a n -hypo-Dirichlet algebra and it is called a Dirichlet algebra when $[A + \bar{A}] = C(X)$, that is, $n = 0$.

If $\dim H < \infty$, Φ is completely bounded for arbitrary uniform algebra A (see [18, Exercises 3.11]). If $\dim H = \infty$ and A is the disc algebra, then there exists a unital bounded homomorphism Φ which is not completely bounded. This was recently shown by G.Pisier [19]. If A is a n -hypo-Dirichlet algebra and Φ is a unital contractive homomorphism then Φ is completely bounded. This was shown by R.G.Douglas and V.I.Paulsen [6]. However we don't know whether Φ is completely contractive or not. They are known solutions for Problem I.

Now we will give known solutions for Problem II when Φ is contractive. If A is the disc algebra then there exists a contractive dilation. This is a famous theorem of B.Sz.-Nagy [10]. T.Ando [2] generalized this to the bidisc algebra. However S.K.Parrot [17] gave an example of Φ which does not have a contractive dilation in the polydisc algebra for $n \geq 3$. If A is a Dirichlet algebra then there exists a contractive dilation (cf. [7]). For a n -hypo-Dirichlet algebra with $n \neq 0$, we don't know whether there exists a contractive dilation or does not. The polydisc algebra for $n \geq 2$ is not a n -hypo-Dirichlet algebra. If A is an annulus algebra, that is, a rational function algebra on an annulus, then there exists a contractive dilation. This was shown by J.Agler [1]. An annulus algebra is a 1-hypo-Dirichlet algebra. If \mathcal{A} is the disc algebra and $A = \{f \in \mathcal{A} ; f(0) = f(1)\}$, then A is also a 1-hypo-Dirichlet algebra. The author [12] proved that Φ has a contractive dilation for this example. Even if $\dim H < \infty$, by an example of S.K.Parrot [17] Φ may not have a contractive dilation for some uniform algebra. The author and the late K.Takahashi [14], and Che-Chen Chu [5] showed that if $\dim H \leq 2$, Φ has a contractive dilation for an arbitrary uniform algebra.

Now we will give more concrete problems than Problem II.

PROBLEM . Suppose Φ is a unital bounded homomorphism of A .

(II-a) Suppose $\|\Phi\| \leq 1$. When A is a n -hypo-Dirichlet algebra and $n \geq 1$, does Φ have a contractive dilation ?

(II-b) Under what conditions on Φ which is $\|\Phi\| > 1$, is Φ completely bounded with $\|\Phi\| = \|\Phi\|_{cb}$ when A is a n -hypo-Dirichlet algebra and $n \geq 0$, or $\dim H \leq 2$?

In this paper, we study Problem (II-a) and (II-b). In §2, we give a new example of a 1-hypo-Dirichlet algebra in which a unital contractive homomorphism has a contractive dilation. In §3, we define a ρ -contractive homomorphism Φ and a ρ -dilation of Φ for $0 < \rho < \infty$. If Φ is a ρ -contractive homomorphism then $\|\Phi\| \leq$

$\max(1, \rho)$ and if it has a ρ -dilation then $\|\Phi\|_{cb} \leq \max(1, \rho)$. In §4, we introduce a δ -homomorphism of A for $-\infty < \delta < 1$. This homomorphism is bounded. In fact, we show more, that is, ' δ -homomorphism' is equivalent to ' $\rho = 1/(1 - \delta)$ -contractive homomorphism'. In §5, we show that a ρ -contractive homomorphism has a ρ -dilation when A is a Dirichlet algebra. In §6, we consider Problem II under conditions on Φ , that is, $\dim A/\ker \Phi = 2$ or a hypothesis on $\ker \Phi$.

2. Third example of a hypo-Dirichlet algebra for Problem II-a

For a n -hypo-Dirichlet algebra with $n \neq 0$, we know only two examples ([1], [11]) in which a unital contractive homomorphism has a contractive dilation, that is, Problem (II-a). They are 1-hypo-Dirichlet algebras. In this section, we give a new example which is also a 1-hypo-Dirichlet algebra. In the proof of Theorem 2.1, a theorem of T.Ando [2] is used essentially. Unfortunately we could not generalize Theorem 2.1 to $A = \{f \in \mathcal{A} ; f'(0) = f''(0) = \dots = f^{(n)}(0) = 0\}$.

THEOREM 2.1. *Let \mathcal{A} be the disc algebra and $A = \{f \in \mathcal{A} ; f'(0) = 0\}$. If Φ is a unital contractive homomorphism of A then Φ has a contractive dilation or equivalently Φ is a completely contractive.*

PROOF. Since $A = \mathbf{C} + z^2\mathcal{A}$, $A_0 = \{f \in A ; f(0) = 0\} = z^2\mathcal{A}$. A_0 has two generators, that is, A_0 is generated by z^2 and z^3 because $2\ell \pm 1$ can be written as the form $2n + 3m$. Let $\Phi(z^2) = S$ and $\Phi(z^3) = T$ then $ST = TS$, $\|S\| \leq 1$ and $\|T\| \leq 1$. By a well known theorem of T.Ando [2], there exist two commuting operators U and V on a Hilbert space K with $H \subset K$ such that

$$S^n T^m = P U^n V^m | H$$

for all nonnegative integers n and m where P is an orthogonal projection from K to H . Any polynomial f in A_0 is written as the following :

$$f = a_{10}z^2 + a_{01}z^3 + \sum_{j, \ell \geq 1} a_{j\ell} z^{2j} z^{3\ell}$$

and so

$$\begin{aligned} \Phi(f) &= a_{10}S + a_{01}T + \sum_{j, \ell \geq 1} a_{j\ell} S^j T^\ell \\ &= P(a_{10}U + a_{01}V + \sum_{j, \ell \geq 1} a_{j\ell} U^j V^\ell) | H \end{aligned}$$

By a theorem of C.R.Putnam and B.Fuglede [20, Corollary 1.19], $U^*V = VU^*$. Hence if we set $\tilde{\Phi}((z^2)^j (z^3)^\ell) = U^j V^\ell$ for any integers j and ℓ then $\tilde{\Phi}$ is a unital contractive homomorphism of $C(X)$ in $L(K)$ and $\Phi = P\tilde{\Phi}|H$ on A_0 . Thus $\tilde{\Phi}$ is a contractive dilation of Φ . \square

3. ρ -Contractive homomorphism

A bounded linear operator T on H is said to be of class C_ρ if there exists a unitary operator U (called a unitary ρ -dilation) on a Hilbert space $K \supset H$ such that $T^n = \rho P U^n | H$ for $n = 1, 2, \dots$ where P is an orthogonal projection from K to H . B.Sz.-Nagy [9] showed that if T is a contraction then it is of class C_1 . If the numerical radius of T is less than equal to one then it is of class C_2 [3]. If T is of class C_ρ then $\|T\| \leq \max(1, \rho)$.

Suppose Φ is a unital algebra homomorphism of A in $L(H)$ and $0 < \rho < \infty$. When $\Phi(f)$ is of class C_ρ for any f in A with $\|f\|_\infty \leq 1$, Φ is called a ρ -contractive homomorphism of A . A 1-contractive homomorphism is equivalent to a contractive homomorphism. A 2-contractive homomorphism Φ is equivalent to that

$$\sup_{\substack{f \in A \\ \|f\|_\infty \leq 1}} \sup_{\substack{y \in H \\ \|y\|=1}} |\langle \Phi(f)y, y \rangle| = 1.$$

If Φ is ρ -contractive then $\|\Phi\| \leq \max(1, \rho)$. We will study Problem (II-b) when Φ is ρ -contractive.

A unital contractive homomorphism $v \rightarrow \tilde{\Phi}(v)$ on a Hilbert space K is called a ρ -dilation of the unital bounded homomorphism $f \rightarrow \Phi(f)$ of A on H if H is a Hilbert subspace of K and

$$\Phi(f) = \rho P \tilde{\Phi}(f)|_H \quad (f \in A_\tau)$$

where P is the orthogonal projection of K onto H , A_τ is the kernel of τ in $M(A)$ and $0 < \rho < \infty$.

If Φ has a ρ -dilation then Φ is a unital completely bounded map. However the converse is not true even for the disc algebra A and $\dim H = 2$. This may be well known. Suppose $T = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$, then $f \rightarrow \Phi(f) = f(T)$ is a unital completely bounded homomorphism but it has not a ρ -dilation for any ρ [15]. If Φ is a unital completely bounded homomorphism, then Φ/ρ is completely contractive but it is not unital. However if Φ has a ρ -dilation then the following is true.

PROPOSITION 3.1. *For a unital bounded homomorphism Φ , Φ has a ρ -dilation with respect to τ if and only if $\Phi(f) = \rho \Phi_0(f)$ ($f \in A_\tau$) where Φ_0 is a unital completely contractive map on A , equivalently Φ_0 has a contractive dilation.*

PROOF. For the 'only if' part, put $\Phi_0(f) = P \tilde{\Phi}(f)|_H$ ($f \in A$). Then Φ_0 has a contractive dilation $\tilde{\Phi}$ and hence it is completely contractive on A . The 'if' part follows from a theorem of W.Arveson (cf.[18, Corollary 6.7]). \square

4. δ -homomorphism

For $-\infty < \delta < 1$, Φ is called a δ -homomorphism for τ in $M(A)$ if Φ is a unital algebra homomorphism of A and $\operatorname{Re} \Phi(f) \geq 0$ whenever f in A , $\tau(f) = 1$ and $\operatorname{Re} f \geq \delta$. In this section we show that Φ is a δ -homomorphism for some τ if and only if Φ is a $\rho = 1/(1 - \delta)$ -contractive homomorphism.

PROPOSITION 4.1. *If a unital algebra homomorphism Φ has a ρ -dilation for τ , then Φ is a $\delta = \left(1 - \frac{1}{\rho}\right)$ -homomorphism for τ .*

PROOF. Suppose that $\Phi(h) = \rho P \tilde{\Phi}(h)|_H$ for $h \in A_\tau$. If $f \in A$, $\tau(f) = 1$ and $\operatorname{Re} f \geq 1 - \frac{1}{\rho}$, since $\tau(f - \tau(f)) = 0$,

$$\Phi(f) - \tau(f)I = \rho P \tilde{\Phi}(f)|_H - \rho \tau(f)I$$

and so

$$\Phi(f) = \rho \left\{ P \tilde{\Phi}(f)|_H + \tau(f) \left(\frac{1}{\rho} - 1 \right) I \right\}.$$

Since $Re f \geq 1 - \frac{1}{\rho}$, $\tilde{\Phi}(Re f) \geq 1 - \frac{1}{\rho}$ and so

$$Re \Phi(f) \geq \rho \left\{ P \tilde{\Phi}(Re f) |H + \left(\frac{1}{\rho} - 1 \right) I \right\} \geq 0.$$

□

PROPOSITION 4.2. Suppose $\delta \neq 0$. Φ is a δ -homomorphism for τ if and only if for any h in A_τ with $\|h\|_\infty < 1$,

$$|\langle \Phi(h)y, y \rangle| \leq \frac{1}{2|\delta|} \|y\|^2 + \frac{2\delta - 1}{2|\delta|} \|\Phi(h)y\|^2 \quad (y \in H)$$

PROOF. Suppose Φ is a δ -homomorphism for τ . Put

$$f = (1 - \delta) \frac{1 + h}{1 - h} + \delta$$

where $h \in A_\tau$ and $\|h\|_\infty < 1$ then $f \in A$, $Re f \geq \delta$ and $\tau(f) = 1$. For $x \in H$, put $x = (I - \Phi(h))y$ then

$$\begin{aligned} & Re \langle \Phi(f)x, x \rangle \\ &= (1 - \delta) Re \left\langle \frac{I + \Phi(h)}{I - \Phi(h)} x, x \right\rangle + \delta \|x\|^2 \\ &= (1 - \delta) Re \langle (I + \Phi(h))y, (I - \Phi(h))y \rangle + \delta \langle (I - \Phi(h))y, (I - \Phi(h))y \rangle \\ &= (1 - \delta) (\|y\|^2 - \|\Phi(h)y\|^2) + \delta (\|y\|^2 + \|\Phi(h)y\|^2 - 2 Re \langle \Phi(h)y, y \rangle) \\ &= \|y\|^2 + (2\delta - 1) \|\Phi(h)y\|^2 - 2\delta \langle \Phi(h)y, y \rangle. \end{aligned}$$

By hypothesis on Φ ,

$$2\delta Re \langle \Phi(h)y, y \rangle \leq \|y\|^2 + (2\delta - 1) \|\Phi(h)y\|^2$$

and so

$$|\langle \Phi(h)y, y \rangle| \leq \frac{1}{2|\delta|} \|y\|^2 + \frac{2\delta - 1}{2|\delta|} \|\Phi(h)y\|^2.$$

The proof is reversible. In fact, if $f \in A$, $Re f \geq \delta$ and $\tau(f) = 1$ then

$$f = (1 - \delta) \frac{1 + h}{1 - h} + \delta$$

for some $h \in A_\tau$ with $\|h\|_\infty \leq 1$. Hence if we put for $0 < \varepsilon < 1$

$$f_\varepsilon = (1 - \delta) \frac{1 + \varepsilon h}{1 - \varepsilon h} + \delta$$

then $f_\varepsilon \rightarrow f$ uniformly as $\varepsilon \rightarrow 1$, $\|\varepsilon h\|_\infty \leq 1$ and $\tau(f_\varepsilon) = 1$. Since $Re \langle \Phi(f_\varepsilon)x, x \rangle \geq 0$, as $\varepsilon \rightarrow 1$, $Re \langle \Phi(f)x, x \rangle \geq 0$ for any $x \in H$. □

THEOREM 4.3. Φ is a δ -homomorphism for some (or any) τ in $M(A)$ if and only if Φ is a $\rho = 1/(1 - \delta)$ -contractive homomorphism.

PROOF. [16, Theorem 2] and Proposition 4.2 imply the theorem, or we can show this by the proof of Proposition 4.2 and [11, Theorem 11.1]. □

5. Condition A

In this section, we show that a ρ -contractive homomorphism has a ρ -dilation when A is a Dirichlet algebra. This is a generalization of a theorem of C.Foias and I.Suciu [7] for $\rho = 1$ and a theorem of B.Sz.Nagy and C.Foias (cf. [11]) for the disc algebra. They give solutions for Problem (II-b).

THEOREM 5.1. *Let A be a Dirichlet algebra and $0 < \rho < \infty$. If Φ is a ρ -contractive homomorphism of A in $L(H)$ then for any τ in $M(A)$ it has a ρ -dilation.*

PROOF. Put $\Phi'(h) = \frac{1}{\rho}\Phi(h) - \tau(h)\left(\frac{1}{\rho} - 1\right)I$ for $h \in A$. By Theorem 4.3 if Φ is ρ -contractive then Φ is a $\delta = \left(1 - \frac{1}{\rho}\right)$ -homomorphism for any $\tau \in M(A)$. Hence if $Reh \geq 0$ then $Re\Phi'(h) \geq 0$. If we extend Φ' to $\tilde{\Phi} : A + \bar{A} \rightarrow L(H)$ by $\tilde{\Phi}(f + \bar{g}) = \Phi'(f) + \Phi'(g)^*$, then $\tilde{\Phi} : C(X) \rightarrow L(H)$ is positive because A is a Dirichlet algebra. By the dilation theorem of M.A.Naimark (cf. [21, Theorem 7.5]) there exists a Hilbert space K , an orthogonal projection $P : K \rightarrow H$ and a multiplicative linear map $u \rightarrow \tilde{\Phi}(u)$ of $C(X)$ in $L(K)$, which satisfies $\tilde{\Phi}(1) = I_K$, $\|\tilde{\Phi}(u)\| \leq \|u\|_\infty$, $u \in C(X)$ and $\Phi'(f) = P\tilde{\Phi}(f)|_H$ for $f \in A$. If $f \in A_\tau$ then $\Phi'(f) = \frac{1}{\rho}\Phi(f)$ and so

$$\Phi(f) = \rho P\tilde{\Phi}(f)|_H.$$

□

PROPOSITION 5.2. *Let A be an arbitrary uniform algebra and $0 < \rho < \infty$. Suppose Φ is a ρ -contractive homomorphism of A in $L(H)$. If $A/\ker \Phi$ is isometrically isomorphic to A/\mathcal{J} where A is a Dirichlet algebra on some compact Hausdorff space Y and \mathcal{J} is a closed ideal in A , then Φ has a ρ -dilation for any τ in $M(A)$ with $\tau = 0$ on $\ker \Phi$.*

PROOF. Let ϕ be an isometric isomorphism from A/\mathcal{J} onto $A/\ker \Phi$. For each $f \in A$, we will write $\phi(f + \mathcal{J}) = \phi(f) + \ker \Phi$ where $\phi(f) \in A$. Moreover we will write Φ again for the map : $f + \ker \Phi \rightarrow \Phi(f)$. Put $\Psi = \Phi \circ \phi$, then Ψ is a unital homomorphism of A/\mathcal{J} in $L(H)$. We will write Ψ again for the map : $f \rightarrow \Psi(f + \mathcal{J})$, then $\mathcal{J} = \ker \Psi$. Since we may assume that τ is a complex homomorphism on $A/\ker \Phi$ by [8, Theorem 6.2 in Chapter I], $\tau \circ \phi$ is a complex homomorphism on A/\mathcal{J} and so we may assume that $\tau \circ \phi \in M(A)$. If $f \in A_{\tau \circ \phi}$, and $\|f\|_\infty \leq 1$, then $\phi(f) \in A_\tau$ and $\|\phi(f) + \mathcal{J}\| \leq 1$. By hypothesis, $\Phi(\phi(f))$ is of class C_ρ and so $\Psi(f) = \Phi \circ \phi(f)$ is of class C_ρ for $f \in A_{\tau \circ \phi}$ with $\|f\|_\infty \leq 1$. Hence Ψ is a ρ -contractive homomorphism of A in $L(H)$ with respect to $\tau \circ \phi$. Since A is a Dirichlet algebra, by Proposition 3.1 and Theorem 5.1 $\Psi = \rho\Psi_0$ on $A_{\tau \circ \phi}$ where Ψ_0 is a unital completely contractive map on A . Put $\Phi_0 = \Psi_0 \circ \phi^{-1}$ then Φ_0 is a unital completely contractive map on A and $\Phi = \rho\Phi_0$ on A_τ . Proposition 3.1 implies the theorem. □

Let A be a n -hypo-Dirichlet algebra and let N_τ be the set of all representing measures of τ in $M(A)$. Then $\dim N_\tau = n$ and there exists a core measure m of N_τ (cf. [8, p106]). Then by [8, Theorem 5.1 in Chapter IV], there is a constant $c > 0$ such that $\nu \leq cm$ for all ν in N_τ . Hence if h is the Radon-Nikodym derivative of ν with respect to m then $\nu = hdm$. Set $N_\tau^m = \{h : \nu = hdm \text{ and}$

$\nu \in N_\tau$, then N_τ^m is a subset of $L^\infty(m)$. Thus N_τ^m can be considered as a subset of $L^\infty(m)$. Many important n -hypo-Dirichlet algebras satisfy a natural condition on $N_\tau^m : N_\tau^m \subset C(X)$. The author showed [13] that if $N_\tau^m \subset C(X)$ then a unital contractive homomorphism Φ of A has a ρ -dilation with respect to τ . It is a long standing open question whether we can choose $\rho = 1$. The motivation of our study in this paper is in this open question. The following Proposition 5.4 implies that if Φ is ρ -contractive for enough small $\rho > 0$ then Φ has a 1-dilation.

LEMMA 5.3. *Let A be a n -hypo-Dirichlet algebra and $N_\tau^m \subset C(X)$. Then there exists a positive linear map T from $C(X)$ to $[A + \bar{A}]$ such that $T(f) = f$ ($f \in A_\tau$) and $T(1)$ is a positive constant ≥ 1 .*

PROOF. This is proved in the proof of [13, Theorem]. \square

PROPOSITION 5.4. *Let A be a n -hypo-Dirichlet algebra and $N_\tau^m \subset C(X)$ for some τ in $M(A)$ where m is a core measure of N_τ . If Φ is a ρ -contractive homomorphism of A in $L(H)$ then it has a ρ' -dilation where $\rho' = \rho T(1)$ and T is a map in Lemma 5.3.*

PROOF. Suppose Φ is ρ -contractive. Then Φ is a $\delta = \left(1 - \frac{1}{\rho}\right)$ -homomorphism for τ by Theorem 4.3. Put

$$\Phi'(f) = \frac{1}{\rho}\Phi(f) - \tau(f) \left(\frac{1}{\rho} - 1\right) I \quad (f \in A),$$

then by the proof of Theorem 5.1, there exists an extension $\tilde{\Phi}'$ of Φ' on $[A + \bar{A}]$ and $\tilde{\Phi}'$ is positive on it. Then $T(1)^{-1}\tilde{\Phi}' \circ T$ is a positive map from $C(X)$ to $L(H)$ and $T(1)^{-1}\tilde{\Phi}' \circ T(1) = I$. By the dilation theorem of M.A.Naimark (cf. [21, Theorem 7.5]) there exists a Hilbert space K , an orthogonal projection P from K to H and a multiplicative linear map $u \rightarrow \tilde{\Phi}(u)$ of $C(X)$ in $L(K)$, which satisfies $\tilde{\Phi}(1) = I_K$, $\|\tilde{\Phi}(u)\| \leq \|u\|_\infty$, $u \in C(X)$ and $\tilde{\Phi}' \circ T(u) = T(1)P\tilde{\Phi}(u)|_H$. Because $T(f) = f$ ($f \in A_\tau$),

$$\Phi(f) = \rho T(1)P\tilde{\Phi}(f)|_H \quad (f \in A_\tau).$$

Suppose $\rho' = \rho T(1)$. \square

6. Condition on Φ

In this section, under conditions on Φ we consider Problem II. We consider ρ -contractive homomorphisms when $A/\ker\Phi$ is of two dimension. The author and the late K.Takahashi [13] showed that Φ has a 1-dilation when $\rho = 1$. We generalize it to any ρ . Proposition 6.3 is a generalization of a result of G.Misra [8] which was shown for a rational uniform algebra on the complex plane and $\rho = 1$.

For x, y in $M(A)$ and a bounded point derivation δ at x , let

$$\sigma_A(x, y) = \sup\{|f(y)| ; f(x) = 0, f \in A \text{ and } \|f\|_\infty \leq 1\}$$

and

$$\omega_A(x, \delta) = \sup\{|\delta(f)| ; f(x) = 0, f \in A \text{ and } \|f\|_\infty \leq 1\}.$$

THEOREM 6.1. *Let A be an arbitrary uniform algebra. If Φ satisfies one of the following conditions (1), (2) and (3), then Φ is completely bounded and $\|\Phi\| = \|\Phi\|_{cb}$.*

1. $\|\Phi(f)^2\| = \|\Phi(f)\|^2 \quad (f \in A)$

2. E is an interpolation set in X and $\ker \Phi = \{f \in A ; f = 0 \text{ on } E\}$
 3. $\|\Phi\| \leq 1$, $\ker \Phi = \{f \in A ; f = 0 \text{ on } E\}$ for some finite set $E \subset M(A)$ and $\sigma_A(x, y) = 1$ for any x, y in E with $x \neq y$.

PROOF. (1) If $\|\Phi(f)^2\| = \|\Phi(f)\|^2$ ($f \in A$) then the closure of $\Phi(A)$ is regarded as a uniform algebra. By [18, Theorem 3.8], $\|\Phi\| = \|\Phi\|_{cb}$.

(2) Since E is an interpolation set in X , $A/\ker \Phi$ is isometrically isomorphic to a subalgebra of $C(E)$ and hence $\Phi(A) \subseteq C(E)$. Again by [18, Theorem 3.8], $\|\Phi\| = \|\Phi\|_{cb}$.

(3) Since $E = \{x_1, \dots, x_n\}$ is a finite set, $\dim A/\ker \Phi = n < \infty$ and $\Phi(A) = \{\sum_{j=1}^n a_j P_j ; a_j \in \mathbf{C}, P_i P_j = \delta_{ij} P_j \text{ and } j = 1, \dots, n\}$. Suppose $\Phi(f_j) = P_j$ and $f_j \in$

A . Then $f_i f_j = \delta_{ij} f_j$ and $f_i(x_j) = \delta_{ij}$. Since $\sigma_A(x_i, x_j) = \delta_{ij}$, there exist $\{g_n^{(i)}\}_{n=1}^\infty$ in A such that $g_n^{(i)}(x_i) \rightarrow 1$ ($n \rightarrow \infty$), $g_n^{(i)}(x_j) = 0$ ($j \neq i$) and $\|g_n^{(i)}\|_\infty \leq 1$. Since

$A/\ker \Phi = \{\sum_{i=1}^n a_i f_i + \ker \Phi ; a_i \in \mathbf{C} \text{ and } i = 1, \dots, n\}$, $g_n^{(i)} - a_{in} f_i \in \ker \Phi$ and

$\Phi(g_n^{(i)}) = a_{in} P_i$. Since Φ is contractive, $|a_{in}| \|P_i\| \leq 1$ and so $\{a_{in}\}$ is bounded. Hence there exists a subsequence $\{a_{in(j)}\}$ such that $a_{in(j)} \rightarrow a_i$ as $j \rightarrow \infty$ for each i . Then $\lim_{j \rightarrow \infty} g_n^{(i)} - a_i f_i \in \ker \Phi$ and $\lim_{j \rightarrow \infty} g_n^{(i)}(x_i) = 1$. Therefore $a_i = 1$ and $\|f_i + \ker \Phi\| \leq 1$, and P_i is selfadjoint for $i = 1, \dots, n$ because Φ is contractive. Thus $\Phi(A)$ is a commutative C^* -algebra. By [18, Theorem 3.8], $\|\Phi\| = \|\Phi\|_{cb}$. \square

THEOREM 6.2. *Suppose Φ is a ρ -contractive homomorphism of A . If $A/\ker \Phi$ is of two dimension then Φ has a ρ -dilation for any τ in $M(A)$ with $\tau = 0$ on $\ker \Phi$.*

PROOF. Suppose $\ker \Phi = \{f \in A ; f(x) = f(y) = 0\}$ where $x, y \in M(A)$ with $x \neq y$. By [13, Lemma 1 and its proof],

$$\Phi(f) = \begin{pmatrix} f(x)I_{H_1} & (f(x) - f(y))C \\ 0 & f(y)I_{H_2} \end{pmatrix} \text{ on } H = H_1 \oplus H_2$$

for all $f \in A$ where C is a bounded linear operator from H_2 to H_1 , and

$$A/\ker \Phi = \{f(x)f_1 + f(y)f_2 + \ker \Phi ; f \in A\}$$

where $f_1(x) = f_2(y) = 1$ and $f_1(y) = f_2(x) = 0$. By [13, Lemma 3], if $\|C\|^2 + 1 = 1/\sigma_A(x, y)^2$ then $\|\Phi(f)\| = \|f + \ker \Phi\|$ for all $f \in A$.

For any $x, y \in M(A)$, there exist a Dirichlet algebra \mathcal{A} and $s, t \in M(\mathcal{A})$ such that $\sigma_A(x, y) = \sigma_A(s, t)$. In fact, we can choose the disc algebra \mathcal{A} . Suppose

$$\Psi(F) = \begin{pmatrix} F(s)I_{H_1} & (F(s) - F(t))B \\ 0 & F(t)I_{H_2} \end{pmatrix}$$

for all $F \in \mathcal{A}$ where B is a bounded linear operator from H_2 to H_1 . Then

$$A/\ker \Psi = \{F(s)F_1 + F(t)F_2 + \ker \Psi ; F \in \mathcal{A}\}$$

where $F_1(s) = F_2(t)$ and $F_1(t) = F_2(s)$, and $\|\Phi(f)\| = \|\Psi(F)\|$ whenever $f(x) = F(s)$ and $f(y) = F(t)$, and $B = C$. If $B = C$ and $\|C\|^2 + 1 = 1/\sigma_A(x, y)^2$, then $\|B\|^2 + 1 = 1/\sigma_A(s, t)^2$ and so $\|f + \ker \Phi\| = \|\Phi(f)\| = \|\Psi(f)\| = \|f + \ker \Psi\|$. Hence for given Φ , we can find a unital homomorphism Ψ on \mathcal{A} such that $A/\ker \Phi \cong A/\ker \Psi$. By Proposition 5.2, Φ has a ρ -dilation for any $\tau \in M(A)$ with $\tau = 0$ on $\ker \Phi$. If $\ker \Phi$ is not the above form, then $\ker \Phi = \{f \in A ; f(x) = \delta(f) = 0\}$

where $x \in M(A)$ and δ is a bounded point derivation at x . By [13, Lemma 1 and its proof],

$$\Phi(f) = \begin{pmatrix} f(x)I_{H_1} & \delta(f)C \\ 0 & f(x)I_{H_2} \end{pmatrix} \text{ on } H = H_1 \oplus H_2$$

for all $f \in A$ where C is a bounded linear operator from H_2 to H_1 , and

$$A/\ker \Phi = \{f(x)1 + \delta(f)f_0 + \ker \Phi ; f \in A\}$$

where $f_0(x) = 0$ and $\delta(f_0) = 1$. By [13, Lemma 3], if $\|C\| = 1/\omega_A(x, \delta)$ then $\|\Phi(f)\| = \|f + \ker \Phi\|$ for all $f \in A$. As in the first part of the proof, by Proposition 5.2 we can show that Φ has a ρ -dilation for any $\tau \in M(A)$. \square

If $\dim H = 2$, then an algebra homomorphism Φ has the following form :

$$\Phi_1(f) = \begin{pmatrix} f(x) & c(f(x) - f(y)) \\ 0 & f(y) \end{pmatrix}$$

where $x, y \in M(A)$ and $x \neq y$ or

$$\Phi_2(f) = \begin{pmatrix} f(x) & c\delta(f) \\ 0 & f(x) \end{pmatrix}$$

where $x \in M(A)$ and δ is a bounded point derivation at x .

PROPOSITION 6.3. *Suppose Φ is a unital bounded homomorphism of A in $L(H)$ and $\dim H = 2$.*

1. *When $\Phi = \Phi_1$, Φ is a ρ -contractive homomorphism if and only if*

$$(1 + |c|^2)|\rho\zeta(f(x) - f(y))|^2 \\ \leq |\{\rho + (1 - \rho)\overline{f(x)\zeta}\}\{\rho + (1 - \rho)f(y)\zeta\} - \overline{f(x)}f(y)|\zeta|^2|$$

for any $f \in A$ with $\|f\|_\infty \leq 1$ and any $\zeta \in D$.

2. *When $\Phi = \Phi_2$, Φ is a ρ -contractive homomorphism if and only if*

$$|c|^2|\delta(f)|^2 \\ \leq (\rho - 2)|f(x)|^2 + 2(1 - \rho)|f(x)| + \rho$$

for any $f \in A$ with $\|f\|_\infty \leq 1$.

PROOF. The author and Okubo [15] gave a necessary and sufficient condition for that a triangle 2×2 matrix is of class C_ρ . By [15, Theorem] $\Phi_1(f)$ is of class C_ρ if and only if

$$|c|^2|f(x) - f(y)|^2 + |f(x) - f(y)|^2 \\ \leq \inf_{\zeta \in D} \left| \frac{\{\rho + (1 - \rho)\overline{f(x)\zeta}\}\{\rho + (1 - \rho)f(y)\zeta\} - \overline{f(x)}f(y)|\zeta|^2}{\rho\zeta} \right|^2$$

and by [15, Remark] $\Phi_2(f)$ is of class C_ρ if and only if

$$|c|^2|\delta(f)|^2 \leq (\rho - 2)|f(x)|^2 + 2(1 - \rho)|f(x)| + \rho.$$

\square

In Proposition 6.3, suppose $\rho = 1$. $\Phi = \Phi_1$ is a 1-contractive homomorphism if and only if $|c|^2 \leq (1 - |f(x)|^2)(1 - |f(y)|^2)/|f(x) - f(y)|^2$ for any $f \in A$ with $\|f\|_\infty \leq 1$. This implies [9, Theorem 1.1]. $\Phi = \Phi_2$ is a 1-contractive homomorphism if and only if $|c|^2 \leq (1 - |f(x)|^2)/|\delta(f)|^2$ for any $f \in A$ with $\|f\|_\infty \leq 1$. Suppose $\rho = 2$. $\Phi = \Phi_1$ is a 2-contractive homomorphism if and only if $1 + |c|^2 \leq$

$\inf_{\zeta \in D} \left| \frac{2 - (\overline{f(x)}\zeta + f(x)\bar{\zeta})}{\zeta(f(x) - f(y))} \right|^2$ for any $f \in A$ with $\|f\|_\infty \leq 1$. $\Phi = \Phi_2$ is a 2-contractive homomorphism if and only if $|c|^2 \leq 2(1 - |f(x)|)/|\delta(f)|^2$ for any $f \in A$ with $\|f\|_\infty \leq 1$.

References

- [1] J. Agler *Rational dilation on an annulus*, Ann. of Math., **121** (1985), 537–564.
- [2] T. Ando *On a pair of commutative contractions*, Acta Sci. Math., **24** (1963), 88–90.
- [3] C. A. Berger *A strange dilation theorem*, Notices Amer. Math. Soc., **12** (1965), 590.
- [4] F. Bonsal and J. Duncan *Complete normed algebras*, Springer-Verlag, New York and Berlin, 1973.
- [5] Che-Chen Chu *Finite dimensional representation of a function algebra*, Ph.D. thesis in the University of Houston, 1992.
- [6] R. G. Douglas and V. I. Paulsen *Completely bounded maps and hypo-Dirichlet algebras*, Acta Sci. Math., **50** (1986), 143–157.
- [7] C. Foias and I. Suciú *Szegő-measures and spectral theory in Hilbert spaces*, Rev. Roum. Math. Pures. et Appl. **11** (1966), 147–159.
- [8] T. Gamelin *Uniform Algebras*, Prentice-Hall, Englewood Cliffs, New Jersey, 1969.
- [9] G. Misra *Curvature inequalities and extremal properties of bundle shifts*, J. Operator Theory **11** (1984), 305–317.
- [10] B. Sz. Nagy *Sur les contractions de l'espace de Hilbert*, Acta Sci. Math. **15** (1953), 87–92.
- [11] B. Sz. Nagy and C. Foias *Harmonic Analysis of Operators on Hilbert Space*, North Holland, New York, 1970.
- [12] T. Nakazi *A spectral dilation of some non-Dirichlet algebra*, Acta Sci. Math., **53** (1989), 119–122.
- [13] T. Nakazi *ρ -dilations and hypo-Dirichlet algebras*, Acta Sci. Math., **56** (1992), 175–181.
- [14] T. Nakazi and K. Takahashi *Two-dimensional representations of uniform algebras*, Proc. Amer. Math. Soc. **123** (1995), 2777–2784.
- [15] T. Nakazi and K. Okubo *ρ -Contraction and 2×2 matrix*, Linear Algebra Appl. **283** (1998), 165–169.
- [16] T. Nakazi and K. Okubo *Generalized numerical radius and unitary ρ -Dilation*, to appear in Math. Japonica.
- [17] S. Parrot *Unitary dilations for commuting contractions*, Pacific J. Math., **34** (1970), 481–490.
- [18] V. I. Paulsen *Completely Bounded Maps and Dilations*, Pitman Research Notes in Math. **146**, Longman, Wiley, New York, 1986.
- [19] G. Pisier *A polynomially bounded operator on Hilbert space which is not similar to a contraction*, J. Amer. Math. Soc. **10** (1997), 351–369.
- [20] H. Radjavi and P. Rosenthal *Invariant Subspaces*, Springer-Verlag, New York, 1973.
- [21] I. Suciú *Function Algebras*, Editura Academiei Republicii Socialiste România, Bucuresti, 1973.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN
E-mail address: nakazi@math.sci.hokudai.ac.jp