



Title	Compact Toeplitz operators with continuous symbols on weighted Bergman spaces
Author(s)	Nakazi, T.; Yoneda, R.
Citation	Hokkaido University Preprint Series in Mathematics, 446, 1-4
Issue Date	1999-2-1
DOI	10.14943/83592
Doc URL	http://hdl.handle.net/2115/69196
Type	bulletin (article)
File Information	pre446.pdf



[Instructions for use](#)

**Compact Toeplitz Operators
With Continuous Symbols
On Weighted Bergman Spaces**

T. Nakazi and R. Yoneda

Series #446. February 1999

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #421 D. Pei and T. Sano, The focal developable and the binormal indicatrix of a nonlightlike curve in Minkowski 3-space, 14 pages. 1998.
- #422 R. Kobayashi, J. A. Warren and W. C. Carter, Modeling grain boundaries using a phase field technique, 12 pages. 1998.
- #423 T. Tsukada, Reticular Legendrian Singularities, 28 pages. 1998.
- #424 A. N. Kirillov and M. Shimozono, A generalization of the Kostka-Foulkes polynomials, 37 pages. 1998.
- #425 M. Nakamura, Strichartz estimates for wave equations in the homogeneous Besov space, 17 pages. 1998.
- #426 A. Arai, On the essential spectra of quantum field Hamiltonians, 18 pages. 1998.
- #427 T. Sano, Bifurcations of affine invariants for one parameter family of generic convex plane curves, 11 pages. 1998.
- #428 F. Hiroshima, Ground states of a model in quantum electrodynamics, 48 pages. 1998.
- #429 F. Hiroshima, Uniqueness of the ground state of a model in quantum electrodynamics: A functional integral approach, 32 pages. 1998.
- #430 J. F. Van Diejen and A. N. Kirillov, Formulas for q -spherical functions using inverse scattering theory of reflectionless Jacobi operators, 33 pages. 1998.
- #431 G. Ishikawa, Determinacy, transversality and Lagrange stability, 13 pages. 1998.
- #432 T. Yoshida, Categorical aspects of generating functions(II): Operations on categories and functors, 65 pages. 1998.
- #433 K. Ito, Loss of convexity of compact hypersurfaces moved by surface diffusion, 20 pages. 1998.
- #434 Y. Shimizu, L^∞ -estimate of first-order space derivatives of Stokes flow in a half space, 22 pages. 1998.
- #435 T. Uemura, Morita-Mumford classes on finite cyclic subgroups of the mapping class group of closed surfaces, 14 pages. 1998.
- #436 A. Inoue and H. Kikuchi, Abel-Tauber theorems for Hankel and Fourier transforms and a problem of Boas, 20 pages. 1998.
- #437 T. Nakazi and T. Osawa, Finite rank intermediate Hankel operators on the Bergman space, 16 pages. 1998.
- #438 R. Yoneda, Compact Toeplitz operators on Bergman spaces, 14 pages. 1998.
- #439 A. Inoue and Y. Kasahara, On the asymptotic behavior of the prediction error of a stationary process, 12 pages. 1998.
- #440 A. Arai, M. Hirokawa and F. Hiroshima, On the absence of eigenvectors of Hamiltonians in a class of massless quantum field models without infrared cutoff, 26 pages. 1998.
- #441 A. N. Kirillov, A. Schilling and M. Shimozono, A bijection between Littlewood-Richardson tableaux and rigged configurations, 66 pages. 1999.
- #442 M. Tsujii, Absolutely continuous invariant measures for expanding piecewise linear maps, 28 pages. 1999.
- #443 T. Nakazi, Some special bounded homomorphisms of a uniform, 10 pages. 1999.
- #444 H. Ishii and T. Mikami, A two dimensional random crystalline algorithm for Gauss curvature flow, 21 pages. 1999.
- #445 A. Arai, Essential spectrum of a self-adjoint operator on an abstract Hilbert space of Fock type and applications to quantum field Hamiltonians, 24 pages. 1999.

Compact Toeplitz Operators With Continuous Symbols On Weighted Bergman Spaces

by

Takahiko Nakazi*

and

Rikio Yoneda

Department of Mathematics
Faculty of Science
Hokkaido University
Sapporo 060-0810, Japan

AMS Subject Classification (1991) ; 47 B 35, 47 B 07

Key Words and Phrases ; Bergman space, Toeplitz operator, continuous function, compact operator, weight

*This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education

Abstract. Let $L_a^2(D, d\sigma d\theta/2\pi)$ be a complete weighted Bergman space on the open unit disc D where $d\sigma$ is a positive finite Borel measure on $[0, 1)$. We show the following : When ϕ is a continuous function on the closed unit disc \bar{D} , T_ϕ is compact if and only if $\phi = 0$ on ∂D .

Let D be the open unit disc and $d\sigma$ be a positive finite Borel measure on $[0, 1)$. Let $L_a^2 = L_a^2(D, d\sigma d\theta/2\pi)$ be a weighted Bergman space on D , that is, L_a^2 consists of analytic functions f in D with

$$\|f\|_2^2 = \int_D |f(re^{i\theta})|^2 d\sigma d\theta/2\pi < \infty.$$

When L_a^2 is closed, P denotes the orthogonal projection from $L^2 = L^2(D, d\sigma d\theta/2\pi)$ onto L_a^2 . For ϕ in $L^\infty = L^\infty(D, d\sigma d\theta/2\pi)$, we consider the Toeplitz operator $T_\phi : L_a^2 \rightarrow L_a^2$ defined by $T_\phi f = P(\phi f)$, $f \in L_a^2$. We show the following theorem in this paper. For the Bergman space (that is, $d\sigma = 2rdr$), Theorem is well known in [5,p107] and [1]. When $d\sigma = (1 - r^2)^\alpha dr$ ($-1 < \alpha < \infty$), Theorem is also known in [3] and [4]. However that argument does not work for the general situation. We need a new idea in order to prove Theorem. $H = H(D)$ denotes the set of all analytic functions on D .

Theorem. *Suppose that $L_a^2 = L_a^2(D, d\sigma d\theta/2\pi)$ is complete. When ϕ is a continuous function on the closed unit disc \bar{D} , T_ϕ is compact if and only if $\phi = 0$ on ∂D .*

In order to prove Theorem, we need three lemmas.

Lemma 1. *L_a^2 is complete if and only if $\sigma([\varepsilon, 1)) > 0$ for some ε with $0 \leq \varepsilon < 1$.*

Proof. For $a \in D$, put

$$s(\mu, a) = \inf \left\{ \int_D |f|^2 d\mu ; f \in H \text{ and } f(a) = 1 \right\}$$

where H is a set of all analytic functions on D and $d\mu = d\sigma d\theta/2\pi$. (1) of Corollary 1 in [2] is valid for $s(\mu, a)$ instead of $S(\mu, a)$. When $(\text{supp}\mu) \cap D$ is a uniqueness set for H , by (1) of Theorem 8 in [2], L_a^2 is complete if and only if for all compact sets K in D $\int_K \log s(\mu, a) r dr d\theta/\pi > -\infty$. If σ is not a zero measure, then $(\text{supp}\mu) \cap D$ is a uniqueness set for H and so these imply the lemma.

Lemma 2. If $\sigma([\varepsilon, 1]) > 0$ for every ε with $0 \leq \varepsilon < 1$, then

$$\lim_{n \rightarrow \infty} \frac{\int_0^\varepsilon r^n d\sigma}{\int_\varepsilon^1 r^n d\sigma} = 0 \quad (0 \leq \varepsilon < 1).$$

Proof. When δ is a positive constant with $\varepsilon + \delta < 1$, the following inequality holds.

$$\begin{aligned} \frac{\int_0^\varepsilon r^n d\sigma}{\int_\varepsilon^1 r^n d\sigma} &\leq \frac{\sigma([0, \varepsilon])}{\int_\varepsilon^1 \left(\frac{r}{\varepsilon}\right)^n d\sigma} \leq \frac{\sigma([0, \varepsilon])}{\int_{\varepsilon+\delta}^1 \left(\frac{r}{\varepsilon}\right)^n d\sigma} \\ &\leq \frac{\sigma([0, \varepsilon])}{\left(\frac{\varepsilon + \delta}{\varepsilon}\right)^n \sigma([\varepsilon + \delta, 1])} \quad (0 < \varepsilon < 1). \end{aligned}$$

Since they are positive and $\lim_{n \rightarrow \infty} \{(\varepsilon + \delta)/\varepsilon\}^n = \infty$, $\lim_{n \rightarrow \infty} (\int_0^\varepsilon r^n d\sigma / \int_\varepsilon^1 r^n d\sigma) = 0$.

Lemma 3. If for every ε with $0 \leq \varepsilon < 1$,

$$\int_\varepsilon^1 r^n d\sigma > 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\int_0^\varepsilon r^n d\sigma}{\int_\varepsilon^1 r^n d\sigma} = 0,$$

then for any nonnegative ℓ

$$\lim_{n \rightarrow \infty} \frac{\int_0^1 r^{n+\ell} d\sigma}{\int_0^1 r^n d\sigma} = 1.$$

Proof. For every ε with $0 \leq \varepsilon < 1$, the following inequality holds.

$$\begin{aligned} 1 &\geq \frac{\int_0^1 r^{n+\ell} d\sigma}{\int_0^1 r^n d\sigma} = \frac{\int_0^\varepsilon r^{n+\ell} d\sigma + \int_\varepsilon^1 r^{n+\ell} d\sigma}{\int_0^\varepsilon r^n d\sigma + \int_\varepsilon^1 r^n d\sigma} \\ &\geq \frac{\varepsilon^\ell \int_\varepsilon^1 r^n d\sigma}{\int_\varepsilon^1 r^n d\sigma + \int_0^\varepsilon r^n d\sigma} \\ &= \frac{\varepsilon^\ell}{1 + \frac{\int_0^\varepsilon r^n d\sigma}{\int_\varepsilon^1 r^n d\sigma}} \end{aligned}$$

because $\int_{\varepsilon}^1 r^n d\sigma > 0$ and $\ell \geq 0$. Thus $\lim_{n \rightarrow \infty} \frac{\int_0^1 r^{n+\ell} d\sigma}{\int_0^1 r^n d\sigma} \geq \varepsilon^\ell$ and as $\varepsilon \rightarrow 1$ we show the lemma.

Proof of Theorem. Suppose $\phi(re^{i\theta}) = \sum_{j=-\infty}^{\infty} \phi_j(r)e^{ij\theta}$ is continuous on \bar{D} where

$$\phi_j(r) = \int_0^{2\pi} \phi(re^{i\theta})e^{-ij\theta} d\theta/2\pi$$

for $j = 0, \pm 1, \pm 2, \dots$. Then $\phi_j(r)$ is continuous on $[0,1]$ for any j . Put

$$\begin{aligned} e_n(re^{i\theta}) &= a_n r^n e^{in\theta} \\ &= r^n e^{in\theta} / \sqrt{\int_0^1 r^{2n} d\sigma} \end{aligned}$$

for $n \geq 0$, then $\{e_n\}$ is an orthonormal basis in L_a^2 . For each j , put

$$\Phi_j(re^{i\theta}) = r^{|j|} e^{-ij\theta} \phi(re^{i\theta}).$$

Then $T_{\Phi_j} = T_{r^{|j|}e^{-ij\theta}}T_\phi$ for $j \geq 0$ and $T_{\Phi_j} = T_\phi T_{r^{|j|}e^{-ij\theta}}$ for $j < 0$. If T_ϕ is compact, then T_{Φ_j} is also compact for any j . For each j , if $n \geq 0$, then

$$|\langle T_{\Phi_j} e_n, e_n \rangle| \leq \|T_{\Phi_j} e_n\|_2 \|e_n\|_2 = \|T_{\Phi_j} e_n\|_2.$$

Since T_{Φ_j} is compact for each j and $e_n \rightarrow 0$ ($n \rightarrow \infty$) weakly, $\|T_{\Phi_j} e_n\|_2 \rightarrow 0$ ($n \rightarrow \infty$) and so $\langle T_{\Phi_j} e_n, e_n \rangle \rightarrow 0$ ($n \rightarrow \infty$). For each j ,

$$\begin{aligned} \langle T_{\Phi_j} e_n, e_n \rangle &= \int_0^{2\pi} \int_0^1 \phi(re^{i\theta}) r^{|j|} e^{-ij\theta} a_n^2 r^{2n} d\sigma d\theta / 2\pi \\ &= a_n^2 \int_0^1 \phi_j(r) r^{|j|+2n} d\sigma \end{aligned}$$

and then $\lim_{n \rightarrow \infty} a_n^2 \int_0^1 \phi_j(r) r^{|j|+2n} d\sigma = 0$. By Lemma 1, $\sigma([\varepsilon, 1]) > 0$ for some ε with $0 \leq \varepsilon < 1$ and hence $\sigma([\varepsilon, 1]) > 0$ for every $\varepsilon < 1$. Hence by Lemma 2,

$$\lim_{n \rightarrow \infty} \frac{\int_0^\varepsilon r^{2n} d\sigma}{\int_\varepsilon^1 r^{2n} d\sigma} = 0 \quad \text{for } (0 \leq \varepsilon < 1).$$

Then by Lemma 3 for any integer j

$$\lim_{n \rightarrow \infty} a_n^2 \int_0^1 r^{|j|+2n} d\sigma = 1.$$

Since $\phi_j(r)$ is continuous on $[0,1]$, we can approximate $\phi_j(r)$ uniformly by polynomials $\sum_{t=0}^k c_t r^t$. Since $\lim_{n \rightarrow \infty} a_n^2 \int_0^1 r^{|j|+2n} d\sigma = 1$ for any j ,

$$\lim_{n \rightarrow \infty} a_n^2 \int_0^1 \left(\sum_{t=0}^k c_t r^t \right) r^{|j|+2n} d\sigma = \sum_{t=0}^k c_t$$

and so

$$\lim_{n \rightarrow \infty} a_n^2 \int_0^1 \phi_j(r) r^{|j|+2n} d\sigma = \phi_j(1).$$

Thus $\phi_j(1) = 0$ for any j because $\lim_{n \rightarrow \infty} a_n^2 \int_0^1 \phi_j(r) r^{|j|+2n} d\sigma = 0$, and hence $\phi = 0$ on ∂D .

Conversely suppose $\phi = 0$ on ∂D . Then we may assume that the support set of ϕ is compact in D . In order to show the compactness of T_ϕ , it is sufficient to show that if $h_n \rightarrow 0$ weakly ($n \rightarrow \infty$) in L_a^2 then $h_n \rightarrow 0$ uniformly on $\text{supp } \phi$. By hypothesis on σ , any point $z \in D$ has a bounded point evaluation for L_a^2 because (1) of Corollary 1 in [1] is valid for $s(\mu, a)$ instead of $S(\mu, a)$ and $r(\mu, a)s(\mu, a) = 1$ ($a \in D$). Hence $h_n(z) \rightarrow 0$. By the boundedness of analytic functions on $\text{supp } \phi$ and the uniform boundedness principle, $h_n \rightarrow 0$ uniformly on $\text{supp } \phi$.

References

1. S.Axler and D.C.Zheng, Compact operators via the Berezin transform, Indiana Univ.Math.J. Vol. 47(1998), 387-400.
2. T.Nakazi and M.Yamada, Riesz's functions in weighted Hardy and Bergman spaces, Can.J.Math. Vol. 48(1996), 930-945.
3. K.Stroethoff, Compact Toeplitz operators on Bergman spaces, Math.Proc.Cambridge Phi.Soc. Vol. 124(1998), 151-160.
4. K.Stroethoff, Compact Toeplitz operators on weighted harmonic Bergman spaces, J. Australian Math.Soc.Series A-Pure Math.and Statistics Vol. 64(1998), 136-148.
5. K.Zhu, Operator Theory in Function Spaces, Dekker, New York, 1990.