Title	Remarks on the uniqueness of bounded solutions of the NavierStokes equations
Author(s)	Giga, Y.; Inui, K.; Kato, J.; Matsui, S.
Citation	Hokkaido University Preprint Series in Mathematics, 497, 1-4
Issue Date	2000-10-1
DOI	10.14943/83643
Doc URL	http://hdl.handle.net/2115/69247
Type	bulletin (article)
File Information	pre497.pdf

[Instructions for use](#)

**Remarks on the uniqueness of bounded solutions
of the Navier-Stokes equations**

**Yoshikazu Giga, Katsuya Inui, Jun Kato
and Shin'ya Matsui**

Series #497. October 2000

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #472 Y. Giga and M.-H. Sato, A level set approach to semicontinuous viscosity solutions for Cauchy problems, 32 pages. 1999.
- #473 R. Yoneda, Characterization of Bloch space and Besov spaces by oscillations, 31 pages. 1999.
- #474 T. Nakazi, Norm inequalities for some singular integral operators, 13 pages. 1999.
- #475 A. Inoue and Y. Kasahara, Asymptotics for prediction errors of stationary processes with reflection positivity, 15 pages. 1999.
- #476 T. Sano, On affine parallels of generic plane curves, 8 pages. 1999.
- #477 T. Nakazi, On an invariant subspace whose common zero set is the zeros of some function, 11 pages. 1999.
- #478 M.-H. Giga and Y. Giga, Generalized motion by nonlocal curvature in the plane, 68 pages. 2000.
- #479 M.-H. Giga and Y. Giga, Crystalline and level set flow – Convergence of a crystalline algorithm for a general anisotropic curvature flow in the plane, 16 pages. 2000.
- #480 A. Arai and M. Hirokawa, Stability of ground states in sectors and its application to the Wigner-Weisskopf model, 16 pages. 2000.
- #481 T. Nakazi, Two dimensional Q -algebras, 11 pages. 2000.
- #482 N. H. Bingham and A. Inoue, Tauberian and Mercerian theorems for systems of kernels, 16 pages. 2000.
- #483 N. H. Bingham and A. Inoue, Abelian, Tauberian and Mercerian theorems for arithmetic sums, 29 pages. 2000.
- #484 I. A. Bogaevski and G. Ishikawa, Lagrange mappings of the first open Whitney umbrella, 22 pages. 2000.
- #485 A. Arai and H. Kawano, A class of deformations of the Schrödinger representation of the Heisenberg commutation relation and exact solution to a Heisenberg equation and a Schrödinger equation, 22 pages. 2000.
- #486 T. Nakazi, Functions in N_+ with the positive real parts on the boundary, 21 pages. 2000.
- #487 Y. Shibukawa, Classification of the R -operator, 36 pages. 2000.
- #488 A. Inoue, Asymptotic behaviour for partial autocorrelation functions of fractional ARIMA processes, 20 pages. 2000.
- #489 S. Ohtani, Construction of unramified Galois extensions over maximal abelian extensions of algebraic number fields, 14 pages. 2000.
- #490 T. Nakazi and T. Yamamoto, The real part of an outer function and a Helson-Szegö weight, 13 pages. 2000.
- #491 A. Yamagami, On Gouvêas conjecture on controlling the conductor, 11 pages. 2000.
- #492 I. Tsuda and M. Hatakeyama, Making sense of internal logic: Theory and a case study, 10 pages. 2000.
- #493 I. Tsuda, Towards an interpretation of dynamic neural activity in terms of chaotic dynamical systems, 73 pages. 2000.
- #494 T. Mikami, Optimal control for absolutely continuous stochastic processes and the mass transportation problem, 17 pages. 2000.
- #495 M. Arisawa and Y. Giga, Anisotropic curvature flow in a very thin domain, 21 pages. 2000.
- #496 T. Nakazi, Backward shift invariant subspaces in the bidisc, 9 pages. 2000.

Remarks on the uniqueness of bounded solutions of the Navier-Stokes equations

Yoshikazu Giga¹, Katsuya Inui², Jun Kato³ and Shin'ya Matsui⁴,

^{1,3} Department of Mathematics, Hokkaido University, Sapporo 060-0810, Japan

² Kozo Keikaku Engineering Inc. ITK Nabeyoko Bldg. 2F 4-30-12, Hon-cho, Nakano-ku, Tokyo, 164-0012 Japan

⁴ Faculty of Business Administration and Information Science, Hokkaido Information University, Nishi-Nopporo, Ebetsu 069-8585, Japan

1 Introduction

This is an announcement of our results on the uniqueness of a bounded solution $u = (u_1, \dots, u_n)$ of the Navier-Stokes equations:

$$(1) \quad u_t - \Delta u + (u, \nabla)u + \nabla p = 0, \quad \operatorname{div} u = 0 \quad \text{in } \mathbf{R}^n \times (0, T).$$

with initial data $u|_{t=0} = u_0$. The detailed argument as well as improvement will be published in [8].

It is by now well-known that for initial data $u_0 \in L^\infty(\mathbf{R}^n)$ the equation (1) admits a unique local-in-time (regular) solution u with

$$(2) \quad p = \sum_{i,j=1}^n R_i R_j u_i u_j,$$

where $R_j = (-\Delta)^{-1/2} \partial / \partial x_j$ is the Riesz operator [1], [11], [2], [6]. (Very recently, it turns out that this solution can be extended globally-in-time when the space dimension is two [7].) It is also well-known that for L^r -initial data ($n \leq r < \infty$) the equation (1) admits a

¹Party supported by the Grant-in-Aid for Scientific Research No.10304010, the Japan Society for the Promotion of Science

⁴Party supported by the Grant-in-Aid for Scientific Research No.11640211, the Japan Society for the Promotion of Science

unique local-in-time solution u with some p [5], [9], [4]... Because of decay at the space infinity of u the relation (2) follow (up to constant) a posteriori for L^r -data ($n \leq r < \infty$).

For L^∞ -initial data the constructed solution u is bounded and may not decay at the space infinity. So even if u solves (1) with some p the relation (2) may not follow. In fact, if we consider $u = g(t)$, then (u, p) always solves (1) with $p = -g'(t) \cdot x$ no matter what function g is. Here $\cdot$ denotes the inner product of $\mathbf{R}^n$. This says the solution u with a constant initial data is not unique without assuming (2). This example suggests that contrary to L^r -case ($n \leq r < \infty$) we need to impose some condition on p to derive uniqueness other than on u .

In this paper we give a new sufficient condition for the uniqueness of solution (u, p) when u is bounded.

Theorem. *Assume that (u, p) is a smooth solution of (1) in $\mathbf{R}^n \times (0, T)$ and that u is bounded. Assume that u is continuous on $[0, T)$ with values in BUC , the space of all bounded uniformly continuous functions. Let u_0 denote the initial data of u . If p is of the form*

$$(3) \quad p = \sum_{1 \leq i, j \leq n} R_i R_j \pi_{ij} + \pi_0$$

with some $\pi_{ij}, \pi_0 \in L^1_{loc}(0, T; L^\infty(\mathbf{R}^n))$, then $(u, \nabla p)$ is uniquely determined by u_0 and p fulfills (2) up to a constant.

Recently, the third author improved the result by weakening the assumption on pressure (3) by

$$(4) \quad p \in L^1_{loc}(0, T; BMO),$$

where BMO is the space of all functions of bounded mean oscillation. By the Calderón-Zygmund inequality (3) implies (4). By a theorem of A. Uchiyama [13] if a function $f \in BMO$ is compactly supported, then it is of the form $f = \sum_{1 \leq i, j \leq n} R_i R_j \eta_{ij} + \eta_0$ with some $\eta_{ij}, \eta_0 \in L^\infty(\mathbf{R}^n)$. Thus the difference of (3) and (4) is small.

The assumption of continuity on u is to determine initial data and can be weakened. The regularity assumption on (u, p) is also can be weakened by using a notion of weak solutions. The reader is referred to a recent paper by the third author [8] for improvement of Theorem.

Let us mention a few known results closely related to our uniqueness results. It was shown in [3] that if $n = 3$ and u with ∇u is bounded in $\mathbf{R}^3 \times (0, T)$, then the uniqueness holds provided that for some $C > 0$ and some $\varepsilon \in (0, 1)$ the inequality

$$(5) \quad |p(x, t)| \leq C(|x|^{1-\varepsilon} + 1)$$

holds for all $(x, t) \in \mathbf{R}^3 \times (0, T)$. Later it was shown in [12], [10] that if $n = 2, 3$ and ∇u is bounded in $\mathbf{R}^3 \times (0, T)$, then the uniqueness holds provided that (5) holds with $\varepsilon = n/2$. (The argument is valid for general n). Our assumptions (3) and (4) do not imply (5) so it is not comparable with those results.

2 Sketch of the proof

The main idea is to prove

$$(6) \quad \nabla p = \sum_{1 \leq i, j \leq n} \nabla R_i R_j u_i u_j$$

which yields (2) so that u solves that integral equation without p whose solution is known [6] to be unique.

A formal idea to prove (6) is to apply $R_i R_j$ to the first j -th equation and add with respect to j . Since $\sum_{j=1}^n R_j f_j = 0$ for divergence free $f = (f_1, \dots, f_n)$, we come to

$$(7) \quad \sum_{1 \leq j, \ell \leq n} R_i R_j (u_\ell \partial_{x_\ell} u_j) + \sum_{j=1}^n R_i R_j \partial_{x_j} p = 0, \quad i = 1, \dots, n,$$

where $\partial_{x_j} = \partial/\partial x_j$. Since $\operatorname{div} u = 0$ and $R_i R_j \partial_{x_\ell} = \partial_{x_i} R_j R_\ell$ the first term of (7) equals $\sum_{j, \ell} \partial_{x_i} R_j R_\ell u_j u_\ell$. Since $\sum_j R_i R_j \partial_{x_j} p = -\partial_{x_i} p$, (7) now yields (6). Unfortunately, the above argument is formal since it is not clear in what way we operate $R_i R_j$ to $\partial_{x_j} p$ when p is merely in BMO .

To overcome this inconvenience we approximate $R_i R_j$ by R_{ij}^ε defined by

$$R_{ij}^\varepsilon f = (\partial_{x_i} \partial_{x_j} k_\varepsilon) * f.$$

Here k_ε is obtained by a truncation of the fundamental solution k of $-\Delta$ and is defined by

$$k_\varepsilon = \psi_\varepsilon \lambda_\varepsilon k$$

with $\psi_\varepsilon(x) = \psi(x/\varepsilon)$, $\lambda_\varepsilon(x) = \lambda(\varepsilon x)$, where ψ is a radial function satisfying $\psi = 0$ for $|x| \leq 1$ and $\psi(x) = 1$ for $|x| \geq 2$ and $\lambda = 1 - \psi$.

Instead of applying $R_i R_j$ we apply R_{ij}^ε to (1) and send $\varepsilon \rightarrow 0$ which realizes the formal idea and obtain (6). A key convergence results is

$$(8) \quad R_{ij}^\varepsilon \partial_{x_j} g \rightarrow -\partial_{x_i} g \quad (1 \leq i \leq n)$$

in the sense of tempered distributions for $g \in BMO$. To prove Theorem it suffices to prove the above convergence for g of the form $g = \sum_{ij} R_i R_j \pi_{ij} + \pi_0$ with some π_{ij} and π_0 . The convergence (8) follows from the observation:

$$(9) \quad (-\Delta) k_\varepsilon * \psi \rightarrow \psi$$

in $\mathcal{H}^1$, the Hardy space for $\psi \in C_0^\infty(\mathbf{R}^n)$ satisfying $\int_{\mathbf{R}^n} \psi dx = 0$. Again to prove (8) for restricted g for Theorem it suffices to prove (9) in L^1 -sense and almost everywhere sense. We won't go into more detail in the present paper and refer to a paper [8] for the detail.

References

- [1] J. R. Cannon and G. H. Knightly, *A note on the Cauchy problem for the Navier-Stokes equations*, SIAM J. Appl. Math **18**(1970), 641-644.
- [2] M. Cannone, *Ondelettes, Panaproducts et Navier-Stokes*, Diclerot Editeur, Arts et Sciences, Paris-New York-Amsterdam (1995).
- [3] G. P. Galdi and P. Maremonti, *A uniqueness theorem for viscous fluid motions in exterior domeins*, Arch. Rational Mech. Anal. **91**(1986), 375-384.
- [4] Y. Giga, *Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes systems*, J. Differential Equations, **62**(1986), 186-212.
- [5] Y. Giga and T. Miyakawa, *Solutions in L_r to the Navier-Stokes initial value problem*, Arch. Rational Mech. Anal. **89**(1985), 267-281.
- [6] Y. Giga, K. Inui and S. Matsui, *On the Cauchy problem for the Navier-Stokes equations with nondecaying initial data*, Quaderni di Matematica, **4**(1999), 28-68.
- [7] Y. Giga, S. Matsui and O. Sawada, *Global existence of two-dimensional Navier-Stokes flow with nondecaying initial velocity*, preprint.
- [8] J. Kato, *On the uniqueness of nondecaying solutions of the Navier-Stokes equations*, preprint.
- [9] T. Kato, *Strong L^p -soulutions of the Navier-Stokes equation in $\mathbf{R}^n$ with applications to weak solutions*, Math. Z. **187**(1984), 471-480.
- [10] N. Kim and D. Chae, *On the uniqueness of the unbounded classical solutions of the Navier-Stokes and associated equations*, J. Math. Anal. Appl. **186**(1994), 91-96.
- [11] G. H. Knightly, *A Cauchy problem for the Navier-Stokes equations*, SIAM J. Math. Anal. **3**(1972), 506-511.
- [12] H. Okamoto, *A uniqueness theorem for the unbounded classical solutions of nonstationary Navier-Stokes equations in $\mathbf{R}^3$* , J. Math. Anal. Appl. **181**(1994), 473-482.
- [13] A. Uchiyama, *A constructive proof of the Fefferman-Stein decomposition of $BMO(\mathbf{R}^n)$* . Acta Math. **148**(1982), 215-241.