



Title	Ground state of the massless Nelson model without infrared cutoff in a non-Fock representation
Author(s)	Arai, A.
Citation	Hokkaido University Preprint Series in Mathematics, 502, 1-19
Issue Date	2000-11-1
DOI	10.14943/83648
Doc URL	http://hdl.handle.net/2115/69252
Type	bulletin (article)
File Information	pre502.pdf



[Instructions for use](#)

**Ground State of the Massless Nelson Model
Without Infrared Cutoff in a Non-Fock
Representation**

Asao Arai

Series #502. November 2000

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #478 M.-H. Giga and Y. Giga, Generalized motion by nonlocal curvature in the plane, 68 pages. 2000.
- #479 M.-H. Giga and Y. Giga, Crystalline and level set flow – Convergence of a crystalline algorithm for a general anisotropic curvature flow in the plane, 16 pages. 2000.
- #480 A. Arai and M. Hirokawa, Stability of ground states in sectors and its application to the Wigner-Weisskopf model, 16 pages. 2000.
- #481 T. Nakazi, Two dimensional Q -algebras, 11 pages. 2000.
- #482 N. H. Bingham and A. Inoue, Tauberian and Mercerian theorems for systems of kernels, 16 pages. 2000.
- #483 N. H. Bingham and A. Inoue, Abelian, Tauberian and Mercerian theorems for arithmetic sums, 29 pages. 2000.
- #484 I. A. Bogaevski and G. Ishikawa, Lagrange mappings of the first open Whitney umbrella, 22 pages. 2000.
- #485 A. Arai and H. Kawano, A class of deformations of the Schrödinger representation of the Heisenberg commutation relation and exact solution to a Heisenberg equation and a Schrödinger equation, 22 pages. 2000.
- #486 T. Nakazi, Functions in N_+ with the positive real parts on the boundary, 21 pages. 2000.
- #487 Y. Shibukawa, Classification of the R -operator, 36 pages. 2000.
- #488 A. Inoue, Asymptotic behaviour for partial autocorrelation functions of fractional ARIMA processes, 20 pages. 2000.
- #489 S. Ohtani, Construction of unramified Galois extensions over maximal abelian extensions of algebraic number fields, 14 pages. 2000.
- #490 T. Nakazi and T. Yamamoto, The real part of an outer function and a Helson-Szegö weight, 13 pages. 2000.
- #491 A. Yamagami, On Gouvêas conjecture on controlling the conductor, 11 pages. 2000.
- #492 I. Tsuda and M. Hatakeyama, Making sense of internal logic: Theory and a case study, 10 pages. 2000.
- #493 I. Tsuda, Towards an interpretation of dynamic neural activity in terms of chaotic dynamical systems, 73 pages. 2000.
- #494 T. Mikami, Optimal control for absolutely continuous stochastic processes and the mass transportation problem, 17 pages. 2000.
- #495 M. Arisawa and Y. Giga, Anisotropic curvature flow in a very thin domain, 21 pages. 2000.
- #496 T. Nakazi, Backward shift invariant subspaces in the bidisc, 9 pages. 2000.
- #497 Y. Giga, K. Inui, J. Kato and S. Matsui, Remarks on the uniqueness of bounded solutions of the Navier-Stokes equations, 4 pages. 2000.
- #498 Y. Giga, M. Paolini and P. Rybka, On the motion by singular interfacial energy, 21 pages. 2000.
- #499 J. Escher and Y. Giga, On a limiting motion and self-interactions of curves moved by the intermediate surface diffusion flow, 12 pages. 2000.
- #500 I. Tsuda and S. Kuroda, Cantor coding in the hippocampus, 20 pages. 2000.
- #501 M. Tsujii, Fat solenoidal attractors, 20 pages. 2000.

Ground State of the Massless Nelson Model Without Infrared Cutoff in a Non-Fock Representation

Asao Arai *

Department of Mathematics
Hokkaido University
Sapporo 060-0810
Japan

November 9, 2000

Abstract

We consider a model of quantum particles coupled to a massless quantum scalar field, called the massless Nelson model, *in a non-Fock representation* of the time-zero fields which satisfy the canonical commutation relations. We show that the model has a ground state for *all* values of the coupling constant *even in the case where no infrared cutoff is made*. The non-Fock representation used is inequivalent to the Fock one if no infrared cutoff is made.

Key words: Nelson's model, massless quantum field, infrared divergence, ground state, canonical commutation relations, non-Fock representation

1 Introduction

We consider a system of N quantum particles ($N \in \mathbb{N}$) moving on the d -dimensional Euclidean space \mathbb{R}^d ($d \in \mathbb{N}$) under the influence of an external potential $V : \mathbb{R}^d \rightarrow \mathbb{R}$ (Borel measurable) and coupled to a massless quantum scalar field. The model we discuss here is the so-called *massless Nelson model* [10]. The problem to which we address ourselves in this paper is that of existence of a ground state of the model *without infrared cutoff*.

It has been shown that the massless Nelson model *with infrared cutoff* has a ground state [14, 7]. A natural question to be asked next is if the model *without infrared cutoff* has a ground state or not.

*Supported by the Grant-In-Aid No. 11440036 for Scientific Research from the Ministry of Education, Science, Sports and Culture, Japan.

We remark that, generally speaking, there is some subtlety on the existence of ground states of massless quantum field models, which is due to possible infrared divergences. Indeed, in a class of models which describe interactions of particles and massless quantum fields, it is proved or suggested that, if the time-zero fields are given by the Fock representation and no infrared cutoff is made in the interactions, then the models have no ground states, although it may depend on the strength of the parameters contained in the Hamiltonians [6, 13, 2, 3]. On the other hand, the Pauli-Fierz model in non-relativistic quantum electrodynamics has a ground state in the Fock representation of time-zero fields even if no infrared cutoff is made. [4, 8].

As for the massless Nelson model without infrared cutoff, Lörinczi, Spohn and Minlos [9] recently proved that, in the case $d = 3$ and $N = 1$, it has no ground state within the Fock space where the time-zero fields are given by the usual Fock representation of the canonical commutation relations (CCR) indexed by $\mathcal{S}_{\text{real}}(\mathbb{R}^d)$, the space of real-valued, rapidly decreasing C^∞ -functions on \mathbb{R}^d .

The purpose of this paper is to point out that, if we consider the massless Nelson model in a non-Fock representation of the CCR for time-zero fields, then it has a ground state even in the case where no infrared cutoff is made. This new representation of the massless Nelson model is inequivalent to the Fock one if no infrared cutoff is made.

This paper is organized as follows. In Section 2 we first briefly review the Nelson model in the standard form in which the time-zero fields are given by the Fock representation of the CCR indexed by $\mathcal{S}_{\text{real}}(\mathbb{R}^d)$. We call it the “standard Nelson model” (SNM). Then we give an algebraic characterization of the Nelson model in a way independent of the choice of representations of the time-zero fields and derive rigorously field equations in a weak sense. To the author’s best knowledge, field equations of the Nelson model has not been rigorously established so far (only formal or classical ones are available).

In Section 3 we define the Nelson model in a non-Fock representation of the time-zero fields.

In the last section, we prove that, under suitable hypotheses, the massless Nelson model introduced in Section 3 has a ground state even in the case where no infrared cutoff is made.

2 Algebraic characterization and field equations of the Nelson model

In this section we first recall the SNM. Then we present an algebraic characterization of the Nelson model in an abstract manner. Finally we derive rigorously field equations of the abstract Nelson model.

2.1 The SNM

The coordinate of the configuration space \mathbb{R}^{dN} of the N particles is denoted $q = (q_1, \dots, q_N) \in \mathbb{R}^{dN}$ with $q_j := (q_{j1}, \dots, q_{jd}) \in \mathbb{R}^d$ ($j = 1, \dots, N$). Let $D_{j\mu}$ ($j = 1, \dots, N$, $\mu = 1, \dots, d$) be the generalized partial differential operator in the variable $q_{j\mu}$, so that the momentum operator of the j -th particle is given by $p_j := (p_{j1}, \dots, p_{jd})$ with $p_{j\mu} := -iD_{j\mu}$.

The Hamiltonian of the particle system is then given by the Schrödinger operator

$$H_p := \sum_{j=1}^N \frac{p_j^2}{2m_j} + V, \quad (2.1)$$

acting on $L^2(\mathbb{R}^{dN})$, where $m_j > 0$ is the mass of the j -th particle and $p_j^2 = \sum_{\mu=1}^d p_{j\mu}^2 = -\Delta_{q_j}$ is the d -dimensional generalized Laplacian in the variable q_j .

For a linear operator T , we denote its domain by $D(T)$.

We assume the following:

(H.1) The operator H_p is self-adjoint on its natural domain $D(H_p) = \cap_{j=1}^N D(\Delta_{q_j}) \cap D(V)$ and bounded from below.

The Hilbert space for state vectors of the quantum scalar field of the SNM is given by

$$\mathcal{F}_b := \bigoplus_{n=0}^{\infty} \otimes_s^n L^2(\mathbb{R}^d), \quad (2.2)$$

the Boson Fock space over $L^2(\mathbb{R}^d)$, where $\otimes_s^n L^2(\mathbb{R}^d)$ is the symmetric tensor product of $L^2(\mathbb{R}^d)$ ($\otimes_s^0 L^2(\mathbb{R}^d) := \mathbb{C}$).

Let ω be a nonnegative Borel measurable function on \mathbb{R}^d such that $0 < \omega(k) < \infty$ for almost everywhere (a.e.) $k \in \mathbb{R}^d$ with respect to the d -dimensional Lebesgue measure and

$$\omega(k) = \omega(-k) \quad \text{a.e. } k. \quad (2.3)$$

For a.e. $k \in \mathbb{R}^d$, $\omega(k)$ physically means the energy of one free boson with momentum k . The function ω defines a nonnegative self-adjoint multiplication operator on $L^2(\mathbb{R}^d)$ which is injective.

Remark 2.1 A physical example for ω is given by

$$\omega_m(k) := \sqrt{k^2 + m^2}, \quad k \in \mathbb{R}^d, \quad (2.4)$$

with $m \geq 0$ a constant denoting the mass of one boson. We are interested in the massless case $m = 0$: $\omega_0(k) = |k|$

The free Hamiltonian of the quantum scalar field is defined by

$$H_b := d\Gamma(\omega), \quad (2.5)$$

the second quantization of ω [11, §X.7].

We denote the annihilation operators on \mathcal{F}_b by $a(f) = \int_{\mathbb{R}^d} a(k) f(k)^* dk$ ($f \in L^2(\mathbb{R}^d)$) [11, §X.7]. The symmetric operator

$$\Phi_s(f) := \frac{1}{\sqrt{2}}(a(f)^* + a(f)), \quad (2.6)$$

called the Segal field operator, is essentially self-adjoint [11, §X.7]. We denote its closure by the same symbol.

Let $\mathcal{S}'_{\text{real}}(\mathbb{R}^d)$ be the set of real tempered distributions on \mathbb{R}^d and, for each $s \in \mathbb{R}$,

$$H_\omega^s := \{f \in \mathcal{S}'_{\text{real}}(\mathbb{R}^d) \mid \omega^s \hat{f} \in L^2(\mathbb{R}^d)\}, \quad (2.7)$$

where

$$\hat{f}(k) := \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} f(x) e^{-ikx} dx \quad (2.8)$$

is the Fourier transform of f .

For $f \in H_\omega^{-1/2}$ and $g \in H_\omega^{1/2}$, we can define

$$\phi_F(f) := \Phi_S \left(\frac{\hat{f}}{\sqrt{\omega}} \right), \quad \pi_F(g) := \Phi_S(i\sqrt{\omega}\hat{g}). \quad (2.9)$$

Let

$$\mathcal{F}_0 := \{\psi = \{\psi^{(n)}\}_{n=0}^\infty \in \mathcal{F}_b \mid \psi^{(n)} = 0 \text{ for all but finitely many } n\text{'s}\}, \quad (2.10)$$

called the subspace of finite-particle vectors. Then, for all $f \in H_\omega^{-1/2}$ and $g \in H_\omega^{1/2}$, $\phi_F(f)$ and $\pi_F(g)$ leave \mathcal{F}_0 invariant satisfying the CCR

$$[\phi_F(f), \pi_F(g)] = i \int_{\mathbb{R}^d} \hat{f}(k) \hat{g}(k) dk, \quad (2.11)$$

$$[\phi_F(f), \phi_F(f')] = 0, \quad [\pi_F(g), \pi_F(g')] = 0, \quad f, f' \in H_\omega^{-1/2}, g, g' \in H_\omega^{1/2}, \quad (2.12)$$

on \mathcal{F}_0 . Namely, $\{\phi_F(f), \pi_F(g) \mid f \in H_\omega^{-1/2}, g \in H_\omega^{1/2}\}$ gives a representation of the CCR. This representation is called the Fock representation of the CCR. In the SNM, $\phi_F(f)$ and $\pi_F(g)$ are taken to be the time-zero fields:

The Hilbert space for the SNM is

$$\mathcal{H} := L^2(\mathbb{R}^{dN}) \otimes \mathcal{F}_b. \quad (2.13)$$

As usual, we freely use the natural identification of \mathcal{H} with $\int_{\mathbb{R}^{dN}}^\oplus \mathcal{F}_b dq$, the constant fibre direct integral with base space (\mathbb{R}^{dN}, dq) and fibre \mathcal{F}_b [12, §XIII.16].

In what follows, for notational simplicity, a decomposable operator $A = \int_{\mathbb{R}^{dN}}^\oplus A(q) dq$ on \mathcal{H} with fibre $A(q)$ (which is an operator on \mathcal{F}_b for each $q \in \mathbb{R}^{dN}$) is denoted $A(q)$ also.

To describe an interaction between the particles and the quantum scalar field, we fix distributions $\rho_j \in \mathcal{S}'_{\text{real}}(\mathbb{R}^d)$, $j = 1, \dots, N$, which satisfy the following:

(H.2) For $j = 1, \dots, N$,

$$\rho_j \in H_\omega^s, \quad s = -1/2, -1. \quad (2.14)$$

The interaction of the particles and the quantum field in the SNM is given by the operator

$$H_I^F := \sum_{j=1}^N \Phi_S \left(e^{-ikq_j} \frac{\hat{\rho}_j^*}{\sqrt{\omega}} \right) \quad (2.15)$$

acting in \mathcal{H} . Formally we have $H_I^F = \sum_{j=1}^N \int_{\mathbb{R}^d} \phi_F(q_j - x) \rho_j(x) dx$, where

$$\phi_F(x) := \int_{\mathbb{R}^d} \frac{1}{\sqrt{2(2\pi)^d \omega(k)}} \{a(k)^* e^{-ikx} + a(k) e^{ikx}\} dk.$$

The total Hamiltonian of the SNM is defined by

$$H_{\text{SNM}} := H_0 + \lambda H_I^{\text{F}}, \quad (2.16)$$

where

$$H_0 := H_p + H_b \quad (2.17)$$

and $\lambda \in \mathbb{R}$ denotes the coupling constant of the model.

Proposition 2.1 *Assume (H.1) and (H.2). Then H_{SNM} is self-adjoint with $D(H_{\text{SNM}}) = D(H_0)$ and bounded from below. Moreover, H_{SNM} is essentially self-adjoint on each core of H_0*

Proof. A simple application of [1, Proposition B.2] which employs the Kato-Rellich theorem (cf. [6]). ■

In summary, the SNM is characterized in terms of the Hamiltonian H_{SNM} and the time-zero-fields $\{\phi_{\text{F}}(f), \pi_{\text{F}}(g) | f \in H_{\omega}^{-1/2}, g \in H_{\omega}^{1/2}\}$.

2.2 An abstract definition of the Nelson model

As is seen above, the SNM uses the Fock representation of the CCR to give its time-zero fields and its Hamiltonian. We want to define the Nelson model in a way independent of the choice of representations of the CCR for time-zero fields. A natural manner for this is to use commutation relations fulfilled by observables, i.e., to find a possible Lie algebraic structure. We shall take commutation relations in a weak sense.

We denote the inner product and the norm of a Hilbert space \mathcal{X} by $\langle \cdot, \cdot \rangle_{\mathcal{X}}$ and $\| \cdot \|_{\mathcal{X}}$ respectively. But, if there is no danger of confusion, then we simply write them as $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$.

Definition 2.2 Let A and B be densely defined linear operators on a Hilbert space \mathcal{X} and \mathcal{D} be a subspace of \mathcal{X} such that $\mathcal{D} \subset D(A) \cap D(B) \cap D(A^*) \cap D(B^*)$. Then we define a quadratic form $[A, B]_{\text{w}}^{\mathcal{D}}$ by

$$[A, B]_{\text{w}}^{\mathcal{D}}(\psi, \phi) := \langle A^* \psi, B \phi \rangle - \langle B^* \psi, A \phi \rangle, \quad \psi, \phi \in \mathcal{D}.$$

(“w” means “weak”.)

Remark 2.2 If A and B are bounded on \mathcal{X} with $D(A) = D(B) = \mathcal{X}$, then, for all dense subspaces \mathcal{D} of \mathcal{X} , $[A, B]_{\text{w}}^{\mathcal{D}}(\psi, \phi) = \langle \psi, [A, B] \phi \rangle$ for all $\psi, \phi \in \mathcal{D}$, where $[A, B] := AB - BA$ (the usual commutator).

Let \mathcal{F} be a Hilbert space and

$$\mathcal{K} := L^2(\mathbb{R}^{dN}) \otimes \mathcal{F} = \int_{\mathbb{R}^{dN}}^{\oplus} \mathcal{F} dq. \quad (2.18)$$

Definition 2.3 Assume (H.1) and (H.2). Let s_0 and s_1 be real constants. A Nelson model is a set $\mathbb{M}_{\text{Nelson}} := \{\mathcal{K}, \mathcal{D}, L_{\lambda}, \{\phi(f), \pi(g) | f \in H_{\omega}^{s_0}, g \in H_{\omega}^{s_1}\}\}$ having the following properties:

(N.1) \mathcal{D} is a dense subspace of \mathcal{K} .

(N.2) L_λ is a symmetric operator on \mathcal{K} and the operator

$$H_{\text{NM}} := H_p + L_\lambda \quad (2.19)$$

is self-adjoint ($D(H_{\text{NM}}) := D(H_p) \cap D(L_\lambda)$). We call H_{NM} the Hamiltonian of the Nelson model $\mathbb{M}_{\text{Nelson}}$.

(N.3) The function ω is such that

$$\mathcal{S}_{\text{real}}(\mathbb{R}^d) \subset H_\omega^{s_0} \cap H_\omega^{s_0+2} \cap H_\omega^{s_1} \cap H_\omega^1 \quad (2.20)$$

and, for all $j = 1, \dots, N, \mu = 1, \dots, d$,

$$D_\mu \rho_j \in H_\omega^{s_0}, \quad (2.21)$$

where D_μ ($\mu = 1, \dots, d$) is the generalized partial differential operator in the μ -th variable x_μ in $x = (x_1, \dots, x_d) \in \mathbb{R}^d$.

(N.4) $\{\mathcal{K}, \mathcal{D}, \{\phi(f), \pi(f) | f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}\}$ is a representation of the CCR indexed by $\mathcal{S}_{\text{real}}(\mathbb{R}^d)$. Namely, for all $f, g \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$, $\phi(f)$ and $\pi(f)$ are self-adjoint operators on \mathcal{K} with $\mathcal{D} \subset D(\phi(f)\pi(g)) \cap D(\pi(f)\phi(g)) \cap D(\phi(f)\phi(g)) \cap D(\pi(f)\pi(g))$ satisfying the CCR

$$[\phi(f), \pi(g)] = i \int_{\mathbb{R}^d} f(x)g(x)dx, \quad (2.22)$$

$$[\phi(f), \phi(g)] = 0, \quad [\pi(f), \pi(g)] = 0, \quad (2.23)$$

on \mathcal{D} and the linearity:

$$\phi(af + bg) = a\phi(f) + b\phi(g), \quad \pi(af + bg) = a\pi(f) + b\pi(g), \quad a, b \in \mathbb{R},$$

on \mathcal{D} .

(N.5) Let $X = q_{j\mu}, p_{j\mu}, Y = \phi(f), \pi(f), j = 1, \dots, N, \mu = 1, \dots, d, f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$. There exists a dense subspace \mathcal{E} of \mathcal{K} such that

$$\mathcal{E} \subset D(X) \cap D(Y) \cap D(L_\lambda) \cap D(H_p) \quad (2.24)$$

and the following relations hold in the sense of quadratic form on \mathcal{E} :

$$[L_\lambda, q_{j\mu}]_{\mathfrak{w}}^{\mathcal{E}} = 0, \quad [L_\lambda, p_{j\mu}]_{\mathfrak{w}}^{\mathcal{E}} = i\lambda\phi(D_\mu \rho_j(q_j - \cdot)),$$

$$[L_\lambda, \phi(f)]_{\mathfrak{w}}^{\mathcal{E}} = -i\pi(f),$$

$$[L_\lambda, \pi(f)]_{\mathfrak{w}}^{\mathcal{E}} = i\phi(\omega(-i\nabla)^2 f) + i\lambda \sum_{j=1}^N (\rho_j * f)(q_j),$$

$$[X, Y]_{\mathfrak{w}}^{\mathcal{E}} = 0, \quad [H_p, Y]_{\mathfrak{w}}^{\mathcal{E}} = 0,$$

where $\nabla = (D_1, \dots, D_d)$ and $\rho_j * f$ is the convolution of ρ_j and f : $(\rho_j * f)(x) := \int_{\mathbb{R}^d} \rho_j(x-y)f(y)dy$.

Remark 2.3 (i) We have

$$(\omega(-i\nabla)^2 f)(x) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} \omega(k)^2 \hat{f}(k) e^{ikx} dk, \quad f \in \mathcal{S}(\mathbb{R}^d).$$

Hence (N.3) implies that

$$\omega(-i\nabla)^2 f \in H_\omega^{s_0},$$

so that $\phi(\omega(-i\nabla)^2 f)$ is defined.

(ii) It follows from (H.2) and (N.3) that, for all $f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$, $\rho_j * f$ is a bounded continuous function on \mathbb{R}^d , so that $\rho_j * f(q_j)$ is a bounded self-adjoint multiplication operator with $D(\rho_j * f(q_j)) = \mathcal{K}$.

Let us show that the SNM is indeed a Nelson model in the above sense. We introduce

$$\mathcal{F}_{\omega, \text{fin}} := \mathcal{L}\{\Omega_0, a(f_1)^* \cdots a(f_n)^* \Omega_0 | n \in \mathbb{N}, f_j \in D(\omega), j = 1, \dots, n\}, \quad (2.25)$$

where $\Omega_0 := \{1, 0, 0, \dots\} \in \mathcal{F}_b$ is the Fock vacuum and $\mathcal{L}\{\dots\}$ means the subspace algebraically spanned by all the vectors in the set $\{\dots\}$, and

$$\mathcal{D}_0 := C_0^\infty(\mathbb{R}^{dN}) \otimes_{\text{alg}} \mathcal{F}_{\omega, \text{fin}}, \quad (2.26)$$

where \otimes_{alg} denotes algebraic tensor product.

Proposition 2.4 Assume (H.1) and (H.2). Suppose that

$$\int_{|q| \leq R} |V(q)|^2 dq < \infty, \quad \forall R > 0, \quad (2.27)$$

$$\mathcal{S}_{\text{real}}(\mathbb{R}^d) \subset H_\omega^{3/2} \cap H_\omega^{-1/2}, \quad (2.28)$$

and, for $j = 1, \dots, N, \mu = 1, \dots, d$,

$$D_\mu \rho_j \in H_\omega^{-1/2}. \quad (2.29)$$

Let

$$L_\lambda^{\text{SNM}} := H_b + \lambda H_I^{\text{F}}. \quad (2.30)$$

Then

$$\mathbb{M}_{\text{SNM}} := \{\mathcal{H}, \mathcal{D}_0, L_\lambda^{\text{SNM}}, \{\phi_{\text{F}}(f), \pi_{\text{F}}(g) | f \in H_\omega^{-1/2}, g \in H_\omega^{1/2}\}\}$$

is a Nelson model.

Proof. (N.1) is obvious. (N.2) follows from Proposition 2.1. (N.3) with $s_0 = -1/2, s_1 = 1/2$ follows from the present assumption. (N.4) follows from (2.11) and (2.12). For (N.5), we take $\mathcal{E} = \mathcal{D}_0$. By condition (2.27), $C_0^\infty(\mathbb{R}^{dN}) \subset D(V)$ so that $C_0^\infty(\mathbb{R}^{dN}) \subset D(H_p)$. Hence (2.24) follows. The commutation relations in (N.5) follow from direct computations using (2.10), (2.11) and

$$[H_b, a(f)] = -a(\omega f), \quad [H_b, a(f)^*] = a(\omega f), \quad (2.31)$$

on $\mathcal{F}_{\omega, \text{fin}} (f, \omega f \in L^2(\mathbb{R}^d))$. ■

2.3 Field equations

We derive field equations for the Nelson model $\mathbb{M}_{\text{Nelson}}$ in a weak sense.

Definition 2.5 Let \mathcal{X} be a Hilbert space and $A(t)$ ($t \in \mathbb{R}$) be a linear operator on \mathcal{X} . Suppose that there exists a dense subspace \mathcal{D} of \mathcal{X} such that $\mathcal{D} \subset D(A(t))$ for all $t \in \mathbb{R}$. We say that $A(t)$ is weakly differentiable on \mathcal{D} if, for all $\psi, \phi \in \mathcal{D}$, the function $\langle \psi, A(t)\phi \rangle$ is differentiable in $t \in \mathbb{R}$. In that case we define a quadratic form $w\text{-}dA(t)/dt|_{\mathcal{D}}$ on \mathcal{D} by

$$w\text{-}\frac{dA(t)}{dt}\Big|_{\mathcal{D}}(\psi, \phi) := \frac{d}{dt} \langle \psi, A(t)\phi \rangle, \quad \psi, \phi \in \mathcal{D}.$$

Definition 2.6 Let \mathcal{X} be a Hilbert space and H be a self-adjoint operator on \mathcal{H} . Let A be a densely defined linear operator on \mathcal{X} . We say that A is in the set \mathbf{A}_H if it satisfies the following (i) and (ii):

- (i) There exists a dense subspace $D_A \subset D(H)$ such that, for all $s \in \mathbb{R}$, $e^{isH}D_A \subset D(A) \cap D(A^*)$.
- (ii) For all $\psi \in D_A$, the \mathcal{X} -valued functions: $s \rightarrow Ae^{isH}\psi$ and $s \rightarrow A^*e^{isH}\psi$ are strongly continuous on \mathbb{R} .

For $A \in \mathbf{A}_H$, we set

$$D_{A,H} := \mathcal{L}\{e^{isH}\psi \mid \psi \in D_A, s \in \mathbb{R}\}. \quad (2.32)$$

Lemma 2.7 Let \mathcal{X} be a Hilbert space and H be a self-adjoint operator on \mathcal{X} . Let A be a densely defined linear operator on \mathcal{X} and set

$$A(t) := e^{itH}Ae^{-itH}, \quad t \in \mathbb{R},$$

Assume that $A \in \mathbf{A}_H$. Then $A(t)$ is weakly differentiable on $D_{A,H}$ and

$$w\text{-}\frac{dA(t)}{dt}\Big|_{D_{A,H}} = i[H, A(t)]_{w}^{D_{A,H}}. \quad (2.33)$$

Proof. Let $\psi, \phi \in D_{A,H}$. Then $\psi, \phi \in D(A(t)) \cap D(A(t)^*)$ for all $t \in \mathbb{R}$. For $h \in \mathbb{R} \setminus \{0\}$,

$$\begin{aligned} \left\langle \psi, \frac{A(t+h) - A(t)}{h} \phi \right\rangle &= \left\langle e^{-itH} \frac{e^{-ihH} - 1}{h} \psi, Ae^{-i(t+h)H} \phi \right\rangle \\ &\quad + \left\langle A(t)^* \psi, \frac{e^{-ihH} - 1}{h} \phi \right\rangle. \end{aligned}$$

Since $\psi \in D(H)$, it follows that

$$\lim_{h \rightarrow 0} \frac{e^{-ihH} - 1}{h} \psi = -iH\psi.$$

It follows from Definition 2.6(ii) that $Ae^{isH}\phi$ is strongly continuous in s . Hence

$$\lim_{h \rightarrow 0} Ae^{-i(t+h)H} \phi = Ae^{-itH} \phi.$$

Hence

$$\lim_{h \rightarrow 0} \left\langle \psi, \frac{A(t+h) - A(t)}{h} \phi \right\rangle = i \left\langle e^{-itH} H \psi, A e^{-itH} \phi \right\rangle - i \left\langle A(t)^* \psi, H \phi \right\rangle.$$

Thus the desired result follows. \blacksquare

The canonical Heisenberg operators of the Nelson model $\mathbb{M}_{\text{Nelson}}$ (Definition 2.3) are defined as follows:

$$q_{j\mu}(t) := e^{itH_{\text{NM}}} q_{j\mu} e^{-itH_{\text{NM}}}, \quad p_{j\mu}(t) := e^{itH_{\text{NM}}} p_{j\mu} e^{-itH_{\text{NM}}}, \quad (2.34)$$

$$\phi(t, f) := e^{itH_{\text{NM}}} \phi(f) e^{-itH_{\text{NM}}}, \quad \pi(t, f) := e^{itH_{\text{NM}}} \pi(t, f) e^{-itH_{\text{NM}}}, \quad (2.35)$$

$$j = 1, \dots, N, \quad \mu = 1, \dots, d, \quad f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d), \quad t \in \mathbb{R}.$$

For a self-adjoint operator T , $Q(T)$ denotes the form domain of the quadratic form associated with T .

Theorem 2.8 Consider the Nelson model $\mathbb{M}_{\text{Nelson}}$ (Definition 2.3). Assume (H.1), (H.2) and the following (A.1)–(A.5):

(A.1) (2.27) holds and H_{p} is essentially self-adjoint on $C_0^\infty(\mathbb{R}^d)$.

(A.2) For all $j = 1, \dots, N$, $\mu = 1, \dots, d$, the distributional partial derivative $D_{j\mu} V$ in the variable $q_{j\mu}$ is a Borel measurable function on \mathbb{R}^d which is a.e. finite with respect to the Lebesgue measure. Moreover, $C_0^\infty(\mathbb{R}^d)$ is a form core of the self-adjoint multiplication operator $D_{j\mu} V$ and $D(H_{\text{p}}) \subset \cap_{j=1}^N \cap_{\mu=1}^d Q(D_{j\mu} V)$.

(A.3) In addition to (2.24),

$$\mathcal{E} \subset D(H_{\text{p}}) \cap \left\{ \cap_{j,l=1}^N \cap_{\mu,\nu=1}^d [D(p_{j\mu} q_{l\nu}) \cap D(q_{l\nu} p_{j\mu})] \right\} \cap \left[\cap_{j=1}^N \cap_{\mu=1}^d Q(D_{j\mu} V) \right].$$

(A.4) For all $s \in \mathbb{R}$, $e^{isH_{\text{NM}}}$ leaves \mathcal{E} invariant: $e^{isH_{\text{NM}}} \mathcal{E} \subset \mathcal{E}$

(A.5) For all $j = 1, \dots, N$, $\mu = 1, \dots, d$ and $\psi \in \mathcal{E}$, the \mathcal{K} -valued function $q_{j\mu} e^{isH_{\text{NM}}} \psi$ is strongly continuous in $s \in \mathbb{R}$.

(A.6) For all $f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$, the \mathcal{K} -valued functions $\phi(f) e^{isH_{\text{NM}}} \psi$ and $\pi(f) e^{isH_{\text{NM}}} \psi$ are strongly continuous in $s \in \mathbb{R}$.

Then the Heisenberg operators $q_{j\mu}(t), p_{j\mu}(t), \phi(t, f)$ and $\pi(t, f)$ ($f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$) are weakly differentiable on \mathcal{E} and

$$\text{w-} \frac{dq_{j\mu}(t)}{dt} \Big|_{\mathcal{E}} = \frac{p_{j\mu}(t)}{m_j}, \quad (2.36)$$

$$\text{w-} \frac{dp_{j\mu}(t)}{dt} \Big|_{\mathcal{E}} = -D_{j\mu} V(q(t)) - \lambda e^{itH_{\text{NM}}} \phi(D_{\mu} \rho_j(q_j - \cdot)) e^{-itH_{\text{NM}}}, \quad (2.37)$$

$$\text{w-} \frac{d\phi(t, f)}{dt} \Big|_{\mathcal{E}} = \pi(t, f), \quad (2.38)$$

$$\text{w-} \frac{d\pi(t, f)}{dt} \Big|_{\mathcal{E}} = -\phi(t, \omega(-i\nabla)^2 f) - \lambda \sum_{j=1}^N (\rho_j * f)(q_j(t)), \quad (2.39)$$

where $q_j(t) := (q_{j1}(t), \dots, q_{jd}(t))$, $q(t) := (q_1(t), \dots, q_N(t))$.

Proof. By (A.3) and the CCR

$$[p_{j\mu}, q_{l\nu}] = -i\delta_{jl}\delta_{\mu\nu} \quad \text{on } D(p_{j\mu}q_{l\nu}) \cap D(q_{l\nu}p_{j\mu}),$$

we have

$$[H_p, q_{j\mu}]_{\mathcal{W}}^{\mathcal{E}} = -i\frac{p_{j\mu}}{m_j},$$

which, together with (N.5), implies that

$$[H_{NM}, q_{j\mu}]_{\mathcal{W}}^{\mathcal{E}} = -i\frac{p_{j\mu}}{m_j}. \quad (2.40)$$

By this fact, (A.4) and (A.5), we can apply Lemma 2.7 to obtain (2.36).

Let

$$h_0 := -\sum_{j=1}^N \frac{\Delta_{q_j}}{2m_j}. \quad (2.41)$$

For all $\psi \in D(h_0)$, we have

$$\begin{aligned} \|p_{j\mu}\psi\|^2 &\leq 2m_j \langle \psi, h_0\psi \rangle \leq 2m_j \|\psi\| \|h_0\psi\| \\ &\leq \varepsilon \|h_0\psi\|^2 + \frac{m_j^2}{\varepsilon} \|\psi\|^2, \end{aligned} \quad (2.42)$$

where $\varepsilon > 0$ is arbitrary. By the self-adjointness of H_p and the closed graph theorem, there exist constants $a \geq 0, b \geq 0$ such that, for all $\psi \in D(H_p) = D(h_0) \cap D(V)$,

$$\|h_0\psi\| + \|V\psi\| \leq a\|H_p\psi\| + b\|\psi\|. \quad (2.43)$$

Hence, by (2.42),

$$\|p_{j\mu}\psi\| \leq a\sqrt{\varepsilon}\|H_p\psi\| + \left(b\sqrt{\varepsilon} + \frac{m_j}{\sqrt{\varepsilon}}\right)\|\psi\|. \quad (2.44)$$

Let $\psi, \phi \in C_0^\infty(\mathbb{R}^{dN})$. Then we have

$$\langle H_p\psi, p_{j\mu}\phi \rangle - \langle p_{j\mu}\psi, H_p\phi \rangle = \langle \psi, i(D_{j\mu}V)\phi \rangle.$$

By (A.1), (A.2) and (2.44), we can extend this equality to all $\psi, \phi \in D(H_p)$. Hence, in particular, we have

$$[H_p, p_{j\mu}]_{\mathcal{W}}^{\mathcal{E}} = iD_{j\mu}V \quad (2.45)$$

in the sense of quadratic form on \mathcal{E} . Hence, by (N.5), we have

$$[H_{NM}, p_{j\mu}]_{\mathcal{W}}^{\mathcal{E}} = iD_{j\mu}V + i\lambda\phi(D_{\mu}\rho_j(q_j - \cdot)). \quad (2.46)$$

in the sense of quadratic form on \mathcal{E} . By this fact, (A.4) and (A.6), we can apply Lemma 2.7 to obtain (2.37).

Similarly (more easily) we can prove (2.38) and (2.39). ■

Now we show that Theorem 2.8 can be applied to the SNM.

Lemma 2.9 *Let S be a self-adjoint operator on a Hilbert space \mathcal{X} and A be a linear operator on \mathcal{X} such that, for some $\alpha > 0$, A is $|S|^\alpha$ -bounded, i.e., $D(|S|^\alpha) \subset D(A)$ and*

$$\|A\psi\| \leq a\||S|^\alpha\psi\| + b\|\psi\|, \quad \psi \in D(|S|^\alpha),$$

where $a, b \geq 0$ are constants. Then, for all $\psi \in D(|S|^\alpha)$, the function: $t \rightarrow Ae^{itS}\psi$ is strongly continuous.

Proof. We have for all $h \in \mathbb{R}$

$$\begin{aligned} \|Ae^{i(t+h)S}\psi - Ae^{itS}\psi\| &\leq a\||S|^\alpha(e^{ihS} - 1)e^{itS}\psi\| + b\|(e^{ihS} - 1)\psi\| \\ &= a\|(e^{ihS} - 1)|S|^\alpha\psi\| + b\|(e^{ihS} - 1)\psi\|, \end{aligned}$$

which implies the strong continuity of the function: $t \rightarrow Ae^{itS}\psi$. ■

For a self-adjoint operator T on a Hilbert space which is bounded from below, we set

$$E_0(T) := \inf \sigma(T), \quad (2.47)$$

the ground state energy of T , where $\sigma(T)$ denotes the spectrum of T .

Under hypotheses (H.1) and (H.2), we set

$$\hat{H}_{\text{SNM}} := H_{\text{SNM}} - E_0(H_{\text{SNM}}), \quad (2.48)$$

which is nonnegative.

Lemma 2.10 *Assume (H.1) and (H.2). Suppose that $Q(V) \subset D(|q|)$ and there exist constants $c_1, c_2 \geq 0$ such that for all $u \in Q(V)$*

$$\||q|u\| \leq c_1\||V|^{1/2}u\| + c_2\|u\|. \quad (2.49)$$

Then, for all $j = 1, \dots, N$, $\mu = 1, \dots, d$ and $\psi \in Q(H_{\text{SNM}})$, the \mathcal{H} -valued function $q_{j\mu}e^{isH_{\text{SNM}}}\psi$ is strongly continuous in $s \in \mathbb{R}$.

Proof. By Proposition 2.1 and the closed graph theorem, there exist constants $a_1, b_1 \geq 0$ such that, for all $\psi \in D(H_{\text{SNM}}) = D(H_0)$,

$$\|H_p\psi\| + \|H_b\psi\| \leq a_1\|H_{\text{SNM}}\psi\| + b_1\|\psi\|. \quad (2.50)$$

By this inequality and (2.43), we have

$$\|V\psi\| \leq aa_1\|H_{\text{SNM}}\psi\| + (ab_1 + b)\|\psi\|,$$

which, via an application [11, Theorem X.18], implies that $Q(H_{\text{SNM}}) \subset Q(V)$ and

$$\||V|^{1/2}\phi\| \leq a_2\|\hat{H}_{\text{SNM}}^{1/2}\phi\| + b_2\|\phi\|, \quad \phi \in Q(H_{\text{SNM}}),$$

with $a_2, b_2 \geq 0$ being constants. Hence, by (2.49), we obtain

$$\|q_{j\mu}\psi\| \leq c_1a_2\|\hat{H}_{\text{SNM}}^{1/2}\psi\| + (c_1b_2 + c_2)\|\psi\|, \quad \psi \in Q(H_{\text{SNM}}).$$

Hence we can apply Lemma 2.9 to obtain the desired result. ■

Lemma 2.11 *Assume (H.1) and (H.2). Suppose that*

$$\mathcal{S}_{\text{real}}(\mathbb{R}^d) \subset H_{\omega}^{-1} \cap H_{\omega}^{1/2}. \quad (2.51)$$

Let $\psi \in Q(H_{\text{SNM}})$. Then, for all $f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$, the \mathcal{H} -valued functions $\phi_{\text{F}}(f)e^{isH_{\text{SNM}}}\psi$ and $\pi_{\text{F}}(f)e^{isH_{\text{SNM}}}\psi$ are strongly continuous in $s \in \mathbb{R}$.

Proof. Using the well-known estimates

$$\|a(f)\psi\| \leq \|\omega^{-1/2}f\|_{L^2(\mathbb{R}^d)} \|H_{\text{b}}^{1/2}\psi\|, \quad (2.52)$$

$$\begin{aligned} \|a(f)^*\psi\| &\leq \|\omega^{-1/2}f\|_{L^2(\mathbb{R}^d)} \|H_{\text{b}}^{1/2}\psi\| + \|f\|_{L^2(\mathbb{R}^d)}, \\ &\psi \in D(H_{\text{b}}^{1/2}), f, \omega^{-1/2}f \in L^2(\mathbb{R}^d), \end{aligned} \quad (2.53)$$

we have

$$\begin{aligned} \|\phi_{\text{F}}(f)\psi\| &\leq \sqrt{2}\|\omega^{-1}\hat{f}\|_{L^2(\mathbb{R}^d)} \|H_{\text{b}}^{1/2}\psi\| + \frac{1}{\sqrt{2}}\|\omega^{-1/2}\hat{f}\|_{L^2(\mathbb{R}^d)}\|\psi\|, \\ &f \in H_{\omega}^{-1} \cap H_{\omega}^{-1/2}, \end{aligned} \quad (2.54)$$

$$\begin{aligned} \|\pi_{\text{F}}(g)\psi\| &\leq \sqrt{2}\|\hat{g}\|_{L^2(\mathbb{R}^d)} \|H_{\text{b}}^{1/2}\psi\| + \frac{1}{\sqrt{2}}\|\omega^{1/2}\hat{g}\|_{L^2(\mathbb{R}^d)}\|\psi\|, \\ &g \in L^2(\mathbb{R}^d) \cap H_{\omega}^{1/2}. \end{aligned} \quad (2.55)$$

By (2.50) and an application [11, Theorem X.18], we have

$$\|H_{\text{b}}^{1/2}\psi\| \leq a_3\|\hat{H}_{\text{SNM}}^{1/2}\psi\| + b_3\|\psi\|, \quad \psi \in Q(H_{\text{SNM}}), \quad (2.56)$$

with $a_3, b_3 \geq 0$ being constants. It follows that, for all $f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$, $\phi_{\text{F}}(f)$ and $\pi_{\text{F}}(f)$ are $\hat{H}_{\text{SNM}}^{1/2}$ -bounded. Hence an application of Lemma 2.9 yields the desired result. \blacksquare

We shall need the following condition too.

(H.3) The function ω is such that

$$\mathcal{S}_{\text{real}}(\mathbb{R}^d) \subset H_{\omega}^{-1} \cap H_{\omega}^{3/2}. \quad (2.57)$$

Theorem 2.12 *Assume (H.1)–(H.3), (A.1), (A.2), (2.29) and (2.49). Suppose that*

$$D(H_{\text{P}}) \subset \bigcap_{j,l=1}^N \bigcap_{\mu,\nu=1}^d [D(p_{j\mu}q_{l\nu}) \cap D(q_{l\nu}p_{j\mu})] \cap \left(\bigcap_{j=1}^N \bigcap_{\mu=1}^d Q(D_{j\mu}V) \right) \quad (2.58)$$

Then the conclusion of Theorem 2.8 holds with $H_{\text{NM}} = H_{\text{SNM}}$, $\phi(f) = \phi_{\text{F}}(f)$, $\pi(f) = \pi_{\text{F}}(f)$ and $\mathcal{E} = D(H_{\text{SNM}})$.

Proof. We need only check that the assumption of Theorem 2.8 is satisfied with $H_{\text{NM}} = H_{\text{SNM}}$, $\phi(f) = \phi_{\text{F}}(f)$, $\pi(f) = \pi_{\text{F}}(f)$ and $\mathcal{E} = D(H_{\text{SNM}})$. (A.3) and (A.4) follow from (2.58). (A.5) and (A.6) follow from Lemmas 2.10 and 2.11 respectively. \blacksquare

3 A Nelson model in a non-Fock representation

In this section we define a Nelson model whose time-zero fields are given by a non-Fock representation of the CCR and show that it has a ground state even if no infrared cutoff is made.

Under hypotheses (H.2) and (H.3), we can define for each $f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)$

$$\tilde{\phi}(f) := \phi_{\text{F}}(f) + \lambda \sum_{j=1}^N \int_{\mathbb{R}^d} \frac{\hat{\rho}_j(k) \hat{f}(k)}{\omega(k)^2} dk. \quad (3.1)$$

We set

$$\tilde{\pi}(f) := \pi_{\text{F}}(f). \quad (3.2)$$

Proposition 3.1 *Assume (H.2) and (H.3). Then:*

- (i) $\{\mathcal{F}_b, \mathcal{F}_0, \{\tilde{\phi}(f), \tilde{\pi}(f) | \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}\}$ is a representation of the CCR indexed by $\mathcal{S}_{\text{real}}(\mathbb{R}^d)$.
- (ii) The representation $\{\tilde{\phi}(f), \tilde{\pi}(f) | \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}$ is unitarily equivalent to the Fock representation $\{\phi_{\text{F}}(f), \pi_{\text{F}}(f) | \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}$ if and only if $\sum_{j=1}^N \rho_j \in H_{\omega}^{-3/2}$.

Proof. Part (i) is obvious. Part (ii) follows from an argument similar to that in [5, Chapter 1, §1-e]. ■

We now consider a Nelson model whose time-zero fields are given by $\{\tilde{\phi}(f), \tilde{\pi}(f) | f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}$. We introduce an $L^2(\mathbb{R}^d)$ -valued function G on \mathbb{R}^{dN} ($G : \mathbb{R}^{dN} \rightarrow L^2(\mathbb{R}^d)$, $G(q) \in L^2(\mathbb{R}^d)$, $q \in \mathbb{R}^{dN}$) by

$$G(q)(k) := \sum_{j=1}^N \frac{\hat{\rho}_j(k)^*}{\sqrt{\omega(k)}} (e^{-iq_j k} - 1) \quad (3.3)$$

and a function

$$W(q) := \sum_{j,l=1}^N \int_{\mathbb{R}^d} \frac{\hat{\rho}_j(k) \hat{\rho}_l(k)^*}{\omega(k)^2} e^{-iq_l k} dk, \quad q \in \mathbb{R}^{dN}. \quad (3.4)$$

By reality of ρ_j and (2.3), W is real-valued.

We define a Hamiltonian by

$$H := H_0 + \lambda \Phi_{\text{S}}(G(q)) - \lambda^2 W + c_0 \lambda^2, \quad (3.5)$$

where

$$c_0 := \frac{1}{2} \left\| \frac{\sum_{j=1}^N \hat{\rho}_j}{\omega} \right\|_{L^2(\mathbb{R}^d)}^2. \quad (3.6)$$

Lemma 3.2 *Assume (H.1)–(H.3). Then H is self-adjoint with $D(H) = D(H_0)$ and bounded from below. Moreover, H is essentially self-adjoint on each core of H_0 .*

Proof. In the same way as in the case of H_{SNM} , we can show that $H' := H_0 + \lambda \Phi_S(G(q))$ is self-adjoint with $D(H') = D(H_0)$, bounded from below, and essentially self-adjoint on each core of H_0 . Since $-\lambda^2 W + c_0 \lambda^2$ is a bounded self-adjoint operator, the Kato-Rellich theorem yields the desired result. \blacksquare

Let

$$L_\lambda^{\text{NF}} := H_b + \lambda \Phi_S(G(q)) - \lambda^2 W + c_0 \lambda^2, \quad (3.7)$$

so that

$$H = H_p + L_\lambda^{\text{NF}}. \quad (3.8)$$

Proposition 3.3 *Assume (H.1)–(H.3). Then*

$$\mathbf{M}_{\text{NF}} := \{\mathcal{H}, \mathcal{D}_0, L_\lambda^{\text{NF}}, \{\tilde{\phi}(f), \tilde{\pi}(g) | f \in H_\omega^{-1}, g \in H_\omega^{1/2}\}\} \quad (3.9)$$

is a Nelson model.

Proof. Similar to the proof of Proposition 2.4. \blacksquare

Proposition 3.4 *Assume (H.1)–(H.3). Then $\{H, \{\tilde{\phi}(f), \tilde{\pi}(g) | f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}\}$ is unitarily equivalent to $\{H_{\text{SNM}}, \{\phi_{\text{F}}(f), \pi_{\text{F}}(g) | f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d)\}\}$ if and only if*

$$\sum_{j=1}^N \rho_j \in H_\omega^{-3/2}. \quad (3.10)$$

Proof. Suppose that there exists a unitary operator Y on \mathcal{H} such that

$$YHY^{-1} = H_{\text{SNM}}, \quad (3.11)$$

$$Y\tilde{\phi}(f)Y^{-1} = \phi_{\text{F}}(f), \quad Y\tilde{\pi}(f)Y^{-1} = \pi_{\text{F}}(f), \quad f \in \mathcal{S}_{\text{real}}(\mathbb{R}^d). \quad (3.12)$$

Then, by Proposition 3.1, (3.10) holds. Conversely, suppose that (3.10) holds. Then we can define a unitary operator U by

$$U := \exp\left(-i\lambda \Phi_S\left(i \frac{\sum_{j=1}^N \hat{\rho}_j^*}{\omega^{3/2}}\right)\right). \quad (3.13)$$

It is easy to check that (3.11) and (3.12) hold with $Y = U$. \blacksquare

Proposition 3.4 shows that, under (H.1)–(H.3) and the condition that $\sum_{j=1}^N \rho_j \in H_\omega^{-3/2}$, the Nelson model \mathbf{M}_{NF} is equivalent to the SNM. In this case, under suitable additional conditions, H_{SNM} has a ground state [7] and so does H . Thus we are interested in the case

$$\sum_{j=1}^N \rho_j \notin H_\omega^{-3/2}. \quad (3.14)$$

This condition is called an *infrared singularity condition*. In this paper, we say that *the Nelson model $\mathbf{M}_{\text{Nelson}}$ has no infrared cutoff* if (3.14) holds. Under condition (3.14), the Nelson model \mathbf{M}_{NF} is not unitarily equivalent to the SNM. Note that (3.14) and the natural condition $\sum_{j=1}^N \rho_j / \sqrt{\omega} \in L^2(\mathbb{R}^d)$ imply

$$\text{ess.inf}_{k \in \mathbb{R}^d} \omega(k) = 0 \quad (3.15)$$

(“ess.inf” means essential infimum), i.e., the quantum scalar field under consideration is “massless”.

4 Existence of a ground state of the Nelson model \mathbb{M}_{NF} without infrared cutoff

Let T be a self-adjoint operator on a Hilbert space and bounded from below. We say that T has a ground state if there exists a non-zero vector $\psi \in D(T)$ such that $T\psi = E_0(T)\psi$. In this case ψ is called a ground state of T .

We show below that, under suitable conditions, the Hamiltonian H of the Nelson model \mathbb{M}_{NF} has a ground state *even in the case where it has no infrared cutoff (i.e., (3.14) holds)*.

We formulate some additional conditions:

(H.4) The function ω is continuous on \mathbb{R}^d satisfying (3.15) and the following conditions:

$$\begin{aligned} D_\mu \omega &\in L^\infty(\mathbb{R}^d), \quad \mu = 1, \dots, d, \\ \lim_{|k| \rightarrow \infty} \omega(k) &= \infty, \end{aligned}$$

(H.5) For $j = 1, \dots, N$,

$$\int_{\mathbb{R}^d} \frac{|k|^2 \hat{\rho}_j(k)|^2}{\omega(k)^3} dk < \infty. \quad (4.1)$$

(H.6) For all $R > 0$, $\int_{|q| \leq R} |V(q)| dq < \infty$ and there exist constants $c_1, c_2 \geq 0$ such that

$$|q|^2 \leq c_1 V(q) + c_2, \quad \text{a.e. } q \in \mathbb{R}^{dN}. \quad (4.2)$$

An infrared-cutoff Hamiltonian of the SNM is defined by

$$H_{\text{SNM}, \sigma} := H_0 + \lambda \sum_{j=1}^N \Phi_S \left(e^{-ikq_j} \frac{\chi_{\omega \geq \sigma} \hat{\rho}_j^*}{\sqrt{\omega}} \right), \quad (4.3)$$

where $\sigma > 0$ is an infrared cutoff parameter and χ_S is a characteristic function of the set S .

Under (H.2), we can define for all $\sigma > 0$ a unitary operator

$$U_\sigma := \exp \left(-i\lambda \Phi_S \left(i \frac{\sum_{j=1}^N \chi_{\omega \geq \sigma} \hat{\rho}_j}{\omega^{3/2}} \right) \right). \quad (4.4)$$

Theorem 4.1 *Assume (H.1), (H.2) and (H.4)–(H.6). Then H has a ground state ψ_0 which has the following property: there exists a sequence $\{\phi_{\sigma_n}\}_{n=1}^\infty$ of unit vectors in $D(H_0)$ such that $\sigma_n > 0$, $n \in \mathbb{N}$, $\lim_{n \rightarrow \infty} \sigma_n = 0$, each ϕ_{σ_n} is a ground state of H_{SNM, σ_n} and*

$$\text{w-}\lim_{n \rightarrow \infty} U_{\sigma_n}^{-1} \phi_{\sigma_n} = \psi_0, \quad (4.5)$$

where “w-lim” means weak limit.

Remark 4.1 Consider the physical case $\omega = \omega_0$ ((2.4) with $m = 0$). Hence (H.4) holds. Assume (H.2) with $\omega = \omega_0$ and that

$$\sum_{j=1}^N \frac{\hat{\rho}_j(\cdot)}{|k|^{3/2}} \notin L^2(\mathbb{R}^d).$$

Then (3.14) and (H.5) hold with $\omega = \omega_0$. Hence Theorem 4.1 holds in the physical case without infrared cutoff.

Remark 4.2 Assume (3.14). Then we have

$$\text{w-}\lim_{\sigma \rightarrow 0} U_\sigma = 0. \quad (4.6)$$

This is an expression of infrared divergence. Hence the relation $\text{w-}\lim_{n \rightarrow \infty} U_{\sigma_n}^{-1} \phi_{\sigma_n} = \psi_0$ in Theorem 4.1 suggests that $\text{w-}\lim_{n \rightarrow \infty} \phi_{\sigma_n} = 0$.

The idea of proof of Theorem 4.1 is to apply [7, Theorem 1].

Let

$$K := H_p - \lambda^2 W + c_0 \lambda^2.$$

Lemma 4.2 Assume (H.1) and (H.2). Then K is self-adjoint and bounded from below.

Proof. This follows from the boundedness of $\lambda^2 W$ and a simple application of the Kato-Rellich theorem. ■

Let

$$\hat{K} := K - E_0(K).$$

Lemma 4.3 Assume (H.1), (H.2) and (H.6). Then $(\hat{K} + i)^{-1}$ is compact.

Proof. By (H.6) and the boundedness of W , we have

$$\lim_{|q| \rightarrow \infty} \{V(q) - \lambda^2 W(q) + c_0 \lambda^2\} = \infty.$$

Hence, applying [12, p.249, Theorem XIII.67], we obtain the desired result. ■

We denote by $\mathbf{B}(L^2(\mathbb{R}^{dN}))$ the set of bounded linear operators on $L^2(\mathbb{R}^{dN})$.

For each $k \in \mathbb{R}^d$, we define a linear operator on $L^2(\mathbb{R}^{dN})$ as follows:

$$v(k) := \begin{cases} G(\cdot)(k) & \omega(k) > 0 \\ 0 & \omega(k) = 0 \end{cases}. \quad (4.7)$$

Remark 4.3 By the original assumption for ω , the Lebesgue measure of the set $\{k \in \mathbb{R}^d | \omega(k) = 0\}$ is equal to zero. Hence, $v(k) = G(\cdot)(k)$ a.e. $k \in \mathbb{R}^d$.

Lemma 4.4 Assume (H.2). Then, for a.e. $k \in \mathbb{R}^{dN}$, $v(k) \in \mathbf{B}(L^2(\mathbb{R}^{dN}))$ with

$$\|v(k)\| \leq 2 \sum_{j=1}^N \frac{|\hat{\rho}_j(k)|}{\sqrt{\omega(k)}}.$$

Proof. This easily follows from (3.3). ■

The following lemma immediately follows from Lemma 4.4.

Lemma 4.5 *Assume (H.1) and (H.2). Then, for a.e. $k \in \mathbb{R}^{dN}$,*

$$v(k)(\hat{K} + 1)^{-1/2} \in \mathbf{B}(L^2(\mathbb{R}^{dN})), \quad (\hat{K} + 1)^{-1/2}v(k) \in \mathbf{B}(L^2(\mathbb{R}^{dN})).$$

Lemma 4.6 *Assume (H.1) and (H.2). Then, for all $u_1, u_2 \in L^2(\mathbb{R}^{dN})$, the functions*

$$(u_2, v(\cdot)(\hat{K} + 1)^{-1/2}u_1), (u_2, (\hat{K} + 1)^{-1/2}v(\cdot)u_1),$$

on \mathbb{R}^d are measurable. Moreover,

$$C(R) := \int_{\mathbb{R}^d} \frac{1}{\omega(k)} \left(\|v(k)(\hat{K} + R)^{-1/2}\|^2 + \|(\hat{K} + R)^{-1/2}v(k)\|^2 \right) dk < \infty$$

and

$$\lim_{R \rightarrow \infty} C(R) = 0. \quad (4.8)$$

Proof. We have

$$(u_2, v(k)(\hat{K} + 1)^{-1/2}u_1) = \sum_{j=1}^N \frac{\hat{\rho}_j(k)^*}{\sqrt{\omega(k)}} g_j(k)$$

with

$$g_j(k) = \int_{\mathbb{R}^d} u_2(q)^* (e^{-iq_j k} - 1) ((\hat{K} + 1)^{-1/2}u_1)(q) dq.$$

It is easy to see that g_j is continuous on \mathbb{R}^d . Hence $(u_2, v(\cdot)(\hat{K} + 1)^{-1/2}u_1)$ is measurable. Similarly we can show that $(u_2, (\hat{K} + 1)^{-1/2}v(\cdot)u_1)$ is measurable.

We have

$$C(R) \leq \frac{2}{R} \int_{\mathbb{R}^d} \frac{1}{\omega(k)} \|v(k)\|^2 dk \leq \frac{8N}{R} \sum_{j=1}^N \int_{\mathbb{R}^d} \frac{|\hat{\rho}_j(k)|^2}{\omega(k)^2} dk.$$

Hence $C(R) < \infty$ and (4.8) holds. ■

Lemma 4.7 *Assume (H.1) and (H.6). Let $\psi \in D(\hat{K}^{1/2})$. Then $\psi \in D(|q_j|)$ and, for all $j = 1, \dots, N$,*

$$\| |q_j| \psi \| \leq c_1 \|\hat{K}^{1/2} \psi\| + (c_1 \lambda^2 \|W\| + c_1 c_0 \lambda^2 + c_1 E_0(K) + c_2) \|\psi\|^2. \quad (4.9)$$

In particular,

$$Q_j := |q_j| (\hat{K} + 1)^{-1/2} \quad (4.10)$$

is bounded.

Proof. Let $\psi \in D(K)$. Then, using (H.6), one can easily prove (4.9). Since $D(K)$ is a core of $\hat{K}^{1/2}$, it follows from a limiting argument that $D(|q_j|) \supset D(\hat{K}^{1/2})$ and (4.9) holds for all $\psi \in D(\hat{K}^{1/2})$. ■

Lemma 4.8 Assume (H.1), (H.2), (H.5) and (H.6). Then

$$\int_{\mathbb{R}^d} \frac{1}{\omega(k)^2} \|v(k)(\hat{K} + 1)^{-1/2}\|^2 dk < \infty. \quad (4.11)$$

Proof. By the elementary inequality

$$|e^{-iq_j k} - 1| \leq |q_j| |k|$$

we have for all $u \in L^2(\mathbb{R}^{dN})$

$$\|v(k)u\| \leq \sum_{j=1}^N \frac{|\hat{\rho}_j(k)| |k|}{\sqrt{\omega(k)}} \| |q_j| u \|.$$

Hence

$$\|v(k)(\hat{K} + 1)^{-1/2}\| \leq \sum_{j=1}^N \frac{|\hat{\rho}_j(k)| |k|}{\sqrt{\omega(k)}} \|Q_j\|,$$

which, by (H.5), implies (4.11). ■

Proof of Theorem 4.1

We have

$$H = K \otimes I + I \otimes H_b + H_I(v)$$

with

$$H_I(v) := \frac{\lambda}{\sqrt{2}} \int_{\mathbb{R}^d} (v(k)a(k)^* + v(k)^*a(k)) dk.$$

Hence H is exactly of the same form as the Hamiltonians considered in [7]. Lemmas 4.2–4.8 show that H satisfies all the assumptions of [7, Theorem 1]. Thus we can apply [7, Theorem 1] to conclude that H has a ground state ψ_0 . From the proof of [7, Theorem 1], we see that ψ_0 is a weak limit of a sequence $\{\psi_{\sigma_n}\}_{n=1}^{\infty}$ of unit vectors in $D(H_0)$ such that $\sigma_n > 0$, $\lim_{n \rightarrow \infty} \sigma_n = 0$ and each ψ_{σ_n} is a ground state of the infrared-cutoff version H_{σ_n} of H :

$$H_{\sigma} := K \otimes I + I \otimes H_b + H_I(v_{\sigma}), \quad \sigma > 0,$$

where $v_{\sigma}(k) := v(k)\chi_{\omega \geq \sigma}(k)$. By Proposition 3.4, $\phi_{\sigma_n} := U_{\sigma_n} \psi_{\sigma_n}$ is a ground state of H_{SNM, σ_n} with $\|\phi_{\sigma_n}\| = 1$. Hence (4.5) holds. ■

References

- [1] A. Arai, Scaling limit for quantum systems of nonrelativistic particles interacting with a Bose field, Hokkaido University Preprint Series in Mathematics #59, 1989.
- [2] A. Arai, M. Hirokawa and F. Hiroshima, On the absence of eigenvectors of Hamiltonians in a class of massless quantum field models without infrared cutoff, J. Funct. Anal. **168**(1999), 470–497.

- [3] A. Arai and M. Hirokawa, Ground states of a general class of quantum field Hamiltonians, Hokkaido University Preprint Series in Mathematics #457, 1999, to be published in *Rev. Math. Phys.*
- [4] V. Bach, J. Fröhlich and I. M. Sigal, Spectral analysis for systems of atoms and molecules coupled to the quantized radiation field, *Commun. Math. Phys.* **207**(1998), 249–290.
- [5] G. G. Emch, Algebraic Methods in Statistical Mechanics and Quantum Field Theory, John Wiley & Sons, 1972.
- [6] J. Fröhlich, On the infrared problem in a model of scalar electrons and massless scalar bosons, *Ann. Inst. Henri Poincaré* **19**(1973), 1–103.
- [7] C. Gérard, On the existence of ground states for massless Pauli-Fierz Hamiltonians, *Ann. Henri Poincaré* **1**(2000), 443–459.
- [8] M. Griesemer, E. H. Lieb and M. Loss, Ground states in non-relativistic quantum electrodynamics, preprint, 2000.
- [9] J. Lörinczi, H. Spohn and R. A. Minlos, The infrared behaviour in Nelson’s model of a quantum particle coupled to a massless scalar field, preprint, 2000.
- [10] E. Nelson, Interaction of nonrelativistic particles with a quantized scalar field, *J. Math. Phys.* **5**(1964), 1190–1197.
- [11] M. Reed and B. Simon, Methods of Modern Mathematical Physics Vol. II: Fourier Analysis, Self-Adjointness, Academic Press, New York, 1975.
- [12] M. Reed and B. Simon, Methods of Modern Mathematical Physics Vol. IV: Analysis of Operators, Academic Press, New York, 1978.
- [13] H. Spohn, Ground state(s) of the spin-boson Hamiltonian, *Commun. Math. Phys.* **123** (1989), 277–304.
- [14] H. Spohn, Ground state of a quantum particle coupled to a scalar Bose field, *Lett. Math. Phys.* **44** (1998), 9–16.