



Title	On the worst conditional expectation
Author(s)	Inoue, A.
Citation	Hokkaido University Preprint Series in Mathematics, 520, 1-10
Issue Date	2001-03
DOI	10.14943/83666
Doc URL	<a href="http://hdl.handle.net/2115/69270">http://hdl.handle.net/2115/69270</a>
Type	bulletin (article)
File Information	pre520.pdf



[Instructions for use](#)

## On the Worst Conditional Expectation

Akihiko Inoue

Series #520. March 2001

HOKKAIDO UNIVERSITY  
PREPRINT SERIES IN MATHEMATICS

- #495 M. Arisawa and Y. Giga, Anisotropic curvature flow in a very thin domain, 21 pages. 2000.
- #496 T. Nakazi, Backward shift invariant subspaces in the bidisc, 9 pages. 2000.
- #497 Y. Giga, K. Inui, J. Kato and S. Matsui, Remarks on the uniqueness of bounded solutions of the Navier-Stokes equations, 4 pages. 2000.
- #498 Y. Giga, M. Paolini and P. Rybka, On the motion by singular interfacial energy, 21 pages. 2000.
- #499 J. Escher, Y. Giga and K. Ito, On a limiting motion and self-interactions of curves moved by the intermediate surface diffusion flow, 12 pages. 2000.
- #500 I. Tsuda and S. Kuroda, Cantor coding in the hippocampus, 20 pages. 2000.
- #501 M. Tsujii, Fat solenoidal attractors, 20 pages. 2000.
- #502 A. Arai, Ground state of the massless Nelson model without infrared cutoff in a non-Fock representation, 19 pages. 2000.
- #503 Y. Giga, S. Matsui and O. Sawada, Global existence of two-dimensional Navier-Stokes flow with nondecaying initial velocity, 19 pages. 2000.
- #504 A. Inoue and Y. Kasahara, Partial autocorrelation functions of the fractional ARIMA processes with negative degree of differencing, 14 pages. 2000.
- #505 T. Nakazi, Interpolation problem for  $\ell^1$  and a uniform algebra, 12 pages. 2000.
- #506 R. Kobayashi and Y. Giga, On anisotropy and curvature effects for growing crystals, 38 pages. 2000.
- #507 A. Arai, Instability in the spectral and the Fredholm properties of an infinite dimensional Dirac operator on the abstract Boson-Fermion Fock space, 6 pages. 2000.
- #508 A. Arai, Supersymmetric methods for constructing soliton-type solutions to multi-component nonlinear Schrödinger and Klein-Gordon equations, 22 pages. 2000.
- #509 Y. Tonegawa, Phase field model with a variable chemical potential, 30 pages. 2000.
- #510 Y. Giga, Shocks and very strong vertical diffusion, 11 pages. 2000.
- #511 S. Izumiya & N. Takeuchi, Special curves and ruled surfaces, 18 pages. 2001.
- #512 S. Izumiya, Generating families of developable surfaces in  $R^3$ , 18 pages. 2001.
- #513 S. Izumiya, K. Maruyama, Transversal topology and singularities of Haefliger foliations, 8 pages. 2001.
- #514 S. Izumiya, D-H. Pei & T. Sano, Singularities of hyperbolic Gauss maps, 27 pages. 2001.
- #515 S. Izumiya, N. Takeuchi, Generic special curves, 12 pages. 2001.
- #516 S. Izumiya, D-H. Pei & T. Sano, Horospherical surfaces of curves in hyperbolic space, 9 pages. 2001.
- #517 R. Yoneda, The composition operators on weighted Bloch space, 8 pages. 2001.
- #518 M. Jinzenji, T. Sasaki,  $N = 4$  supersymmetric Yang-Mills theory on orbifold  $T^4/\mathbf{Z}_2$ , 18 pages. 2001.
- #519 Y. Giga, Viscosity solutions with shocks, 58 pages. 2001.

# ON THE WORST CONDITIONAL EXPECTATION

AKIHIKO INOUE

Department of Mathematics  
Faculty of Science  
Hokkaido University  
Sapporo 060-0810, Japan  
E-mail: inoue@math.sci.hokudai.ac.jp

ABSTRACT. We study continuous coherent risk measures on  $L^p$ , in particular, the worst conditional expectations. We show some representation theorems for them, extending the results of Artzner, Delbaen, Eber, Heath, and Kusuoka.

## 1. INTRODUCTION

The worst conditional expectations are important examples of coherent risk measures. Both concepts were introduced by Artzner *et al.* [1], [2]. There they *defined* risks (such as market risks) axiomatically and provided a unified framework for the analysis of them. In these two papers, the underlying probability space was supposed to be finite. Subsequently Delbaen [3] extended the theory of coherent risk measures to general probability spaces.

In [3], the space  $L^\infty$  of all bounded real random variables or the space  $L^0$  of all real random variables was taken as the space of risks  $X$  to be measured. However, from the viewpoint of application, the assumption  $X \in L^\infty$  seems to be inconvenient, while the space  $L^0$  seems to be too large for a simple theory. In this paper, we take the intermediate spaces  $L^p$ , in particular  $L^1$ , as the spaces of risks. We can develop a simple theory at the cost of restricting ourselves to the *continuous* coherent risk measures on  $L^p$ .

Among such coherent risk measures, we are especially interested in the law invariant ones. The worst conditional expectations are again important examples. In fact, Kusuoka [4] proved that if the probability space was standard and nonatomic, then all the law invariant coherent risk measures on  $L^\infty$  with the “Fatou property” could be represented by the worst conditional expectations. We extend this result to the continuous, law invariant, coherent risk measures on  $L^p$ .

We refer to Nakano [5] where he also considers coherent risk measures on  $L^1$ . He uses them to measure the shortfall risks that appear in hedging contingent claims under constraints on the initial capital.

Let  $(\Omega, \mathcal{F}, P)$  be a probability space. For  $1 \leq p \leq \infty$ , we write  $L^p$  for  $L^p(\Omega, \mathcal{F}, P)$  and  $\|\cdot\|_p$  for its norm. A mapping  $\rho : L^p \rightarrow \mathbf{R}$  is called a *coherent risk measure* if the following conditions are satisfied:

- (1) if  $X \geq 0$ , then  $\rho(X) \leq 0$ ;

---

*Date:* March 2, 2001.

*Key words and phrases.* coherent risk measure, worst conditional expectation, Neyman-Pearson lemma.

- (2)  $\rho(X_1 + X_2) \leq \rho(X_1) + \rho(X_2)$ ;
- (3)  $\rho(\lambda X) = \lambda \rho(X)$  for  $\lambda \geq 0$ ;
- (4)  $\rho(a + X) = \rho(X) - a$  for  $a \in \mathbf{R}$

(cf. [1], [2], and [3]). We say that  $\rho$  is *continuous* if it is so in the norm  $\|\cdot\|_p$ , or, equivalently, there exists  $C \in (0, \infty)$  such that  $|\rho(X)| \leq C\|X\|_p$  (see Lemma 2.1). From [3, Theorem 3.2] (see also [2, Proposition 4.1]), we obtain the following characterization:

**Theorem 1.1.** *Let  $1 \leq p < \infty$  and  $(1/p) + (1/q) = 1$  ( $q = \infty$  if  $p = 1$ ). Then, for a mapping  $\rho : L^p \rightarrow \mathbf{R}$ , the following conditions are equivalent:*

- (1) *the mapping  $\rho$  is a continuous coherent risk measure;*
- (2) *there exists a set  $G$  of nonnegative random variables  $g$  with  $E[g] = 1$  such that*

$$\sup_{g \in G} \|g\|_q < \infty, \quad (1.1)$$

$$\rho(X) = \sup_{g \in G} E[(-X)g] \quad (X \in L^p). \quad (1.2)$$

We refer to [5, Theorem 1.2] for an analogue of this theorem for the lower semi-continuous coherent risk measures on  $L^1$ .

Let  $\alpha \in (0, 1]$ . There are several possible definitions of the worst conditional expectation  $\rho_\alpha$ . In this paper, suggested by [3, Example 4.2], we adopt the following one:

$$\rho_\alpha(X) := \sup_{g \in G(\alpha)} E[(-X)g] \quad (X \in L^1), \quad (1.3)$$

where

$$G(\alpha) := \left\{ g : \begin{array}{l} g \text{ is a nonnegative random variable on } (\Omega, \mathcal{F}, P) \\ \text{such that } E[g] = 1, \|g\|_\infty \leq 1/\alpha \end{array} \right\}. \quad (1.4)$$

Notice that

$$\rho_1(X) = E[-X] \quad (X \in L^1).$$

We easily see that, for  $1 \leq p \leq \infty$  and  $\alpha \in (0, 1]$ , the worst conditional expectation  $\rho_\alpha$  defines a continuous coherent risk measure on  $L^p$ .

For  $0 < \alpha < 1$ , the worst conditional expectation  $\text{WCE}_\alpha$  in [2] is defined by

$$\text{WCE}_\alpha(X) := \sup \left\{ \frac{1}{P(A)} E[(-X)I_A] : A \in \mathcal{F}, P(A) > \alpha \right\}.$$

We easily see that  $\text{WCE}_\alpha$  also defines a continuous coherent risk measure on  $L^p$  with  $1 \leq p \leq \infty$ . In fact, if  $P$  is atomless, then, for  $0 < \alpha < 1$ ,  $\text{WCE}_\alpha$  and  $\rho_\alpha$  coincide on  $L^\infty$  (see [3, Example 4.2]), whence on  $L^1$ .

To give an explicit representation of  $\rho_\alpha(X)$  for  $X \in L^1$ , we use an idea in statistical hypothesis testing. Let  $X \in L^1$  and  $\alpha \in (0, 1)$ . We define two constants  $k = k(X, \alpha) \in \mathbf{R}$  and  $\gamma = \gamma(X, \alpha) \in [0, 1]$  by

$$k := \inf\{z \in \mathbf{R} : P(X > z) \leq \alpha\}$$

and

$$\gamma := \begin{cases} \frac{\alpha - P(X > k)}{P(X = k)} & \text{if } P(X = k) > 0, \\ 0 & \text{if } P(X = k) = 0, \end{cases}$$

respectively. We define the random variable  $\phi_{(X,\alpha)}$  (called the *most powerful test*) by

$$\phi_{(X,\alpha)} := I_{(X>k)} + \gamma I_{(X=k)}. \quad (1.5)$$

Notice that  $E[\phi_{(X,\alpha)}] = \alpha$ . Applying the Neyman-Pearson lemma, we prove the following theorem:

**Theorem 1.2.** *Let  $\alpha \in (0, 1)$ . Then, for  $X \in L^1$ ,*

$$\rho_\alpha(-X) = \frac{1}{\alpha} E[X\phi_{(X,\alpha)}]. \quad (1.6)$$

We say that a coherent risk measure  $\rho$  on  $L^p$  is *law invariant* if  $\rho(X) = \rho(Y)$  for every pair  $(X, Y)$  of  $X, Y \in L^p$  with the same distribution. Since the expectation on the right-hand side of (1.6) depends only on the distribution of  $X$  (and  $\alpha$ ), we immediately obtain the following corollary:

**Corollary 1.3.** *For  $\alpha \in (0, 1]$ , the worst conditional expectation  $\rho_\alpha$  is law invariant on  $L^1$ , hence on  $L^p$  with  $1 < p \leq \infty$ .*

It should be noticed that, in Corollary 1.3, there is no restriction on  $(\Omega, \mathcal{F}, P)$ . In particular, it applies to those with atoms, such as discrete probability spaces. We may regard this as one advantage of the definition (1.3) with (1.4).

For a random variable  $X$ , we write  $F_X$  for the distribution function of  $X$ :

$$F_X(x) := P(X \leq x) \quad (x \in \mathbf{R}).$$

We denote by  $Z_X$  the following Skorokhod representation of  $X$ :

$$Z_X(x) := \inf\{z \in \mathbf{R} : F_X(z) > x\} \quad (0 < x < 1).$$

Then  $Z_X$  is a non-decreasing, right continuous function on  $(0, 1)$ . We write Leb for the Lebesgue measure on  $((0, 1), \mathcal{B}(0, 1))$ . As is well known (see, e.g., Williams [7, §3.12]), as a random variable on  $((0, 1), \mathcal{B}(0, 1), \text{Leb})$ ,  $Z_X$  has the same distribution as  $X$ . Using Theorem 1.2, we prove the following theorem:

**Theorem 1.4.** *Let  $\alpha \in (0, 1]$ . Then, for  $X \in L^1$ ,*

$$\rho_\alpha(-X) = \frac{1}{\alpha} \int_{1-\alpha}^1 Z_X(x) dx. \quad (1.7)$$

Eq. (1.7) is used as the definition of  $\rho_\alpha$  on  $L^\infty$  in Kusuoka [4]. Following his method, we prove the following analogue of [4, Theorem 4]:

**Theorem 1.5.** *Let  $1 \leq p < \infty$  and  $(1/p) + (1/q) = 1$ . We assume that*

$$(\Omega, \mathcal{F}, P) \text{ is a standard nonatomic probability space.} \quad (1.8)$$

*Then, for a mapping  $\rho : L^p \rightarrow \mathbf{R}$ , the following conditions are equivalent:*

- (1) *the mapping  $\rho$  is a continuous, law invariant coherent risk measure;*
- (2) *there exists a set  $M$  of probability measures on  $((0, 1], \mathcal{B}(0, 1])$  such that*

$$\sup_{m \in M} \int_{(0,1]} \frac{1}{\alpha} m(d\alpha) < \infty \quad \text{if } p = 1, \quad (1.9)$$

$$\sup_{m \in M} \int_0^1 \left\{ \int_{[1-t,1]} \frac{1}{\alpha} m(d\alpha) \right\}^q dt < \infty \quad \text{if } 1 < p < \infty, \quad (1.10)$$

$$\rho(X) = \sup_{m \in M} \int_{(0,1]} \rho_\alpha(X) m(d\alpha) \quad (X \in L^p). \quad (1.11)$$

We prove Theorems 1.1, 1.2, and 1.4 in §2. In §3, we prove Theorem 1.5, and give an example showing that this assertion does not hold in general without the assumption (1.8).

## 2. PROOFS OF THEOREMS 1.2 AND 1.4

**Lemma 2.1.** *Let  $1 \leq p \leq \infty$ . Then, for a coherent risk measure  $\rho : L^p \rightarrow \mathbf{R}$ , the following conditions are equivalent:*

- (1) *the coherent risk measure  $\rho$  is continuous;*
- (2) *there exists  $C \in (0, \infty)$  such that  $|\rho(X) - \rho(Y)| \leq C\|X - Y\|_p$  for  $X, Y \in L^p$ ;*
- (3) *there exists  $C \in (0, \infty)$  such that  $\rho(X) \leq C\|X\|_p$  for  $X \in L^p$ ;*
- (4) *there exists  $C \in (0, \infty)$  such that  $|\rho(X)| \leq C\|X\|_p$  for  $X \in L^p$ .*

*Proof.* It is clear that (2) implies (1), and (4) implies (3) trivially. Since  $\rho(0) = 0$ , one can prove the implication (1)  $\Rightarrow$  (4) in the same way as the standard proof of boundedness of continuous linear transformations. Suppose (3). Then  $\rho(X) - \rho(Y) \leq \rho(X - Y) \leq C\|X - Y\|_p$ , and similarly  $\rho(Y) - \rho(X) \leq C\|X - Y\|_p$ . Hence (2).  $\square$

*Proof of Theorem 1.1.* (2)  $\Rightarrow$  (1) By Hölder's inequality, we have

$$\rho(X) \leq \sup\{\|g\|_q : g \in G\} \cdot \|X\|_p.$$

Using this, we can easily prove (1).

(1)  $\Rightarrow$  (2) Let  $X \in L^\infty$  and let  $X_n$  be a uniformly bounded sequence that decreases to  $X$  a.s. Then  $X_n \rightarrow X$  in  $L^p$ , and so, by the continuity of  $\rho$ , we have  $\rho(X_n) \rightarrow \rho(X)$ . This implies that the restriction of  $\rho$  on  $L^\infty$  has the Fatou property (see [3]). Therefore, by [3, Theorem 3.2], there exists a set  $G$  of nonnegative random variables  $g$  with  $E[g] = 1$  such that  $\rho(X) = \mu(X)$  for  $X \in L^\infty$ , where

$$\mu(X) := \sup_{g \in G} E[(-X)g]. \tag{2.1}$$

From this and Lemma 2.1, we see that there exists  $C \in (0, \infty)$  such that

$$E\|X|g| \leq C\|X\|_p \quad (g \in G, X \in L^\infty).$$

Following the standard method, we obtain (1.1) from this. Eq. (2.1) now defines a continuous coherent risk measure  $\mu$  on  $L^1$ , whence  $\rho$  and  $\mu$  coincide on  $L^1$ .  $\square$

*Proof of Theorem 1.2.* First we assume that  $X \geq C$  for some  $C \in \mathbf{R}$ . Clearly we may assume  $P(X > C) > 0$ . Set  $Y := X - C$ . Then  $\rho_\alpha(-X) = \rho_\alpha(-Y) + C$ . We introduce the following class of "randomized tests"  $\phi$ :

$$\Phi(\alpha) := \left\{ \phi : \begin{array}{l} \phi \text{ is a random variable on } (\Omega, \mathcal{F}, P) \text{ such that} \\ 0 \leq \phi \leq 1 \text{ (} P \text{ a.s.)}, \quad E[\phi] = \alpha \end{array} \right\}.$$

Then from the definition of  $\rho_\alpha$  it follows that

$$\rho_\alpha(-Y) = \frac{E[Y]}{\alpha} \sup \{ E^{P_1}[\phi] : \phi \in \Phi(\alpha) \},$$

where  $E^{P_1}$  stands for expectation with respect to the probability measure  $P_1$  given by

$$\frac{dP_1}{dP} = \frac{Y}{E[Y]}.$$

The Neyman-Pearson lemma (cf. [6, Chap. III, Sect. 3]) now gives

$$\sup \{ E^{P_1}[\phi] : \phi \in \Phi(\alpha) \} = E^{P_1}[\phi_{(Y, \alpha)}]$$

or

$$\rho_\alpha(-Y) = \frac{1}{\alpha} E[\phi_{(Y,\alpha)} Y].$$

We have

$$\begin{aligned} k(Y, \alpha) &= \inf\{z : P(Y > z) \leq \alpha\} \\ &= \inf\{u : P(X > u) \leq \alpha\} - C \\ &= k(X, \alpha) - C. \end{aligned}$$

So

$$\begin{aligned} \{X > k(X, \alpha)\} &= \{Y > k(Y, \alpha)\}, \\ \{X = k(X, \alpha)\} &= \{Y = k(Y, \alpha)\}, \end{aligned}$$

whence  $\phi_{(X,\alpha)} = \phi_{(Y,\alpha)}$ . Combining,

$$\begin{aligned} \rho_\alpha(-X) &= \frac{1}{\alpha} E[Y\phi_{(Y,\alpha)}] + C = \frac{1}{\alpha} E[Y\phi_{(X,\alpha)}] + \frac{C}{\alpha} E[\phi_{(X,\alpha)}] \\ &= \frac{1}{\alpha} E[X\phi_{(X,\alpha)}]. \end{aligned}$$

We now assume that  $X$  is an arbitrary element of  $L^1$ . Write

$$X_n := XI_{(X \geq -n)} - nI_{(X < -n)} \quad (n = 1, 2, \dots).$$

If  $a > -n$ , then

$$\{X_n > a\} = \{X \geq -n\} \cap \{X > a\} = \{X > a\}.$$

Thus if  $n > -k(X, \alpha)$ , then  $\phi_{(X,\alpha)} = \phi_{(X_n,\alpha)}$ , and so

$$\rho_\alpha(-X_n) = \frac{1}{\alpha} E[X_n \phi_{(X,\alpha)}].$$

Since  $X_n \rightarrow X$  as  $n \rightarrow \infty$  in  $L^1$  and  $0 \leq \phi_{(X,\alpha)} \leq 1$ , we have

$$E[X_n \phi_{(X,\alpha)}] \rightarrow E[X \phi_{(X,\alpha)}] \quad (n \rightarrow \infty);$$

also the continuity of  $\rho_\alpha$  implies that  $\rho_\alpha(-X_n)$  tends to  $\rho_\alpha(-X)$  as  $n \rightarrow \infty$ . Thus (1.6) follows.  $\square$

*Proof of Theorem 1.4.* We may (and shall) assume  $0 < \alpha < 1$ . For  $X \in L^1$ , write

$$Y_X(x) := \inf\{z \in \mathbf{R} : F_X(z) \geq x\} \quad (0 < x < 1).$$

Then  $\text{Leb}(Z_X = Y_X) = 1$  (see [7, p. 35]). So, instead of (1.7), we may prove

$$\rho_\alpha(-X) = \frac{1}{\alpha} \int_{1-\alpha}^1 Y_X(x) dx. \quad (2.2)$$

Since, as a random variable on  $((0, 1), \mathcal{B}(0, 1), \text{Leb})$ ,  $Y_X$  has the same distribution as  $X$ , Theorem 1.2 yields

$$\rho_\alpha(-X) = \frac{1}{\alpha} \int_0^1 Y_X(x) \phi_{(Y_X,\alpha)}(x) dx.$$

Now

$$k(Y_X, \alpha) = k(X, \alpha) = \inf\{z \in \mathbf{R} : F_X(z) \geq 1 - \alpha\} = Y_X(1 - \alpha).$$

Define  $c \in [1 - \alpha, 1]$  by

$$c := \inf\{x \in (0, 1) : Y_X(x) > Y_X(1 - \alpha)\} \quad (:= 1 \text{ if the set is empty}).$$



Then since  $Y_X$  is nondecreasing on  $(0, 1)$ , we have

$$\{x \in (0, 1) : Y_X(x) > Y_X(1 - \alpha)\} = \begin{cases} (c, 1) \text{ or } [c, 1) & \text{if } c < 1, \\ \emptyset & \text{if } c = 1, \end{cases}$$

whence

$$\alpha - \text{Leb}\{x \in (0, 1) : Y_X(x) > Y_X(1 - \alpha)\} = c - (1 - \alpha).$$

We also have

$$Y_X(x) = Y_X(1 - \alpha) \quad (1 - \alpha \leq x < c);$$

in particular,  $c = 1 - \alpha$  if  $\text{Leb}\{x \in (0, 1) : Y_X(x) = Y_X(1 - \alpha)\} = 0$ . Thus

$$\begin{aligned} & \int_0^1 Y_X(x) \phi_{(Y_X, \alpha)}(x) dx \\ &= Y_X(1 - \alpha) \gamma(Y_X, \alpha) \text{Leb}\{x \in (0, 1) : Y_X(x) = Y_X(1 - \alpha)\} + \int_c^1 Y_X(x) dx \\ &= Y_X(1 - \alpha) \{c - (1 - \alpha)\} + \int_c^1 Y_X(x) dx = \int_{1-\alpha}^1 Y_X(x) dx, \end{aligned}$$

whence (2.2). □

### 3. PROOF OF THEOREMS 1.5

**Proposition 3.1.** *Let  $1 \leq p < \infty$  and  $X \in L^p$ . Write*

$$X_n := -nI_{(X < -n)} + XI_{(-n \leq X \leq n)} + nI_{(X > n)} \quad (n = 1, 2, \dots). \quad (3.1)$$

Then

$$\int_0^1 |Z_X(t) - Z_{X_n}(t)|^p dt \rightarrow 0 \quad (n \rightarrow \infty).$$

*Proof.* We have

$$P(X_n \leq z) = \begin{cases} 0 & (-\infty < z < -n), \\ P(X \leq z) & (-n \leq z < n), \\ 1 & (n \leq z < \infty). \end{cases}$$

Suppose  $0 < x < P(X \leq -n)$ . Then  $Z_X(x) \leq -n$ . On the other hand, since  $x < P(X \leq -n) = P(X_n \leq -n)$ , we have  $Z_{X_n}(x) \leq -n$ , hence  $= -n$ . Thus

$$|Z_X(x) - Z_{X_n}(x)| \leq |Z_X(x)| \quad (0 < x < P(X \leq -n)).$$

Next we suppose  $P(X \leq -n) \leq x < P(X < n)$ . Then since

$$\lim_{z \uparrow n} P(X \leq z) = P(X < n),$$

we see that  $P(X \leq n - \epsilon) > x$  for  $\epsilon > 0$  small enough. So

$$Z_X(x) = \inf\{-n < z < n : P(X \leq z) > x\}.$$

On the other hand,

$$P(X_n \leq -n) = P(X \leq -n) \leq x < P(X < n) = P(X_n < n),$$

so that, similarly

$$Z_{X_n}(x) = \inf\{-n < z < n : P(X_n \leq z) > x\}.$$

Since  $P(X \leq z) = P(X_n \leq z)$  for  $-n < z < n$ , it follows that

$$Z_X(x) = Z_{X_n}(x) \quad (P(X \leq -n) \leq x < P(X < n)).$$

Finally we suppose  $P(X < n) \leq x < 1$ . Then, for any  $\epsilon > 0$ , we have

$$x \geq P(X < n) = P(X_n < n) \geq P(X_n \leq n - \epsilon),$$

so that  $Z_{X_n}(x) \geq n - \epsilon$ . Since  $\epsilon$  is arbitrary, we see that  $Z_{X_n}(x) \geq n$ , hence  $= n$ . Similarly, we have  $Z_X(x) \geq n$ . Thus

$$|Z_X(x) - Z_{X_n}(x)| \leq |Z_X(x)| \quad (P(X < n) \leq x < 1).$$

Combining,

$$\begin{aligned} & \int_0^1 |Z_X(x) - Z_{X_n}(x)|^p dx \\ & \leq \int_0^{P(X \leq -n)} |Z_X(x)|^p dx + \int_{P(X < n)}^1 |Z_X(x)|^p dx \rightarrow 0 \quad (n \rightarrow \infty). \end{aligned}$$

Thus we obtain the proposition.  $\square$

By assumption (1.8), we may (and shall) assume that

$$(\Omega, \mathcal{F}, P) \text{ is equal to the Lebesgue space } ((0, 1), \mathcal{B}(0, 1), \text{Leb}). \quad (3.2)$$

Then, for a random variable  $X$  on  $(\Omega, \mathcal{F}, P)$ , the Skorokhod representation  $Z_X$  is again a random variable on  $(\Omega, \mathcal{F}, P)$ . Now if  $X$  is nondecreasing and right continuous, then

$$X(t) = Z_X(t) \quad (0 < t < 1). \quad (3.3)$$

Indeed, from the definition of  $Z_X$ , we find that

$$P(X \leq Z_X(t)) \geq t, \quad P(X < Z_X(t)) \leq t \quad (0 < t < 1).$$

The latter implies  $X(s) \geq Z_X(t)$  for  $t < s < 1$ , whence  $X(t) \geq Z_X(t)$ . Similarly, the former implies  $X(s) \leq Z_X(t)$  for  $0 < s < t$ , whence  $X(t-) \leq Z_X(t)$ . Since  $X(t-) = X(t)$  a.s., we have  $X(t) = Z_X(t)$  a.s. However, both  $X$  and  $Z_X$  are right continuous, whence (3.3).

For  $1 \leq q \leq \infty$ , we define

$$G^q := \{g \in L^q : g \geq 0 \text{ (} P \text{ a.s.)}, E[g] = 1\}.$$

Recall that, for a random variable  $X$ , we denote by  $F_X$  its distribution function.

**Proposition 3.2.** *Let  $1 \leq p < \infty$  and  $(1/p) + (1/q) = 1$ . Then, for  $X \in L^p$  and  $g \in G^q$ ,*

$$E[Z_X Z_g] = \sup\{E[Xf] : f \in G^q, F_f = F_g\}.$$

*Proof.* Write

$$\mu(X) := \sup\{E[(-X)f] : f \in G^q, F_f = F_g\} \quad (X \in L^p).$$

Then since  $\|f\|_q = \|g\|_q$  if  $F_f = F_g$ , the mapping  $\mu : L^p \rightarrow \mathbf{R}$  defines a continuous coherent risk measure on  $L^p$ . Define  $X_n$  by (3.1). Then  $X_n \in L^\infty$ , and so [4, Proposition 14] implies  $E[Z_{X_n} Z_g] = \mu(-X_n)$ . Now  $\|X - X_n\|_p \rightarrow 0$ , so that the continuity of  $\mu$  implies  $\mu(-X_n) \rightarrow \mu(-X)$ . On the other hand, by Proposition 3.1, we have

$$\lim_{n \rightarrow \infty} E[Z_{X_n} Z_g] = E[Z_X Z_g].$$

Thus the proposition follows.  $\square$

**Proposition 3.3.** *Let  $1 \leq p < \infty$  and  $(1/p) + (1/q) = 1$ . Then, for a mapping  $\rho : L^p \rightarrow \mathbf{R}$ , the following conditions are equivalent:*

- (1) *the mapping  $\rho$  is a continuous, law invariant coherent risk measure;*

(2) there exists a set  $H$  of nondecreasing, right continuous probability density functions on  $(0, 1)$  such that

$$\sup_{h \in H} \|h\|_q < \infty, \quad (3.4)$$

$$\rho(-X) = \sup_{h \in H} E[Z_X h] \quad (X \in L^p). \quad (3.5)$$

*Proof.* (2)  $\Rightarrow$  (1) Clearly  $\rho$  is law invariant. Define  $G$  by

$$G := \{g \in G^q : Z_g \in H\}.$$

Then since  $Z_h = h$  for  $h \in H$ , we see that  $\{Z_g : g \in G\} = H$ . Thus

$$\rho(-X) = \sup_{g \in G} E[Z_X Z_g].$$

Now if  $g \in G$  and  $F_f = F_g$  for  $f \in G^q$ , then  $f \in G$ . So by Proposition 3.2 we have

$$\sup_{g \in G} E[Z_X Z_g] = \sup_{g \in G} \sup_{f \in G^q, F_f = F_g} \{E[Xf] : f \in G^q, F_f = F_g\} = \sup_{g \in G} E[Xg].$$

Moreover,

$$\sup_{g \in G} \|g\|_q = \sup_{g \in G} \|Z_g\|_q = \sup_{h \in H} \|h\|_q < \infty.$$

Thus  $\rho$  is a continuous coherent risk measure on  $L^p$ .

(1)  $\Rightarrow$  (2) The restriction of  $\rho$  on  $L^\infty$  defines a law invariant, coherent risk measure on  $L^\infty$  with the Fatou property (see the proof of Theorem 1.1). So, by [4, Proposition 16], there exists a set  $H$  of nondecreasing, right continuous probability density functions on  $(0, 1)$  such that

$$\rho(-X) = \sup_{h \in H} E[Z_X h] \quad (X \in L^\infty). \quad (3.6)$$

By Lemma 2.1, we can take  $C > 0$  such that  $\rho(-X) \leq C\|X\|_p$  for  $X \in L^p$ .

Let  $h \in H$ . Then since  $Z_h = h$ , it follows from [4, Proposition 14] that, for any nonnegative  $X \in L^\infty$ ,

$$E[Xh] \leq E[Z_X Z_h] = E[Z_X h] \leq \rho(-X) \leq C\|X\|_p.$$

This implies

$$E[|X|h] \leq C\|X\|_p \quad (h \in H, X \in L^\infty).$$

From this, we obtain (3.4). So if we write

$$\mu(X) := \sup_{h \in H} E[(-X)h] \quad (X \in L^p),$$

then  $\mu$  defines a continuous coherent risk measure on  $L^p$ . For  $X \in L^p$ , we define  $X_n$  by (3.1). Then (3.6) implies  $\rho(-X_n) = \mu(-Z_{X_n})$  for  $n = 1, 2, \dots$ . If we let  $n \rightarrow \infty$ , then by Proposition 3.1 we obtain  $\rho(-X) = \mu(Z_X)$  or (3.5).  $\square$

*Proof of Theorem 1.5.* We prove the implication (1)  $\Rightarrow$  (2). Let  $H$  be as in Proposition 3.3. For  $h \in H$ , we define

$$f(t) = \begin{cases} 0 & (-\infty < t < 0), \\ h(0+) & (t = 0), \\ h(t) & (0 < t < 1). \end{cases} \quad (3.7)$$

Then, as in the proof of [4, Theorem 4], it follows from Theorem 1.4 that, for  $X \in L^p$ ,

$$\begin{aligned} E[Z_X h] &= \int_{[0,1)} \left\{ \int_x^1 Z_X(t) dt \right\} df(x) = \int_{[0,1)} \rho_{1-x}(-X)(1-x) df(x) \\ &= \int_{(0,1]} \rho_\alpha(-X) m(d\alpha), \end{aligned}$$

where  $m$  is the probability measure on  $(0, 1]$  defined by

$$m(d\alpha) := \alpha df \circ \phi \tag{3.8}$$

with  $\phi(t) := 1 - t$ . Now if  $1 < p < \infty$ , then

$$\sup_{m \in M} \int_0^1 \left\{ \int_{[1-t,1]} \frac{1}{\alpha} m(d\alpha) \right\}^q dt = \int_0^1 h(t)^q dt = \|h\|_q^q,$$

while if  $p = 1$ , then

$$\int_{(0,1]} \frac{1}{\alpha} m(d\alpha) = h(1-) = \|h\|_\infty.$$

Thus (2) follows from Proposition 3.3.

The proof of the implication (2)  $\Rightarrow$  (1) is similar. For  $m \in M$ , we define the nondecreasing, right continuous function  $h$  on  $(0, 1)$  so that (3.7) and (3.8) hold. We prove the representation (3.5) with (3.4) from (2), and apply Proposition 3.3 to obtain (1).  $\square$

**Example 3.4.** In this example, we show that Theorem 1.5 does not hold in general without (1.8). We set  $\Omega = \{\omega_1, \omega_2\}$  and  $\mathcal{F} = \{\emptyset, \Omega, \{\omega_1\}, \{\omega_2\}\}$ . Choose  $p_1, p_2$  so that  $0 < p_1 < p_2 < 1$ ,  $p_1 + p_2 = 1$ , and define the probability measure  $P$  on  $(\Omega, \mathcal{F})$  by  $P\{\omega_1\} = p_1$ ,  $P\{\omega_2\} = p_2$ . Then two random variables on  $\Omega$  have the same distribution if and only if they are identical. Thus, on this probability space, all the coherent risk measures are law invariant. Take  $g_1 > 0$  and  $g_2 > 0$  so that  $g_1 p_1 + g_2 p_2 = 1$ ,  $g_1 p_1 > g_2 p_2$ . We define  $\rho$  by

$$\rho(-X) := E[Xg] = X_1 g_1 p_1 + X_2 g_2 p_2,$$

where  $X(\omega_i) = X_i$ ,  $g(\omega_i) = g_i$  for  $i = 1, 2$ . Now suppose Theorem 1.5 (2) holds. Then as in the proof of (2)  $\Rightarrow$  (1) in Theorem 1.5, we get the representation (3.5). If we take  $(0, 1)$  and  $(1, 0)$  as  $(X_1, X_2)$ , then we obtain

$$g_2 p_2 = \sup_{h \in H} \int_{p_1}^1 h(t) dt, \quad g_1 p_1 = \sup_{h \in H} \int_{p_2}^1 h(t) dt.$$

However this contradicts the assumption  $g_1 p_1 > g_2 p_2$  since  $p_1 < p_2$  and  $h \geq 0$  for  $h \in H$ .

**Acknowledgements** I should like to thank Yumiharu Nakano for valuable discussions.

#### REFERENCES

- [1] Ph. Artzner, F. Delbaen, J.-M. Eber, and D. Heath, Thinking coherently, *Risk* **10** (1999), November, 68–71.
- [2] Ph. Artzner, F. Delbaen, J.-M. Eber, and D. Heath, Coherent measures of risk, *Math. Finance* **9** (1999), 203–228.
- [3] F. Delbaen, Coherent risk measures on general probability spaces, preprint.

- [4] S. Kusuoka, On law invariant coherent risk measures, preprint.
- [5] Y. Nakano, Efficient hedging with coherent risk measure, preprint.
- [6] L. Schmetterer, "Introduction to mathematical statistics," Springer-Verlag, New York, 1974.
- [7] D. Williams, "Probability with martingales." Cambridge Univ. Press, Cambridge, 1991.