



Title	Covariance kernel and the central limit theorem in the total variation distance
Author(s)	Mikami, T.
Citation	Hokkaido University Preprint Series in Mathematics, 523, 1-12
Issue Date	2001-04
DOI	10.14943/83669
Doc URL	http://hdl.handle.net/2115/69273
Type	bulletin (article)
File Information	pre523.pdf



[Instructions for use](#)

Covariance kernel and the central limit theorem
in the total variation distance

Toshio Mikami

Series #523, April 2001

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #498 Y. Giga, M. Paolini and P. Rybka, On the motion by singular interfacial energy, 21 pages. 2000.
- #499 J. Escher, Y. Giga and K. Ito, On a limiting motion and self-interactions of curves moved by the intermediate surface diffusion flow, 12 pages. 2000.
- #500 I. Tsuda and S. Kuroda, Cantor coding in the hippocampus, 20 pages. 2000.
- #501 M. Tsujii, Fat solenoidal attractors, 20 pages. 2000.
- #502 A. Arai, Ground state of the massless Nelson model without infrared cutoff in a non-Fock representation, 19 pages. 2000.
- #503 Y. Giga, S. Matsui and O. Sawada, Global existence of two-dimensional Navier-Stokes flow with nondecaying initial velocity, 19 pages. 2000.
- #504 A. Inoue and Y. Kasahara, Partial autocorrelation functions of the fractional ARIMA processes with negative degree of differencing, 14 pages. 2000.
- #505 T. Nakazi, Interpolation problem for ℓ^1 and a uniform algebra, 12 pages. 2000.
- #506 R. Kobayashi and Y. Giga, On anisotropy and curvature effects for growing crystals, 38 pages. 2000.
- #507 A. Arai, Instability in the spectral and the Fredholm properties of an infinite dimensional Dirac operator on the abstract Boson-Fermion Fock space, 6 pages. 2000.
- #508 A. Arai, Supersymmetric methods for constructing soliton-type solutions to multi-component nonlinear Schrödinger and Klein-Gordon equations, 22 pages. 2000.
- #509 Y. Tonegawa, Phase field model with a variable chemical potential, 30 pages. 2000.
- #510 Y. Giga, Shocks and very strong vertical diffusion, 11 pages. 2000.
- #511 S. Izumiya & N. Takeuchi, Special curves and ruled surfaces, 18 pages. 2001.
- #512 S. Izumiya, Generating families of developable surfaces in \mathbb{R}^3 , 18 pages. 2001.
- #513 S. Izumiya, K. Maruyama, Transversal topology and singularities of Haefliger foliations, 8 pages. 2001.
- #514 S. Izumiya, D-H. Pei & T. Sano, Singularities of hyperbolic Gauss maps, 27 pages. 2001.
- #515 S. Izumiya, N. Takeuchi, Generic special curves, 12 pages. 2001.
- #516 S. Izumiya, D-H. Pei & T. Sano, Horospherical surfaces of curves in hyperbolic space, 9 pages. 2001.
- #517 R. Yoneda, The composition operators on weighted Bloch space, 8 pages. 2001.
- #518 M. Jinzenji, T. Sasaki, $N = 4$ supersymmetric Yang-Mills theory on orbifold- T^4/\mathbb{Z}_2 , 18 pages. 2001.
- #519 Y. Giga, Viscosity solutions with shocks, 58 pages. 2001.
- #520 A. Inoue, On the worst conditional expectation, 10 pages. 2001.
- #521 Yumiharu Nakano, Efficient hedging with coherent risk measure, 10 pages. 2001.
- #522 T. Nakazi, Toeplitz operators and weighted norm inequalities on the bidisc, 15 pages. 2001.

Covariance kernel and the central limit theorem in the total variation distance

Toshio Mikami
Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan

April 2, 2001

Abstract

We introduce covariance kernels for Borel probability measures on \mathbf{R}^d , and study the relation between the central limit theorem in the total variation distance and the convergence of covariance kernels.

1 Introduction.

Cacoullos and Papathanasiou (1989) introduced a function, called a covariance kernel or ω -function $\omega(x)$, for a probability density function $f(x)$ on \mathbf{R} to study the characterization of probability distributions. It is known that $f(x)$ is normal if and only if $\omega(x) \equiv 1$ (see Cacoullos, Papathanasiou and Utev (1992)). Cacoullos, Papathanasiou and Utev (1994) proved that the convergence, as $n \rightarrow \infty$ in $L^1(\mathbf{R}, dx)$, of a sequence of probability density functions $\{f_n\}_{n=1}^\infty$ with interval supports on \mathbf{R} to $g_1(x) \equiv (2\pi)^{-1/2} \exp(-x^2/2)$ is equivalent to that of $\{\omega_n f_n - f_n\}_{n=1}^\infty$ to 0, where ω_n denotes a ω -function of f_n .

Mikami (1998) generalized their result, by a different method, to the case where probability measures under consideration are Borel probability measures on \mathbf{R} .

Cacoullos and Papathanasiou (1992) introduced a covariance kernel for a probability density function $f(x)$ on \mathbf{R}^d for $d \geq 2$. Papathanasiou (1996)

used it to show that $L^1(\mathbf{R}^d, dx)$ -norm of $f - g_d$ ($g_d(x) \equiv \prod_{i=1}^d g_1(x_i)$ for $x = (x_i)_{i=1}^d \in \mathbf{R}^d$) is dominated by that of $(\omega_f^i f - f)_{i=1}^d$, where $(\omega_f^i)_{i=1}^d$ denotes a covariance kernel (vector) of f . Papadatos and Papathanasiou (1998) studied the relation between $L^1(\mathbf{R}^d, dx)$ -norm of $f_1 - f_2$ and covariance kernels of f_1 and f_2 and of their marginals for two probability density functions f_1 and f_2 on \mathbf{R}^d .

In these papers they assumed that

$$\sigma_k \equiv \left(\int_{\mathbf{R}^d} y_i y_j f_k(y) dy - \int_{\mathbf{R}^d} y_i f_k(y) dy \int_{\mathbf{R}^d} y_j f_k(y) dy \right)_{i,j=1}^d \quad (1.1)$$

is positive definite for $k = 1, 2$, and that the following holds:

$$\left(\int_{\mathbf{R}^d} (\sigma_k^{-1} y)_i f_k(y) dy \right) \left(\int_{\mathbf{R}} f_k(x) dx_i \right) = \int_{\mathbf{R}} (\sigma_k^{-1} x)_i f_k(x) dx_i \quad (1.2)$$

for all $i = 1, \dots, d$ and $k = 1, 2$, and that f_1 and f_2 have convex supports since they used an identity in Cacoullos and Papathanasiou (1992). They also considered the discrete case under a similar condition.

In this paper we modify the definition of a covariance kernel so that it can be defined for any Borel probability measure on \mathbf{R}^d . We also show, without such a restriction as above, that the convergence, as $n \rightarrow \infty$ in the total variation distance, of a sequence of Borel probability measures $\{P_n\}_{n \geq 1}$ on \mathbf{R}^d to a standard normal distribution is equivalent to that of $\mathbf{W}(P_n) - Id \times P_n$ to 0 (see section 2 for definition), where Id denotes an $d \times d$ -identity matrix. Our proof is different from that of Mikami (1998), and our result in this paper generalizes it to a multi-dimensional case.

In section 2 we state our main result which will be proved in section 3.

2 Main Result.

Before we introduce a modified definition of a covariance kernel, we give some notations.

For a Borel probability measure P on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$ and any set S and $S' \subset \{1, \dots, d\}$ for which $S \cap S' = \emptyset$ and for which $S' \neq \{1, \dots, d\}$, put

$$P_{(x_j)_{j \in S'}}^S (\prod_{i \in S} dx_i) \quad (2.1)$$

$$\equiv \begin{cases} \int_{\{(x_j)_{j \notin S \cup S'} \in \mathbf{R}^{d-\#(S \cup S')}\}} P_{(x_j)_{j \in S'}}(\Pi_{i \notin S'} dx_i) & \text{if } S \neq \emptyset, S \cup S' \neq \{1, \dots, d\}, \\ P_{(x_j)_{j \in S'}}(\Pi_{i \notin S'} dx_i) & \text{if } S \cup S' = \{1, \dots, d\}, \\ 1 & \text{if } S = \emptyset, \end{cases}$$

$$P^S(\Pi_{i \in S} dx_i) \equiv P_{(x_j)_{j \in \emptyset}}^S(\Pi_{i \in S} dx_i). \quad (2.2)$$

Here $\#(S)$ denotes a cardinal number of the set S , and $P_{(x_j)_{j \in S'}}(\Pi_{i \notin S'} dx_i)$ denotes a regular conditional probability of P given $(x_j)_{j \in S'}$ (see Shiryaev (1995)). When it is not confusing, we write $\{i\} \equiv i$, $(x_j)_{\{j:j < i\}} \equiv (x_j)_{j < i}$, $(x_j)_{\{j:j \neq i\}} \equiv (x_j)_{j \neq i}$, etc, for the sake of simplicity.

The following definition is a modification of covariance kernels in Cacoullos and Papathanasiou (1992), and generalizes that in Mikami (1998) to a multi-dimensional case.

Definition 2.1 For a Borel probability measure P on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$ such that $\int_{\mathbf{R}^d} |y|^2 P(dy) < \infty$, put for $i = 1, \dots, d$,

$$\mathbf{W}^i(P)(dx) \equiv P^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) dx_i \quad (2.3)$$

$$\times \int_{-\infty}^{x_i} \left\{ \int_{\mathbf{R}} z P_{(x_j)_{j \neq i}}(dz) - y \right\} P_{(x_j)_{j \neq i}}(dy),$$

$$\mathbf{W}(P)(dx) \equiv (\delta_{ij} \mathbf{W}^i(P)(dx))_{i,j=1}^d. \quad (2.4)$$

Here we put $\delta_{ij} = 1$ if $i = j$, and $= 0$ if $i \neq j$ ($1 \leq i, j \leq d$). When $\mathbf{W}^i(P)(dx)$ is absolutely continuous with respect to dx , we put $\mathbf{W}^i(P)(dx)/dx \equiv W^i(P)(x)$ and $\mathbf{W}(P)(dx)/dx \equiv W(P)(x)$.

Remark 2.1 Suppose that $\int_{\mathbf{R}^d} x^i P(dx) = 0$ and $\int_{\mathbf{R}^d} |x^i|^2 P(dx) = 1$ for $i = 1, \dots, d$. Then $\mathbf{W}^i(P)$ is a probability measure only when $d = 1$ (see (3.12)). Suppose also that $P(dx)/dx \equiv p(x)$ exists. Then the covariance kernel ω_p^i in Cacoullos and Papathanasiou (1992) is equal to $W^i(P)(x)/p(x)$ if $P(dx)$ is a product measure.

For two finite measures P and Q on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$, let

$$\rho(P(dx), Q(dx)) \equiv \sup \left\{ \left| \int_{\mathbf{R}^d} \varphi(x) (P(dx) - Q(dx)) \right| \right. \quad (2.5)$$

$$\left. : \varphi \text{ is Borel measurable from } \mathbf{R}^d \text{ to } [-1, 1] \right\}$$

denote the total variation distance between them.

Remark 2.2 For two probability measures P and Q on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$,

$$\rho(P(dx), Q(dx)) = 2 \sup_{A \in \mathbf{B}(\mathbf{R}^d)} |P(A) - Q(A)| \quad (2.6)$$

(see Shiryaev (1995, p. 360, Lemma 1)).

The following is our main result.

Theorem 2.1 Suppose that $\{P_n\}_{n \geq 1}$ is a sequence of Borel probability measures on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$ such that $\int_{\mathbf{R}^d} |y_i|^2 P_n(dy) = 1$ ($1 \leq i \leq d, 1 \leq n$). Then the following (I) and (II) are equivalent.

$$(I). \quad \lim_{n \rightarrow \infty} \rho(P_n(dx), g_d(x)dx) = 0.$$

$$(II). \quad \lim_{n \rightarrow \infty} \sum_{i=1}^d \rho(P_n(dx), W^i(P_n)(dx)) = 0.$$

Roughly speaking, Theorem 2.1 means that the central limit theorem in the total variation distance is equivalent to the convergence of nonnegative definite matrices, to an identity matrix, which are coefficients of the second order differential operators of the second order PDEs that are satisfied by probability measures under consideration.

In fact, when $P(dx)/dx \equiv p(x)$ exists, $W(P)(x)$ is a nonnegative definite matrix and the following holds: for any $\varphi \in C_o^\infty(\mathbf{R}^d; \mathbf{R})$,

$$\begin{aligned} & \int_{\mathbf{R}^d} \sum_{i,j=1}^d (\delta_{ij} W^i(P)(x)/p(x)) (\partial^2 \varphi(x) / \partial x_i \partial x_j) p(x) dx \quad (2.7) \\ &= - \int_{\mathbf{R}^d} \sum_{i=1}^d \left(\int_{\mathbf{R}} z P_{(x_j)_{j \neq i}}(dz) - x_i \right) (\partial \varphi(x) / \partial x_i) p(x) dx. \end{aligned}$$

If (I) or (II) in Theorem 2.1 holds, then

$$\lim_{n \rightarrow \infty} \sum_{i=1}^d \int_{\mathbf{R}^d} \left| \int_{\mathbf{R}} z (P_n)_{(x_j)_{j \neq i}}(dz) \right|^2 P_n(dx) = 0 \quad (2.8)$$

(see Lemmas 3.2 and 3.3).

3 Proof.

Before we prove Theorem 2.1, we state and prove technical lemmas.

Lemma 3.1 *For any Borel probability measure P on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$,*

$$\begin{aligned} & \rho(P(dx), g_d(x)dx) \tag{3.1} \\ & \leq \sum_{i=1}^d \rho(P(dx), g_1(x_i)dx_i P^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j)) \\ & \leq 2d\rho(P(dx), g_d(x)dx). \end{aligned}$$

(Proof). When $d = 1$, (3.1) is true (see (2.1)). Suppose that $d > 1$. Then one can show the following by induction in d :

$$\begin{aligned} & P(dx) - g_d(x)dx \\ & = \sum_{i=1}^d \Pi_{1 \leq k \leq i-1} g_1(x_k) dx_k (P^{(j)_{j \geq i}}(\Pi_{j \geq i} dx_j) - g_1(x_i) dx_i P^{(j)_{j > i}}(\Pi_{j > i} dx_j)), \end{aligned}$$

where we put $\Pi_{1 \leq k \leq 0} g_1(x_k) dx_k \equiv 1$. This together with the following proves the first inequality in (3.1): for $i = 2, \dots, d$,

$$\begin{aligned} & P^{(j)_{j \geq i}}(\Pi_{j \geq i} dx_j) - g_1(x_i) dx_i P^{(j)_{j > i}}(\Pi_{j > i} dx_j) \\ & = \int_{\{(x_j)_{j < i} \in \mathbf{R}^{i-1}\}} (P(dx) - g_1(x_i) dx_i P^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j)). \end{aligned}$$

The second inequality in (3.1) can be shown by the following: for $i = 1, \dots, d$,

$$\begin{aligned} & P(dx) - g_1(x_i) dx_i P^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \\ & = P(dx) - g_d(x) dx + g_1(x_i) dx_i \int_{\{x_i \in \mathbf{R}\}} (g_d(x) dx - P(dx)). \end{aligned}$$

Q. E. D.

Lemma 3.2 *Suppose that $d > 1$, and that a sequence of Borel probability measures $\{P_n\}_{n \geq 1}$ on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$ satisfies the following: for some $i \in \{1, \dots, d\}$,*

$$\lim_{n \rightarrow \infty} \rho(P_n(dx), g_1(x_i) dx_i (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j)) = 0, \quad (3.2)$$

and $\int_{\mathbf{R}^d} |x_i|^2 P_n(dx) = 1$ for all $n \geq 1$. Then the following holds:

$$\lim_{n \rightarrow \infty} \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{\mathbf{R}} x_i (P_n)^i_{(x_j)_{j \neq i}}(dx_i) \right|^2 = 0. \quad (3.3)$$

(Proof). For $R > 0$,

$$\begin{aligned} & \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{\mathbf{R}} x_i (P_n)^i_{(x_j)_{j \neq i}}(dx_i) \right|^2 \\ & \leq 2 \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{-R}^R x_i (P_n)^i_{(x_j)_{j \neq i}}(dx_i) \right|^2 \\ & \quad + 2 \int_{\{x \in \mathbf{R}^d: |x_i| \geq R\}} |x_i|^2 P_n(dx). \end{aligned} \quad (3.4)$$

The first part of the right hand side of (3.4) can be shown to converge to zero as $n \rightarrow \infty$ by the following:

$$\left| \int_{-R}^R x_i (P_n)^i_{(x_j)_{j \neq i}}(dx_i) \right| \leq R, \quad (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) - a.s., \quad (3.5)$$

and

$$\begin{aligned} & \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{-R}^R x_i (P_n)^i_{(x_j)_{j \neq i}}(dx_i) \right| \\ & = \sum_{k=1}^2 (-1)^k \int_{A_{n,k}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \\ & \quad \times \left\{ R((P_n)^i_{(x_j)_{j \neq i}}((-\infty, R]) - \int_{-\infty}^R g_1(x_i) dx_i) \right. \\ & \quad + (P_n)^i_{(x_j)_{j \neq i}}((-\infty, -R]) - \int_{-\infty}^{-R} g_1(x_i) dx_i \\ & \quad \left. - \int_{-R}^R ((P_n)^i_{(x_j)_{j \neq i}}((-\infty, x_i]) - \int_{-\infty}^{x_i} g_1(y) dy) dx_i \right\} \\ & \leq 8R \rho(P_n(dx), g_1(x_i) dx_i (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j)), \end{aligned} \quad (3.6)$$

where we put

$$A_{n,1} \equiv \{(x_j)_{j \neq i} \in \mathbf{R}^{d-1} : \int_{-R}^R x_i (P_n)_{(x_j)_{j \neq i}}^i(dx_i) < 0\},$$

$$A_{n,2} \equiv \{(x_j)_{j \neq i} \in \mathbf{R}^{d-1} : \int_{-R}^R x_i (P_n)_{(x_j)_{j \neq i}}^i(dx_i) \geq 0\}.$$

In (3.6) we used the following:

$$\int_{-R}^R x_i g_1(x_i) dx_i = 0.$$

The second part of the right hand side of (3.4) can be shown to converge to zero as $n \rightarrow \infty$ by the following: by (3.2),

$$\int_{\{x \in \mathbf{R}^d : |x_i| \geq R\}} |x_i|^2 P_n(dx) \xrightarrow{n \rightarrow \infty} 1 - \int_{-R}^R |y|^2 g_1(y) dy \xrightarrow{R \rightarrow \infty} 0. \quad (3.7)$$

Q. E. D.

Lemma 3.3 *Suppose that $d > 1$, and that a sequence of Borel probability measures $\{P_n\}_{n \geq 1}$ on $(\mathbf{R}^d, \mathbf{B}(\mathbf{R}^d))$ satisfies the following: for some $i \in \{1, \dots, d\}$,*

$$\lim_{n \rightarrow \infty} \rho(W^i(P_n)(dx), P_n(dx)) = 0, \quad (3.8)$$

and that $\int_{\mathbf{R}^d} |x_i|^2 P_n(x) dx = 1$ for all $n \geq 1$. Then the following holds:

$$\lim_{n \rightarrow \infty} \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{\mathbf{R}} y (P_n)_{(x_j)_{j \neq i}}^i(dy) \right|^2 = 0, \quad (3.9)$$

and for any $R > 0$,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sup_{A \in \mathbf{B}(\mathbf{R}^d)} \left| \int_{\{(x_j)_{j \neq i}, y\} \in A : |y| \leq R} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) g_1(y) dy \right. \\ & \quad \left. \times \left(\int_{-\infty}^y -z (P_n)_{(x_j)_{j \neq i}}^i(dz) / g_1(y) - \int_{-\infty}^0 -z (P_n)_{(x_j)_{j \neq i}}^i(dz) / g_1(0) \right) \right| = 0, \end{aligned} \quad (3.10)$$

and

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sup_{A \in \mathbf{B}(\mathbf{R}^{d-1})} \left| \int_A (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) \right. \\ & \quad \left. \times \left(\int_{-\infty}^0 -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(0) - 1 \right) \right| = 0. \end{aligned} \quad (3.11)$$

(Proof). (3.9) can be proved by (3.8) and by the following:

$$\begin{aligned} & \int_{\mathbf{R}^d} W^i(P_n)(dx) \\ = & \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) \\ & \times \left\{ \int_{-\infty}^0 (-y) \left(\int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) - y \right) (P_n)_{(x_j)_{j \neq i}}^i(dy) \right. \\ & \left. - \int_0^{\infty} y \left(\int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) - y \right) (P_n)_{(x_j)_{j \neq i}}^i(dy) \right\} \\ = & - \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}} ((x_j)_{j \neq i}) \Pi_{j \neq i} dx_j \left| \int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) \right|^2 + 1, \end{aligned} \quad (3.12)$$

where we used the following:

$$\int_{\mathbf{R}} \left(\int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) - y \right) (P_n)_{(x_j)_{j \neq i}}^i(dy) = 0. \quad (3.13)$$

(3.10) can be proved by (3.8)-(3.9) and by the following: for $y \in [-R, R]$,

$$\begin{aligned} & \int_{-\infty}^y -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(y) - \int_{-\infty}^0 -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(0) \\ = & \int_0^y x_i g_1(x_i)^{-1} \left(\int_{-\infty}^{x_i} -z(P_n)_{(x_j)_{j \neq i}}^i(dz) dx_i - (P_n)_{(x_j)_{j \neq i}}^i(dx_i) \right), \end{aligned} \quad (3.14)$$

and

$$\begin{aligned} & (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) \int_{-\infty}^{x_i} -z(P_n)_{(x_j)_{j \neq i}}^i(dz) dx_i - P_n(dx) \\ = & -(P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) \int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) (P_n)_{(x_j)_{j \neq i}}^i((-\infty, x_i]) dx_i \\ & + W^i(P_n)(dx) - P_n(dx). \end{aligned} \quad (3.15)$$

(3.11) can be proved by (3.8)-(3.10) and by the following: for $R > 0$,

$$\begin{aligned}
& (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left(1 - \int_{-\infty}^0 -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(0)\right) \quad (3.16) \\
= & (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left\{ \int_{\{x_i \in \mathbf{R}: |x_i| > R\}} ((P_n)_{(x_j)_{j \neq i}}^i(dx_i) \right. \\
& - \left(\int_{-\infty}^0 -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(0) \right) g_1(x_i) dx_i \\
& + \int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) \int_{-R}^R (P_n)_{(x_j)_{j \neq i}}^i((-\infty, x_i]) dx_i \\
& + \int_{-R}^R \left(\int_{-\infty}^y -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(y) \right. \\
& \left. - \int_{-\infty}^0 -z(P_n)_{(x_j)_{j \neq i}}^i(dz)/g_1(0) \right) g_1(y) dy \left. \right\} \\
& + \int_{\{x_i \in \mathbf{R}: |x_i| \leq R\}} (P_n(dx) - W^i(P_n)(dx)).
\end{aligned}$$

Q. E. D.

Finally we prove Theorem 2.1.

(Proof of Theorem 2.1). We only have to prove the case where $d > 1$. In fact, when $d = 1$, the proof is done from the case where $d > 1$, by considering probability measures $\{P_n(dx) \times g_1(y) dy\}_{n \geq 1}$ on $(\mathbf{R}^2, \mathbf{B}(\mathbf{R}^2))$.

We assume that $d > 1$ from here on.

Suppose that (I) in Theorem 2.1 holds. Then the following which will be proved later holds: for any $i \in \{1, \dots, d\}$

$$\lim_{n \rightarrow \infty} \rho(W^i(P_n)(dx), (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) g_1(x_i) dx_i) = 0, \quad (3.17)$$

which implies (II) by Lemma 3.1.

(3.17) is true. Indeed, by (3.13),

$$\begin{aligned}
& W^i(P_n)(dx) - (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) g_1(x_i) dx_i \quad (3.18) \\
= & (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) dx_i \\
& \times \left\{ 1_{(-\infty, 0]}(x_i) \left(\int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) (P_n)_{(x_j)_{j \neq i}}^i((-\infty, x_i]) \right. \right. \\
& \left. \left. - \int_{-\infty}^{x_i} y((P_n)_{(x_j)_{j \neq i}}^i(dy) - g_1(y) dy) \right) \right\}
\end{aligned}$$

$$\begin{aligned}
& -1_{(0,\infty)}(x_i) \left(\int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) (P_n)_{(x_j)_{j \neq i}}^i((x_i, \infty)) \right. \\
& \quad \left. - \int_{x_i}^{\infty} y((P_n)_{(x_j)_{j \neq i}}^i(dy) - g_1(y)dy) \right),
\end{aligned}$$

where $1_A(x)$ denotes an indicator function of the set A . We also have

$$\begin{aligned}
& \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) \right| \tag{3.19} \\
& \quad \times \left(\int_{-\infty}^0 dx_i (P_n)_{(x_j)_{j \neq i}}^i((-\infty, x_i]) + \int_0^{\infty} dx_i (P_n)_{(x_j)_{j \neq i}}^i((x_i, \infty)) \right) \\
& = \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) \right| \int_{\mathbf{R}} |y| (P_n)_{(x_j)_{j \neq i}}^i(dy) \\
& \leq \left(\int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \left| \int_{\mathbf{R}} z(P_n)_{(x_j)_{j \neq i}}^i(dz) \right|^2 \right)^{1/2} \rightarrow 0,
\end{aligned}$$

as $n \rightarrow \infty$ by Lemma 3.2. Decompose the integral in (3.18) as follows: for $R > 0$,

$$\begin{aligned}
1_{(-\infty, 0]}(x_i) \int_{-\infty}^{x_i} dy &= 1_{(-\infty, -R]}(x_i) \int_{-\infty}^{x_i} dy + 1_{(-R, 0]}(x_i) \left(\int_{-\infty}^{-R} dy + \int_{-R}^{x_i} dy \right), \\
1_{(0, \infty)}(x_i) \int_{x_i}^{\infty} dy &= 1_{(R, \infty)}(x_i) \int_{x_i}^{\infty} dy + 1_{(0, R]}(x_i) \left(\int_R^{\infty} dy + \int_{x_i}^R dy \right).
\end{aligned}$$

Then the following (3.20)-(3.21) completes the proof of (3.17).

$$\begin{aligned}
& \int_{\mathbf{R}^{d-1}} (P_n)^{(j)_{j \neq i}}(\Pi_{j \neq i} dx_j) \tag{3.20} \\
& \quad \times \left\{ \int_{-\infty}^{-R} dx_i \int_{-\infty}^{x_i} |y| \left((P_n)_{(x_j)_{j \neq i}}^i(dy) + g_1(y)dy \right) \right. \\
& \quad + \int_R^{\infty} dx_i \int_{x_i}^{\infty} |y| \left((P_n)_{(x_j)_{j \neq i}}^i(dy) + g_1(y)dy \right) \\
& \quad + R \left(\int_{-\infty}^{-R} |y| \left((P_n)_{(x_j)_{j \neq i}}^i(dy) + g_1(y)dy \right) \right. \\
& \quad \left. \left. + \int_R^{\infty} |y| \left((P_n)_{(x_j)_{j \neq i}}^i(dy) + g_1(y)dy \right) \right) \right\} \\
& = \int_{\{x \in \mathbf{R}^d: |x_i| \geq R\}} |x_i|^2 P_n(dx) + \int_{\{y \in \mathbf{R}: |y| \geq R\}} |y|^2 g_1(y)dy \rightarrow 0,
\end{aligned}$$

as $n \rightarrow \infty$ and then $R \rightarrow \infty$ (see (3.7)), and for any Borel measurable $\varphi : \mathbf{R}^d \mapsto [-1, 1]$,

$$\begin{aligned}
& \left| \int_{\mathbf{R}^d} \varphi(x) (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) dx_i \right. & (3.21) \\
& \times \{ 1_{(-R, 0]}(x_i) \int_{-R}^{x_i} y (-P_n)_{(x_j)_{j \neq i}}^i(dy) + g_1(y) dy \\
& \quad \left. + 1_{(0, R]}(x_i) \int_{x_i}^R y (P_n)_{(x_j)_{j \neq i}}^i(dy) - g_1(y) dy \} \right| \\
& \leq 2R^2 \rho(P_n(dx), (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) g_1(x_i) dx_i).
\end{aligned}$$

Suppose that (II) in Theorem 2.1 holds. Then the following which will be proved later holds: for all $i \in \{1, \dots, d\}$,

$$\lim_{n \rightarrow \infty} \rho(P_n(dx), (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) g_1(x_i) dx_i) = 0, \quad (3.22)$$

which implies (I) in Theorem 2.1 by Lemma 3.1.

We prove (3.22) to complete the proof. For any $R > 0$, by Chebychev's inequality,

$$\begin{aligned}
& \int_{\{x \in \mathbf{R}^d : |x_i| \geq R\}} (P_n(dx) + (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) g_1(x_i) dx_i) & (3.23) \\
& \leq R^{-2} \left(\int_{\{x \in \mathbf{R}^d : |x_i| \geq R\}} |x_i|^2 P_n(dx) + \int_{\{y \in \mathbf{R} : |y| \geq R\}} |y|^2 g_1(y) dy \right) < 2/R^2,
\end{aligned}$$

and for any $A \in \mathbf{B}(\mathbf{R}^d)$

$$\begin{aligned}
& \int_{\{((x_j)_{j \neq i}, x_i) \in A : |x_i| \leq R\}} (P_n(dx) - (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) g_1(x_i) dx_i) & (3.24) \\
& = \int_{\{((x_j)_{j \neq i}, x_i) \in A : |x_i| \leq R\}} (P_n(dx) - W^i(P_n)(dx)) \\
& \quad + \int_{\{((x_j)_{j \neq i}, x_i) \in A : |x_i| \leq R\}} (P_n)^{(j)_{j \neq i}} (\Pi_{j \neq i} dx_j) dx_i \\
& \quad \times \left\{ \int_{\mathbf{R}} z (P_n)_{(x_j)_{j \neq i}}^i(dz) \int_{-\infty}^{x_i} (P_n)_{(x_j)_{j \neq i}}^i(dz) \right. \\
& \quad \left. + \left(\int_{-\infty}^{x_i} -z (P_n)_{(x_j)_{j \neq i}}^i(dz) / g_1(x_i) - 1 \right) g_1(x_i) \right\},
\end{aligned}$$

which completes the proof of (3.22) by Lemma 3.3.

Q. E. D.

References

- [1] Cacoullos, T. and Papathanasiou, V. (1989). Characterizations of distributions by variance bounds. *Statist. Probab. Lett.* **7**, 351-356.
- [2] Cacoullos, T. and Papathanasiou, V. (1992). Lower variance bounds and a new proof of the central limit theorem. *J. Multivariate Anal.* **43**, 173–184.
- [3] Cacoullos, T., Papathanasiou, V. and Utev, S. A. (1992). Another characterization of the normal law and a proof of the central limit theorem connected with it. *Theory Probab. Appl.* **37**, 581-588.
- [4] Cacoullos, T., Papathanasiou, V. and Utev, S. A. (1994). Variational inequalities with examples and an application to the central limit theorem. *Ann. Probab.* **22**, 1607-1618.
- [5] Mikami, T. (1998). Equivalent conditions on the central limit theorem for a sequence of probability measures on \mathbf{R} . *Statist. Probab. Lett.* **37**, 237-242.
- [6] Papadatos, N. and Papathanasiou, V. (1998). Variational inequalities for arbitrary multivariate distributions. *J. Multivariate Anal.* **67**, 154–168.
- [7] Papathanasiou, V. (1996). Multivariate variational inequalities and the central limit theorem. *J. Multivariate Anal.* **58**, 189–196.
- [8] Shiryaev, A. N. (1995). Probability. Springer-Verlag, Berlin, Heidelberg, New York, Tokyo.