



Title	On the two-dimensional nonstationary vorticity equations
Author(s)	Giga, Y.
Citation	Hokkaido University Preprint Series in Mathematics, 538, 1-12
Issue Date	2001-09
DOI	10.14943/83683
Doc URL	<a href="http://hdl.handle.net/2115/69287">http://hdl.handle.net/2115/69287</a>
Type	bulletin (article)
File Information	pre538.pdf



[Instructions for use](#)

On the two-dimensional nonstationary vorticity  
equations

Yoshikazu Giga

Series #538 . September 2001

**HOKKAIDO UNIVERSITY**  
**PREPRINT SERIES IN MATHEMATICS**

- #509 Y. Tonegawa, Phase field model with a variable chemical potential, 30 pages. 2000.
- #510 Y. Giga, Shocks and very strong vertical diffusion, 11 pages. 2000.
- #511 S. Izumiya & N. Takeuchi, Special curves and ruled surfaces, 18 pages. 2001.
- #512 S. Izumiya, Generating families of developable surfaces in  $R^3$ , 18 pages. 2001.
- #513 S. Izumiya, K. Maruyama, Transversal topology and singularities of Haefliger foliations, 8 pages. 2001.
- #514 S. Izumiya, D-H. Pei & T. Sano, Singularities of hyperbolic Gauss maps, 27 pages. 2001.
- #515 S. Izumiya, N. Takeuchi, Generic special curves, 12 pages. 2001.
- #516 S. Izumiya, D-H. Pei & T. Sano, Horospherical surfaces of curves in hyperbolic space, 9 pages. 2001.
- #517 R. Yoneda, The composition operators on weighted Bloch space, 8 pages. 2001.
- #518 M. Jinzenji, T. Sasaki,  $N = 4$  supersymmetric Yang-Mills theory on orbifold- $T^4/\mathbf{Z}_2$ , 18 pages. 2001.
- #519 Y. Giga, Viscosity solutions with shocks, 58 pages. 2001.
- #520 A. Inoue, On the worst conditional expectation, 10 pages. 2001.
- #521 Yumiharu Nakano, Efficient hedging with coherent risk measure, 10 pages. 2001.
- #522 T. Nakazi, Toeplitz operators and weighted norm inequalities on the bidisc, 15 pages. 2001.
- #523 T. Mikami, Covariance kernel and the central limit theorem in the total variation distance, 80 pages. 2001.
- #524 K. Yamaguchi and T. Yatsui, Geometry of higher order differential equations of finite type associated with symmetric spaces, 43 pages. 2001.
- #525 T. Suwa, Residues of Chern classes, 20 pages. 2001.
- #526 V. Anh and A. Inoue, Dynamic models of asset prices with long memory, 21 pages. 2001.
- #527 T. Izawa and T. Suwa, Multiplicity of functions on singular varieties, 21 pages. 2001.
- #528 T. Nakazi and T. Yamamoto, Two dimensional commutative Banach algebras and von Neumann inequality, 18 pages. 2001.
- #529 Y. Giga, N. Ishimura and Y. Kohsaka, Spiral solutions for a weakly anisotropic curvature flow equation, 16 pages. 2001.
- #530 Y. Giga and P. Rybka, Quasi-static evolution of 3-D crystals grown from supersaturated vapor, 16 pages. 2001.
- #531 Y. Tonegawa, Remarks on convergence of the Allen-Cahn equation, 18 pages. 2001.
- #532 T. Suwa, Characteristic classes of singular varieties, 26 pages. 2001.
- #533 J. Escher, Y. Giga and K. Ito, On a limiting motion and self-intersections for the intermediate surface diffusion flow, 20 pages. 2001.
- #534 Y.-H. R. Tsai, Y. Giga and S. Osher, A level set approach for computing discontinuous solutions of a class of Hamilton-Jacobi equations, 30 pages. 2001.
- #535 A. Yamagami, On Gouvêa's conjecture in the unobstructed case, 19 pages. 2001.
- #536 A. Inoue, What does the partial autocorrelation function look like for large lags, 27 pages. 2001.
- #537 T. Nakazi and T. Yamamoto, Norm of a linear combination of two operators of a Hilbert space, 16 pages. 2001.

# On the two-dimensional nonstationary vorticity equations

Yoshikazu Giga  
Department of Mathematics  
Hokkaido University  
Sapporo 060-0810, Japan

## 1 Introduction

It is my great honor to contribute to the proceedings of the international conference on ‘Tosio Kato’s method and principle for evolution equations in mathematical physics’. Tosio Kato established mathematical foundations for various important partial differential equations in physics. For example, he established a method to solve the Navier-Stokes equations by a smart application of semigroup theory in his seminal paper with H. Fujita [17], [5]. These papers are very influential in development of mathematical analysis on the Navier-Stokes equations. For example, my research career started by extending their  $L^2$  type Hilbert theory to  $L^p$ -theory [6], [7], [10]. Of course, Tosio Kato often came back to the Navier-Stokes equations and established several fundamental and interesting results based on his smart iteration method eg. [13], [14], [18], [19], [15].

In this paper we survey analytic results on the two-dimensional nonstationary vorticity equations for the Navier-Stokes flow. This topic was studied in his last paper on the Navier-Stokes equations [15]. We consider the Navier-Stokes equations

$$u_t - \Delta u + (u, \nabla)u + \nabla \pi = 0 \text{ in } \mathbf{R}^n \times (0, T), \quad (1.1)$$

$$\operatorname{div} u = 0 \text{ in } \mathbf{R}^n \times (0, T), \quad (1.2)$$

where  $u = u(x, t)$  represents the velocity field and  $\pi = \pi(x, t)$  represents the pressure field. The kinematic viscosity and density are normalized to one;  $(u, \nabla) = \sum_{j=1}^n u^j \partial / \partial x_j$  and  $u_t = \partial_t u = \partial u / \partial t$  where  $u = (u^1, \dots, u^n)$ . If  $n = 2$ , we take curl of (1.1) to get

$$\omega_t - \Delta \omega + (u, \nabla)\omega = 0 \text{ in } \mathbf{R}^2 \times (0, T) \quad (1.3)$$

with the vorticity  $\omega = \operatorname{curl} u$  which is regarded as a scalar function.

Since

$$-\Delta v = \nabla^\perp \operatorname{curl} v - \nabla \operatorname{div} v$$

for a vector field  $v$  in the plane, (1.2) implies

$$-\Delta u = \nabla^\perp \omega,$$

where  $\nabla^\perp \omega = (\partial\omega/\partial x_2, -\partial\omega/\partial x_1)$ . The last relation is formally equivalent to the Biot-Savart law

$$u = \nabla^\perp (-\Delta)^{-1} \omega. \quad (1.4)$$

The system (1.3)-(1.4) is called the vorticity equations and it is formally equivalent to the Navier-Stokes system (1.1)-(1.2).

We consider the initial-value problem of (1.3), (1.4) for the vorticity especially when initial data  $\omega_0$  is merely a finite Radon measure or  $L^1$  function. The global existence result is first proved by [11] and later improved by [15], [1]. This does not follow from a famous result of J. Leray [22] since the initial kinetic energy may not be finite. The uniqueness is still open unless point mass part of  $\omega_0$  is small [11], [15]. Large time behaviour

$$\omega(x, t) \sim m \frac{1}{4\pi} e^{-|x|^2/4t}, m = \int_{\mathbf{R}^2} \omega_0 \quad (t \rightarrow \infty)$$

is established for small total variation of  $\omega_0$  [9] and later for small  $m$  [3]. It is well-known that solution  $\omega$  becomes smooth for  $t > 0$  even if  $\omega_0$  is just a measure since the problem is parabolic. However, no one quantified this effect until Tosio Kato [15] derived the following smoothing rate estimate for  $L^p$ -norms of derivatives:

$$\sup_{t>0} t^{k+1+\frac{|\alpha|}{2}-\frac{1}{p}} \|\partial_t^k \partial_x^\alpha \omega\|_p(t) < \infty, \quad 1 < p < \infty, \quad k = 0, 1, 2, \dots \quad (1.5)$$

for multi-index  $\alpha = (\alpha_1, \alpha_2)$ ,  $|\alpha| = \alpha_1 + \alpha_2$ ,  $\partial_x^\alpha = (\partial/\partial x_1)^{\alpha_1} (\partial/\partial x_2)^{\alpha_2}$ . This is also viewed as a decay estimate of derivatives which is new. His proof is rather sophisticated by using estimates on fractional derivatives of products of functions (e.g. [20], [21]). Here we give an elementary proof for (1.5) including  $p = 1$  and  $p = \infty$  by deriving a new form of the Gronwall inequality. We shall also apply the inequality to derive smoothing rate estimate for the Navier-Stokes equation. Except the last application the contents of this paper is taken from the book [8].

## 2 Existence

We consider the initial value problem for the vorticity equations (1.3), (1.4). Let  $M(\mathbf{R}^2)$  denote the space of all finite Radon measures on  $\mathbf{R}^2$  equipped with the total variation norm  $\|\cdot\|_1$ . The space  $L^1(\mathbf{R}^2)$  is regarded as a closed subspace of  $M(\mathbf{R}^2)$ .

**Theorem 2.1.** *Let  $\omega_0 \in M(\mathbf{R}^2)$ . Then there exists a global solution  $(\omega, u)$  of (1.3), (1.4) such that*

(i)  $\omega \in C((0, \infty); L^q(\mathbf{R}^2))$  for  $1 \leq q \leq \infty$  satisfying

$$t^{1-1/q} \|\omega\|_q(t) \leq \kappa^{-(1-1/q)} \|\omega_0\|_1, \quad 1 \leq q \leq \infty. \quad (2.1)$$

with a numerical constant  $\kappa$ ;

(ii)  $\omega(t) \rightarrow \omega_0$  in the weak \* topology of  $M(\mathbf{R}^2)$  as  $t \rightarrow 0$ ;

(iii)  $\omega(x, t)$  is smooth for  $t > 0$ .

The existence of a smooth solution for  $\omega_0 \in M(\mathbf{R}^2)$  satisfying (i), (ii), (iii) goes back to [11] where the constant  $\kappa^{-(1-1/q)}$  in (2.1) is replaced by a constant  $C_o$  depending on  $q$  and  $\|\omega_0\|_1$ . The key to prove the global existence is to prove a priori estimate (2.1). In [15] and [1] improved a priori estimate of [11]. They prove the estimate of solution  $W$  of a perturbed heat equation

$$W_t - \Delta W + (v, \nabla)W = 0$$

with  $\operatorname{div} v = 0$  of form

$$t^{1-1/q} \|W\|_q(t) \leq \kappa^{-(1-1/q)} \|W|_{t=0}\|_1$$

with the best possible constant  $\kappa$  in the Nash inequality

$$\|\nabla \psi\|_2^2 \|\psi\|_1^2 \geq \kappa \|\psi\|_2^4.$$

Their method differ from each other. For  $W$  one is able to prove that  $\|W\|_q(t)$  is nonincreasing in time  $t$ . Thus  $\|\omega\|_q(t)$  in Theorem 2.1 is also nonincreasing in  $t$ . We refer [8] for more details as well as original papers [15], [1].

The existence of a global weak solution of (1.3), (1.4) with  $\omega_0 \in M(\mathbf{R}^2)$  has been proved in [4].

**Open problem.** Existence of solutions for measure initial vorticity is not known when we consider the Navier-Stokes equations (1.1), (1.2) on the half space with Dirichlet boundary condition  $u = 0$ .

### 3 Uniqueness

The situation of uniqueness is unsatisfactory.

**Theorem 3.1.** For each  $q \in [4/3, 2)$  there exists a constant  $\varepsilon_q > 0$  such that if

$$\overline{\lim}_{t \rightarrow 0} t^{1-\frac{1}{q}} \|e^{t\Delta} \omega_0\|_q < \varepsilon_q/2, \quad (3.1)$$

then it is possible to construct a solution  $\omega$  in Theorem 2.1 satisfying

$$\overline{\lim}_{t \rightarrow 0} t^{1-\frac{1}{q}} \|\omega\|_q(t) < \varepsilon_q. \quad (3.2)$$

The solution  $\omega$  satisfying (3.2) is unique provided that (i), (ii) of Theorem 2.1 is fulfilled where (2.1) is replaced with boundedness of  $t^{1-1/q} \|\omega\|_q^{(t)}$  on  $[0, T)$ ,  $T > 0$ . Here  $e^{t\Delta} \omega_0 = G_t * \omega_0$ ,  $G_t(x) = (4\pi t)^{-1} \exp(-|x|^2/4t)$  and  $*$  denotes the convolution on  $\mathbf{R}^2$ . Such a kind of uniqueness goes back to [11]. It is shown that if point mass part of  $\omega_0$  is small then (3.1) is fulfilled, so the uniqueness holds for example for  $\omega_0 \in L^1(\mathbf{R}^2)$ . In [15] Tosio Kato clarified the explicit value of  $\varepsilon_q$ . Especially, he proved that  $\varepsilon_{4/3} \approx 0.4922$ . If  $\|\omega_0\|_1$  is small, the uniqueness (among small solution) is easy to prove (cf. [4]).

**Open problem.** Unfortunately, the uniqueness of solution is not known even when  $\omega_0 = m\delta(x)$ , a constant multiple of Dirac's delta function when  $|m|$  is not small. For such an initial data

$$\omega(t, x) = m(4\pi t)^{-1} \exp(-|x|^2/4t)$$

is a solution of (1.3) and (1.4) satisfying (3.2). According to [15] if  $|m| < \varepsilon_q/2c_q$  with  $c_q = (4\pi)^{-(1-1/q)} q^{-1/q}$  then such a solution is unique and satisfies (3.2) with  $q = 4/3$ . (The value  $\varepsilon_q/2c_q \approx 0.5749$  for  $q = 4/3$ ) A typical question is whether  $\omega$  is a unique solution when  $|m|$  is large.

**Remark 3.2.** In Theorem 3.1 the uniqueness is actually proved for solutions of the integral equation

$$\omega(t) = e^{t\Delta} \omega_0 - \int_0^t e^{(t-s)\Delta} (u, \nabla) \omega ds \quad (3.3)$$

with (1.4). This is formally equivalent to (1.3), (1.4) with initial data  $\omega_0$ . Here  $\omega(t)$  is interpreted as a function on  $[0, \infty)$  with values in some space of functions of  $x$ -variables. Even if  $\omega_0 \in L^1(\mathbf{R}^2)$  the method (originally due to [17]) provides uniqueness for (3.3) among the class of  $\omega$  satisfying

$$\omega \in C([0, T), L^1(\mathbf{R}^2)) \text{ and } t^{1-1/q} \omega \in L^\infty(0, T; L^q(\mathbf{R}^2))$$

with

$$\lim_{t \rightarrow 0} t^{1-1/q} \|\omega\|_q(t) = 0 \quad (3.4)$$

for some  $q \in [4/3, 2)$ . The last assumption (3.4) is very convenient when the space of initial data is scaling-invariant under the rescaling  $\omega_{0\lambda}(x) = \lambda^2 \omega_0(x)$ ,  $\lambda > 0$ . The extra assumption (3.4) turns to be removed by a remark of Brezis [2]. See also [16].

## 4 Large time behaviour

If  $\omega_0$  is radially symmetric, then  $G_t * \omega_0$  is a solution of (1.3), (1.4) with initial data and for large  $t$  it is asymptotically like  $\int \omega_0 dx G_t$ . For the nonradial  $\omega_0$  we still have such a result provided that  $\|\omega_0\|_1$  is small. Let  $\omega$  be a solution of (1.3), (1.4) satisfying (2.1) with  $\omega|_{t=0} = \omega_0$ .

**Theorem 4.1.** *For each  $\delta \in (0, 1/2)$  there exists  $\varepsilon = \varepsilon(\delta) > 0$  such that if  $\|\omega_0\|_1 < \varepsilon$  then*

$$\begin{aligned} \|\omega - mG_t\|_p(t) &\leq CNt^{-1+1/p-\delta}, \quad m = \int_{\mathbf{R}^2} \omega_0 \\ \|\omega - e^{t\Delta}\omega_0\|_p(t) &\leq CNt^{-1+1/p-\delta}, \quad 1 \leq p \leq \infty, \quad t > 0 \end{aligned}$$

with a universal constant  $C$  and  $N = \| |x| \omega_0 \|_1$

This result is first proved in [9]. Later the smallness assumption is weakened by the smallness of  $|m|$  in [3] as follows.

**Theorem 4.2.** *Let  $\omega$  be a solution of (1.3), (1.4) satisfying (2.1) with  $\omega|_{t=0} = \omega_0$ . If  $|m| < \varepsilon_q/2c_q$  with  $c_q = (4\pi)^{-(1-1/q)}q^{-1/q}$  for some  $q \in [4/3, 2)$ , then*

$$\lim_{t \rightarrow \infty} t^{1-1/p} \|\omega - mG_t\|_p(t) = 0.$$

The constant  $\varepsilon_q$  is taken as in Theorem 3.1.

We apply the perturbation argument to prove Theorem 4.1 while we use the scaling argument to prove Theorem 4.2. Indeed we consider

$$\omega_\lambda(x, t) = \lambda^2 \omega(\lambda x, \lambda^2 t), \quad u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t)$$

and study the behaviour of  $(\omega_\lambda, u_\lambda)$  as  $\lambda \rightarrow \infty$ . The limit satisfies the vorticity equation with initial data  $m\delta(x)$ . If the uniqueness result apply, then the limit must equal  $mG_t$ . This yields the convergence. Unfortunately, the proof in [3] needs more explanation since the estimate (3.2) is not proved for the limit. We have given a full proof in [8] at least  $p = \infty$ .



## 5 Smoothing rate estimates

**Theorem 5.1.** *Let  $(\omega, u)$  be a solution of (1.3), (1.4) in Theorem 2.1 with initial data  $\omega_0 \in M(\mathbf{R}^2)$ . For multi-index  $\beta$  and  $b = 0, 1, 2$ , there is a constant  $C = C(\beta, b, \|\omega_0\|_1)$  such that*

$$t^{b + \frac{|\beta|}{2} + 1 - \frac{1}{p}} \|\partial_t^b \partial_x^\beta \omega\|_p(t) \leq C \|\omega_0\|_1, \quad 1 \leq p \leq \infty. \quad (5.1)$$

*There is a constant  $C' = C'(\beta, b, p, \|\omega_0\|_1)$  such that*

$$t^{b + \frac{|\beta|}{2} + \frac{1}{2} - \frac{1}{p}} \|\partial_t^b \partial_x^\beta u\|_p(t) \leq C' \|\omega_0\|_1, \quad 2 < p \leq \infty. \quad (5.2)$$

*The constants  $C, C'$  can be taken so that it is nondecreasing in  $\|\omega_0\|_1$ .*

The estimates (5.1) and (5.2) are first proved by Tosio Kato [15] excluding the case  $p = 1$  and  $p = \infty$ . His method is based on a sophisticated application of interpolation theory and fractional derivatives estimates of product of functions. We shall give an elementary proof based on the next Gronwall type inequality.

## 6 A new Gronwall type inequality

**Lemma 6.1.** *Let  $\psi$  be a continuous function on  $(0, T)$ , where  $T \in (0, \infty]$ . Let  $\alpha, \gamma, \delta$  be real numbers such that  $\gamma > 0, \delta > 0, \gamma + \delta = 1$  and  $0 < \gamma < 1$ . Let  $b_\varepsilon > 0$  be a number depending on  $\varepsilon \in (0, 1)$ . (For simplicity, assume that  $\varepsilon \mapsto b_\varepsilon$  is nonincreasing. Assume that there is  $\sigma > 0$  such that*

$$0 \leq \psi(t) \leq \sigma \left( b_\varepsilon t^{-\alpha} + \int_{t(1-\varepsilon)}^t \frac{\psi(s)}{(t-s)^\gamma s^\delta} ds \right) \quad (6.1)$$

*for all  $t \in (0, T)$ ,  $\varepsilon \in (0, 1)$ . Then there exists a constant  $C = C(\sigma, \alpha, \delta, \gamma, b_\varepsilon)$  such that*

$$\psi(t) \leq C \sigma t^{-\alpha} \text{ for all } t \in (0, T). \quad (6.2)$$

*The constant  $C$  can be taken so that it is nondecreasing in  $\sigma$ .*

**Proof.** By (6.1) we have

$$\psi(t)t^\alpha \leq \sigma \left( b_\varepsilon + t^\alpha \int_{t(1-\varepsilon)}^t \frac{\psi(s)s^\alpha}{(t-s)^\delta s^{\delta+\alpha}} ds \right), \quad t \in (0, T).$$

For  $\eta \in (0, T)$  we set

$$\varphi(t) = \sup_{\eta \leq \tau \leq t} \tau^\alpha \psi(\tau).$$

(Note that  $\sup_{0 \leq \tau \leq t} \tau^\alpha \psi(\tau)$  may be infinite so we truncate near  $\tau = 0$ .) Then for  $t > \eta/(1 - \varepsilon)$  we observe that

$$\begin{aligned} \varphi(t) &\leq \sigma \left( b_\varepsilon + t^\alpha \varphi(t) \int_{t(1-\varepsilon)}^t \frac{ds}{(t-s)^\gamma s^{\delta+\alpha}} \right) \\ &= \sigma \left( b_\varepsilon + \varphi(t) \int_{1-\varepsilon}^1 \frac{d\tau}{(1-\tau)^\gamma \tau^{\delta+\alpha}} \right) \end{aligned}$$

since  $\gamma + \delta = 1$  by changing the variable  $s = t\tau$ . Since  $\delta \in (0, 1)$ ,

$$I(\varepsilon) = \int_{1-\varepsilon}^1 \frac{d\tau}{(1-\tau)^\gamma \tau^{\delta+\alpha}}$$

converges for  $\varepsilon \in (0, 1)$ . Moreover  $I(\varepsilon)$  is continuous on  $[0, 1)$  and  $I(0) = 0$ . Since  $\varepsilon \mapsto I(\varepsilon)$  is increasing, there is a unique  $\varepsilon = \varepsilon(\sigma) < 1$  such that

$$I(\varepsilon) = \min \left( \frac{1}{2\sigma}, I(1) \right).$$

(The value  $I(1)$  may be  $+\infty$ .) For such an  $\varepsilon(\sigma)$

$$\varphi(t) \leq \sigma b_{\varepsilon(\sigma)} + \frac{1}{2} \varphi(t), \quad t \in (\eta/(1 - \varepsilon(\sigma)), T).$$

Sending  $\eta \rightarrow 0$  to get

$$t^\alpha \psi(t) \leq 2\sigma b_{\varepsilon(\sigma)}, \quad t \in (0, T).$$

This estimate yields (6.2) with  $c = 2b_{\varepsilon(\sigma)}$ .  $\square$

Although there are various type of the Gronwall type estimate, truncated integral in (6.1) is not usual. The estimate (6.2) also gives the decay estimate by taking  $T = \infty$ . The Gronwall type inequality yielding decay is known for example in [12, Lemma2,1]. It says that

$$0 \leq \psi(t) \leq (1+t)^{-\gamma} + a \int_0^t \frac{\psi(s)}{(t-s)^\gamma (1+s)^\delta} ds, \quad t \geq 0$$

implies  $\psi(t) \leq C(1+t)^{-\gamma}$  for  $a > 0$ ,  $\gamma \in [0, 1)$ ,  $\gamma + \delta > 1$  with  $C = C(a, \gamma, \delta)$ . However, clearly, it is different from ours, since decay rate  $\gamma$  is restricted. The author is grateful to Professor Tohru Ozawa for pointing [12, Lemma 2.1].

## 7 Indication of the proof of smoothing rate estimates

Instead of giving the full proof of Theorem 5.1 we only indicate its strategy by showing a typical situation:  $|\beta| = 1$ ,  $b = 0$ ,  $2 \leq p \leq \infty$ . In this situation  $\|\nabla \omega\|_p(t)$  is expected to be nonintegrable near  $t = 0$  so usual argument fails to work. By (2.1) it is standard to

observe that  $\|\partial_t^b \partial_x^\beta \omega\|_p(t)$  is bounded in any interval  $(\delta, T)$  if  $T > \delta > 0$ . The proof for  $p = \infty$  is actually written in [11] but the proof for other  $p \geq 1$  is the essentially the same. So we may assume that  $\|\nabla \omega\|_p(t)$  is continuous on  $(0, \infty)$  as a function of time  $t$ .

We first note that

$$\|u\|_\infty(t) \leq C_1 \|\omega\|_1 t^{-1/2} \quad (7.1)$$

by applying the Gagliardo-Nirenberg inequality

$$\|u\|_\infty \leq C_2 \|u\|_r^{1-2/r} \|\nabla u\|_r^{2/r}, \quad 2 < r < \infty,$$

the Calderón-Zygmund inequality

$$\|\nabla u\|_r \leq C_3 \|\omega\|_r, \quad 1 < r < \infty$$

the Hardy-Littlewood-Sobolev inequality

$$\|u\|_r \leq C_4 \|\omega\|_q, \quad 1/r = 1/q - 1/2, \quad 1 < q < 2$$

and (2.1) with  $C_q$  independent of  $u, \omega, \omega_0$  and  $t$ . Our  $\omega$  solves the integral equation

$$\omega(t) = e^{t\Delta} \omega_0 - \int_0^t e^{(t-s)\Delta} (u, \nabla) \omega \, ds.$$

We differentiate in space and estimate its  $L^p$ -norm to get

$$\|\nabla \omega\|_p(t) \leq \|\nabla e^{t\Delta} \omega_0\|_p + \int_0^t \|\nabla e^{(t-s)\Delta} (u, \nabla) \omega\|_p \, ds := I(t) + J(t). \quad (7.2)$$

By  $L^p - L^1$  estimate of derivate for the heat equation we observe

$$I(t) \leq C_5 t^{-\alpha} \|\omega_0\|_1, \quad \alpha = \frac{1}{2} + 1 - \frac{1}{p}. \quad (7.3)$$

To estimate  $J(t)$  we divide the interval of integration into two parts  $(0, t(1-\varepsilon))$ ,  $(t(1-\varepsilon), t)$ ,  $\varepsilon \in (0, 1)$ :

$$J(t) = \int_0^{t(1-\varepsilon)} \dots + \int_{t(1-\varepsilon)}^t \dots := J_1(t) + J_2(t). \quad (7.4)$$

To estimate  $J_1$  we use the property  $(u, \nabla) \omega = \operatorname{div} (u, \omega)$  to get

$$J_1(t) = \int_0^{t(1-\varepsilon)} \|\operatorname{div} \nabla e^{(t-s)\Delta} u \omega\|_p \, ds \leq \int_0^{t(1-\varepsilon)} \frac{C_6}{(t-s)^{1+1-1/p}} \|u \omega\|_1 \, ds$$

by  $L^p - L^1$  estimate  $\|\partial_x^\beta e^{t\Delta} f\|_p \leq C_6 t^{-|\beta|-1+1/p} \|f\|_1$ . We now apply (7.1) and (2.1) to get

$$J_1(t) \leq C_7 \int_0^{t(1-\varepsilon)} \frac{ds}{s^{1/2}(t-s)^{2-1/p}} \|\omega_0\|_1^2 = A_\varepsilon t^{-\alpha} \|\omega_0\|_1^2 \quad (7.5)$$

with  $A_\varepsilon$  depending only on  $\varepsilon$  and  $p$ . Contrary to  $J_1$  the singularity of  $\|\nabla\omega\|_p(s)$  near  $s = 0$  is excluded in  $J_2$  from the interval of integration. We thus directly estimate  $J_2$  to get

$$\begin{aligned} J_2(t) &\leq \int_{t(1-\varepsilon)}^t \frac{C_8}{(t-s)^{1/2}} \|u\|_\infty \|\nabla\omega\|_p ds \\ &\leq C_9 \int_{t(1-\varepsilon)}^t \frac{1}{(t-s)^{1/2} s^{1/2}} \|\nabla\omega\|_p(s) ds \|\omega_0\|_1 \end{aligned} \quad (7.6)$$

by (2.1). Combining (7.2)-(7.6) yields

$$\|\nabla\omega\|_p(t) \leq C_{10} \|\omega_0\|_1 \left( B_\varepsilon t^{-\alpha} \|\omega_0\|_1 + \int_t^{t(1-\varepsilon)} \frac{1}{(t-s)^{1/2} s^{1/2}} \|\nabla\omega\|_p(s) ds \right). \quad (7.7)$$

For all  $t > 0$ ,  $\varepsilon \in (0, 1)$  with  $B_\varepsilon$  depending only on  $p$  and  $\varepsilon$ . (It is easy to see that  $A_\varepsilon$  is nonincreasing in  $\varepsilon$  so is  $B_\varepsilon$ ). We now apply Lemma 6.1 to (7.7) to get (5.1) for  $|\beta| = 1$ ,  $b = 0$ ,  $2 \leq p \leq \infty$ .

In a similar way to derive (7.2) we estimate  $\|\nabla u\|_\infty(t)$ . Using this and estimate for  $\|\nabla\omega\|_p$  we just obtained, we are able to estimate  $\|\partial_x^\beta \omega\|_p(t)$  with  $|\beta| = 1$  like the estimate of  $\|\nabla\omega\|_p$ . The remaining estimates for  $b = 0$  can be proved inductively. The estimates with  $b > 0$  easily follows from (5.1), (5.2) with  $b = 0$  and the vorticity equation (1.3). The reader is referred to [8] for more detailed argument.

## 8 Smoothing rate estimate for the Navier-Stokes flow

In [13] it has been shown that for  $u_0 \in L^n(\mathbf{R}^n)$  (with  $\operatorname{div} u_0 = 0$ ), there is a unique global smooth solution of the Navier-Stokes equation (1.1), (1.2) with some  $\pi$  for initial velocity  $u_0$  provided that  $\|u_0\|_n$  is small. The solution  $u$  is continuous in  $[0, \infty)$  with values in  $L^n(\mathbf{R}^n)$ . Without size restriction of  $\|u_0\|_n$  we only have a local solution. (When  $n = 2$ , there exists always global smooth solution for  $L^2(\mathbf{R}^2)$  data as shown in [22].) We note our argument applies to get a smoothing rate estimate for velocity.

**Theorem 8.1.** *For  $u_0 \in L^n(\mathbf{R}^n)$  (with  $\operatorname{div} u_0 = 0$ ) let  $u \in C([0, T], L^n(\mathbf{R}^n))$  be a smooth solution of the Navier-Stokes equations (1.1), (1.2) with  $u|_{t=0} = u_0$ . Assume that  $\sup_{0 \leq t < T} \|u\|_n(t) := M < \infty$ . Then there is a constant  $C = C(M)$  such that*

$$\|\partial_x^\beta u\|_p(t) \leq C(M) \|u_0\|_n t^{-\frac{n}{2} \left( \frac{1}{n} - \frac{1}{p} \right) - \frac{|\beta|}{2}}, t \in (0, T) \quad (8.1)$$

for all  $n \leq p \leq \infty$ , multi-index  $\beta$ .

For the proof we use the integral equation

$$u(t) = e^{t\Delta} u_0 - \int_0^t e^{(t-s)\Delta} \mathbf{P}(u, \nabla) u ds$$

for  $u$ , where  $\mathbf{P}$  is the orthogonal projection to the divergence-free vector space and its explicit form is

$$(\mathbf{P})_{ij} = \delta_{ij} + R_i R_j \quad (1 \leq i, j \leq n)$$

with the Riesz operator  $R_i = \partial_{x_i}(-\Delta)^{-1/2}$ . By the way the idea converting to be original system to this integral equations for time-dependent functions with values in spaces of functions of spatial variables goes back to [17]. Tosio Kato derived various important results by using this integral equations [13]. We estimate  $u$  in a similar way to estimate  $\nabla\omega$  to get the desired result. Here note that  $\mathbf{P}$  is bounded in  $L^p$  to  $L^p$  for  $1 < p < \infty$  by the Calderón-Zygmund inequality. The detail as well as various extension of this estimate will be discussed elsewhere.

**Remark 8.2.** If the solution  $u$  is a global-in-time solution with  $\lim_{t \rightarrow \infty} \|u\|_n(t) = 0$ , our estimate (8.1) implies the decay estimate. In [23] for  $n = 2$  it has been proved that if  $u_0 \in H^m \cap L^2$ , then

$$\begin{aligned} \|\partial_x^\beta u\|_2(t) &\leq C(t+1)^{-(|\beta|+1)/2}, \quad t \geq 1, \\ \|\partial_x^\beta u\|_\infty(t) &\leq C(t+1)^{-(|\beta|+1/2)}, \quad t \geq 1. \end{aligned}$$

Our estimates are not comparable to theirs but it is likely that our method would give such estimates.

## References

- [1] M. Ben-Artzi, Global solutions of two-dimensional Navier-Stokes and Euler equations, Arch. Rational Mech. Anal., **128**(1994), 329-358.
- [2] H. Brezis, Remarks on the preceding paper by M. Ben-Artzi: "Global solutions of two-dimensional Navier-Stokes and Euler equations", Arch. Rational Mech. Anal. **128**(1994), 329-358 (see also MR 96h:35148; <http://www.ams.org/mathscinet>).
- [3] A. Carpio, Asymptotic behavior for the vorticity equations in dimension two and three, Commun. in Partial Differential Equations, **19**(1994), 827-872.
- [4] G.-H. Cottet, Équations de Navier-Stokes dans le plan avec tourbillon initial mesure, C. R. Acad. Sci. Ser. I, **303**(1986), 105-108.
- [5] H. Fujita and T. Kato, On the Navier-Stokes initial value problem I. Arch. Rational Mech. Anal., **16**(1964), 269-315.

- [6] Y. Giga, Analyticity of the semigroup generated by the Stokes operator in  $L_r$  spaces, *Math. Z.*, **178**(1981), 297-329.
- [7] Y. Giga, Domains of fractional powers of the Stokes operator in  $L_r$  spaces, *Arch. Rational Mech. Anal.*, **89**(1985), 251-265.
- [8] Y. Giga and M.-H. Giga, Nonlinear Partial Differential Equations- Asymptotic behaviour of solutions and self-similar solutions, Kyoritsu (1999) (in Japanese), English version (in preparation).
- [9] Y. Giga and T. Kambe, Large time behavior of the vorticity of two dimensional viscous flow and its application to vortex formation, *Comm. Math. Phys.*, **117**(1988), 549-568.
- [10] Y. Giga and T. Miyakawa, Solutions in  $L_r$  of the Navier-Stokes initial value problem, *Arch. Rational Mech. Anal.*, **89**(1985), 267-281.
- [11] Y. Giga, T. Miyakawa and H. Osada, Two-dimensional Navier-Stokes flow with measures as initial vorticity, *Arch. Rational Mech. Anal.*, **104**(1988), 223-250.
- [12] J. Ginibre and G. Velo, Conformal invariance and time decay for non linear wave equations II, *Ann. Inst. Henri Poincaré*, **47**(1987), 263-276.
- [13] T. Kato, Strong  $L^p$ -solutions of the Navier-Stokes equation in  $\mathbf{R}^m$  with applications to weak solutions, *Math. Z.*, **187**(1984), 471-480.
- [14] T. Kato, Strong solutions of the Navier-Stokes equation in Morrey spaces, *Bol. Soc. Bras. Mat.* **22**(1992), 127-155.
- [15] T. Kato, The Navier-Stokes equation for an incompressible fluid in  $\mathbf{R}^2$  with a measure as the initial vorticity, *Diff. Integral Equations*, **7**(1994), 949-966.
- [16] T. Kato, Well-posedness, *Sūgaku* **48**(1996), 298-300, (in Japanese).
- [17] T. Kato and H. Fujita, On the nonstationary Navier-Stokes system, *Rend. Sem. Mat. Univ. Padova*, **32**(1962), 243-260.
- [18] T. Kato and G. Ponce, Well-posedness of the Euler and Navier-Stokes equations in the Lebesgue spaces  $L^p_s(\mathbf{R}^2)$ , *Revista Mat. Iberoamericana* **2**(1986), 73-88.
- [19] T. Kato and G. Ponce, The Navier-Stokes equations with weak initial data, *Internat. Math. Res. Notices* (1994), no. **10**, 435ff., approx. 10pp. (electronic).

- [20] C. E. Kenig, G. Ponce and L. Vega, Well-posedness of the initial value problem for the Korteweg-des Vries equations, *J. Amer. Math. Soc.* **4**(1991), 323-347.
- [21] C. E. Kenig, G. Ponce and L. Vega, Well-posedness and scattering results for the generalized Korteweg-de Vries equation via contraction principle, *Comm. Pure Appl. Math.*, **46**(1993), 527-620.
- [22] J. Leray, Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l' Hydrodynamique, *J. Math. Pures Appl.*, **12**(1933), 1-82.
- [23] M. Schonbek, Large time behaviour of solutions to the Navier-Stokes equations in  $H^m$  spaces, *Commun. in Partial Differential Equations*, **20**(1995), 103-117.