



Title	Properties of a Rubin's orthogonal function which is a linear combination of two inner functions
Author(s)	Nakazi, T.; Watanabe, T.
Citation	Hokkaido University Preprint Series in Mathematics, 550, 1-9
Issue Date	2002-04
DOI	10.14943/83695
Doc URL	http://hdl.handle.net/2115/69299
Type	bulletin (article)
File Information	pre550.pdf



[Instructions for use](#)

Properties of A Rudin's Orthogonal Function Which
Is A Linear Combination of Two Inner Functions

Takahiko Nakazi and Toru Watanabe

Series #550. April 2002

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #524 K. Yamaguchi and T. Yatsui, Geometry of higher order differential equations of finite type associated with symmetric spaces, 43 pages. 2001.
- #525 T. Suwa, Residues of Chern classes, 20 pages. 2001.
- #526 V. Anh and A. Inoue, Dynamic models of asset prices with long memory, 21 pages. 2001.
- #527 T. Izawa and T. Suwa, Multiplicity of functions on singular varieties, 21 pages. 2001.
- #528 T. Nakazi and T. Yamamoto, Two dimensional commutative Banach algebras and von Neumann inequality, 18 pages. 2001.
- #529 Y. Giga, N. Ishimura and Y. Kohsaka, Spiral solutions for a weakly anisotropic curvature flow equation, 16 pages. 2001.
- #530 Y. Giga and P. Rybka, Quasi-static evolution of 3-D crystals grown from supersaturated vapor, 16 pages. 2001.
- #531 Y. Tonegawa, Remarks on convergence of the Allen-Cahn equation, 18 pages. 2001.
- #532 T. Suwa, Characteristic classes of singular varieties, 26 pages. 2001.
- #533 J. Escher, Y. Giga and K. Ito, On a limiting motion and self-intersections for the intermediate surface diffusion flow, 20 pages. 2001.
- #534 Y.-H. R. Tsai, Y. Giga and S. Osher, A level set approach for computing discontinuous solutions of a class of Hamilton-Jacobi equations, 30 pages. 2001.
- #535 A. Yamagami, On Gouvêa's conjecture in the unobstructed case, 19 pages. 2001.
- #536 A. Inoue, What does the partial autocorrelation function look like for large lags, 27 pages. 2001.
- #537 T. Nakazi and T. Yamamoto, Norm of a linear combination of two operators of a Hilbert space, 16 pages. 2001.
- #538 Y. Giga, On the two-dimensional nonstationary vorticity equations, 12 pages. 2001.
- #539 M. Jinzenji, Gauss-Manin system and the virtual structure constants, 25 pages. 2001.
- #540 H. Ishii and T. Mikami, Motion of a graph by R -curvature, 28 pages. 2001.
- #541 M. Jinzenji and T. Sasaki, $\mathcal{N} = 4$ supersymmetric Yang-Mills theory on orbifold- T^4/\mathbb{Z}_2 : higher rank case, 17 pages. 2001.
- #542 T. Nakazi, The Nevanlinna counting functions for Rudin's orthogonal functions, 7 pages. 2001.
- #543 K. Sugano, On H-separable extensions of QF-3 rings, 7 pages. 2001.
- #544 A. Arai, Non-relativistic limit of a Dirac-Maxwell operator in relativistic quantum electrodynamics, 27 pages. 2001.
- #545 O. Sawada, On time-local solvability of the Navier-Stokes equations in Besov spaces, 30 pages. 2001.
- #546 C. M. Elliott, Y. Giga, and S. Goto, Dynamic boundary conditions for Hamilton-Jacobi equations, 27 pages. 2001.
- #547 Y. Nakano, Minimizing coherent risk measures of shortfall in discrete-time models with cone constraints, 22 pages. 2002.
- #548 K. Tachizawa, A generalization of the Lieb-Thirring inequalities in low dimensions, 13 pages. 2002.
- #549 T. Nakazi, Absolute values and real parts for functions in the Smirnov class, 8 pages. 2002.

Properties of A Rudin's Orthogonal Function Which Is A Linear Combination of Two Inner Functions

by

Takahiko Nakazi*

and

Toru Watanabe

*This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education

2000 Mathematics Subject Classification : Primary 30 D 55, 46 J 15

Key words and phrases : Hardy space, inner function, orthogonal, polynomial

Abstract. ϕ is called a Rudin's (orthogonal) function if ϕ is a function in H^∞ and the different nonnegative powers of ϕ are orthogonal in H^2 . When ϕ is a multiple of an inner function and $\phi(0) = 0$, ϕ is a Rudin's function. Sundberg and Bishop showed that a Rudin's function is not necessarily a multiple of an inner function. We study a Rudin's function which is a linear combination of two inner functions or a polynomial of an inner function.

§1. Introduction.

For $1 \leq p \leq \infty$, H^p denotes the usual Hardy space on the unit circle T and σ is a normalized Lebesgue measure on T . ϕ is called a Rudin's (orthogonal) function if ϕ is a function in H^∞ and the different nonnegative powers of ϕ are orthogonal in H^2 . Two inner functions f and g are called (statistically) independent if

$$\int_T f^\ell \bar{g}^s d\sigma = 0$$

for all nonnegative integers $\ell, s, |\ell| + |s| > 0$. W.Rudin posed two problems (see [3]) : R1. Is a Rudin's (orthogonal) function necessarily a multiple of an inner function ? R2. Do there exist two (statistically) independent inner functions ? R1 has been negatively solved by Sundberg [6] and Bishop [1]. It is shown in [3] that if R1 is valid, then so is R2. We don't know whether the converse is true or not. Under some conditions, R1 has been positively solved. That is, when ϕ is a univalent function [2] and ϕ is in the disc algebra with boundary function in Lip α for some $\alpha > 1/2$ [3].

In §2, we give a few properties two (statistically) independent inner functions. In §3, we study a Rudin's (independent) function which is a linear combination of two inner functions. In §4, we show that a Rudin's (orthogonal) function is a multiple of an inner function when it is a polynomial of an inner function.

§2. Statistically independent inner functions

In this section, we give few necessary conditions for two independent inner functions.

Proposition 1. *If f and g are independent inner functions, then neither f nor g is a finite Blaschke product.*

Proof. Let $H^2[g]$ be the closure of the set of all polynomials of g in H^2 , then $f^j H^2[g]$ is orthogonal to $f^\ell H^2[g]$ if $j \neq \ell$ because f and g are independent inner functions. Set

$$M = \sum_{j=0}^{\infty} \oplus f^j H^2[g],$$

then M is a closed invariant subspace of H^2 under the multiplication by f . It is known (cf. [5, p11]) that the dimension of $M \ominus fM$ is less than or equal to that of $H^2 \ominus fH^2$. This implies that f can not be a finite Blaschke product.

Proposition 2. *If f and g are independent inner functions, then the set of all polynomials of f and g is not dense in H^2 .*

Proof. Suppose the set of all polynomials of f and g is dense in H^2 . Then

$$H^2 = \sum_{j=0}^{\infty} \oplus f^j H^2[g] = H^2[g] \oplus f H^2$$

because f and g are independent. Hence for any $n \geq 1$, g^n is orthogonal to $f H^2$ and so $f \bar{g}^n \in H_0^2 = L^2 \ominus \bar{H}^2$. Therefore $f = 0$ because $f \in \bigcap_{n=0}^{\infty} g^n H^2 = \{0\}$. This contradiction shows the proposition.

Proposition 3. *If f and g are independent inner functions, then there exists a positive integer n such that $z^n f$ and $z^n g$ are not independent.*

Proof. Suppose $z^n f$ and $z^n g$ are independent for any positive integer n . For any integer $\ell \geq 1$, f^ℓ is orthogonal to $z^n g^{\ell+1}$ for all $n \geq 0$. Put $\phi = \bar{f}g$, then $\phi^\ell g$ belongs to H^2 for all integer $\ell \geq 0$. By [3, p177], ϕ belongs to H^2 . Similarly we can show that $\bar{\phi}$ belongs to H^2 . Therefore ϕ is constant and so $g = \alpha f$ for some constant α with absolute value 1. This contradicts that f and g are independent.

§3. A linear combination of two inner functions

Let f and g be inner functions with $f(0) = g(0) = 0$ and let a and b complex numbers. Put

$$\phi(a, b) = af + bg.$$

If f and g are independent, then for an arbitrary pair (a, b) $\phi(a, b)$ is a Rudin's function. Theorem 4 shows that the converse is not true formally.

Lemma 1. *For some pair (a, b) , if $\phi(a, b)$ is a Rudin's function, then for any positive integer $\ell \geq 2$*

$$b^\ell \bar{a} \int_T \bar{f} g^\ell d\sigma + a^\ell \bar{b} \int_T f^\ell \bar{g} d\sigma = 0.$$

Proof. Since $(\phi^\ell, \phi) = 0$ for $\ell \geq 2$,

$$\sum_{j=0}^{\ell} \binom{\ell}{j} a^{\ell-j} \bar{a} b^j \int_T f^{\ell-j-1} g^j d\sigma + \sum_{j=0}^{\ell} \binom{\ell}{j} a^{\ell-j} b^j \bar{b} \int_T f^{\ell-j} g^{j-1} d\sigma = 0$$

because $\phi^\ell = \sum_{j=0}^{\ell} \binom{\ell}{j} a^{\ell-j} b^j f^{\ell-j} g^j$. If $1 \leq j \leq \ell - 1$, then $f^{\ell-j-1} g^j \in H^\infty$ and $f^{\ell-j} g^{j-1} \in H^\infty$. This implies that $\int_T f^{\ell-j-1} g^j d\sigma = \int_T f^{\ell-j} g^{j-1} d\sigma = 0$ because $f(0) = g(0) = 0$. Therefore

$$\bar{a} b^\ell \int_T \bar{f} g^\ell d\sigma + a^\ell \bar{b} \int_T f^\ell \bar{g} d\sigma = 0.$$

Corollary 1. For some pair (a, b) , if $\phi(a, b) = az + bg$ is a Rudin's function where g is an inner function with $g(0) = 0$, then $a = 0$, $b = 0$ or $g = cz$ for some complex number c .

Proof. By Lemma 1,

$$a^{\ell}\bar{b} \int_T z^{\ell} \bar{g} d\sigma = 0 \quad (\ell \geq 2)$$

and so if $a^{\ell}\bar{b} \neq 0$, then $g = cz$ for some complex number c .

Theorem 4. For arbitrary pair (a, b) , $\phi(a, b)$ is a Rudin's function if and only if

$$\int_T \bar{f}^s g^t d\sigma = \int_T f^t \bar{g}^s d\sigma = 0$$

for all non-negative integers t, s , $t \geq s + 1$.

Proof. When $\ell > k$,

$$\begin{aligned} (\phi^{\ell}, \phi^k) &= \left(\sum_{j=0}^{\ell} \binom{\ell}{j} a^{\ell-j} b^j f^{\ell-j} g^j, \sum_{i=0}^k \binom{k}{i} a^{k-i} b^i f^{k-i} g^i \right) \\ &= \sum_{j=0}^{\ell} \sum_{i=0}^k \binom{\ell}{j} \binom{k}{i} a^{\ell-j} \bar{a}^{k-i} b^j \bar{b}^i \int_T f^{\ell-k+i-j} g^{j-i} d\sigma \\ &= \sum_{-k \leq j-i < 0} \sum F(\ell, k, i, j) + \sum_{\ell-k < j-i \leq \ell} \sum F(\ell, k, i, j) \end{aligned}$$

where $F(\ell, k, i, j) = \binom{\ell}{j} \binom{k}{i} a^{\ell-j} \bar{a}^{k-i} b^j \bar{b}^i \int_T f^{\ell-k+i-j} g^{j-i} d\sigma$. Because if $j - i = 0$ or $0 \leq j - i \leq \ell - k$, then $f^{\ell-k+i-j} g^{j-i} \in H^{\infty}$ and so $F(\ell, k, i, j) = 0$. In the last line, note the following. When $-k \leq j - i < 0$, $t = \ell - k + i - j \underset{\neq}{>} s$ if $s = -(j - i)$. When $\ell - k \underset{\neq}{<} j - i \leq \ell$, $t = j - i \underset{\neq}{>} s$ if $s = -(\ell - k + i - j)$. Hence

$$\int_T f^{\ell-k+i-j} g^{j-i} d\sigma = \int_T f^t \bar{g}^s d\sigma \quad \text{or} \quad \int_T \bar{f}^s g^t d\sigma$$

where $t \geq s + 1$.

The 'if' part of the theorem is clear by the fact noted above. We will prove the 'only if' part by induction about s . If $s = 1$, by Lemma 1 it holds because (a, b) is arbitrary. Suppose it holds for $2 \leq s \leq n - 1$, that is, for $t \geq s + 1$

$$\int_T \bar{f}^s g^t d\sigma = \int_T f^t \bar{g}^s d\sigma = 0.$$

Since ϕ is a Rudin's function, $(\phi^{\ell}, \phi^n) = 0$ if $\ell > n$. By the fact noted above,

$$(\phi^{\ell}, \phi^n) = \sum_{-n \leq j-i < 0} \sum F(\ell, n, i, j) + \sum_{\ell-n < j-i \leq \ell} \sum F(\ell, n, i, j).$$

By the hypothesis on induction,

$$\begin{aligned} (\phi^\ell, \phi^n) &= \sum_{j-i=-n} \sum F(\ell, n, i, j) + \sum_{j-i=\ell} \sum F(\ell, n, i, j) \\ &= F(\ell, n, n, 0) + F(\ell, n, 0, \ell). \end{aligned}$$

In fact, when $-(n-1) \leq j-i < 0$, $t = \ell - n + i - j > s = -(j-i)$ and so $F(\ell, n, i, j) = 0$.

When $\ell - n < j-i \leq -1$, $t = j-i > s = -(\ell - n + i - j) \leq n-1$ and so $F(\ell, n, i, j) = 0$.

Thus

$$\binom{\ell}{0} \binom{n}{n} a^\ell \bar{a}^0 b^0 \bar{b}^n \int_T f^\ell \bar{g}^n d\sigma + \binom{\ell}{\ell} \binom{n}{0} a^0 \bar{a}^n b^\ell \bar{b}^0 \int_T \bar{f}^n g^\ell d\sigma = 0.$$

Since (a, b) is arbitrary,

$$\int_T f^\ell \bar{g}^n d\sigma = \int_T \bar{f}^n g^\ell d\sigma = 0.$$

Question. If $\phi(a, b) = af + bg$ is a Rudin's function for arbitrary pair (a, b) , then are f and g independent inner function ?

Theorem 5. Let q and Q be inner functions with $q(0) = 0$ and $Q(0) \neq 0$. If $f = q^s$ and $g = q^m Q$ where $m \geq s+1$, $s \geq 1$ and $\phi(a, b) = af + bg$ is a Rudin's function, then $a = 0$ or $b = 0$.

Proof. We will prove the following claim :

$$a^\ell \bar{b}^{\ell-k} \int_T f^\ell \bar{g}^{\ell-k} d\sigma = 0 \quad (\ell \geq k).$$

Put $k = m - s$ and $\ell = m$, then

$$\int_T f^\ell \bar{g}^{\ell-k} d\sigma = \int_T q^{s\ell - m(\ell-k)} \bar{Q}^{\ell-k} d\sigma = \int_T \bar{Q}^s d\sigma \neq 0$$

because $s\ell - m(\ell-k) = 0$. This implies $a^\ell \bar{b}^{\ell-k} = 0$.

We will show the claim by induction about ℓ . When $\ell = k$, it is clear because $f(0) = 0$. Suppose it holds for $k \leq \ell \leq n-1$. Since ϕ is a Rudin's function.

$$(\phi^n, \phi^{n-k}) = \sum_{j=0}^n \sum_{i=0}^{n-k} \binom{n}{j} \binom{n-k}{i} a^{n-j} \bar{a}^{n-k-i} b^j \bar{b}^i \int_T f^{n-(n-k)+i-j} \bar{g}^{i-j} d\sigma = 0.$$

When $i-j \leq 0$,

$$\int_T f^{n-(n-k)+i-j} \bar{g}^{i-j} d\sigma = 0$$

because

$$f^{n-(n-k)+i-j} \bar{g}^{i-j} = q^{s\{n-(n-k)+i-j\}} \bar{q}^{m(i-j)} Q^{j-i} = q^{sk+(m-s)(j-i)} Q^{j-i}.$$

When $0 \leq i - j \leq (n - 1) - k$, by the hypothesis on induction,

$$\int_T f^{n-(n-k)+i-j} \bar{g}^{i-j} d\sigma = 0.$$

Because if we put $i - j = s - k$ and $k \leq s \leq n - 1$, then $n - (n - k) + i - j = s$. Thus

$$\begin{aligned} 0 &= (\phi^n, \phi^{n-k}) \\ &= \sum_{j=0}^n \sum_{i=0}^{n-k} \binom{n}{j} \binom{n-k}{i} a^{n-j} \bar{a}^{n-k-i} b^j \bar{b}^i \int_T f^{n-(n-k)+i-j} \bar{g}^{i-j} d\sigma \\ &= \binom{n}{0} \binom{n-k}{n-k} a^n \bar{a}^0 b^0 \bar{b}^{n-k} \int_T f^n \bar{g}^{n-k} d\sigma. \end{aligned}$$

§4. Rudin's orthogonality function.

In this section, we study a Rudin's function when ϕ is a polynomial of an inner function. Theorem 5 determines a Rudin's function when $\phi = aq^s + bq^m$, $m \geq s + 1$ and q is an inner function with $q(0) = 0$. On the other hand, Theorem 6 solves affirmatively R1 when ϕ is a polynomial of an inner function. Proposition 7 gives another proof of Corollary 3 in [2] and another one of Proposition 3 in §2.

Theorem 6. *Let ϕ_0 be a Rudin's function and $\phi = \sum_{j=1}^n a_j \phi_0^j$ with $a_n \neq 0$. If ϕ is a Rudin's function, then $\phi = a_n \phi_0^n$.*

Proof. We may assume that $\phi = \sum_{j=1}^n a_j \phi_0^j$, $a_n = 1$ and $n > 1$. By induction, we will show that $a_\ell = 0$ for $1 \leq \ell \leq n - 1$. Suppose $\ell = 1$. Since

$$\phi^n = \sum_{i=0}^n \binom{n}{i} (a_1 \phi_0)^{n-i} \left(\sum_{k=2}^n a_k \phi_0^k \right)^i,$$

the smallest degree of ϕ^n is n because the smallest one of $\binom{n}{i} (a_1 \phi_0)^{n-i} \left(\sum_{k=2}^n a_k \phi_0^k \right)^i$ is $n + i$. On the other hand, the degree of ϕ is n . Hence if ϕ is a Rudin's function,

$$(\phi^n, \phi) = a_1^n = 0$$

because ϕ_0 is a Rudin's function.

Suppose $a_1 = a_2 = \cdots = a_\ell = 0$ for $\ell < n - 1$. Then since

$$\phi^n = \sum_{i=0}^n \binom{n}{i} (a_{\ell+1} \phi_0^{\ell+1})^{(n-i)} \left(\sum_{k=\ell+2}^n a_k \phi_0^k \right)^i,$$

the smallest degree of ϕ^n is $n(\ell+1)$ because the smallest one of $\binom{n}{i} (a_{\ell+1} \phi_0^{\ell+1})^{(n-i)} \left(\sum_{k=\ell+2}^n a_k \phi_0^k \right)^i$ is $(\ell+1)(n-i) + (\ell+2)i = i + n(\ell+1)$. On the other hand, since

$$\phi^{\ell+1} = \sum_{j=0}^{\ell+1} \binom{\ell+1}{j} \phi_0^{n(\ell+1-j)} \left(\sum_{k=\ell+1}^{n-1} a_k \phi_0^k \right)^j,$$

the largest degree of $\phi^{\ell+1}$ is $n(\ell+1)$ because the largest one of $\binom{\ell+1}{j} \phi_0^{n(\ell+1-j)} \left(\sum_{k=\ell+1}^{n-1} a_k \phi_0^k \right)^j$ is $(n-1)j + n(\ell+1-j) = n(\ell+1) - j$. Hence if ϕ is a Rudin's function,

$$(\phi^n, \phi^{\ell+1}) = (a_{\ell+1})^{n(\ell+1)} = 0$$

because ϕ_0 is a Rudin's function.

Proposition 7. *If $z^n \phi$ is a Rudin's function for all $n \geq 0$, then ϕ is a multiple of an inner function.*

Proof. Fix a positive integer k . For all $n \geq 0$, $z^{nk} \phi^k$ is orthogonal to $z^{n(k+1)} \phi^{k+1}$ because $z^n \phi$ is a Rudin's function. Thus ϕ^k is orthogonal to $\{z^n \phi^{k+1}\}_{n=0}^{\infty}$. Put $\phi = qh$ where q is inner and h is outer. Then by the Beurling's theorem [5, p11], ϕ^k is orthogonal to $q^{k+1} H^2$ and so $\bar{h}^{k+1} q \in H^2$ for all $k \geq 0$. By [4, p177], h is constant and so ϕ is inner.

References

1. C.Bishop, Orthogonal functions in H^∞ , in preprint
2. P.S.Bourdon, Rudin's orthogonality problem and the Nevanlinna counting function, Proc.Amer.Math.Soc. 125(1997), 1187-1192
3. J.A.Cima, B.Korenblum and M.Stessin, Composition isometries and Rudin's problems, in preprint
4. T.Gamelin, Uniform Algebras, Prentice-Hall, Englewood Cliffs, New Jersey, 1969
5. M.Rosenblum and J.Rovnyak, Hardy Classes and Operator Theory, Oxford Math. Monographs, 1985
6. C.Sundberg, Measures induced by analytic functions and a problem of Walter Rudin, in preprint

Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan

nakazi @ math.sci.hokudai.ac.jp