



Title	Backward shift invariant subspaces in the bidisc
Author(s)	Izuchi, K.; Nakazi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 572, 1-8
Issue Date	2002-11
DOI	10.14943/83717
Doc URL	http://hdl.handle.net/2115/69321
Type	bulletin (article)
File Information	pre572.pdf



[Instructions for use](#)

Backward Shift Invariant Subspaces in the Bidisc

Keiji Izuchi and Takahiko Nakazi

Series #572. November 2002

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #546 C. M. Elliott, Y. Giga, and S. Goto, Dynamic boundary conditions for Hamilton-Jacobi equations, 27 pages. 2001.
- #547 Y. Nakano, Minimizing coherent risk measures of shortfall in discrete-time models with cone constraints, 22 pages. 2002.
- #548 K. tachizawa, A generalization of the Lieb-Thirring inequalities in low dimensions, 13 pages. 2002.
- #549 T. Nakazi, Absolute values and real parts for functions in the Smirnov class, 8 pages. 2002.
- #550 T. Nakazi and T. Watanabe, Properties of a Rubin's orthogonal function which is a linear combination of two inner functions, 9 pages. 2002.
- #551 T. Ohtsuka, A level set method for spiral crystal growth, 24 pages. 2002.
- #552 M.-H. Giga and Y. Giga, Minimal vertical singular diffusion preventing overturning for the Burgers equation, 18 pages. 2002.
- #553 Y. Giga and P. Rybka, Berg's effect, 12 pages. 2002.
- #554 Y. Tonegawa, Domain dependent monotonicity formula for a singular perturbation problem, 14 pages. 2002.
- #555 S. Izumiya and N. Takeuchi, New special curves and developable surfaces, 10 pages. 2002.
- #556 S. Izumiya, K. Saji and N. Takeuchi, Singularities of line congruences, 17 pages. 2002.
- #557 S. Izumiya, D. Pei and M. C. Romero Fuster, The lightcone Gauss map of a spacelike surface in Minkowski 4-space, 21 pages. 2002.
- #558 S. Izumiya, D. Pei and M. C. Romero Fuster, Umbilicity of spacelike submanifolds of Minkowski space, 14 pages. 2002.
- #559 S. Izumiya, D. Pei and M. Takahashi, Curves and surfaces in Hyperbolic space, 16 pages. 2002.
- #560 J. Kato, On the uniqueness of nondecaying solutions for the Navier-Stokes equations, 19 pages. 2002.
- #561 M. Jinzenji and T. Sasaki, An Approach to $\mathcal{N} = 4$ ADE gauge Theory on $K3$, 29 pages. 2002.
- #562 T.Nakazi and T.Yamamoto, Norms of some singular integral operators on weighted L^2 spaces, 27 pages. 2002.
- #563 A.Harris and Y.Tonegawa, A $\bar{\partial}\partial$ -poincaré lemma for forms near an isolated complex singularity, 8 pages. 2002.
- #564 M.Takahashi, Bifurcations of ordinary differential equations of Clairaut type, 23 pages. 2002.
- #565 G.Ishikawa, Classifying singular Legendre curves by contactomorphisms, 17 pages. 2002.
- #566 G.Ishikawa, Perturbations of Caustics and fronts, 17 pages. 2002.
- #567 Y.Giga, On regularizing-decay rate estimates for solutions to the Navier-Stokes initial value problem, 12 pages. 2002.
- #568 T.Miyao, Strongly supercommuting self-adjoint operators, 34 pages. 2002.
- #569 Jun-Muk Hwang and K.Yamaguchi, Characterization of Hermitian symmetric spaces by fundamental forms, 10 pages. 2002.
- #570 H.Ishii and T.Mikami, Convexified Gauss curvature flow of bounded open sets in an anisotropic external field: a stochastic approximation and PDE, 37 pages. 2002.
- #571 Y.Nakano, Minimization of shortfall risk in a jump-diffusion model, 10 pages. 2002.

Backward Shift Invariant Subspaces in the Bidisc

by

Keiji Izuchi*

and

Takahiko Nakazi*

*This research was partially supported by Grant-in-Aid for Scientific Research,
Ministry of Education

2000 Mathematics Subject Classification : Primary 47 A 15, 46 J 15, Secondary
47 A 20.

Key words and phrases : bidisc, Hardy space, backward shift, invariant subspace,
double commuting

Abstract. Suppose that T_ϕ is a Toeplitz operator with a symbol ϕ on the Hardy space H^2 on the bidisc. Let N be a backward shift invariant subspace of H^2 , that is, N is an invariant subspace under T_z^* and T_w^* . Let P be the orthogonal projection from H^2 onto N . For ϕ in H^∞ , put $S_\phi = PT_\phi|N$. In this paper, we give a characterization of a backward shift invariant subspace which satisfies $S_z S_w^* = S_w^* S_z$.

§1. Introduction

Let T^2 be the torus that is the Cartesian product of two unit circles T in \mathbf{C} . Let $p = 1$ or $p = \infty$. The usual Lebesgue spaces, with respect to the Haar measure m on T^2 , are denoted by $L^p = L^p(T^2)$, and $H^p = H^p(T^2)$ is the space of all f in L^p whose Fourier coefficients

$$\hat{f}(j, \ell) = \int_{T^2} f(z, w) \bar{z}^j \bar{w}^\ell dm(z, w)$$

are 0 as soon as at least one component of (j, ℓ) is negative. Then H^p is called the Hardy space. As $T^2 = (z, T) \times (w, T)$, $H^p(z, T)$ and $H^p(w, T)$ denote the one variable Hardy spaces.

Let P_{H^2} be the orthogonal projection from L^2 onto H^2 . For ϕ in L^∞ , the Toeplitz operator T_ϕ is defined by

$$T_\phi f = P_{H^2}(\phi f) \quad (f \in H^2).$$

A closed subspace N of H^2 is said to be backward shift invariant if

$$T_z^* N \subset N \quad \text{and} \quad T_w^* N \subset N.$$

A closed subspace M of H^2 is said to be shift invariant if $T_z M \subset M$ and $T_w M \subset M$. The orthogonal complement of N is shift invariant. Let P_N and P_M be the orthogonal projections from H^2 onto N and M , respectively. For ϕ in H^∞ , put

$$S_\phi = P_N T_\phi P_N|N \quad \text{and} \quad V_\phi = P_M T_\phi P_M|M.$$

It is known in [2] that $V_z V_w^* = V_w^* V_z$ if and only if $M = qH^2$ for some inner function q in H^∞ . In this paper, we are interested in backward shift invariant subspaces N which satisfy $S_z S_w^* = S_w^* S_z$. Let $M = H^2 \ominus N$. We will write $P = P_N$ and $Q = I - P_N$, where I is the identity operator on H^2 . In this paper, we also study two operators

$$A = QT_z P \quad \text{and} \quad B = PT_w^* Q.$$

In §2, we show that $AB|M = V_w^* V_z - V_z V_w^*$ and $BA|N = S_z S_w^* - S_w^* S_z$. Then $AB = 0$ is equivalent to $V_z V_w^* = V_w^* V_z$, and $BA = 0$ is equivalent to $S_z S_w^* = S_w^* S_z$. Moreover we determine backward shift invariant subspaces satisfying $A = 0$ or $B = 0$. In §3, we give a characterization of backward shift invariant subspaces satisfying $BA = 0$, equivalently

$S_z S_w^* = S_w^* S_z$. And we give simple sufficient conditions to be $S_z S_w^* = S_w^* S_z$. In §4, we give a conjecture, that is, the sufficient condition is also necessary one.

Throughout this paper, for a subset H of H^2 , $[H]_2$ denotes the closed linear span of H and $[H]$ the linear span of H .

§2. Invariant subspace with $A = 0$ or $B = 0$

Let N be a backward shift invariant subspace and M be the orthogonal complement of N in H^2 . Put $P = P_N$ and $Q = I - P_N$, then Q is the orthogonal projection from H^2 onto M .

Lemma 2.1.

- (1) $AB = QT_w^* QT_z Q - QT_z QT_w^* Q$ and so $AB|M = V_w^* V_z - V_z V_w^*$.
- (2) $BA = PT_z PT_w^* P - PT_w^* PT_z P$ and so $BA|N = S_z S_w^* - S_w^* S_z$
- (3) $\ker A = \{f \in N : T_z f \in N\} \oplus M$.
- (4) $\ker B = \{f \in M : T_w^* f \in M\} \oplus N$.

Proof. (1) Since $T_z Q = QT_z Q$ and $T_z T_w^* = T_w^* T_z$,

$$\begin{aligned} AB &= QT_z PT_w^* Q \\ &= QT_z T_w^* Q - QT_z QT_w^* Q \\ &= QT_w^* QT_z Q - QT_z QT_w^* Q. \end{aligned}$$

(2) Since $T_w^* P = PT_w^* P$ and $T_w^* T_z = T_z T_w^*$,

$$\begin{aligned} BA &= PT_w^* QT_z P \\ &= PT_w^* T_z P - PT_w^* PT_z P \\ &= PT_z PT_w^* P - PT_w^* PT_z P. \end{aligned}$$

The properties (3) and (4) are clear.

Theorem 2.2.

- (1) $A = 0$ if and only if $N = H^2$ or $N = H^2 \ominus qH^2$ where q is a one variable inner function with $q = q(w)$.
- (2) $B = 0$ if and only if $M = [0]$ or $M = qH^2$ where q is a one variable inner function with $q = q(z)$.
- (3) $A = B = 0$ if and only if $N = [0]$ or $N = H^2$.

Proof. (2) follows from (1). We will show (1). We have $H^2 = N \oplus M$ and $T_z M \subset M$. Suppose $A = 0$. By Lemma 2.1 (3), $T_z N \subset N$. Put $N_0 = N \ominus T_z N$ and

$M_0 = M \ominus T_z M$. Then

$$H^2 = \sum_{n=0}^{\infty} \oplus (N_0 \oplus M_0) z^n = \sum_{n=0}^{\infty} \oplus H^2(w, T) z^n$$

because $zH^2 = zN \oplus zM$ and so $N_0 \oplus M_0 = H^2(w, T)$. By Lemma 2.1 (1), $V_w^* V_z = V_z V_w^*$ and so $V_z^* V_w = V_w V_z^*$ because $AB = 0$. Hence $V_w(\ker V_z^*) \subseteq \ker V_z^*$ and $\ker V_z^* = M_0$. Therefore by a theorem of Beurling [1], if $M_0 \neq [0]$, $M_0 = qH^2(w, T)$ and q is a one variable inner function with $q = q(w)$. Hence $M = qH^2$ and so $N = H^2 \ominus qH^2$. If $M_0 = [0]$, then $M = [0]$, and so $N = H^2$.

§3. Invariant subspace with $AB = 0$ or $BA = 0$

Suppose that N is a backward shift invariant subspace and $M = H^2 \ominus N$. By Lemma 2.1, $AB = 0$ if and only if $V_w^* V_z = V_z V_w^*$, and $BA = 0$ if and only if $S_z S_w^* = S_w^* S_z$. Hence we know (see [2],[3],[4]) that $AB = 0$ if and only if $M = qH^2$ for some inner function q . In this section, we study N when $BA = 0$, that is, $S_z S_w^* = S_w^* S_z$.

Lemma 3.1.

$$[\text{ran } A]_2 = \{M \ominus zM\} \ominus \{H^2(w, T) \cap M\}$$

and

$$\ker B = \{H^2(z, T) \cap M\} \oplus wM \oplus N.$$

Proof. Since $(T_w^* f, g) = (f, wg)$ if $f, g \in H^2$,

$$\{f \in M ; T_w^* f \in M\} = M \cap \{H^2 \ominus wN\} = \{H^2(z, T) \cap M\} \oplus wM,$$

because $H^2 \ominus wN = (H^2 \ominus wH^2) \oplus w(H^2 \ominus N)$ and $N = H^2 \ominus M$. Hence by Lemma 2.1 (4), $\ker B = \{f \in M ; T_w^* f \in M\} \oplus N = \{H^2(z, T) \cap M\} \oplus wM \oplus N$. By the same argument, $\ker A^* = \{H^2(w, T) \cap M\} \oplus zM \oplus N$ and so

$$\begin{aligned} [\text{ran } A]_2 &= H^2 \ominus \ker A^* \\ &= \{M \ominus zM\} \ominus \{H^2(w, T) \cap M\}. \end{aligned}$$

Lemma 3.2.

- (1) $A = 0$ if and only if $M = \{H^2(w, T) \cap M\} \oplus zM$.
- (2) $B = 0$ if and only if $M = \{H^2(z, T) \cap M\} \oplus wM$.
- (3) $BA = 0$ if and only if $\{H^2(z, T) \cap M\} \oplus wM \supseteq \{M \ominus zM\} \ominus \{H^2(w, T) \cap M\}$.

Proof. These follow from Lemma 3.1.

For a subset H of H^2 , let $H_k = \sum_{i+j=k} z^i w^j H$ for $k \geq 0$.

Theorem 3.3. *Let N be a backward shift invariant subspace of H^2 and M its orthogonal complement. Suppose $N \neq H^2$.*

(1) $S_z S_w^* = S_w^* S_z$ if and only if $M = H + M_1$ and if and only if $M = \sum_{j=0}^{k-1} H_j + M_k$ for any $k \geq 1$, where $H = H_0 = H^2(z, T) \cap M + H^2(w, T) \cap M$. If $S_z S_w^* = S_w^* S_z$, then $H \neq [0]$.

(2) When $M \cap H^2(z, T) = [0]$ or $M \cap H^2(w, T) = [0]$, $S_z S_w^* = S_w^* S_z$ if and only if $M = qH^2 + M_k$ for any $k \geq 1$ where q is a one variable inner function such that $M \cap H^2(z, T) = qH^2(z, T)$ or $M \cap H^2(w, T) = qH^2(w, T)$.

(3) When $M \cap H^2(z, T) \neq [0]$ and $M \cap H^2(w, T) \neq [0]$, $S_z S_w^* = S_w^* S_z$ if and only if $M = q_1 H^2 + q_2 H^2 + M_k$ for any $k \geq 1$ where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one variable inner functions such that $M \cap H^2(z, T) = q_1 H^2(z, T)$ and $M \cap H^2(w, T) = q_2 H^2(w, T)$.

Proof. (1) Since $S_z S_w^* = S_w^* S_z$ is equivalent to $BA = 0$, $S_z S_w^* = S_w^* S_z$ if and only if $M = H + M_1$ by Lemma 3.2 (3). It is easy to see that $M = H + M_1 = \sum_{j=0}^{k-1} H_j + M_k$ for any $k \geq 1$. If $H = [0]$, then $M = M_k$ and hence $M = [0]$. This contradicts $N \neq H^2$.

(2) We may assume that $M \cap H^2(z, T) = [0]$ and $M \cap H^2(w, T) \neq [0]$. By a theorem of Beurling [1], $M \cap H^2(w, T) = qH^2(w, T)$ for some one variable inner function $q = q(w)$. By (1), $S_z S_w^* = S_w^* S_z$ if and only if $M = qH^2(w, T) + M_1$ if and only if $M = q \sum_{j=0}^{k-1} \oplus H^2(w, T) z^j + M_k$ for any $k \geq 1$. This is equivalent to $M = qH^2 + M_k$ for any $k \geq 1$. For, $M_k \supseteq qz^k H^2$.

(3) By a theorem of Beurling, $M \cap H^2(z, T) = q_1 H^2(z, T)$ and $M \cap H^2(w, T) = q_2 H^2(w, T)$ where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one variable inner functions. By (1), $S_z S_w^* = S_w^* S_z$ if and only if $M = q_1 H^2(z, T) + q_2 H^2(w, T) + M_1$ if and only if

$$M = q_1 \sum_{j=0}^{k-1} \oplus H^2(z, T) w^j + q_2 \sum_{j=0}^{k-1} \oplus H^2(w, T) z^j + M_k$$

for any $k \geq 1$. This is equivalent to $M = q_1 H^2 + q_2 H^2 + M_k$ for any $k \geq 1$. For, $M_k \supseteq q_1 w^k H^2 + q_2 z^k H^2$.

Corollary 3.4.

(1) $AB = BA = 0$ if and only if $A = 0$ or $B = 0$.

(2) If $N = H^2 \ominus qH^2$ and q is an inner function and $S_z S_w^* = S_w^* S_z$, then q is a one variable.

Proof. (1) If $AB = BA = 0$, then by Lemma 2.1 (1) $V_w^* V_z = V_z V_w^*$ and so $M = qH^2$ for some inner function q (see [2], [4]). On the other hand, by Theorem 3.3 (1), $M \cap H^2(z, T) \neq [0]$ or $M \cap H^2(w, T) \neq [0]$ because $S_z S_w^* = S_w^* S_z$. Hence q is one variable. By Theorem 2.2, $A = 0$ or $B = 0$.

(2) is clear by (1).

Corollary 3.5. *Let N be a backward shift invariant subspace and $N \neq H^2$.*

(1) *If $S_z S_w^* = S_w^* S_z$, then $N \subseteq H^2 \ominus qH^2$ for some one variable inner function q .*

(2) *If $N = H^2 \ominus qH^2$ for some one variable inner function q , then $S_z S_w^* = S_w^* S_z$.*

Proof. (1) By Theorem 3.3, if $S_z S_w^* = S_w^* S_z$ then $M \supseteq qH^2$ for some one variable inner function q . Hence $N \subseteq H^2 \ominus qH^2$. (2) is clear by Theorem 3.3 (3).

Corollary 3.6. *Suppose that $A \neq 0$ and $B \neq 0$.*

(1) *If $S_z S_w^* = S_w^* S_z$, then $N \subseteq (H^2 \ominus q_1 H^2) \cap (H^2 \ominus q_2 H^2)$ where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one variable inner functions.*

(2) *If $N = (H^2 \ominus q_1 H^2) \cap (H^2 \ominus q_2 H^2)$ where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one variable inner functions, then $S_z S_w^* = S_w^* S_z$.*

Proof. By Theorem 2.2, we can prove (1) as in the proof of Corollary 3.5 (1). (2) Since $q_1 H^2 + q_2 H^2 = [q_1, q_2] + (q_1 H_1^2 + q_2 H_1^2)$, $M = [q_1, q_2] + (zM + wM) = q_1 H^2 + q_2 H^2 + M_k$ for any $k \geq 1$. It is easy to see that $M \cap H^2(z, T) = q_1 H^2(z, T)$ and $M \cap H^2(w, T) = q_2 H^2(w, T)$. Hence by Theorem 3.3 (3) $S_z S_w^* = S_w^* S_z$.

§4. Conjecture

By Corollary 3.5 (2) and Corollary 3.6, if $N = H^2$, $N = H^2 \ominus qH^2$ for some one variable inner function q or $N = (H^2 \ominus q_1 H^2) \cap (H^2 \ominus q_2 H^2)$ for some one variable inner functions $q_1 = q_1(z)$ and $q_2 = q_2(w)$, then $S_z S_w^* = S_w^* S_z$. Because of Theorem 3.3, we have the following conjecture. In this section, we study this conjecture.

Conjecture. *If $S_z S_w^* = S_w^* S_z$, then $N = H^2$, $N = H^2 \ominus qH^2$ for some one variable inner function q or $N = (H^2 \ominus q_1 H^2) \cap (H^2 \ominus q_2 H^2)$, where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one variable inner functions.*

Proposition 4.1. *If $M = q_1 H^2 + q_2 H^2 + M_k$ for any $k \geq 1$, where $M \cap H^2(z, T) = q_1 H^2(z, T)$ and $M \cap H^2(w, T) = q_2 H^2(w, T)$, then $M = q_1 (H^2 \ominus w^k H^2) + q_2 (H^2 \ominus w^k H^2) + w^k M$ for any $k \geq 1$. The converse is also true.*

Proof. Since $wM \supseteq q_2 w H^2(w, T)$, by Lemma 3.2 (3) and Theorem 3.3 (3),

$$q_1 H^2(z, T) \oplus wM \supseteq K_2 \oplus q_2 w H^2(w, T),$$

where $M \ominus zM = K_2 \oplus q_2 H^2(w, T)$. Thus

$$q_1 H^2(z, T) \oplus wM \supseteq \sum_{j=0}^{\infty} \oplus \{K_2 \oplus q_2 w H^2(w, T)\} z^j.$$

Since

$$M = \sum_{j=0}^{\infty} \oplus (M \ominus zM)z^j = \sum_{j=0}^{\infty} \oplus \{K_2 \oplus q_2H^2(w, T)\}z^j,$$

we have

$$q_1H^2(z, T) + q_2H^2(z, T) + wM \supseteq M.$$

Hence $M = q_1H^2(z, T) + q_2H^2(z, T) + wM$. This leads our assertion.

Corollary 4.2. *If $M = qH^2 + M_k$ for any $k \geq 1$ where $M \cap H^2(z, T) = qH^2(z, T)$ and $M \cap H^2(w, T) = [0]$, then $M = qH^2$.*

Proof. By Proposition 4.1 and its proof, $M = q(H^2 \ominus w^kH^2) + w^kM$ for any $k \geq 1$. This implies that $M = qH^2$, because $q(H^2 \ominus w^kH^2)$ is orthogonal to w^kM .

It is not difficult to prove that $q_1H^2 + q_2H^2$ is closed when $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one variable. Hence our conjecture is equivalent to the following one. If $S_zS_w^* = S_w^*S_z$, then $M = [0]$, $M = qH^2$ or $M = q_1H^2 + q_2H^2$. Even if N is of finite dimension, $S_zS_w^* \neq S_w^*S_z$ may happen. In fact, $N = \{1, z, w\}$ is such an example.

References

1. A. Beurling, On two problems concerning linear transformations in Hilbert space, Acta Math. 81(1949), 239-255.
2. V. Mandrekar, The validity of Beurling theorems in polydiscs, Proc. Amer. Math. Soc. 103(1988), 145-148.
3. T. Nakazi, Certain invariant subspaces of H^2 and L^2 on a bidisc. Can. J. Math. XL(1988), 1272-1280.
4. T. Nakazi, Invariant subspaces in the bidisc and commutators, J. Austral. Math. Soc. (Series A) 56(1994), 232-242.

K. Izuchi
Department of Mathematics
Faculty of Science
Niigata University
Niigata 950-2181, Japan

izuchi@scux.sc.niigata-u.ac.jp

T. Nakazi
Department of Mathematics
Faculty of Science
Hokkaido University
Sapporo 060-0810, Japan

nakazi@math.sci.hokudai.ac.jp