



Title	Exposed points and extremal problems in H_1 on a bidisc
Author(s)	Nakazi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 590, 1-11
Issue Date	2003-04
DOI	10.14943/83735
Doc URL	http://hdl.handle.net/2115/69339
Type	bulletin (article)
File Information	pre590.pdf



[Instructions for use](#)

EXPOSED POINTS AND EXTREMAL
PROBLEMS IN H^1 ON A BIDISC

Takahiko Nakazi

Series #590. April 2003

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #564 M. Takahashi, Bifurcations of ordinary differential equations of Clairaut type, 23 pages. 2002.
- #565 G. Ishikawa, Classifying singular Legendre curves by contactomorphisms, 17 pages. 2002.
- #566 G. Ishikawa, Perturbations of Caustics and fronts, 17 pages. 2002.
- #567 Y. Giga and O. Sawada, On regularizing–decay rate estimates for solutions to the Navier–Stokes initial value problem, 12 pages. 2002.
- #568 T. Miyao, Strongly supercommuting self-adjoint operators, 34 pages. 2002.
- #569 J.M. Hwang and K. Yamaguchi, Characterization of Hermitian symmetric spaces by fundamental forms, 10 pages. 2002.
- #570 H. Ishii and T. Mikami, Convexified Gauss curvature flow of bounded open sets in an anisotropic external field: a stochastic approximation and PDE, 37 pages. 2002.
- #571 Y. Nakano, Minimization of shortfall risk in a jump-diffusion model, 10 pages. 2002.
- #572 K. Izuchi and T. Nakazi, Backward shift invariant subspaces in the bidisc, 8 pages. 2002.
- #573 S. Izumiya, D. Pei and M. C. Romero-Fuster, The horospherical geometry of surfaces in Hyperbolic 4-space, 17 pages. 2002.
- #574 S. Izumiya and M. C. Romero-Fuster, The hyperbolic Gauss-Bonnet type theorem, 10 pages. 2002.
- #575 S. Izumiya and S. Janeczko, A symplectic framework for multiplane gravitational lensing, 19 pages. 2002.
- #576 S. Izumiya, M. Kossowski, D. Pei and M. C. Romero-Fuster, Singularities of C^∞ -lightlike hypersurfaces in Minkowski 4-space, 18 pages. 2002.
- #577 S. Izumiya, D. Pei and M. Takahashi, Evolutes of hypersurfaces in Hyperbolic space, 21 pages. 2002.
- #578 Y. Giga, S. Matsui and S. Sasayama, Blow up rate for semilinear heat equation with subcritical nonlinearity, 29 pages. 2002.
- #579 M. Tsujii, Physical measures for partially hyperbolic surface endomorphisms, 71 pages. 2003.
- #580 Y. Giga and K. Yamada, On viscous Burgers-like equations with linearly growing initial data, 19 pages. 2003.
- #581 T. Nakazi and T. Osawa, Spectra of Toeplitz Operators and Uniform Algebras, 9 pages. 2003.
- #582 Y. Daido, M. Ikehata and G. Nakamura, Reconstruction of Inclusions for the Inverse Boundary Value Problem with Mixed Type Boundary Condition, 18 pages. 2003.
- #583 Y. Daido and G. Nakamura, Reconstruction of Inclusions for the Inverse Boundary Value Problem with Mixed Type Boundary Condition and Source Term, 26 pages. 2003.
- #584 M.-H. Giga and Y. Giga, A PDE approach for motion of phase-boundaries by a singular interfacial energy, 19 pages. 2003.
- #585 A.A. Davydov, G. Ishikawa, S. Izumiya and W.-Z. Sun, Generic singularities of implicit systems of first order differential equations on the plane, 28 pages. 2003.
- #586 K. Yamauchi, On an underlying structure for the consistency of viscosity solutions, 12 pages. 2003.
- #587 T. Miyao, Momentum operators with a winding gauge potential, 15 pages. 2003.
- #588 Y. Giga and R. Kobayashi, On constrained equations with singular diffusivity, 35 pages. 2003.
- #589 O. Sawada, On analyticity rate estimates of the solutions to the Navier-Stokes equations in Bessel-potential spaces, 13 pages. 2003.

EXPOSED POINTS AND EXTREMAL PROBLEMS IN H^1 ON A BIDISC

BY TAKAHIKO NAKAZI*

Kurztitel : Extremal problems in H^1

2000 Mathematics Subject Classification : 32 A 35, 42 B 30

Key words and phrases : Hardy space, rational function, bidisc, exposed point, linear functional, extremal problem

nakazi@math.sci.hokudai.ac.jp

*This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education

Abstract. An essential bounded function ϕ gives a continuous linear functional on the Hardy space H^1 on the bitorus. In this paper, we consider extremal problems on H^1 when ϕ is a rational function, ϕ is a product of one variable functions or $\phi = |f|/f$ for some outer function f in H^1 such that $f(z, w)$ has a good property with respect to w for a.e. z .

1. Introduction

Let m be the normalized Lebesgue measure of the bitorus T^2 , the distinguished boundary of the open unit bidisc D^2 in the space \mathbf{C}^2 of two complex variables (z, w) . Let Z be the set of all integers, Z_+ the set of all nonnegative integers, $Z^2 = Z \times Z$ and $Z_+^2 = Z_+ \times Z_+$. For $1 \leq p \leq \infty$, $L^p = L^p(T^2, m)$ denotes the Lebesgue space and $H^p = H^p(T^2, m) = \{f \in L^p ; \hat{f}(\ell, n) = 0 \text{ if } (\ell, n) \notin Z_+^2\}$, that is, H^p denotes the usual Hardy space on T^2 . Any function f in H^p has an analytic extension in D^2 . m_z and m_w denote the normalized Lebesgue measures on the unit circle $T = T_z$ and $T = T_w$, $T^2 = T \times T$ and $m = m_z \times m_w$. For $1 \leq p \leq \infty$, $H^p(T)$ denotes the usual Hardy space on T . Let $K_0^p = \{f \in L^p ; \hat{f}(\ell, n) = 0 \text{ if } -(\ell, n) \in Z_+^2\}$, then $K_0^p = \{f \in L^p ; \int_{T^2} fg dm = 0 \text{ if } g \in H^q\}$ where $1/p + 1/q = 1$. Let $C = C(T^2)$ be the set of all continuous functions on T^2 , $K_0 = K_0^p \cap C$ and $A = H^p \cap C$. A is usually called the bidisc algebra. When $\alpha = (\alpha_1, \alpha_2) \in Z_+^2$ and $\zeta = (z, w) \in \mathbf{C}^2$, we write $\zeta^\alpha = z^{\alpha_1} w^{\alpha_2}$ and $|\alpha| = \alpha_1 + \alpha_2$. For $\alpha = (\alpha_1, \alpha_2)$ and $\beta = (\beta_1, \beta_2)$ in Z_+^2 , $0 \leq \alpha \leq \beta$ means that $0 \leq \alpha_1 \leq \beta_1$ and $0 \leq \alpha_2 \leq \beta_2$.

If ϕ is a function in L^∞ , we denote by B_ϕ the functional defined on H^1 by

$$B_\phi(f) = \int_{T^2} f(\zeta)\phi(\zeta)dm(\zeta).$$

The norm of B_ϕ is $\|B_\phi\| = \sup\{|B_\phi(f)| ; f \in S\}$ and let S_ϕ denote the set of all f in S for which $B_\phi(f) = \|B_\phi\|$. Here S is the unit sphere of H^1 . The functions in S_ϕ are called extremal functions. By the duality relation

$$\|B_\phi\| = \|\phi + K_0^\infty\|.$$

A function $\psi = \phi + k_0$ ($k_0 \in K_0^\infty$) for which $\|B_\phi\| = \|\psi\|_\infty$ is called an extremal kernel. If S_ϕ is not empty and ψ is an extremal kernel with $\|\psi\|_\infty = 1$, then $\psi = |f|/f$ for some function f in H^1 . This can be proved as in one variable case (see [D, §8.1 and §8.2], [Ha]). Let $\langle S_\phi \rangle$ denote the linear span of S_ϕ . Yabuta [Y1], [Y2] and Hasumi [Ha] and Wiegerinck [W] studied S_ϕ when $\langle S_\phi \rangle$ is of dimension one. For example, Yabuta [Y2] showed that if f is not identically zero and $\text{Re} f \geq 0$ a.e. on T^2 then $S_\phi = \{\gamma f ; \gamma > 0, \|\gamma f\|_1 = 1\}$. However, S_ϕ has not been studied when $\langle S_\phi \rangle$ is of finite dimension. In this paper we will study such cases. In Section 2, we show that if ϕ is a continuous function on T^2 , then S_ϕ is not empty. Moreover three important problems in this area are given. In Section 3, we show that if ϕ is a rational extremal kernel, then $\langle S_\phi \rangle$ is of finite dimension. In Section 4, S_ϕ is studied when ϕ is a product of one variable functions. In Sections 5 and 6, S_ϕ is studied when $\phi = \bar{\zeta}^\alpha |f_0|/f_0$ and $S_{|f_0|/f_0} = \{\gamma f_0 ; \gamma > 0, \|\gamma f_0\|_1 = 1\}$.

The function h in H^p is called an outer function if

$$\int_{T^2} \log |h| dm = \log \left| \int_{T^2} h dm \right| > -\infty$$

and a weakly outer function if

$$\int_{T^2} \log |h| dm = \int_T \log \left| \int_T h dm_z \right| dm_w = \int_T \log \left| \int_T h dm_w \right| dm_z > -\infty.$$

When $\phi = |f|/f$ and f is a function in H^1 , f is called a strongly outer function if

$$S_\phi = \{\gamma f ; \gamma > 0, \|\gamma f\|_1 = 1\}$$

and f is called a w -strongly outer function if for a.e. $z \in T$

$$S_{\phi_z} = \{\gamma f_z ; \gamma > 0, \|\gamma f_z\|_1 = 1\}$$

where $\phi_z(w) = \phi(z, w)$ and $f_z(w) = f(z, w)$. The following are known from [Ha]. If f is w -strongly outer and z -strongly outer, then f is strongly outer. However there exists a weakly outer function which is not outer. For example, $f(z, w) = z - w$. In the one variable case, a strongly outer function is outer but the converse is not true. In the two variable case, a strongly outer function may not be outer. In fact, it is easy to see that $f(z, w) = z - w$ is strongly outer but not outer.

2. Extremal kernel and problems

For a unimodular function ϕ in L^∞ , put $\mathcal{P}_\phi = \{f \in H^1 ; \phi f \text{ is nonnegative on } T^2\}$ and $\mathcal{R}_\phi = \{f \in H^1 ; \phi f \text{ is real on } T^2\}$. Then $\mathcal{P}_\phi \subseteq \mathcal{R}_\phi$. When ϕ is an extremal kernel and $S_\phi \neq \emptyset$, $\phi = \|B_\phi\| |f|/f$ for some function f in H^1 (see Introduction) and so $S_\phi \subset \mathcal{P}_\phi$.

Proposition 1. *If ϕ is a continuous function on T^2 , then S_ϕ is not empty and S_ϕ is weak $*$ compact.*

Proof : By a theorem of Bochner [B, p717], $(C/K_0)^* = H^1$ and so if ϕ is in C , then $S_\phi \neq \emptyset$ and S_ϕ is weak $*$ compact because the unit ball of H^1 is weak $*$ compact. \square

Lemma 1. *Let ϕ be a unimodular function in L^∞ . Then*

$$\mathcal{R}_\phi + i\mathcal{R}_\phi = H^1 \cap \bar{\phi}^2 \bar{H}^1 \text{ and } \mathcal{R}_\phi = \left\{ \frac{f + \bar{\phi}^2 \bar{f}}{2} ; f \in H^1 \cap \bar{\phi}^2 \bar{H}^1 \right\}.$$

Proof : If $f \in \mathcal{R}_\phi$, then $\phi^2 f^2$ is non-negative and so $\phi^2 f = \bar{f}$. Hence $f = \bar{\phi}^2 \bar{f}$ and $if = \bar{\phi}^2 \overline{(-if)}$. This implies that $\mathcal{R}_\phi + i\mathcal{R}_\phi \subseteq H^1 \cap \bar{\phi}^2 \bar{H}^1$. Conversely if $f \in H^1 \cap \bar{\phi}^2 \bar{H}^1$, then $f = \bar{\phi}^2 \bar{g}$ for some $g \in H^1$ and so $g = \bar{\phi}^2 \bar{f}$. Put $F = (f + \bar{\phi}^2 \bar{f})/2$, then $\phi^2 F = \bar{F}$ and so ϕF is real-valued. Hence $F \in \mathcal{R}_\phi$. Put $G = -i(f - \bar{\phi}^2 \bar{f})/2$, $\phi^2 G = \bar{G}$ and so ϕG is real-valued. Hence $G \in \mathcal{R}_\phi$. Since

$$F + iG = \frac{f + \bar{\phi}^2 \bar{f}}{2} + \frac{f - \bar{\phi}^2 \bar{f}}{2} = f,$$

$\mathcal{R}_\phi + i\mathcal{R}_\phi \supseteq H^1 \cap \bar{\phi}^2 \bar{H}^1$. This implies the first equality. The second equality follows from the proof of the first one. In fact, if $f \in \mathcal{R}_\phi$ then $f = (f + \bar{\phi}^2 \bar{f})/2$. Conversely if $F = (f + \bar{\phi}^2 \bar{f})/2$ and $f \in H^1 \cap \bar{\phi}^2 \bar{H}^1$, then ϕF is real-valued and $F \in \mathcal{R}_\phi$. \square

Proposition 2. *If ϕ is a unimodular function in L^∞ and ϕ is an extremal kernel, then $\{f^2 ; f \in \mathcal{R}_{\sqrt{\phi}} \cap H^2\} \subset S_\phi \subset \mathcal{R}_\phi$ and so $\langle S_\phi \rangle \subseteq H^1 \cap \bar{\phi}^2 \bar{H}^1$.*

Proof : Lemma 1 implies the proposition. \square

Now we give three important problems in this area. These have been solved affirmatively in one variable case by the author [N]. In the one variable case, $S_{\bar{\zeta}^\alpha} = \left\{ \lambda \prod_{j=1}^{\alpha} (\zeta - a_j)(1 - \bar{a}_j \zeta) \in S ; |a_j| \leq 1 (j = 1, \dots, \alpha) \text{ and } \lambda > 0 \right\}$ (see [D, p143]).

Problem 1. *If S_ϕ is a non-empty weak $*$ compact set, then is $\langle S_\phi \rangle$ of finite dimension ?*

Problem 2. *If $\phi = \bar{\zeta}^\alpha |f_0|/f_0$, $\alpha \in \mathbf{Z}_+^2$ and f_0 is a strongly outer function in H^1 , then is $\langle S_\phi \rangle$ of finite dimension ?*

Problem 3. *If $\langle S_\phi \rangle$ is of finite dimension, then*

$$S_\phi = (\{\gamma ; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times f_0) \cap S$$

where $\alpha \in \mathbf{Z}_+^2$ and f_0 is a strongly outer function ?

In this paper, we solve these problems in some special cases.

3. Rational extremal kernel

If ϕ is a rational function in L^∞ , then ϕ is continuous on T^2 and so by Proposition 1 S_ϕ is not empty. When ϕ is a rational function, unlike the one variable case. We don't know whether the extremal kernel is also rational. In this section, we show that $\langle S_\phi \rangle$ is of finite dimension when ϕ is a rational extremal kernel. If q is a rational inner function, then on T^2 $q = \zeta^\alpha \bar{g}/g$ where $\alpha = (\alpha_1, \alpha_2) \in \mathbf{Z}_+^2$, $\zeta^\alpha = z^{\alpha_1} w^{\alpha_2}$, g is an analytic polynomial without zeros in the bidisc D^2 . If q is continuous, then g is an invertible function in H^∞ (see [R, p112]).

Lemma 2. *If ψ is a rational function in L^∞ , then $H^1 \cap \psi\bar{H}^1$ is of finite dimension.*

Proof : Put $\psi = P/Q$ where P and Q are analytic polynomials, then

$$H^1 \cap \psi\bar{H}^1 \subseteq Q^{-1}(H^1 \cap P\bar{H}^1).$$

If $P = P(\zeta) = \sum_{|\alpha|=0}^n a_\alpha \zeta^\alpha$ where $\alpha \in Z_+^2$ and $\zeta \in \mathbf{C}^2$, then

$$H^1 \cap P\bar{H}^1 \subseteq \bigcup_{|\alpha|=0}^n H^1 \cap \zeta^\alpha \bar{H}^1 = \bigcup_{|\alpha|=n} H^1 \cap \zeta^\alpha \bar{H}^1.$$

Since $H^1 \cap \zeta^\alpha \bar{H}^1$ is of finite dimension, $H^1 \cap \psi\bar{H}^1$ is of finite dimension. \square

Theorem 2. *Let ϕ be a rational function and an extremal kernel. Then $\langle S_\phi \rangle$ is of finite dimension.*

Proof : We may assume that $\|\phi + K_0^\infty\| = 1$. Since ϕ is an extremal kernel, ϕ is unimodular rational function and $S_\phi = \{f \in H^1 ; \|f\|_1 = 1 \text{ and } \phi f \text{ is non-negative on } T^2\}$ (see Introduction). Hence $S_\phi \subset \mathcal{P}_\phi$. By Lemma 1 and Proposition 2, $S_\phi \subset H^1 \cap \bar{\phi}^2 \bar{H}^1$. Lemma 2 implies that $\langle S_\phi \rangle$ is of finite dimension. \square

Theorem 3. *Let q be a rational inner function. That is, suppose $q = \zeta^\alpha \bar{g}/g$, $\alpha \in Z_+^2$ and g is an analytic polynomial with no zeros in D^2 . If $\phi = \bar{q}$, then*

$$S_\phi = (\{\gamma ; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times g^{-2}) \cap S$$

and

$$\left\{ \left(\sum_{0 \leq \beta \leq \alpha} b_\beta \zeta^\beta \right)^2 \in S ; b_{\alpha-\beta} = \bar{b}_\beta \right\} \subset S_{\bar{\zeta}^\alpha} \subset \left\{ \sum_{0 \leq \beta \leq 2\alpha} b_\beta \zeta^\beta \in S ; b_{2\alpha-\beta} = \bar{b}_\beta \right\}.$$

Proof : If $f \in \mathcal{P}_\phi$, then $\bar{\zeta}^\alpha \frac{g}{\bar{g}} f \geq 0$ and so $\bar{\zeta}^\alpha g^2 f \geq 0$. Hence $g^2 \mathcal{P}_\phi \subseteq \mathcal{P}_{\bar{\zeta}^\alpha}$. If $F \in \mathcal{P}_{\bar{\zeta}^\alpha}$, then $\bar{\zeta}^\alpha F \geq 0$ and so $\bar{\zeta}^\alpha \frac{g}{\bar{g}} g^{-2} F \geq 0$. Hence $\mathcal{P}_\phi = (g^{-2} \mathcal{P}_{\bar{\zeta}^\alpha}) \cap H^1$ and so $S_\phi = \{\gamma F g^{-2} \in S ; \gamma > 0, F \in S_{\bar{\zeta}^\alpha}\}$. By Lemma 1

$$\mathcal{R}_{\bar{\zeta}^\alpha} + i\mathcal{R}_{\bar{\zeta}^\alpha} = H^1 \cap \zeta^{2\alpha} \bar{H}^1 = \left\{ \sum_{0 \leq \beta \leq 2\alpha} a_\beta \zeta^\beta \right\}$$

and

$$\mathcal{R}_{\bar{\zeta}^\alpha} = \left\{ \frac{1}{2} \left(\sum_{0 \leq \beta \leq 2\alpha} a_\beta \zeta^\beta + \zeta^{2\alpha} \sum_{0 \leq \beta \leq 2\alpha} \bar{a}_\beta \bar{\zeta}^\beta \right) \right\} = \left\{ \sum_{0 \leq \beta \leq 2\alpha} b_\beta \zeta^\beta ; b_{2\alpha-\beta} = \bar{b}_\beta \right\}.$$

Hence by Proposition 2 $S_{\bar{\zeta}^\alpha} \subset \left\{ \sum_{0 \leq \beta \leq 2\alpha} b_\beta \zeta^\beta \in S ; b_{2\alpha-\beta} = \bar{b}_\beta \right\}$. Observe that $\mathcal{R}_{\bar{\zeta}^{\alpha/2}} = \mathcal{R}_{\bar{\zeta}^{\alpha/2}} \cap H^2$ and by Lemma 1

$$\mathcal{R}_{\bar{\zeta}^{\alpha/2}} + i\mathcal{R}_{\bar{\zeta}^{\alpha/2}} = H^2 \cap \zeta^\alpha \bar{H}^2 = \left\{ \sum_{0 \leq \beta \leq \alpha} a_\beta \zeta^\beta \right\}.$$

Since $\mathcal{R}_{\bar{\zeta}^{\alpha/2}} = \left\{ \sum_{0 \leq \beta \leq \alpha} b_\beta \zeta^\beta ; b_{\alpha-\beta} = \bar{b}_\beta \right\}$, by Proposition 2 $\left\{ \left(\sum_{0 \leq \beta \leq \alpha} b_\beta \zeta^\beta \right)^2 \in S ; b_{\alpha-\beta} = \bar{b}_\beta \right\} \subset S_{\bar{\zeta}^\alpha}$. \square

Corollary 1. *Let $q = \zeta^\alpha \bar{g}/g$ be a rational inner function. If $\phi = \bar{q}$, then the dimension of $\langle S_\phi \rangle$ is less than or equal to $2|\alpha| + 1$. If ϕ is continuous, then the dimension of $\langle S_\phi \rangle$ is $2|\alpha| + 1$.*

Proof : The first statement is clear by Theorem 3 if the dimension of $\langle S_{\bar{\zeta}^\alpha} \rangle = 2|\alpha| + 1$. For the second statement, if ϕ is continuous, by [R, p112] g is invertible in H^∞ and so the dimension of $\langle S_\phi \rangle$ is equal to the dimension of $\langle S_{\bar{\zeta}^\alpha} \rangle$. Now we will prove that the dimension of $\langle S_{\bar{\zeta}^\alpha} \rangle = 2|\alpha| + 1$. For any $a \in \mathbf{C}$ with $|a| \leq 1$,

$$\zeta^\beta (\zeta^\gamma + a)(1 + \bar{a}\zeta^\gamma) \in \mathcal{P}_{\bar{\zeta}^\alpha}$$

where $\beta + \gamma = \alpha$, $\beta \in Z_+^2$ and $\gamma \in Z_+^2$. An elementary calculation implies that

$$\zeta^{\alpha+\beta} \in \langle \mathcal{P}_{\bar{\zeta}^\alpha} \rangle \text{ and } \zeta^{\alpha-\beta} \in \langle \mathcal{P}_{\bar{\zeta}^\alpha} \rangle$$

for any β with $\alpha - \beta \in Z_+^2$. Hence the dimension of $\langle \mathcal{P}_{\bar{\zeta}^\alpha} \rangle$ is $2|\alpha| + 1$ and $\langle S_{\bar{\zeta}^\alpha} \rangle = \langle \mathcal{R}_{\bar{\zeta}^\alpha} \rangle$. \square

In one variable case, if q is an irreducible inner function (that is, $q = z$) then $\langle S_{\bar{q}} \rangle$ is of dimension three. In the two variable case, this is not true. In fact, if $q(z, w) = \frac{zw - a}{1 - \bar{a}zw}$ for some $a \in D$, then $S_{\bar{q}} = S_{\bar{z}\bar{w}} \times (1 - \bar{a}zw)^{-2}$ and so $\langle S_{\bar{q}} \rangle$ is of dimension nine because $\langle S_{\bar{z}\bar{w}} \rangle = H^1 \cap z^2 w^2 \bar{H}^1$.

4. Kernels of products of one variable functions

In general, even if ϕ is a rational function, ϕ can not be factorized to $\phi_1 \phi_2$ where $\phi_1 = \phi_1(z)$ and $\phi_2 = \phi_2(w)$. In this section, we assume that ϕ can be factorized to $\phi_1 \phi_2$.

Theorem 4. Suppose $\phi = \phi_1\phi_2$ where $\phi_1 = \phi_1(z) \in L^\infty(T_z)$ and $\phi_2 = \phi_2(w) \in L^\infty(T_w)$. If S_{ϕ_j} is not empty for $j = 1, 2$, then S_ϕ is not empty and $S_\phi \supset S_{\phi_1} \times S_{\phi_2}$.

Proof : We may assume that $\|B_{\phi_j}\| = 1$ for $j = 1, 2$. Then $\|\phi_1 + zH^\infty(T_z)\| = \|\phi_2 + wH^\infty(T_w)\| = 1$. By the one variable theory, there exist unimodular functions $\psi_1 \in L^\infty(T_z)$ and $\psi_2 \in L^\infty(T_w)$ such that $\psi_1 = \phi_1 + g_1$ and $\psi_2 = \phi_2 + g_2$ where $g_1 \in zH^\infty(T_z)$ and $g_2 \in wH^\infty(T_w)$. Put $\psi = \psi_1\psi_2$, then ψ is a unimodular function in L^∞ and $B_\psi = B_\phi$ because

$$\psi = (\phi_1 + g_1)(\phi_2 + g_2) = \phi + g_1\phi_2 + g_2\phi_1 + g_1g_2$$

belongs to $\phi + K_0^\infty$. It is easy to see that $S_\psi \supseteq S_{\psi_1} \times S_{\psi_2} = S_{\phi_1} \times S_{\phi_2}$, and so $S_\psi \neq \emptyset$.

Lemma 4. Suppose $q = q_1q_2$ is an inner function where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one-variable inner functions. If q_1 is a finite Blaschke product, then

$$H^1 \cap q\bar{H}^1 = (H^1 \cap q_1\bar{H}^1) \times (H^1 \cap q_2\bar{H}^1),$$

and $H^1 \cap q_1\bar{H}^1 = H^1(T_z) \cap q_1\bar{H}^1(T_z)$ and $H^1 \cap q_2\bar{H}^1 = H^1(T_w) \cap q_2\bar{H}^1(T_w)$.

Proof : Since q_1 is a one variable finite Blaschke product, $q_1(z) = z^n\bar{g}/g$ where g is an invertible function in H^∞ . Hence

$$H^1 \cap q_1q_2\bar{H}^1 = g^{-1}(H^1 \cap z^nq_2\bar{H}^1).$$

If $f = z^nq_2\bar{h} \in H^1 \cap z^nq_2\bar{H}^1$, then $q_2\bar{h}$ is conjugate analytic in z and $z^nq_2\bar{h}$ is analytic in z . Hence $h = \sum_{j=0}^n h_j(w)z^j$ and so $f = z^nq_2\bar{h} = \sum_{j=0}^n q_2\bar{h}_jz^{n-j}$. Therefore

$$\begin{aligned} H^1 \cap z^nq_2\bar{H}^1 &= \sum_{j=0}^n (H^1 \cap q_2\bar{H}^1)z^j = \sum_{j=0}^n (H^1(T_w) \cap q_2\bar{H}^1(T_w))z^j \\ &= (H^1 \cap z^n\bar{H}^1) \times (H^1 \cap q_2\bar{H}^1) \end{aligned}$$

because $H^1 \cap z^n\bar{H}^1 = H^1(T_z) \cap z^n\bar{H}^1(T_z)$ and $H^1 \cap q_2\bar{H}^1 = H^1(T_w) \cap q_2\bar{H}^1(T_w)$. This implies the lemma. \square

Proposition 5. Suppose $q = q_1q_2$ is an inner function where $q_1 = q_1(z)$ and $q_2 = q_2(w)$ are one-variable finite Blaschke products. If $\phi = \bar{q}$, $\phi_1 = \bar{q}_1$ and $\phi_2 = \bar{q}_2$, then

$$\langle S_\phi \rangle = \langle S_{\phi_1} \rangle \times \langle S_{\phi_2} \rangle$$

and $S_{\phi_1} = S_{\phi_1} \cap H^1(T_z)$ and $S_{\phi_2} = S_{\phi_2} \cap H^1(T_w)$.

Proof : By Lemma 1,

$$S_\phi \subset \mathcal{P}_\phi \subset \mathcal{R}_\phi \subset H^1 \cap \bar{\phi}^2\bar{H}^1$$

and by Lemma 4,

$$\begin{aligned}
H^1 \cap \bar{\phi}^2 \bar{H}^1 &= H^1 \cap q_1^2 q_2^2 \bar{H}^1 \\
&= (H^1 \cap q_1^2 \bar{H}^1) \times (H^1 \cap q_2^2 \bar{H}^1) \\
&= (H^1(T_z) \cap q_1^2 \bar{H}^1(T_z)) \times (H^1(T_w) \cap q_2^2 \bar{H}^1(T_w)).
\end{aligned}$$

It is easy to see that $\langle S_{\phi_1} \rangle = H^1(T_z) \cap q_1^2 \bar{H}^1(T_z)$ and $\langle S_{\phi_2} \rangle = H^1(T_w) \cap q_2^2 \bar{H}^1(T_w)$ because q_1 and q_2 are finite Blaschke products. Thus $\langle S_\phi \rangle = \langle S_{\phi_1} \rangle \times \langle S_{\phi_2} \rangle$. \square

$$5. \quad \phi = \bar{\zeta}^\alpha |f_0|/f_0 \text{ for } f_0^{-1} \in H^1$$

If q is a rational inner function and $\phi = \bar{q}$, then $\phi = \bar{\zeta}^\alpha |g^{-2}|/g^{-2}$ where $\alpha \in \mathbf{Z}_+^2$ and g is an analytic polynomial with no zeros in D^2 . Then g^{-2} belongs to the class N_* (see [R, p44]) but we don't know whether g^{-2} is in H^1 . If q is continuous, it is known that g^{-2} belongs to H^∞ (see [R, p112]). In general, if $\phi = \bar{\zeta}^\alpha |f_0|/f_0$, $f_0 \in H^1$ and $f_0^{-1} \in H^\infty$, then by the proof of Theorem 3 it is easy to see that $S_\phi = (\{\gamma; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times f_0) \cap S$. In this section, we assume that f_0^{-1} belongs to H^1 but we don't assume that $f_0^{-1} \in H^\infty$.

Lemma 5. *If $\alpha \in \mathbf{Z}_+^2$ and $f \in H^{1/2}$ such that $\bar{\zeta}^\alpha f$ is non-negative on T^2 , then f belongs to H^1 .*

Proof : If $\alpha = (\ell, m)$, $\ell \geq 0$ and $m \geq 0$, then by a result of one variable Hardy space [HS] $\bar{z}^\ell \bar{w}^m f(z, w) = \sum_{j=-\ell}^{\ell} a_j(w) z^j$ a.e. w and $\bar{z}^\ell \bar{w}^m f(z, w) = \sum_{j=-m}^m b_j(z) w^j$ a.e. z . \square

Proposition 6. *Suppose $\alpha \in \mathbf{Z}_+^2$ and f_0 is a function in H^1 whose inverse is in H^1 . If $\phi = \bar{\zeta}^\alpha |f_0|/f_0$, then*

$$S_\phi = (\{\gamma; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times f_0) \cap S.$$

Proof : It is clear that $(\{\gamma; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times f_0) \cap S \subseteq S_\phi$. If $f \in S_\phi$, then $\bar{\zeta}^\alpha f/f_0 \geq 0$ a.e. on T^2 and $f f_0^{-1} \in H^{1/2}$. Lemma 5 implies that $f f_0^{-1}$ belongs to H^1 and so $f f_0^{-1} \in \{\gamma; \gamma > 0\} \times S_{\bar{\zeta}^\alpha}$. Thus $f \in (\{\gamma; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times f_0) \cap S$. \square

6. $\phi = \bar{\zeta}^\alpha |f_0|/f_0$ and $\operatorname{Re} f_0 \geq 0$

We call a function f in H^1 w -strongly outer when $f(z, w)$ is strongly outer with respect to w for a.e. z in T . It is easy to see that if f is w -strongly outer and z -strongly outer, then f is strongly outer [Ha]. The converse is not true [Ha]. When f is a function in H^1 with $f^{-1} \in H^1$ or with $\operatorname{Re} f \geq 0$, f is w -strongly outer and z -strongly outer (see [N]). In this section, we consider an easier problem than Problems 2 and 3 in §2. Suppose f_0 is w -strongly outer and z -strongly outer, $\alpha \in Z_+^2$ and $\phi = \bar{\zeta}^\alpha |f_0|/f_0$. We can ask whether

$$S_\phi = (\{\gamma ; \gamma > 0\} \times S_{\bar{\zeta}^\alpha} \times f_0) \cap S.$$

In this section, we show that it is true when $\alpha = (\alpha_1, 0)$ or $\alpha = (0, \alpha_2)$. In particular, if $\operatorname{Re} f_0 \geq 0$, then it is true. ϕ_z is defined by $\phi_z(w) = \phi(z, w)$ for a.e. z and ϕ_w is defined similarly.

Theorem 6. *Suppose f is a weakly outer function in H^1 and a w -strongly outer function, and $\phi = |f|/f$.*

- (1) $\langle S_{\phi_z} \rangle$ is of finite dimension one for a.e. $z \in T$.
- (2) If $\langle S_{\phi_w} \rangle$ is of finite dimension for a.e. w , then there exist a non-negative integer n and a strongly outer function f_0 in H^1 such that

$$S_\phi = (\{\gamma ; \gamma > 0\} \times S_{z^n} \times f_0) \cap S.$$

Proof: If $g \in S_\phi$ and $\alpha_g = g/f$, then α_g is a non-negative function in the class N_* (see [R, p112]) because f is weakly outer and $\alpha_g = \alpha_g(z)$ because f is w -strongly outer. The first statement is clear. We will show the second statement. If $\langle S_{\phi_w} \rangle$ is of finite dimension for all $w \in E$ with $m_w(T \setminus E) = 0$, put $E_k = \{w \in E ; \langle S_{\phi_w} \rangle \text{ has dimension } k\}$ for $1 \leq k < \infty$. By [N, Theorem 2], we may assume that $k = 2n + 1$ and $0 \leq n < \infty$. Then there exist the smallest integer n such that $k = 2n + 1$ and $E_k \neq \emptyset$. Since $\langle S_{\phi_w} \rangle$ is of finite dimension $k = 2n + 1$ for all $w \in E_k$, by [N] for any $g \in S_\phi$

$$\alpha_g(z) = \frac{\prod_{j=1}^n (z - a_j)(1 - \bar{a}_j z)}{\prod_{j=1}^n (z - b_j)(1 - \bar{b}_j z)}$$

where $\{a_j\}_{j=1}^n$ and $\{b_j\}_{j=1}^n$ are in \bar{D} . Moreover there exists a function $g \in S_\phi$ such that $\{a_j\}_{j=1}^n \cap \{b_j\}_{j=1}^n = \emptyset$ in the above representation of α_g . We will show that $E_k = E$. If $E_k \neq E$ and $k = 2n + 1$, then there exists an integer ℓ such that $n < \ell < \infty$ and $E_{2\ell+1} \neq \emptyset$. Hence, again by [N] there exists a function $h \in S_\phi$ such that

$$\alpha_h(z) = \frac{\prod_{j=1}^\ell (z - c_j)(1 - \bar{c}_j z)}{\prod_{j=1}^\ell (z - d_j)(1 - \bar{d}_j z)}$$

where $\{c_j\}_{j=1}^\ell$ and $\{d_j\}_{j=1}^\ell$ are in \bar{D} , and $\{c_j\}_{j=1}^\ell \cap \{d_j\}_{j=1}^\ell = \emptyset$. This contradicts that $\alpha_h(z) = \prod_{j=1}^n (z - a_j)(1 - \bar{a}_j z) / \prod_{j=1}^n (z - b_j)(1 - \bar{b}_j z)$ with $n < \ell$. Hence $E_k = E$. Thus $\langle S_{\phi_w} \rangle$ is of finite dimension $2n + 1$ for a.e. w . By the above argument, there exists a function $g \in S_\phi$ such that

$$g(z, w) = \frac{\prod_{j=1}^n (z - a_j)(1 - \bar{a}_j z)}{\prod_{j=1}^n (z - b_j)(1 - \bar{b}_j z)} f(z, w)$$

where $\{a_j\}_{j=1}^n$ and $\{b_j\}_{j=1}^n$ are two disjoint sets in \bar{D} . Since g is in H^1 ,

$$f_0(z, w) = f(z, w) / \prod_{j=1}^n (z - b_j)(1 - \bar{b}_j z)$$

belongs to H^1 because $\{a_j\}_{j=1}^n \cap \{b_j\}_{j=1}^n = \emptyset$. Then $f_0(z, w)$ is strongly outer. For, if not, there exists a function k in the unit ball of H^1 such that $\beta(z) = k(z, w) / f_0(z, w)$ is non-negative and non-constant on T^2 because $f_0(z, w)$ is w -strongly outer. Since $\langle S_{\phi_w} \rangle$ is of finite dimension for a.e. w , $\beta(z) = \prod_{j=1}^s (z - c_j)(1 - \bar{c}_j z) / \prod_{j=1}^s (z - d_j)(1 - \bar{d}_j z)$, where $1 \leq$

$s < \infty$, $\{c_j\}_{j=1}^s$ and $\{d_j\}_{j=1}^s$ are two disjoint set in \bar{D} . Then $f_0(z, w) / \prod_{j=1}^s (z - d_j)(1 - \bar{d}_j z)$

belongs to H^1 . This contradicts that $\langle S_{\phi_w} \rangle$ is of finite dimension n for a.e. w . Hence $f_0(z, w)$ is strongly outer. Thus $S_\phi = (\{\gamma ; \gamma > 0\} \times S_{\bar{z}^n} \times f_0) \cap S$.

Corollary 3. *Suppose $f_0 \in H^1$ is w -strongly outer and z -strongly outer and $\phi = \bar{z}^n |f_0| / f_0$ for some positive integer n . Then*

$$S_\phi = (\{\gamma ; \gamma > 0\} \times S_{\bar{z}^n} \times f_0) \cap S.$$

Proof : Let $f = \gamma \prod_{j=1}^n (z - a_j)(1 - \bar{a}_j z) f_0$ where $|a_j| = 1$ ($1 \leq j \leq n$), $\gamma > 0$ and $\|f\|_1 = 1$. Then $\phi = |f| / f$, f is a weakly outer function in H^1 and w -strongly outer. Apply Theorem 6.

Corollary 4. *Suppose f_0 is a nonzero function in H^1 with $\operatorname{Re} f_0 \geq 0$ and $\phi = \bar{z}^n |f_0| / f_0$ for some positive integer n . Then*

$$S_\phi = (\{\gamma ; \gamma > 0\} \times S_{\bar{z}^n} \times f_0) \cap S.$$

Proof : If $\operatorname{Re} f_0 \geq 0$ and $f_0 \not\equiv 0$, then f_0 is w -strongly outer and z -strongly outer.

References

- [B] S.Bochner, Boundary values of analytic functions in several variables and of almost periodic functions, Ann. of Math. 45(1944), 708-722.
- [D] P.L.Duren, Theory of H^p Spaces, Academic Press Inc, New York and London 1970.
- [Ha] M.Hasumi, Extreme points and unicity of extremum problems in H^1 on the polydiscs, Pacific J.Math.44(1973), 523-535.
- [N] T.Nakazi, Exposed points and extremal problems in H^1 , J.Funct.Anal. 53(1983), 224-230.
- [HS] H.Helson and D.Sarason, Past and future, Math.Scand. 21(1967), 5-16.
- [Y1] K.Yabuta, Unicity of the extremum problems in $H^1(U^n)$, Proc.Amer.Math.Soc. 28(1971), 181-184.
- [Y2] K.Yabuta, Some uniqueness theorems for $H^p(U^n)$ functions, Tohoku Math.J. 24(1972), 353-357.
- [R] W.Rudin, Function Theory in Polydiscs, Benjamin, New York, 1969.
- [W] J.Wiegerinck, A characterization of strongly exposed points of the unit ball of H^1 , Indag.Mathem., n.s. 4(509-519).

Hokkaido University
Department of Mathematics
Sapporo 060-0810,
Japan