



Title	Invariant subspaces of finite codimension and uniform algebras
Author(s)	Nakazi, T.; Osawa, T.
Citation	Hokkaido University Preprint Series in Mathematics, 595, 1-6
Issue Date	2003-07
DOI	10.14943/83740
Doc URL	http://hdl.handle.net/2115/69344
Type	bulletin (article)
File Information	pre595.pdf



[Instructions for use](#)

Invariant Subspaces Of Finite
Codimension And Uniform Algebras

Takahiko Nakazi and Tomoko Osawa

Series #595. July, 2003

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #569 J.M. Hwang and K. Yamaguchi, Characterization of Hermitian symmetric spaces by fundamental forms, 10 pages. 2002.
- #570 H. Ishii and T. Mikami, Convexified Gauss curvature flow of bounded open sets in an anisotropic external field: a stochastic approximation and PDE, 37 pages. 2002.
- #571 Y. Nakano, Minimization of shortfall risk in a jump-diffusion model, 10 pages. 2002.
- #572 K. Izuchi and T. Nakazi, Backward shift invariant subspaces in the bidisc, 8 pages. 2002.
- #573 S. Izumiya, D. Pei and M. C. Romero-Fuster, The horospherical geometry of surfaces in Hyperbolic 4-space, 17 pages. 2002.
- #574 S. Izumiya and M. C. Romero-Fuster, The hyperbolic Gauss-Bonnet type theorem, 10 pages. 2002.
- #575 S. Izumiya and S. Janeczko, A symplectic framework for multiplane gravitational lensing, 19 pages. 2002.
- #576 S. Izumiya, M. Kossowski, D. Pei and M. C. Romero-Fuster, Singularities of C^∞ -lightlike hypersurfaces in Minkowski 4-space, 18 pages. 2002.
- #577 S. Izumiya, D. Pei and M. Takahashi, Evolutes of hypersurfaces in Hyperbolic space, 21 pages. 2002.
- #578 Y. Giga, S. Matsui and S. Sasayama, Blow up rate for semilinear heat equation with subcritical nonlinearity, 29 pages. 2002.
- #579 M. Tsujii, Physical measures for partially hyperbolic surface endomorphisms, 71 pages. 2003.
- #580 Y. Giga and K. Yamada, On viscous Burgers-like equations with linearly growing initial data, 19 pages. 2003.
- #581 T. Nakazi and T. Osawa, Spectra of Toeplitz Operators and Uniform Algebras, 9 pages. 2003.
- #582 Y. Daido, M. Ikehata and G. Nakamura, Reconstruction of Inclusions for the Inverse Boundary Value Problem with Mixed Type Boundary Condition, 18 pages. 2003.
- #583 Y. Daido and G. Nakamura, Reconstruction of Inclusions for the Inverse Boundary Value Problem with Mixed Type Boundary Condition and Source Term, 26 pages. 2003.
- #584 M.-H. Giga and Y. Giga, A PDE approach for motion of phase-boundaries by a singular interfacial energy, 19 pages. 2003.
- #585 A.A. Davydov, G. Ishikawa, S. Izumiya and W.-Z. Sun, Generic singularities of implicit systems of first order differential equations on the plane, 28 pages. 2003.
- #586 K. Yamauchi, On an underlying structure for the consistency of viscosity solutions, 12 pages. 2003.
- #587 T. Miyao, Momentum operators with a winding gauge potential, 15 pages. 2003.
- #588 Y. Giga and R. Kobayashi, On constrained equations with singular diffusivity, 35 pages. 2003.
- #589 O. Sawada, On analyticity rate estimates of the solutions to the Navier-Stokes equations in Bessel-potential spaces, 13 pages. 2003.
- #590 T. Nakazi, Exposed points and extremal problems in H^1 on a bidisc, 11 pages. 2003.
- #591 Y. Tsai and Y. Giga, A numerical study of anisotropic crystal growth with bunching under very singular vertical diffusion, 9 pages. 2003.
- #592 K. Izuchi, T. Nakazi and M. Seto, Backward shift invariant subspaces in the bidisc II, 17 pages. 2003.
- #593 S. Jimbo, Singular perturbation of domains and Semilinear elliptic equations III, 26 pages. 2003.
- #594 Y. Giga, S. Matsui and S. Sasayama, On blow up rate for sign-changing solutions in a convex domain, 12 pages. 2003.

Invariant Subspaces Of Finite Codimension And Uniform Algebras

by

Takahiko Nakazi*

and

Tomoko Osawa

* This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education.

2000 Mathematics Subject Classification : 46 J 15, 46 J 20

Key words and phrases : uniform algebra, Q -algebra, finite dimension, invariant subspace, ideal

Abstract. Let A be a uniform algebra on a compact Hausdorff space X and m a probability measure on X . Let $H^p(m)$ be the norm closure of A in $L^p(m)$ with $1 \leq p < \infty$ and $H^\infty(m)$ the weak $*$ closure in $L^\infty(m)$. In this paper, we describe a closed ideal of A and a closed invariant subspace of $H^p(m)$ which is of finite codimension.

§1. Introduction

Let A be a uniform algebra on a compact Hausdorff space X . $M(A)$ denotes the maximal ideal space of A . Let m be a probability measure on X . $H^p(m)$ denotes the norm closure of A in $L^p(m)$ with $1 \leq p < \infty$ and $H^\infty(m)$ denotes the weak $*$ closure of A in $L^\infty(m)$. $H^p(m)$ is called an abstract Hardy space. When A is a disc algebra, if m is the normalized Lebesgue measure on the unit circle, $H^p(m)$ is the usual Hardy space and if m is the normalized area measure on the unit disc, $H^p(m)$ is the usual Bergman space.

Let I be a closed ideal of A . In this paper, we are interested in I with $\dim A/I < \infty$. Then A/I is called a Q-algebra. Two dimensional Q-algebras can be described easily, that is, $I = \{f \in A ; \phi_1(f) = \phi_2(f) = 0\}$ where $\phi_j \in M(A)$ ($j = 1, 2$), or $I = \{f \in A ; \phi(f) = D_\phi(f) = 0\}$ where $\phi \in M(A)$ and D_ϕ is a bounded point derivation at ϕ . One of the authors [3] showed that a two dimensional operator algebra on a Hilbert space is a Q-algebra. It seems to be worthwhile to describe a finite dimensional Q-algebra. In Section 2, we describe an ideal I with $\dim A/I < \infty$ using a theorem of T.W.Gamelin [2]. As a result, a finite dimensional Q-algebra is described.

When M is a closed subspace of $H^p(m)$ and $AM \subset M$, M is called an invariant subspace. In this paper, we are interested in M with $\dim H^p(m)/M < \infty$. When A is the polydisc algebra on T^n and m is the normalized Lebesgue measure on T^n , a finite codimensional invariant subspace M in $H^p(m)$ was described by P.Ahern and D.N.Clark [1] using the ideals in the polynomial ring $\mathcal{C}[z_1, \dots, z_n]$ of finite codimension whose zero sets are contained in the polydisk, D^n . In Section 3, for arbitrary uniform algebra A we describe a finite codimensional invariant subspace M in $H^p(m)$ using the result in Section 2.

§2. Finite codimensional ideal

Let $\phi \in M(A)$. A closed subalgebra H of A is a (ϕ, k) -subalgebra if there is a sequence of closed subalgebras $A = A_0 \supset A_1 \supset \dots \supset A_k = H$ such that A_j is the kernel of a continuous point derivation D_j of A_{j-1} at ϕ . If H is a (ϕ, k) -subalgebra of A , then H has finite codimension in A and $M(H) = M(A)$ by [2, Lemma 9.1].

If I is a closed ideal of A and A/I is of finite dimension, $B = \mathcal{C} + I$ is a closed subalgebra of A and A/B is of finite dimension. By a theorem of T. W. Gamelin [2, Theorem 9.8], we can describe B and so I . Since B is a special closed subalgebra of A we can describe I more explicitly.

Theorem 1. *If I is a closed ideal of A and A/I is of finite dimension, then there exists a closed subalgebra $E = E(I)$ of A such that $E = \{f \in A : \phi_1(f) = \dots = \phi_n(f)\}$, $1 \leq n < \infty$, $\{\phi_j\} \subset M(A)$ and*

$$I = H_\phi^E \cap \ker \phi$$

where $\phi = \phi_j|_E$, $1 \leq j \leq n$ and H_ϕ^E is a (ϕ, k) -subalgebra with respect to E for some k .

Proof. Put $H = I + \mathcal{C}$, then A/H is of finite dimension. By a theorem of T.W.Gamelin [2, Theorem 9.8], H can be obtained from A in two steps.

(i) There exist pairs of points $\psi_j, \psi'_j, 1 \leq j \leq \ell$, in $M(A)$ such that if E consists of the $f \in A$ such that $\psi_j(f) = \psi'_j(f), 1 \leq j \leq \ell$, then $H \subset E \subset A$.

(ii) There exist distinct points $\theta_j \in M(E)$ and θ_j -subalgebras H_j of $E, 1 \leq j \leq k$, such that $H = H_1 \cap \dots \cap H_k$.

Put $\tilde{\psi}_j = \psi_j|_E = \psi'_j|_E$ for $1 \leq j \leq \ell$, then $\tilde{\psi}_j$ belongs to $M(E)$. Since I is an ideal of $A, I \subset \bigcap_{j=1}^{\ell} \ker \tilde{\psi}_j$. For let $f \in A$ such that $\psi_j(f) \neq \psi'_j(f)$. If $g \in I$, then $fg \in I$ but $\psi_j(fg) \neq \psi'_j(fg)$ when $\tilde{\psi}_j(g) \neq 0$. This contradicts that $fg \in E$. Thus $\tilde{\psi}_j(g) = 0$. Hence $I \subset \bigcap_{j=1}^{\ell} \ker \tilde{\psi}_j$ and so $H \subseteq \bigcap_{j=1}^{\ell} \ker \tilde{\psi}_j + \mathcal{C}$. By definition of $E, \tilde{\psi}_1 = \dots = \tilde{\psi}_{\ell}$. Therefore E has the form : $E = \{f \in A ; \phi_1(f) = \dots = \phi_n(f)\}, 1 \leq n < \infty$ and $\{\phi_j\} \subset M(A)$.

For each j with $1 \leq j \leq k, H_j$ is a θ_j -subalgebra of E for $\theta_j \in M(E)$. Hence there is a sequence of closed subalgebras $E = E_{j0} \supset E_{j1} \supset \dots \supset E_{j\ell_j} = H_j$ such that E_{jt} is the kernel of a continuous point derivation D_{jt} of $E_{j,t-1}$ at θ_j . We will write

$E_{j\ell_j} = \ker D_{\theta_j}$ where D_{θ_j} is a derivation on $E_{j(\ell_j-1)}$. Then $H = \bigcap_{j=1}^k \ker D_{\theta_j}$ and so

$I = \left\{ \bigcap_{j=1}^k \ker D_{\theta_j} \right\} \cap \ker \theta$ for some $\theta \in M(H)$. Suppose g is an arbitrary function in I .

For any $j(1 \leq j \leq k)$, there exists a function $f \in E_{j(\ell_j-1)}$ such that $f \notin E_{j\ell_j} = \ker D_{\theta_j}$. Since $fg \in I$ and $D_{\theta_j}(g) = 0, D_{\theta_j}(fg) = \theta_j(g)D_{\theta_j}(f) = 0$ because D_{θ_j} is a derivation on

$E_{j(\ell_j-1)}$. This implies that $\theta_j(g) = 0$. Hence $I \subset \bigcap_{j=1}^k \ker \theta_j$. Therefore by definition of

$E, \theta_1 = \dots = \theta_k \in M(E)$ and so $H_1 = \dots = H_k$. Thus $\theta_1|_H = \theta$ and $I = (\ker D_{\theta_1}) \cap \ker \theta_1$.

Since $I \subset \bigcap_{j=1}^n \ker \phi_j, I \subset (\ker \phi_1) \cap (\ker D_{\theta_1} \cap \ker \theta_1)$ and so $\phi_1|_E = \theta_1$.

Corollary 1. *If I is a closed ideal of A and A/I is of finite dimension 2, then $I = \{f \in A ; \phi_1(f) = \phi_2(f) = 0\}$ where $\phi_j \in M(A) (j = 1, 2)$ and $\phi_1 \neq \phi_2$, or $I = \{f \in A ; \phi(f) = D_{\phi}(f) = 0\}$ where $\phi \in M(A)$ and D_{ϕ} is a bounded point derivation at ϕ .*

Proof. When $\dim A/I = 2$, by Theorem 1 $E = A$ or $E = \{f \in A ; \phi_1(f) = \phi_2(f)\}$. If $E = A$, then $H_{\phi}^E = \{f \in A ; D_{\phi}(f) = 0\}$ and if $E = \{f \in A ; \phi_1(f) = \phi_2(f)\}$, then $H_{\phi}^E = E$. For $\dim A/H_{\phi}^E = 1$ because $H_{\phi}^E = I + \mathcal{C}$. This implies the corollary.

Corollary 2. *If B is a finite dimensional Q -algebra and $B_0 = \text{rad } B$ is its radical, then there exist subalgebras B_1, B_2, \dots, B_{k+1} in B_0 such that $B_{k+1} = \{0\}, \dim B_j/B_{j+1} = 1$ and B_{j+1} is an ideal of B_j for $j = 0, 1, \dots, k$. Hence $\text{rad } B$ has a basis $\{f_0, f_1, \dots, f_k\}$ such that $(f_j)^{2^{(k+1)-j}} = 0$ for $j = 0, 1, \dots, k$.*

Proof. Since B is a Q -algebra, $B = A/I$ for some uniform algebra A and some closed ideal I of A . Since B is of finite dimension, we can apply Theorem 1 to A and I . In the notation of Theorem 1, $\text{rad } B = \{f \in E ; \phi(f) = 0\}/I$. Since H_ϕ^E is a ϕ -subalgebra with respect to E , there exists a sequence of closed subalgebras $E = E_0 \underset{\neq}{\supset} E_2 \underset{\neq}{\supset} \cdots \underset{\neq}{\supset} E_{k+1} = H_\phi^E$ such that E_j is the kernel of a continuous point derivation D_j of E_{j-1} at ϕ . Hence $E_{j+1} \cap \ker \phi$ is an ideal of $E_j \cap \ker \phi$ and $\dim\{E_j \cap \ker \phi / E_{j+1} \cap \ker \phi\} = 1$. Put $B_j = (E_j \cap \ker \phi)/I$. Then $\dim B_j / B_{j+1} = 1$ and B_{j+1} is an ideal of B_j for $j = 0, 1, \dots, k$ and $B_{k+1} = \{0\}$. For each j , there exists f_j such that $B_j = \langle f_j \rangle + B_{j+1}$ and then $\{f_0, f_1, \dots, f_k\}$ is a basis of $\text{rad } B = B_0$. Since f_j^2 belongs to B_{j+1} because $E_{j+1} = \ker D_{j+1}$. Thus $(f_j)^{2(k+1-j)} = 0$.

§3. Finite codimensional invariant subspace

Let m be a probability measure on X . For $1 \leq p < \infty$, $H^p(m)$ denotes the norm closure of A in $L^p(m)$ and $H^\infty(m)$ denotes the weak $*$ closure of A in $L^\infty(m)$. When M is a closed subspace of $H^p(m)$ and $AM \subset M$, M is called an invariant subspace. For a subset S of $H^p(m)$, $[S]_p$ denotes the closure of S in $H^p(m)$.

Theorem 2. *If M is an invariant subspace of H^p with $\dim H^p/M = n < \infty$, then there exists a closed ideal of A such that $\dim A/I = n$, $[I]_p = M$ and $I = M \cap A$. If H_ϕ^E is a (ϕ, k) -subalgebra with respect to $E = E(I)$, then $[E_j]_p \underset{\neq}{\supset} [E_{j+1}]_p$ for any $j (0 \leq j \leq k-1)$ and $\dim H^p/[E]_p = \dim A/E$. Conversely if $\dim A/I = n < \infty$ then $\dim H^p/[I]_p \leq n$. If $[E_j]_p \underset{\neq}{\supset} [E_{j+1}]_p$ for any $j (0 \leq j \leq k-1)$ and $\dim H^p/[E]_p = \dim A/E$, then $\dim H^p/[I]_p = n$ and $[I]_p \cap A = I$.*

Proof. Suppose that M is an invariant subspace of $H^p(m)$ and $\dim H^p(m)/M = n < \infty$. Then there exist n linear independent linear functionals ψ_1, \dots, ψ_n in $(H^p)^*$ such that $\psi_j = 0$ on M for $1 \leq j \leq n$. Put $\phi_j = \psi_j|_A$ for $1 \leq j \leq n$ and $I = M \cap A$, then $I = \bigcap_{j=1}^n \ker \phi_j$ and so $\dim A/I = n$. For ϕ_1, \dots, ϕ_n are independent linear functionals in A^* because A is dense in $H^p(m)$. If $M \underset{\neq}{\supset} [I]_p$, then there exists $\psi_{n+1} \in (H^p)^*$ such that $\psi_{n+1} = 0$ on $[I]_p$ and $\psi_1, \dots, \psi_n, \psi_{n+1}$ are independent linear functionals in $(H^p)^*$. If put $\phi_{n+1} = \psi_{n+1}|_A$, then $\phi_1, \dots, \phi_n, \phi_{n+1}$ are independent linear functionals in A^* and $I \subseteq \bigcap_{j=1}^{n+1} \ker \phi_j$. This contradiction implies that $M = [I]_p$. Note that $\dim H^p/[E_k] = \dim H^p/[I]_p - 1 = \dim A/I - 1 = \dim A/E_k$. If $\dim H^p/[E_0]_p < \dim A/E_0$ where $E_0 = E$ or $[E_j]_p = [E_{j+1}]_p$ for some $j (0 \leq j \leq k-1)$, then this contradicts that $\dim H^p/[E_k]_p = \dim A/E_k$. The converse is clear.

Corollary 3. *If M is an invariant subspace of H^p with $\dim H^p/M = 2$, then $M = \{f \in H^p ; \Phi_1(f) = \Phi_2(f) = 0\}$ where $\Phi_j \in (H^p)^*$, and $\Phi_j(fg) = \Phi_j(f)\Phi_j(g)$ for*

$f \in H^p$ and $g \in A$, or $M = \{f \in H^p ; \Phi(f) = D_\phi(f) = 0\}$ where $\Phi, D_\Phi \in (H^p)^*$, and $\Phi(fg) = \Phi(f)\Phi(g)$ and $D_\Phi(fg) = \Phi(f)D_\Phi(g) + \Phi(g)D_\Phi(f)$ for $f \in H^p$ and $g \in A$.

Proof. This is a result of Corollary 1 and Theorem 2.

Corollary 4. *If M is an invariant subspace of H^p with $\dim H^p/M = n < \infty$, then there exist f_1, \dots, f_n in A such that $\{f_j + M\}_{j=1}^n$ is a basis in H^p/M .*

Proof. By Theorem 2, if $I = M \cap A$ then $\dim A/I = n$ and $M = [I]_p$. Hence there exist f_1, \dots, f_n in A such that $\{f_j + I\}_{j=1}^n$ is a basis in A/I . If f_j belongs to M , then f_j also belongs to $M \cap A = I$ and so f_j does not belong to M . This shows the corollary.

References

1. P.Ahern and D.N.Clark, Invariant subspaces and analytic continuation in several variables, J.Math.Mech. 19(1969/1970), 963-969.
2. T.W.Gamelin, Embedding Riemann surfaces in maximal ideal spaces, J.Funct.Anal. 2(1968), 123-146.
3. T.Nakazi, Two-dimensional Q -algebras, Linear algebra Appl. 315(2000), 197-205.

Takahiko Nakazi
Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan

E-mail : nakazi@math.sci.hokudai.ac.jp

Tomoko Osawa
Mathematical and Scientific Subjects
Asahikawa National College of Technology
Asahikawa 071-8142, Japan

E-mail : ohsawa@asahikawa-nct.ac.jp