Title	Steady solution and its stability to Navier-Stokes equations with general Navier slip boundary condition
Author(s)	Itoh, Shigeharu; Tanaka, Naoto; Tani, Atusi
Citation	Hokkaido University Preprint Series in Mathematics, 643, 1-23
Issue Date	2004
DOI	10.14943/83796
Doc URL	http://hdl.handle.net/2115/69450
Type	bulletin (article)
File Information	pre643.pdf

[Instructions for use](#)

Steady solution and its stability to Navier-Stokes equations with general Navier slip boundary condition

S.Itoh, N.Tanaka and A.Tani

Dedicated to the memory of Olga Ladyzenskaya

Shigeharu Itoh

Hirosaki University, Department of Mathematics, Hirosaki 036-8560, Japan,
sitoh@cc.hirosaki-u.ac.jp

Naoto Tanaka

Fukuoka University, Department of Applied Mathematics, Fukuoka 814-0180,
Japan, naoto@cis.fukuoka-u.ac.jp

Atusi Tani

Keio University, Department of Mathematics, Yokohama 223-8522, Japan,
tani@math.keio.ac.jp

Abstract. Steady solution and asymptotic behaviour of corresponding non-steady solution are studied for the Navier-Stokes equations under general Navier slip boundary condition. It is proved that the existence of a unique stationary solution and that this solution is asymptotically stable under some restrictions of the data.

Mathematics Subject Classification (2000). 35B40, 35Q30, 76D03, 76D05.

Keywords. incompressible Navier-Stokes equations, general Navier slip boundary condition, steady solution, stability of steady solution.

1 Introduction

The motion of an incompressible viscous fluid in a bounded domain Ω in $\mathbb{R}^3$ is described by the Navier-Stokes equations :

$$\frac{\partial v}{\partial t} + (v \cdot \nabla)v - \nabla \cdot \mathbf{P}(v, p) = f, \quad \nabla \cdot v = 0. \quad (1.1)$$

Here $v = v(x, t)$ is the velocity vector field, $p = p(x, t)$ is the pressure, $\mathbf{P}(v, p) = -p\mathbf{I}_3 + 2\nu\mathbf{D}(v)$ is the stress tensor, $\mathbf{I}_3$ is an identity matrix of degree 3, $\mathbf{D}(v)$ is the velocity deformation tensor with the elements $D_{ij} = \frac{1}{2}(\frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i})$, $\nu > 0$ is a constant coefficient of viscosity and $f = f(x, t)$ is a given vector field of external forces.

Equations (1.1) is considered under the initial condition

$$v|_{t=0} = v_0(x) \quad (1.2)$$

and the boundary conditions

$$v \cdot n = 0, \quad v \cdot \tau = K\mathbf{P}n \cdot \tau, \quad (1.3)$$

or equivalently,

$$v \cdot n = 0, \quad v = K[\mathbf{P}n - (\mathbf{P}n \cdot n)n], \quad (1.4)$$

where n and τ are a unit inward normal and a unit tangential vectors to a smooth boundary Γ of Ω , respectively, such that $n \times \tau = 1$ and $K = K(x, t)$ is assumed to be a nonnegative function defined on $\Gamma_\infty = \Gamma \times (0, \infty)$. Dividing both sides of (1.4)₂ by $1 + \nu K$ and using the same letter K in place of $1/(1 + \nu K)$, we have

$$v \cdot n = 0, \quad 2(1 - K)\Pi\mathbf{D}(v)n - Kv = 0, \quad (1.5)$$

where $0 \leq K \leq 1$ and $\Pi w = w - (w \cdot n)n$. Condition (1.3) implies that the fluid particles are partially slipping on a solid boundary (see Navier [8], Goldstein [4], Serrin [10], Sect. 64, and references therein). We note that if $K \equiv 1$, then condition (1.5) reduces the well-investigated adherence one.

The aim of the present paper is to prove the existence of a unique solution $(\bar{v}(x), \bar{p}(x))$ to the stationary problem

$$\begin{cases} (\bar{v} \cdot \nabla)\bar{v} - \nabla \cdot \mathbf{P}(\bar{v}, \bar{p}) = \bar{f}, & \nabla \cdot \bar{v} = 0 \quad \text{in } \Omega, \\ \bar{v} \cdot n = 0, \quad 2(1 - \bar{K})\Pi\mathbf{D}(\bar{v})n - \bar{K}\bar{v} = 0 & \text{on } \Gamma \end{cases} \quad (1.6)$$

and to study its stability with respect to the corresponding nonstationary problem (1.1)-(1.2), (1.5). In §§3 – 4, we prove the following existence theorem to stationary problem (1.6). Throughout Theorems 1.1-1.4 we always assume that the boundary Γ belongs to $W_2^{\frac{5}{2}+l}$ ($\frac{1}{2} < l < 1$) (as for function spaces, see §2).

Throughout this paper we denote by c_i ($i = 1, 2, 3, \dots$) positive constants that will be used later. Otherwise we do not distinguish them and use the same symbol c .

Theorem 1.1 *Let $\bar{K} \in W_2^{\frac{1}{2}+l}(\Gamma)$ ($0 \leq \bar{K} \leq 1$). If $\bar{f} \in W_2^l(\Omega)$ satisfies condition (4.6), then there exists a unique solution $(\bar{v}, \nabla \bar{p}) \in (W_2^{2+l}(\Omega) \cap H(\Omega)) \times W_2^l(\Omega)$ of (1.6) satisfying the inequality*

$$\|\bar{v}\|_{W_2^{2+l}(\Omega)} + \|\nabla \bar{p}\|_{W_2^l(\Omega)} \leq c \|\bar{f}\|_{W_2^l(\Omega)}. \quad (1.7)$$

The proof of Theorem 1.1 depends on the investigation of the linearized problem for (1.6) and the contraction mapping principle. In studying the linearized problem we follow the general theory of elliptic boundary value problems developed by [1, 11, 16]. But our problem is not included completely in the framework of known theory [1, 11, 16], because in boundary condition $2(1-K)\Pi\mathbf{D}(v)n - Kv = 0$ we must regard both terms $2(1-K)\Pi\mathbf{D}(v)n$ and Kv are principal, since $0 \leq K \leq 1$. To overcome this difficulty we make some devices. The most important one is a partition of unity of a domain which was originally introduced by Tani [15] for the study of time-dependent compressible Navier-Stokes equations under condition (1.5), and later used by the authors [5, 14] for incompressible case. We follow this idea with some natural modifications for stationary case (see §3.4).

Now let us turn to nonstationary problem (1.1)-(1.2), (1.5). First of all existence of a temporally local solution was established in [14].

Theorem 1.2 ([14]) *Suppose $K \in W_2^{\frac{1}{2}+l, \frac{1}{4}+\frac{l}{2}}(\Gamma_\infty)$, $f \in W_2^{l, \frac{l}{2}}(Q_\infty)$ ($Q_\infty = \Omega \times (0, \infty)$), $v_0 \in W_2^{1+l}(\Omega)$ and v_0 satisfies the compatibility conditions. Then problem (1.1)-(1.2), (1.5) has a unique solution $(v, \nabla p) \in W_2^{2+l, 1+\frac{l}{2}}(Q_{T_1}) \times W_2^{l, \frac{l}{2}}(Q_{T_1})$ for some $T_1 \in (0, \infty)$ such that*

$$\|v\|_{W_2^{2+l, 1+\frac{l}{2}}(Q_{T_1})} + \|\nabla p\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \leq c_1 (\|v_0\|_{W_2^{1+l}(\Omega)} + \|f\|_{W_2^{l, \frac{l}{2}}(Q_\infty)}) \equiv c_1 E. \quad (1.8)$$

Moreover, the number T_1 increases unboundedly as E tends to zero.

Theorem 1.2 and the a priori estimates proved in §5 yield the following temporally global existence theorem.

Theorem 1.3 *In addition to the assumptions of Theorem 1.2, suppose $f \in W_2^{2l, l}(Q_\infty)$. If $E_0 = \|v_0\|_{W_2^{1+l}(\Omega)} + \|f\|_{W_2^{2l, l}(Q_\infty)} < \varepsilon_0$ with a sufficiently small positive number ε_0 , then the solution $(v, \nabla p)$ of Theorem 1.2 exists for all $t > 0$ and satisfies the inequality*

$$\sup_{t \geq t_1 > 0} (\|v(t)\|_{W_2^{2+l}(\Omega)} + \|\nabla p(t)\|_{W_2^l(\Omega)}) \leq c E_0 \quad (1.9)$$

for each $t_1 > 0$.

Finally stability of stationary solution from Theorem 1.1 is discussed in §6.

Theorem 1.4 *Let $(v(x, t), p(x, t))$ and $(\bar{v}(x), \bar{p}(x))$ be the solution of nonstationary problem (1.1)-(1.2), (1.5) from Theorem 1.3 and of stationary problem (1.6) from Theorem 1.1, respectively. Suppose that $\frac{K - \bar{K}}{(1 - K)(1 - \bar{K})} \in L_2(\hat{\Gamma}(t))$ ($\hat{\Gamma}(t) = \{x \in \Gamma \mid K(x, t) \neq 1, \bar{K}(x) \neq 1\}$). Then the difference $u = v - \bar{v}$ obeys the inequality*

$$\|u(t)\|_{L_2(\Omega)}^2 \leq e^{-Mt} (\|u(0)\|_{L_2(\Omega)}^2 + \int_0^t e^{Ms} F(s) ds) \quad (1.10)$$

for any $t > 0$, where M and $F(t)$ are defined by (6.5) and (6.6), respectively.

In conclusion let us mention some previous works about Navier-Stokes and Stokes equations under slip boundary conditions. In the case of perfect slip, *i.e.*, $K \equiv 0$, stationary problem for incompressible Stokes equations was discussed by Solonnikov-Sčadilov [13], while for compressible heat-conductive Navier-Stokes equations by Farwig [2]. Note that the problem with perfect slip condition is closely resemble to that of stationary motion with a free boundary (see *e.g.*, [9]).

On the other hand, the time-dependent problem (1.1)-(1.2), (1.5), besides our previous work [14] mentioned above, was also investigated in [5]. In [5], it was proved that the solution exists for a small time interval in Hölder class of functions and that this solution exists for all time without restriction of smallness of the data provided the space dimension is two. The existence of a temporally global solution for non-homogeneous fluid was also established in [6].

Finally the initial value problem for viscous compressible heat conducting fluid with general slip boundary condition was studied by Tani [15] in Hölder class of functions.

2 Preliminaries

2.1. Function spaces

Throughout this paper we use the Sobolev-Slobodetskiĭ spaces defined as follows. Let Ω be a domain in $\mathbb{R}^n$ and $l > 0$ be a noninteger with an integral part $[l]$ and a nonintegral part $\{l\}$. By $W_2^l(\Omega)$ we mean the space of functions

$u(x)$, $x \in \Omega$, equipped with the norm

$$\begin{aligned} \|u\|_{W_2^l(\Omega)}^2 &= \sum_{|\alpha|=0}^{[l]} \|D_x^\alpha u\|_{L_2(\Omega)}^2 + \sum_{|\alpha|=[l]} \int_{\Omega} \int_{\Omega} \frac{|D_x^\alpha u(x) - D_y^\alpha u(y)|^2}{|x-y|^{n+2\{l\}}} dx dy \\ &\equiv \|u\|_{W_2^{[l]}(\Omega)}^2 + \|u\|_{\dot{W}_2^l(\Omega)}^2, \end{aligned}$$

where

$$D_x^\alpha u(x) = \frac{\partial^{|\alpha|} u(x)}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \dots \partial x_n^{\alpha_n}}, \quad \alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$$

denotes a distributional derivative of $u(x)$ of order $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$. Now we define anisotropic spaces $W_2^{l, \frac{l}{2}}(Q_T)$ of functions $u(x, t)$ in a cylindrical domain $Q_T = \Omega \times (0, T)$ ($0 < T \leq \infty$) as $W_2^{l, \frac{l}{2}}(Q_T) = L_2(0, T; W_2^l(\Omega)) \cap L_2(\Omega; W_2^{\frac{l}{2}}(0, T))$ and introduce in this space the norm

$$\begin{aligned} \|u\|_{W_2^{l, \frac{l}{2}}(Q_T)}^2 &= \int_0^T \|u(\cdot, t)\|_{W_2^l(\Omega)}^2 dt + \int_{\Omega} \|u(x, \cdot)\|_{W_2^{\frac{l}{2}}(0, T)}^2 dx \\ &\equiv \|u\|_{W_2^{l, 0}(Q_T)}^2 + \|u\|_{W_2^{0, \frac{l}{2}}(Q_T)}^2. \end{aligned}$$

Here the norm in $W_2^{\frac{l}{2}}(0, T)$ is defined by

$$\begin{aligned} \|u(x, \cdot)\|_{W_2^{\frac{l}{2}}(0, T)}^2 &= \sum_{j=0}^{[\frac{l}{2}]} \|\partial_t^j u(x, \cdot)\|_{L_2(0, T)}^2 \\ &\quad + \int_0^T dt \int_0^t \left| \partial_t^{[\frac{l}{2}]} u(x, t) - \partial_t^{[\frac{l}{2}]} u(x, t - \tau) \right|^2 \frac{d\tau}{\tau^{1+2\{\frac{l}{2}\}}} \\ &\equiv \|u\|_{W_2^{[\frac{l}{2}]}(0, T)}^2 + \|u\|_{\dot{W}_2^{\frac{l}{2}}(0, T)}^2, \end{aligned}$$

where $\partial_t = \frac{\partial}{\partial t}$.

For a smooth manifold $\partial\Omega = \Gamma$, the space $W_2^l(\Gamma)$ of functions defined on Γ is introduced in a standard manner by means of the local coordinates

and the partition of unity, and $W_2^{l, \frac{l}{2}}(\Gamma_T)$ can be defined in the same way as above.

The spaces of vector fields whose components belong to $W_2^l(\Omega)$, $W_2^{l, \frac{l}{2}}(Q_T)$ etc. are denoted by the same notation as the scalar case $W_2^l(\Omega)$, $W_2^{l, \frac{l}{2}}(Q_T)$ etc. and their norms are supposed to be equal to the sum of norms of all its components.

2.2. Auxiliary lemmas

In this subsection, we present auxiliary inequalities which will be frequently used in later sections. Hereafter, we assume that Ω is a bounded domain in $\mathbb{R}^3$ with a boundary $\Gamma \in W_2^{\frac{5}{2}+l}$ ($l > \frac{1}{2}$).

Lemma 2.1 ([7]) *Suppose $u \in W_2^1(\Omega)$. Then $u \in L_4(\Gamma)$ and*

$$\|u\|_{L_4(\Gamma)} \leq c_2 \|u\|_{W_2^1(\Omega)}.$$

Lemma 2.2 ([3]) *Let u be a vector function in $W_2^1(\Omega)$ satisfying $u \cdot n = 0$ on Γ . Then we have*

$$\|u\|_{L_2(\Omega)} \leq c_3 \|\nabla u\|_{L_2(\Omega)}.$$

Next we state Korn's inequality discussed in [13]. For vectors $u, v \in W_2^1(\Omega)$, we introduce

$$E(u, v) = \int_{\Omega} \sum_{i,j=1}^3 \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right) \left(\frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} \right) dx.$$

We recall that the vectors satisfying $E(u, u) = 0$ form a finite-dimensional affine space of vectors of the form

$$u = A + B \times x,$$

where A and B are constant vectors. Let us define $\tilde{H}(\Omega) = \{ u \in W_2^1(\Omega) \mid E(u) \equiv E(u, u) < \infty, u \cdot n = 0 \text{ on } \Gamma \}$. If Ω is a region obtained by rotation about a vector B , we denote by $H(\Omega)$ the space of functions in $\tilde{H}(\Omega)$ satisfying the condition

$$\int_{\Omega} u \cdot (B \times x) dx = 0.$$

Otherwise we set $H(\Omega) = \tilde{H}(\Omega)$.

Lemma 2.3 ([13]) *The inequality*

$$\|\nabla u\|_{L_2(\Omega)}^2 \leq c_4 E(u)$$

is valid for each $u \in H(\Omega)$.

3 Stationary Stokes problem

In this section we consider the problem

$$\begin{cases} -\nu\Delta u + \nabla q = \bar{f}, & \nabla \cdot u = \rho \quad \text{in } \Omega, \\ 2(1 - \bar{K})\Pi\mathbf{D}(u)n - \bar{K}u = b_\tau, & u \cdot n = b_n \quad \text{on } \Gamma. \end{cases} \quad (3.1)$$

We prove

Theorem 3.1 *Let $\frac{1}{2} < l < 1$. Suppose $\Gamma \in W^{\frac{5}{2}+l}_2$, $\bar{K} \in W^{\frac{1}{2}+l}_2(\Gamma)$,*

$$\begin{cases} \bar{f} \in W^l_2(\Omega), & \rho \in W^{1+l}_2(\Omega), \\ b_\tau \in W^{\frac{1}{2}+l}_2(\Gamma) \cap W^{\frac{3}{2}+l}_2(\gamma), & \gamma = \{x \in \Gamma \mid \bar{K}(x) = 1\}, \\ b_n \in W^{\frac{3}{2}+l}_2(\Gamma) \end{cases} \quad (3.2)$$

and

$$\int_{\Omega} \rho \, dx = - \int_{\Gamma} b_n \, dS. \quad (3.3)$$

Then problem (3.1) has a unique solution $(u, \nabla q)$ such that

$$(u, \nabla q) \in V_l \equiv W^{2+l}_2(\Omega) \times W^l_2(\Omega) \quad (3.4)$$

and

$$\begin{aligned} \|(u, \nabla q)\|_{V_l} &\equiv \|u\|_{W^{2+l}_2(\Omega)} + \|\nabla q\|_{W^l_2(\Omega)} \\ &\leq c_5 \left(\|\bar{f}\|_{W^l_2(\Omega)} + \|\rho\|_{W^{1+l}_2(\Omega)} + \|b_\tau\|_{W^{\frac{1}{2}+l}_2(\Gamma)} + \|b_\tau\|_{W^{\frac{3}{2}+l}_2(\gamma)} + \|b_n\|_{W^{\frac{3}{2}+l}_2(\Gamma)} \right) \\ &\equiv c_5 \|(\bar{f}, \rho, b_\tau, b_n)\|_{H_l}. \end{aligned} \quad (3.5)$$

As usual we start with the study of a model problem.

3.1. Half-space problem for the homogeneous system

First of all, let us consider the boundary value problem for the homogeneous Stokes system in a half space $\mathbb{R}^3_+ = \{x = (x', x_3) \mid x' \in \mathbb{R}^2, x_3 > 0\}$:

$$\begin{cases} -\nu\Delta u + \nabla q = 0, & \nabla \cdot u = 0 \quad \text{in } \mathbb{R}^3_+, \\ (1 - k) \left(\frac{\partial u_3}{\partial x_j} + \frac{\partial u_j}{\partial x_3} \right) - k u_j \Big|_{x_3=0} = b_j \quad (j = 1, 2), \\ u_3|_{x_3=0} = 0 \quad \text{on } \mathbb{R}^2, \end{cases} \quad (3.6)$$

where $0 \leq k \leq 1$ is a constant and b_1, b_2 are given functions on $\mathbb{R}^2$. Applying to (3.6) the Fourier transform with respect to $x' = (x_1, x_2)$:

$$\hat{f}(\xi, x_3) = F[f] = \int_{\mathbb{R}^2} e^{-ix' \cdot \xi} f(x', x_3) dx', \quad (3.7)$$

where $\xi = (\xi_1, \xi_2)$, $x' \cdot \xi = x_1\xi_1 + x_2\xi_2$. Then we have the following system of ordinary differential equations:

$$\begin{cases} \nu \left(\xi^2 - \frac{d^2}{dx_3^2} \right) \hat{u}_j + i\xi_j \hat{q} = 0 & (j = 1, 2), \\ \nu \left(\xi^2 - \frac{d^2}{dx_3^2} \right) \hat{u}_3 + \frac{d\hat{q}}{dx_3} = 0, \\ i\xi_1 \hat{u}_1 + i\xi_2 \hat{u}_2 + \frac{d\hat{u}_3}{dx_3} = 0, & (\hat{u}, \hat{q}) \longrightarrow 0 \quad (x_3 \longrightarrow +\infty), \end{cases} \quad (3.8)$$

$$\begin{cases} (1 - k) \left(i\xi_j \hat{u}_3 + \frac{d\hat{u}_j}{dx_3} \right) - k\hat{u}_j \Big|_{x_3=0} = \hat{b}_j & (j = 1, 2), \\ \hat{u}_3|_{x_3=0} = 0. \end{cases} \quad (3.9)$$

We seek the solution to (3.8), (3.9) in the form

$$\begin{aligned} \begin{pmatrix} \hat{u}_1 \\ \hat{u}_2 \\ \hat{u}_3 \\ \hat{q} \end{pmatrix} &= a_1 \left[\begin{pmatrix} -|\xi| \\ 0 \\ -(i\xi_1 + 1) \\ -2\nu|\xi| \end{pmatrix} e^{-|\xi|x_3} + \begin{pmatrix} i\xi_1 \\ i\xi_2 \\ -|\xi| \\ 0 \end{pmatrix} x_3 e^{-|\xi|x_3} \right] \\ &+ a_2 \left[\begin{pmatrix} 0 \\ -|\xi| \\ -(i\xi_2 + 1) \\ -2\nu|\xi| \end{pmatrix} e^{-|\xi|x_3} + \begin{pmatrix} i\xi_1 \\ i\xi_2 \\ -|\xi| \\ 0 \end{pmatrix} x_3 e^{-|\xi|x_3} \right] \\ &+ a_3 \begin{pmatrix} i\xi_1 \\ i\xi_2 \\ -|\xi| \\ 0 \end{pmatrix} e^{-|\xi|x_3}. \end{aligned} \quad (3.10)$$

The coefficient (a_1, a_2, a_3) is determined by substituting (3.10) into (3.9). We have

$$\begin{pmatrix} \hat{u}_1 \\ \hat{u}_2 \\ \hat{u}_3 \\ \hat{q} \end{pmatrix} = \begin{pmatrix} G_{11} & G_{12} & 0 \\ G_{21} & G_{22} & 0 \\ 0 & 0 & 0 \\ G_{41} & G_{42} & 0 \end{pmatrix} \begin{pmatrix} \hat{b}_1 \\ \hat{b}_2 \\ 0 \end{pmatrix} e^{-|\xi|x_3} + \begin{pmatrix} H_{11} & H_{12} & 0 \\ H_{21} & H_{22} & 0 \\ H_{31} & H_{32} & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \hat{b}_1 \\ \hat{b}_2 \\ 0 \end{pmatrix} x_3 e^{-|\xi|x_3}, \quad (3.11)$$

where

$$\begin{aligned} G_{mj} &= -\frac{1}{(1-k)|\xi|+k} \left(\delta_{mj} + \frac{(1-k)}{2(1-k)|\xi|+k} \frac{(i\xi_m)(i\xi_j)}{|\xi|} \right) \quad (m, j = 1, 2), \\ G_{4j} &= 2\nu \frac{i\xi_j}{2(1-k)|\xi|+k} \quad (j = 1, 2), \\ H_{mj} &= -\frac{1}{2(1-k)|\xi|+k} \frac{(i\xi_m)(i\xi_j)}{|\xi|} \quad (m, j = 1, 2), \\ H_{3j} &= \frac{i\xi_j}{2(1-k)|\xi|+k} \quad (j = 1, 2). \end{aligned} \quad (3.12)$$

In order to estimate (3.11) in $W_2^l(\mathbb{R}_+^3)$, we make use of Parseval's equality. Indeed, we find

Lemma 3.1 *Let us introduce the norms*

$$\begin{aligned} \|u\|_{l, \mathbb{R}^2}^2 &= \int_{\mathbb{R}^2} |\hat{u}(\xi)|^2 |\xi|^{2l} d\xi, \\ \|u\|_{l, \mathbb{R}_+^3}^2 &= \sum_{k < l} \int_{\mathbb{R}^2} \left\| \left(\frac{d}{dx_3} \right)^k \hat{u}(\xi, \cdot) \right\|_{L_2(\mathbb{R}_+)}^2 |\xi|^{2(l-k)} d\xi + \int_{\mathbb{R}^2} \|\hat{u}(\xi, \cdot)\|_{\dot{W}_2^l(\mathbb{R}_+)}^2 d\xi. \end{aligned}$$

Then the norms $\|u\|_{l, \mathbb{R}^2}$ and $\|u\|_{l, \mathbb{R}_+^3}$ are equivalent to $\|u\|_{\dot{W}_2^l(\mathbb{R}^2)}$ and $\|u\|_{\dot{W}_2^l(\mathbb{R}_+^3)}$, respectively.

As for the estimates of $e^{-|\xi|x_3}$ and $x_3 e^{-|\xi|x_3}$, we obtain the following lemma by direct calculations.

Lemma 3.2 *Let $k \geq 0$ be an integer and $\alpha \in (0, 1)$. Then we have*

$$\int_0^\infty \left| \left(\frac{d}{dx_3} \right)^k e^{-|\xi|x_3} \right|^2 dx_3 \leq c|\xi|^{2k-1},$$

$$\int_0^\infty \int_0^\infty \left| \left(\frac{d}{dx_3} \right)^k e^{-|\xi|(x_3+y_3)} - \left(\frac{d}{dx_3} \right)^k e^{-|\xi|x_3} \right|^2 \frac{dx_3 dy_3}{y_3^{1+2\alpha}} \leq c|\xi|^{2(k+\alpha)-1},$$

$$\int_0^\infty \left| \left(\frac{d}{dx_3} \right)^k x_3 e^{-|\xi|x_3} \right|^2 dx_3 \leq c|\xi|^{2k-3},$$

$$\int_0^\infty \int_0^\infty \left| \left(\frac{d}{dx_3} \right)^k (x_3 + y_3) e^{-|\xi|(x_3+y_3)} - \left(\frac{d}{dx_3} \right)^k x_3 e^{-|\xi|x_3} \right|^2 \frac{dx_3 dy_3}{y_3^{1+2\alpha}} \leq c|\xi|^{2(k+\alpha)-3}$$

with a constant c independent of $|\xi|$.

From (3.12) and Lemmas 3.1 and 3.2, we obtain

Lemma 3.3 *Let $b = (b_1, b_2) = (1 - k)b^N - kb^D$ with $b^N \in \dot{W}_2^{\frac{1}{2}+l}(\mathbb{R}^2)$, $b^D \in \dot{W}_2^{\frac{3}{2}+l}(\mathbb{R}^2)$, $l \geq 0$. Then solution (3.11) to problem (3.6) satisfies the inequality*

$$\|u\|_{\dot{W}_2^{2+l}(\mathbb{R}_+^3)} + \|\nabla q\|_{\dot{W}_2^l(\mathbb{R}_+^3)} \leq c \left(\|b^N\|_{\dot{W}_2^{\frac{1}{2}+l}(\mathbb{R}^2)} + \|b^D\|_{\dot{W}_2^{\frac{3}{2}+l}(\mathbb{R}^2)} \right). \quad (3.13)$$

3.2. Non-homogeneous system

Now we consider the non-homogeneous system

$$\begin{cases} -\nu \Delta u + \nabla q = \bar{f}, & \nabla \cdot u = \rho & \text{in } \mathbb{R}_+^3, \\ (1 - k) \left(\frac{\partial u_3}{\partial x_j} + \frac{\partial u_j}{\partial x_3} \right) - k u_j \Big|_{x_3=0} = b_j & (j = 1, 2), \\ u_3|_{x_3=0} = b_3 & \text{on } \mathbb{R}^2. \end{cases} \quad (3.14)$$

We prove

Lemma 3.4 *Let $b = (b_1, b_2)$ be as in Lemma 3.3. Suppose that $\bar{f} \in \dot{W}_2^l(\mathbb{R}_+^3)$, $\rho \in \dot{W}_2^{1+l}(\mathbb{R}_+^3)$, $b_3 \in \dot{W}_2^{\frac{3}{2}+l}(\mathbb{R}^2)$ and the condition $\int_{\mathbb{R}_+^3} \rho dx = -\int_{\mathbb{R}^2} b_3 dx'$ is satisfied. Then the solution $(u, \nabla q)$ to problem (3.14) satisfies the estimate*

$$\begin{aligned} & \|u\|_{\dot{W}_2^{2+l}(\mathbb{R}_+^3)} + \|\nabla q\|_{\dot{W}_2^l(\mathbb{R}_+^3)} \\ & \leq c \left(\|\bar{f}\|_{\dot{W}_2^l(\mathbb{R}_+^3)} + \|\rho\|_{\dot{W}_2^{1+l}(\mathbb{R}_+^3)} + \|b^N\|_{\dot{W}_2^{\frac{1}{2}+l}(\mathbb{R}^2)} + \|b^D\|_{\dot{W}_2^{\frac{3}{2}+l}(\mathbb{R}^2)} + \|b_3\|_{\dot{W}_2^{\frac{3}{2}+l}(\mathbb{R}^2)} \right). \end{aligned} \quad (3.15)$$

Proof. We seek the solution of (3.14) in the form $(u, q) = (u^{(1)} + u^{(2)} + u^{(3)}, \nu\rho' + q^{(3)})$. Here $u^{(1)}$ is a solution of Dirichlet problem:

$$-\nu\Delta u^{(1)} = \bar{f} \quad \text{in } \mathbb{R}_+^3, \quad u^{(1)}|_{x_3=0} = 0 \quad \text{on } \mathbb{R}^2. \quad (3.16)$$

While $u^{(2)} = \nabla\phi$, where ϕ is a solution of Neumann problem

$$\Delta\phi = \rho - \nabla \cdot u^{(1)} \equiv \rho' \quad \text{in } \mathbb{R}_+^3, \quad \frac{\partial\phi}{\partial x_3}\bigg|_{x_3=0} = b_3 \quad \text{on } \mathbb{R}^2, \quad (3.17)$$

and $(u^{(3)}, q^{(3)})$ is a solution of problem (3.6) with $b = (b_1, b_2)$ replaced by $d = (d_1, d_2)$, where

$$\begin{aligned} d_j &= b_j - (1-k) \left(\frac{\partial}{\partial x_j} (u_3^{(1)} + u_3^{(2)}) + \frac{\partial}{\partial x_3} (u_j^{(1)} + u_j^{(2)}) \right) + k (u_j^{(1)} + u_j^{(2)}) \bigg|_{x_3=0} \\ &\equiv (1-k)d_j^N - kd_j^D \quad (j = 1, 2). \end{aligned}$$

From the known estimates to problem (3.16) and (3.17) combined with Lemma 3.3 yields (3.15). $\blacksquare$

3.3. Uniqueness of the solution

Before proving the normal solvability to problem (3.1), we discuss uniqueness of the solution.

Lemma 3.5 *The solution to problem (3.1) is unique. (Here uniqueness of q means within an additive constant.)*

Proof. Let (u, q) be a solution of (3.1) with $\bar{f} = \rho = b_\tau = b_n = 0$. Then we have

$$\begin{aligned} 0 &= - \int_{\Omega} (\nabla \cdot \mathbf{P}(u, q)) \cdot u \, dx = \int_{\Gamma} \mathbf{P}(u, q)n \cdot u \, dS + \frac{\nu}{2} E(u) \\ &= \int_{\Gamma} 2\nu \mathbf{D}(u)n \cdot u \, dS + \frac{\nu}{2} E(u) \\ &= \nu \int_{\Gamma^*} \frac{\bar{K}}{1 - \bar{K}} |u|^2 \, dS + \frac{\nu}{2} E(u), \end{aligned}$$

where $\Gamma^* = \{x \in \Gamma \mid \bar{K}(x) \neq 1\}$. Therefore the boundary condition $u \cdot n = 0$ implies $u \equiv 0$. $\blacksquare$

3.4. Proof of Theorem 3.1

Here and in what follows we simply write $\|\cdot\|_{L_2(\Omega)}$ as $\|\cdot\|$.

We decompose a solution (3.1) in a similar manner as in §3.2. Namely $(u, q) = (u^{(1)} + u^{(2)} + v, \nu\rho' + p)$, where $u^{(1)}, u^{(2)} = \nabla\phi$ and (v, p) satisfy the following equations, respectively:

$$-\nu\Delta u^{(1)} = \bar{f} \quad \text{in } \Omega, \quad u^{(1)} = 0 \quad \text{on } \Gamma, \quad (3.18)$$

$$-\nu\Delta\phi = \rho - \nabla \cdot u^{(1)} \equiv \rho' \quad \text{in } \Omega, \quad \frac{\partial\phi}{\partial n} = b_n \quad \text{on } \Gamma, \quad (3.19)$$

$$\begin{cases} -\nu\Delta v + \nabla p = 0, & \nabla \cdot v = 0 \quad \text{in } \Omega, \\ 2(1 - \bar{K})\Pi\mathbf{D}(v)n - \bar{K}v \\ \quad = b_\tau - 2(1 - \bar{K})\Pi\mathbf{D}(u^{(1)} + \nabla\phi) + \bar{K}(u^{(1)} + \nabla\phi)|_\Gamma \equiv d_\tau, \\ v \cdot n = 0 \quad \text{on } \Gamma. \end{cases} \quad (3.20)$$

Since problems (3.18) and (3.19) are well investigated, we have only to consider problem (3.20). The solvability of (3.20) will be proved by the method of the regularizer (cf., [11, 16]), which necessitates to introduce two systems of covering $\{\omega^{(k)}\}$ and $\{\Omega^{(k)}\}$ of $\bar{\Omega}$. As was mentioned in introduction, we make some devices for $\{\omega^{(k)}\}$ and $\{\Omega^{(k)}\}$ because of boundary condition $2(1 - \bar{K})\Pi\mathbf{D}(v)n - \bar{K}v = d_\tau$ ($0 \leq \bar{K} \leq 1$).

For arbitrary small positive number λ , $\{\omega^{(k)}\}$ and $\{\Omega^{(k)}\}$ are constructed as follows :

For $k = k'$ satisfying $\omega^{(k')} \cap \Gamma = \emptyset$, $\{\omega^{(k')}\}$ and $\{\Omega^{(k')}\}$ are the cubes with the same center and with the length of their edges, in a parallel direction with axes, equal to $\lambda/2$ and λ , respectively.

For $k = k''$ such that $\xi^{(k'')} \in \Gamma - \gamma$ ($\gamma = \{x \in \Gamma \mid \bar{K}(x) = 1\}$), we define by the local rectangular coordinate system $\{y\}$:

$$\begin{aligned} \omega^{(k'')} &= \Pi_x^y \{ |y_j| \leq \frac{1}{2} \beta_1 \lambda \quad (j = 1, 2), \quad 0 \leq y_3 - F(y'; \xi^{(k'')}) \leq \beta_1 \lambda \}, \\ \Omega^{(k'')} &= \Pi_x^y \{ |y_j| \leq \beta_1 \lambda \quad (j = 1, 2), \quad 0 \leq y_3 - F(y'; \xi^{(k'')}) \leq 2\beta_1 \lambda \}. \end{aligned}$$

Here the equation $y_3 = F(y'; \xi^{(k'')})$ ($y' = (y_1, y_2)$) represents the boundary Γ in the neighborhood of the point $\xi^{(k'')}$, Π_x^y is the transformation from $\{y\}$ to $\{x\}$ and β_1 is some positive constant independent of λ . If γ is covered by $\cup_{k''}(\omega^{(k'')} \cap \Gamma)$, then it is clear that $\bar{\Omega}$ is covered by $\{\omega^{(k)}\}$ and $\{\Omega^{(k)}\}$ constructed above.

Otherwise (in this case we shall denote $k = k'''$), we define $\{\omega^{(k''')}\}$ and

$\{\Omega^{(k''')}\}$ by the same way as $\{\omega^{(k'')}\}$ and $\{\Omega^{(k'')}\}$ with another positive constant $\beta_2(\leq \beta_1)$ also independent of λ so that $\gamma - \cup_{k''}(\omega^{(k'')} \cap \Gamma) \subset \cup_{k'''}(\Omega^{(k''')} \cap \Gamma) \subset \gamma$.

Once we introduce the system of coverings as above, the rest of the proof is carried out in line with the general theory of Solonnikov [11]. Hence we only describe it briefly.

Now we consider two families of smooth functions $\{\zeta^{(k)}(x)\}$ and $\{\eta^{(k)}(x)\}$ associated with the coverings $\{\omega^{(k)}\}$ and $\{\Omega^{(k)}\}$: $\zeta^{(k)}(x) = 1$ if $x \in \omega^{(k)}(t)$, $\zeta^{(k)}(x) = 0$ if $x \in \bar{\Omega} - \Omega^{(k)}$, $0 \leq \zeta^{(k)}(x) \leq 1$, $|D_x^r \zeta^{(k)}(x)| \leq c\lambda^{-|r|}$, $\eta^{(k)}(x) \equiv \frac{\zeta^{(k)}(x)}{\sum_k (\zeta^{(k)}(x))^2}$. Obviously, $\{\eta^{(k)}(x)\}$ are smooth functions such that $\eta^{(k)}(x) = 0$ if $x \in \bar{\Omega} - \Omega^{(k)}$, $\sum_k \eta^{(k)}(x) \zeta^{(k)}(x) = 1$ and

$$|D_x^r \eta^{(k)}(x)| \leq c\lambda^{-|r|}. \quad (3.21)$$

We note that $\Gamma \in W^{\frac{5}{2}+l}$ means that $F(y'; \xi^{(k)}) \in W^{\frac{5}{2}+l}(B_r)$, $B_r \equiv \{y' \in \mathbb{R}^2 \mid |y'| < r\}$, has the properties $F(0) = 0$, $\nabla' F(0) = 0$, $\|F\|_{W^{\frac{5}{2}+l}(B_r)} \leq N$ with the constants r and N being independent of y' . We take λ small enough so that $\beta_1 \lambda \leq \frac{r}{2}$ holds. Clearly,

$$\begin{aligned} |F(y')| &= |F(y') - F(0)| \leq N|y'|, \\ |\nabla' F(y')| &= |\nabla' F(y') - \nabla' F(0)| \leq N|y'|. \end{aligned} \quad (3.22)$$

We define operator $\mathcal{R}$ by

$$\begin{aligned} \mathcal{R}h &= \sum_{k'', k'''} \eta^{(k)}(x) \Pi_x^z \mathcal{R}^{(k)} \Pi_z^x (h\zeta^{(k)})(x) \\ &= \sum_{k'', k'''} \eta^{(k)}(x) \Pi_x^z (\bar{v}^{(k)}, \nabla \bar{p}^{(k)})(z) \\ &= \sum_{k'', k'''} \eta^{(k)}(x) (v^{(k)}, \nabla p^{(k)})(x) \end{aligned}$$

where $h = (0, 0, d_\tau, 0)$, the local coordinate system $\{z\}$ connected with $\{y\}$ is given by the relation $z' = y'$, $z_3 = y_3 - F(y'; \xi^{(k'')})$, Π_x^z is the transformation from $\{z\}$ to $\{x\}$, Π_z^x is its inverse and $\mathcal{R}^{(k)} \Pi_z^x (h\zeta^{(k)})(x) = (\bar{v}^{(k)}, \nabla \bar{p}^{(k)})(z)$, $(\bar{v}^{(k)}, \nabla \bar{p}^{(k)})(z)$ is the solution of following problem in the half space $\mathbb{R}_+^{3(k)} =$

$\Pi_x^z \Omega^{(k)}$:

$$\begin{cases} -\nu \Delta \bar{v}^{(k)} + \nabla \bar{p}^{(k)} = 0, & \nabla \cdot \bar{v}^{(k)} = 0 \quad \text{in } \mathbb{R}_+^{3(k)}, \\ (1 - \bar{K}(\xi^{(k)}, 0)) \left(\frac{\partial \bar{v}_3^{(k)}}{\partial x_j} + \frac{\partial \bar{v}_j^{(k)}}{\partial x_3} \right) - \bar{K}(\xi^{(k)}, 0) \bar{v}_j^{(k)} \Big|_{z_3=0} = \Pi_x^z \zeta^{(k)} d_{\tau j} \quad (j = 1, 2), \\ \bar{v}_3^{(k)} \Big|_{z_3=0} = 0 \quad \text{on } \mathbb{R}^{2(k)} = \Pi_x^z \partial \Omega^{(k)} \cap \{z_3 = 0\}. \end{cases}$$

Then one can easily show that $\mathcal{R}h = (v', \nabla p')(x)$ satisfies

$$\begin{cases} -\nu \Delta v' + \nabla p' = \mathcal{M}_1 h, & \nabla \cdot v' = \mathcal{M}_2 h \quad \text{in } \Omega, \\ 2(1 - \bar{K}) \Pi \mathbf{D}(v') n - \bar{K} v' = d_\tau + \mathcal{M}_3 h, \\ v' \cdot n = \mathcal{M}_4 h \quad \text{on } \Gamma. \end{cases} \quad (3.23)$$

Here the operator $\mathcal{M} = (\mathcal{M}_1, \mathcal{M}_2, \mathcal{M}_3, \mathcal{M}_4)$ is defined on $H_l = \{(\bar{f}, \rho, b_\tau, b_n) \mid (\bar{f}, \rho, b_\tau, b_n) \text{ has the smoothness property in (3.2)}\}$ equipped with the norm $\|(\bar{f}, \rho, b_\tau, b_n)\|_{H_l}$ (cf.(3.5)), and is represented as follows:

$$\begin{aligned} \mathcal{M}_1 h &= \sum_{k'', k'''} (-\nu (\Delta(\eta^{(k)} v^{(k)}) - \eta^{(k)} \Delta v^{(k)}) + \nabla(\eta^{(k)} p^{(k)}) - \eta^{(k)} \nabla p^{(k)}) \\ &\quad + \sum_{k'', k'''} \eta^{(k)} \Pi_x^z (-\nu (\Delta^{(k)} - \Delta) \bar{v}^{(k)} + (\nabla^{(k)} - \nabla) \bar{p}^{(k)}) \\ &\equiv \mathcal{T}_1 h + \mathcal{K}_1 h, \\ \mathcal{M}_2 h &= \sum_{k'', k'''} (\nabla \cdot (\eta^{(k)} v^{(k)}) - \eta^{(k)} \nabla \cdot v^{(k)}) + \sum_{k'', k'''} \eta^{(k)} \Pi_x^z (\nabla^{(k)} - \nabla) \cdot \bar{v}^{(k)} \\ &\equiv \mathcal{T}_2 h + \mathcal{K}_2 h, \\ \mathcal{M}_3 h &= \sum_{k''} 2(1 - \bar{K}(x)) \Pi(\mathbf{D}(\eta^{(k)} v^{(k)}) n - \eta^{(k)} \mathbf{D}(v^{(k)}) n) \\ &\quad + \sum_{k''} \left(\eta^{(k)} (\bar{K}(\xi^{(k)}) - \bar{K}(x)) (2 \Pi \mathbf{D}(v^{(k)}) n + v^{(k)}) \right. \\ &\quad \left. + 2 \eta^{(k)} (1 - \bar{K}(\xi^{(k)})) \Pi_x^z (\Pi \mathbf{D}^{(k)}(\bar{v}^{(k)}) n - \Pi_0 \mathbf{D}(\bar{v}^{(k)}) n_0) \right) \\ &\equiv \mathcal{T}_3 h + \mathcal{K}_3 h, \\ \mathcal{M}_4 h &= \sum_{k'', k'''} \eta^{(k)} v^{(k)} \cdot (n - n_0) \equiv \mathcal{K}_4 h, \end{aligned}$$

where $\nabla^{(k)} = \Pi_z^x \nabla_x = {}^t(\frac{\partial x_i}{\partial z_j})^{-1} \nabla_z \equiv \mathbf{g} \nabla_z$, $\Delta^{(k)} = \nabla^{(k)} \cdot \nabla^{(k)}$, $\mathbf{D}^{(k)} = \Pi_z^x \mathbf{D}$, $n_0 = n(\xi^{(k)}) = (0, 0, 1)^t$ and $\Pi_0 w = w - (w \cdot n_0) n_0$.
If we denote by $\mathcal{A}$ the differential operators in the left hand side of (3.23), then we find

$$\mathcal{A} \mathcal{R} h = h + \mathcal{T} h + \mathcal{K} h, \quad (3.24)$$

where $\mathcal{T} = (\mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_3, 0)$ and $\mathcal{K} = (\mathcal{K}_1, \mathcal{K}_2, \mathcal{K}_3, \mathcal{K}_4)$.

To estimate $\|\mathcal{T} h\|_{H_l}$ and $\|\mathcal{K} h\|_{H_l}$, we introduce the norm depending on parameter λ :

$$\langle u \rangle_{l, \Omega^{(k)}}^2 = \sum_{j=0}^{[l]} \frac{1}{\lambda^{l-j}} \|u\|_{W_2^j(\Omega^{(k)})}^2 + \|u\|_{\dot{W}_2^l(\Omega^{(k)})}^2.$$

Certainly the norms $\|u\|_{W_2^l(\Omega^{(k)})}$ and $\langle u \rangle_{l, \Omega^{(k)}}$ are equivalent for each $\lambda > 0$, and the interpolation inequality implies

$$\|u\|_{W_2^m(\Omega^{(k)})} \leq c \lambda^{l-m} \langle u \rangle_{l, \Omega^{(k)}}$$

for $0 \leq m < l$. By making use of (3.21), (3.22) and the smoothness of $\bar{K}(x)$, one can show that $\mathcal{K}$ is a contraction operator on H_l for small λ , and $\mathcal{T}$ is a compact operator on H_l for each λ , since the imbedding operator from H_l into H_{l-m} is compact for a bounded Ω . Therefore, (3.24) implies that existence of the right regularizer.

Now let us consider $\mathcal{R} \mathcal{A} u$. Similar calculations yield

$$\mathcal{R} \mathcal{A} u = u + \mathcal{S} u + \mathcal{Q} u, \quad (3.25)$$

where $u = (v, \nabla p) \in V_l$, $\mathcal{S} = (\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3, 0)$, $\mathcal{Q} = (\mathcal{Q}_1, \mathcal{Q}_2, \mathcal{Q}_3, \mathcal{Q}_4)$,

$$\begin{aligned} \mathcal{S}_1 u &= \sum_{k'', k'''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} \Pi_z^x \left(-\nu \left(\zeta^{(k)} \Delta v - \Delta(\zeta^{(k)} v) \right) + \zeta^{(k)} \nabla p - \nabla(\zeta^{(k)} p) \right), \\ \mathcal{Q}_1 u &= \sum_{k'', k'''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} \left(-\nu(\Delta^{(k)} - \Delta) \bar{v}^{(k)} + (\nabla^{(k)} - \nabla) \bar{p}^{(k)} \right), \\ \mathcal{S}_2 u &= \sum_{k'', k'''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} \Pi_z^x \left(\zeta^{(k)} \nabla \cdot v - \nabla \cdot (\zeta^{(k)} v) \right), \\ \mathcal{Q}_2 u &= \sum_{k'', k'''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} (\nabla^{(k)} - \nabla) \cdot \bar{v}^{(k)}, \\ \mathcal{S}_3 u &= \sum_{k''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} \Pi_z^x 2 \left(1 - \bar{K}(x) \right) \Pi \left(\zeta^{(k)} \mathbf{D}(v) n - \mathbf{D}(\zeta^{(k)} v) n \right), \end{aligned}$$

$$\begin{aligned}
\mathcal{Q}_3 u &= \sum_{k''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} \Pi_z^x ((\bar{K}(\xi^{(k)}) - \bar{K}(x)) (2\Pi\mathbf{D}(\zeta^{(k)}v)n + \zeta^{(k)}v) \\
&\quad + \sum_{k''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} 2(1 - \bar{K}(\xi^{(k)})) (\Pi\mathbf{D}^{(k)}(\bar{v}^{(k)})n - \Pi_0\mathbf{D}(\bar{v}^{(k)})n_0) \\
\mathcal{Q}_4 u &= \sum_{k'', k'''} \eta^{(k)} \Pi_x^z \mathcal{R}^{(k)} \Pi_z^x \zeta^{(k)}v \cdot (n - n_0).
\end{aligned}$$

By exactly the same way as $\mathcal{K}$ and $\mathcal{T}$, one can show that $\mathcal{Q}$ is a contraction operator on V_l and that $\mathcal{S}$ is a compact operator on V_l , which together with (3.25) imply the existence of the left regularizer. By combining these and uniqueness of a solution from Lemma 3.5, Theorem 3.1 is proved. $\blacksquare$

4 Proof of Theorem 1.1

We solve (1.6) by the method of successive approximations. Let $(\bar{v}^{(0)}, \nabla \bar{p}^{(0)}) = (0, 0)$ and $(\bar{v}^{(m)}, \nabla \bar{p}^{(m)}) \in \bar{X}(\Omega) \equiv \{ (\bar{v}^{(m)}, \nabla \bar{p}^{(m)}) \in V_l \mid \|(\bar{v}^{(m)}, \nabla \bar{p}^{(m)})\|_{V_l} \leq 2c_5 \|\bar{f}\|_{W_2^l(\Omega)} \}$ ($m = 1, 2, 3, \dots$). We define $(\bar{v}^{(m+1)}, \nabla \bar{p}^{(m+1)})$ as a solution to the linear problem

$$\begin{cases} -\nu \Delta \bar{v}^{(m+1)} + \nabla \bar{p}^{(m+1)} = \bar{f} - (\bar{v}^{(m)} \cdot \nabla) \bar{v}^{(m)}, \\ \nabla \cdot \bar{v}^{(m+1)} = 0 \quad \text{in } \Omega, \\ \bar{v}^{(m+1)} \cdot n = 0, \quad 2(1 - \bar{K})\Pi\mathbf{D}(\bar{v}^{(m+1)})n - \bar{K}\bar{v}^{(m+1)} = 0 \quad \text{on } \Gamma. \end{cases} \quad (4.1)$$

By Theorem 3.1 problem (4.1) has a unique solution $(\bar{v}^{(m+1)}, \nabla \bar{p}^{(m+1)}) \in V_l$ satisfying

$$\begin{aligned}
\|(\bar{v}^{(m+1)}, \nabla \bar{p}^{(m+1)})\|_{V_l} &\leq c_5 (\|\bar{f}\|_{W_2^l(\Omega)} + \|(\bar{v}^{(m)} \cdot \nabla) \bar{v}^{(m)}\|_{W_2^l(\Omega)}) \\
&\leq c_5 (\|\bar{f}\|_{W_2^l(\Omega)} + c_6 \|\bar{v}^{(m)}\|_{W_2^{1+l}(\Omega)}^2) \\
&\leq c_5 (\|\bar{f}\|_{W_2^l(\Omega)} + c_6 (2c_5 \|\bar{f}\|_{W_2^l(\Omega)})^2) \\
&\leq c_5 (1 + 4c_5^2 c_6 \|\bar{f}\|_{W_2^l(\Omega)}) \|\bar{f}\|_{W_2^l(\Omega)}. \quad (4.2)
\end{aligned}$$

Hence we find $(\bar{v}^{(m+1)}, \nabla \bar{p}^{(m+1)}) \in \bar{X}(\Omega)$ provided

$$4c_5^2 c_6 \|\bar{f}\|_{W_2^l(\Omega)} \leq 1. \quad (4.3)$$

Now let us prove the convergence of the successive approximations. Subtracting from (4.1) the similar equations for $(\bar{v}^{(m)}, \nabla \bar{p}^{(m)})$ and setting $(\bar{V}^{(m+1)}, \nabla \bar{P}^{(m+1)}) = (\bar{v}^{(m+1)} - \bar{v}^{(m)}, \nabla \bar{p}^{(m+1)} - \nabla \bar{p}^{(m)})$, we obtain

$$\begin{cases} -\nu \Delta \bar{V}^{(m+1)} + \nabla \bar{P}^{(m+1)} = -(\bar{v}^{(m)} \cdot \nabla) \bar{V}^{(m)} - (\bar{V}^{(m)} \cdot \nabla) \bar{v}^{(m-1)}, \\ \nabla \cdot \bar{V}^{(m+1)} = 0 \quad \text{in } \Omega, \\ \bar{V}^{(m+1)} \cdot n = 0, \quad 2(1 - \bar{K}) \Pi \mathbf{D}(\bar{V}^{(m+1)}) n - \bar{K} \bar{V}^{(m+1)} = 0 \quad \text{on } \Gamma. \end{cases} \quad (4.4)$$

By virtue of Theorem 3.1, there exists a unique solution $(\bar{V}^{(m+1)}, \nabla \bar{P}^{(m+1)}) \in V_l$ of (4.4), which satisfies

$$\begin{aligned} \|(\bar{V}^{(m+1)}, \nabla \bar{P}^{(m+1)})\|_{V_l} &\leq c_5 \left(\|(\bar{v}^{(m)} \cdot \nabla) \bar{V}^{(m)}\|_{W_2^l(\Omega)} + \|(\bar{V}^{(m)} \cdot \nabla) \bar{v}^{(m-1)}\|_{W_2^l(\Omega)} \right) \\ &\leq c_5 c_6 \left(\|\bar{v}^{(m)}\|_{W_2^{1+l}(\Omega)} + \|\bar{v}^{(m-1)}\|_{W_2^{1+l}(\Omega)} \right) \|\bar{V}^{(m)}\|_{W_2^{1+l}(\Omega)} \\ &\leq 4c_5^2 c_6 \|\bar{f}\|_{W_2^l(\Omega)} \|(\bar{V}^{(m)}, \nabla \bar{P}^{(m)})\|_{V_l}. \end{aligned} \quad (4.5)$$

Therefore if we assume

$$4c_5^2 c_6 \|\bar{f}\|_{W_2^l(\Omega)} < 1, \quad (4.6)$$

then we see that the sequence $(\bar{v}^{(m)}, \nabla \bar{p}^{(m)})$ converges to some $(\bar{v}, \nabla \bar{p}) \in \bar{X}(\Omega)$ as $m \rightarrow \infty$, which is our desired solution to (1.6). The uniqueness of the solution follows from the estimate similar to (4.5). $\blacksquare$

5 Proof of Theorem 1.3

We begin with the conservation of energy for (1.1)-(1.2), (1.5).

Lemma 5.1 *The estimate*

$$\|v(t)\| \leq \|v_0\| + \int_0^t \|f(s)\| ds \quad (t > 0) \quad (5.1)$$

is true for the solution $(v, \nabla p)$ to problem (1.1)-(1.2), (1.5).

Proof. Multiplying v for (1.1)₁ and integrating it over Ω , we have the equality

$$\frac{1}{2} \frac{d}{dt} \|v\|^2 + \frac{\nu}{2} E(v) = - \int_{\Gamma} \mathbf{P}(v, p) n \cdot v \, dS + \int_{\Omega} f \cdot v \, dx.$$

Similarly in the proof of Lemma 3.5, we find

$$\frac{1}{2} \frac{d}{dt} \|v\|^2 + \frac{\nu}{2} E(v) + \nu \int_{\Gamma^*(t)} \frac{K}{1-K} |v|^2 dS = \int_{\Omega} f \cdot v dx \leq \|f\| \|v\|, \quad (5.2)$$

where $\Gamma^*(t) = \{x \in \Gamma \mid K(x, t) \neq 1\}$. ■

For the estimates of higher derivatives of the solution, we follow Solonnikov [12].

Lemma 5.2 *Let the solution from Theorem 1.2 satisfy the condition*

$$\|v\|_{W_2^{2+l, 1+\frac{l}{2}}(Q_{T_1})} \leq \delta \quad (5.3)$$

with a sufficiently small number $\delta > 0$. Then

$$U(\lambda) = \|v\|_{W_2^{2+l, 1+\frac{l}{2}}(Q(\lambda))} + \|\nabla p\|_{W_2^{l, \frac{l}{2}}(Q(\lambda))} \leq c(\lambda^{-\frac{l}{2}} \|v\|_{L_2(Q(0))} + \|f\|_{W_2^{l, \frac{l}{2}}(Q_{\infty})}), \quad (5.4)$$

where $\lambda \in (0, 1)$, $Q(\lambda) = \Omega \times (2t_0 + \lambda, T_1)$, $t_0 > 0$, $2t_0 + \lambda < T_1$. Furthermore,

$$\sup_{t \in (t_1, T_1)} (\|v(t)\|_{W_2^{2+l}(\Omega)} + \|\nabla p(t)\|_{W_2^l(\Omega)}) \leq c(\|v\|_{L_2(Q(0))} + \|f\|_{W_2^{2l, l}(Q_{\infty})}) \quad (5.5)$$

is valid for each $t_1 \in (2t_0, T_1)$.

Proof. Let $\zeta_{\lambda}(t)$ be a smooth function of $t \in \mathbb{R}$ which vanishes for $t \leq t_0 + \frac{\lambda}{2}$, equals to 1 for $t \geq t_0 + \lambda$ and satisfies $0 \leq \zeta_{\lambda}(t) \leq 1$, $|\zeta_{\lambda}^{(k)}(t)| \leq c\lambda^{-k}$. Then it is easily seen that $(v_{\lambda}, \nabla p_{\lambda}) = (\zeta_{\lambda}v, \zeta_{\lambda}\nabla p)$ satisfies the equations

$$\begin{cases} \frac{\partial v_{\lambda}}{\partial t} - \nabla \cdot \mathbf{P}(v_{\lambda}, p_{\lambda}) = \zeta_{\lambda}f + \zeta_{\lambda}(v \cdot \nabla)v - \zeta'_{\lambda}v, & \nabla \cdot v_{\lambda} = 0 \quad \text{in } Q(\lambda), \\ v_{\lambda}|_{t=0} = 0 & \text{on } \Omega, \\ v_{\lambda} \cdot n = 0, \quad 2(1-K)\Pi\mathbf{D}(v_{\lambda})n - Kv_{\lambda} = 0 & \text{on } \Gamma(\lambda) = \Gamma \times (2t_0 + \lambda, T_1). \end{cases} \quad (5.6)$$

Applying Theorem 3.4 in [14], we obtain

$$\begin{aligned} & \|v_{\lambda}\|_{W_2^{2+l, 1+\frac{l}{2}}(Q_{T_1})} + \|\nabla p_{\lambda}\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\ & \leq c(\|\zeta_{\lambda}f\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} + \|\zeta_{\lambda}(v \cdot \nabla)v\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} + \|\zeta'_{\lambda}v\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})}) \\ & \leq c(\|f\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} + \delta\|v_{\lambda}\|_{W_2^{1+l, \frac{1}{2}+\frac{l}{2}}(Q_{T_1})} + \lambda^{-1}\|v\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})}). \end{aligned}$$

Therefore,

$$U(\lambda) \leq c(\|f\|_{W_2^{l, \frac{l}{2}}(Q_\infty)} + \lambda^{-1}\|v\|_{W_2^{l, \frac{l}{2}}(Q(\frac{\lambda}{2}))}). \quad (5.7)$$

Using the interpolation inequality (see [12]), we find

$$c\lambda^{-1}\|v\|_{W_2^{l, \frac{l}{2}}(Q(\frac{\lambda}{2}))} \leq \varepsilon_1\|v\|_{W_2^{2+l, 1+\frac{l}{2}}(Q(\frac{\lambda}{2}))} + \left(\frac{\varepsilon_1\lambda}{c}\right)^{-\frac{l}{2}}\|v\|_{L_2(Q(0))}, \quad (5.8)$$

where $\varepsilon_1 > 0$ is an arbitrary small number. Substituting (5.8) into (5.7) leads to

$$U(\lambda) \leq \varepsilon_1 U\left(\frac{\lambda}{2}\right) + \left(\frac{\varepsilon_1\lambda}{c}\right)^{-\frac{l}{2}}\|v\|_{L_2(Q(0))} + c\|f\|_{W_2^{l, \frac{l}{2}}(Q_\infty)}. \quad (5.9)$$

If $\varepsilon_1 > 0$ is so small that $\frac{\varepsilon_1}{2^{\frac{l}{2}}} < 1$, then using (5.9) recursively, we get (5.4).

Next consider the difference $(v^{(s)}(x, t), \nabla p^{(s)}(x, t)) = (v_\lambda(x, t) - v_\lambda(x, t-s), \nabla p_\lambda(x, t) - \nabla p_\lambda(x, t-s))$, $0 < s < t_0$. Subtracting from (5.6) the similar equations for $(v_\lambda(x, t-s), \nabla p_\lambda(x, t-s))$, we obtain the system of equations for $(v^{(s)}(x, t), \nabla p^{(s)}(x, t))$. We apply to it Theorem 3.4 in [14] once again. Then the following inequality holds :

$$\begin{aligned} & \|v^{(s)}\|_{W_2^{2+l, 1+\frac{l}{2}}(Q_{T_1})} + \|\nabla p^{(s)}\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\ & \leq c(\|(\zeta_\lambda(v \cdot \nabla)v)(t) - (\zeta_\lambda(v \cdot \nabla)v)(t-s)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\ & \quad + \|(\zeta'_\lambda v)(t) - (\zeta'_\lambda v)(t-s)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} + \|(\zeta_\lambda f)(t) - (\zeta_\lambda f)(t-s)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})}). \end{aligned} \quad (5.10)$$

We calculate the right hand side of (5.10), for example, as follows.

$$\begin{aligned} & \|(\zeta_\lambda(v \cdot \nabla)v)(t) - (\zeta_\lambda(v \cdot \nabla)v)(t-s)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\ & \leq \|(v^{(s)} \cdot \nabla)v(t)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} + \|(v(t-s) \cdot \nabla)v^{(s)}\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\ & \quad + \|(\zeta_\lambda(t-s) - \zeta_\lambda(t))(v(t-s) \cdot \nabla)v(t)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\ & \leq c(\delta\|v^{(s)}\|_{W_2^{1+l, \frac{1}{2}+\frac{l}{2}}(Q_{T_1})} + \lambda^{-1}\delta s\|v\|_{W_2^{1+l, \frac{1}{2}+\frac{l}{2}}(Q(\lambda))}), \end{aligned}$$

$$\begin{aligned}
& \|(\zeta'_\lambda v)(t) - (\zeta'_\lambda v)(t-s)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\
& \leq \|(\zeta'_\lambda(t) - \zeta'_\lambda(t-s))v(t)\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} + \|\zeta'_\lambda(t-s)(v(t) - v(t-s))\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \\
& \leq c(\lambda^{-1} + \lambda^{-2}) s \|v\|_{W_2^{2+l, 1+\frac{l}{2}}(Q(\lambda))}.
\end{aligned}$$

Therefore we conclude from (5.10) that

$$\|v^{(s)}\|_{W_2^{2+l, 1+\frac{l}{2}}(Q_{T_1})} + \|\nabla p^{(s)}\|_{W_2^{l, \frac{l}{2}}(Q_{T_1})} \leq c(\|v\|_{L_2(Q(0))} + \|f\|_{W_2^{2l, l}(Q_\infty)}) s^l. \quad (5.11)$$

Since $l > \frac{1}{2}$, (5.11) together with well-known imbedding theorem yields (5.5). $\blacksquare$

Proof of Theorem 1.3. Let ε_0 be a number such that if $E_0 < \varepsilon_0$, then solution $(v, \nabla p)$ from Theorem 1.2 exists on $[0, 1]$. This solution satisfies

$$\|v\|_{W_2^{2+l, 1+\frac{l}{2}}(Q_1)} + \|\nabla p\|_{W_2^{l, \frac{l}{2}}(Q_1)} \leq c_1 E_0,$$

where $Q_1 = \Omega \times (0, 1)$. Then condition (5.3) is true provided $c_1 E_0 < \delta$. On the other hand

$$\sup_{t \in (t_1, 1)} \|v(\cdot, t)\|_{W_2^{2+l}(\Omega)} \leq c_7 E_0$$

holds by virtue of Lemmas 5.1 and 5.2. Therefore assuming $c_7 E_0 < \varepsilon_0$, we have

$$\|v(\cdot, 1)\|_{W_2^{2+l}(\Omega)} \leq \varepsilon_0,$$

which implies that Theorem 1.2 is applicable for the initial time $t = 1$. This means that the solution exists on $[1, 2]$. Hence if $E_0 < \min\{\varepsilon_0, \frac{\delta}{c_1}, \frac{\varepsilon_0}{c_7}\}$, then we can repeat above argument infinitely many times. This completes the proof. $\blacksquare$

6 Proof of Theorem 1.4

It is easy to see that $(u, \nabla q) = (v - \bar{v}, \nabla p - \nabla \bar{p})$ satisfies the equations

$$\begin{cases} \frac{\partial u}{\partial t} + (u \cdot \nabla)u - \nabla \cdot \mathbf{P}(u, q) = -((\bar{v} \cdot \nabla)u + (u \cdot \nabla)\bar{v}) + (f - \bar{f}), \\ \nabla \cdot u = 0 \quad x \in \Omega, \quad t > 0, \\ u|_{t=0} = v_0(x) - \bar{v}(x) \equiv u_0(x) \quad x \in \Omega, \\ u \cdot n = 0, \\ 2(1 - \bar{K})\Pi\mathbf{D}(u)n - \bar{K}u = (K - \bar{K})(2\Pi\mathbf{D}(v)n + v) \quad x \in \Gamma, \quad t > 0. \end{cases} \quad (6.1)$$

Multiplying the first equation of (6.1) by u and integrating it over Ω , we find

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u\|^2 + \frac{\nu}{2} E(u) \\ &= - \int_{\Gamma} \mathbf{P}(u, q) n \cdot u \, dS - \int_{\Omega} (u \cdot \nabla) \bar{v} \cdot u \, dx + \int_{\Omega} (f - \bar{f}) \cdot u \, dx, \end{aligned} \quad (6.2)$$

from which

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u\|^2 + \frac{\nu}{2} E(u) + \nu \int_{\Gamma^*} \frac{\bar{K}}{1 - \bar{K}} |u|^2 \, dS \\ &= -\nu \int_{\hat{\Gamma}(t)} \frac{K - \bar{K}}{(1 - K)(1 - \bar{K})} u \cdot v \, dS - \int_{\Omega} (u \cdot \nabla) \bar{v} \cdot u \, dx + \int_{\Omega} (f - \bar{f}) \cdot u \, dx \\ &\leq \nu \left\| \frac{K - \bar{K}}{(1 - K)(1 - \bar{K})} \right\|_{L_2(\hat{\Gamma}(t))} \|u\|_{L_4(\Gamma)} \|v\|_{L_4(\Gamma)} + \|\nabla \bar{v}\|_{L_{\infty}(\Omega)} \|u\|^2 + \|f - \bar{f}\| \|u\| \\ &\leq \varepsilon \nu E(u) + \|\nabla \bar{v}\|_{L_{\infty}(\Omega)} \|u\|^2 + c_8 \left(\left\| \frac{K - \bar{K}}{(1 - \bar{K})(1 - K)} \right\|_{L_2(\hat{\Gamma}(t))}^2 \|v\|_{W_2^1(\Omega)}^2 + \|f - \bar{f}\|^2 \right), \end{aligned} \quad (6.3)$$

where $0 < \varepsilon < \frac{1}{2}$. Inequality (6.3) can be written as

$$\frac{d}{dt} \|u\|^2 + M \|u\|^2 \leq F(t) \quad (6.4)$$

with

$$M = c_3^{-2} c_4^{-1} \nu (1 - 2\varepsilon) - 2 \|\nabla \bar{v}\|_{L_{\infty}(\Omega)}, \quad (6.5)$$

$$F(t) = 2c_8 \left(\left\| \frac{K - \bar{K}}{(1 - \bar{K})(1 - K)} \right\|_{L_2(\hat{\Gamma}(t))}^2 \|v\|_{W_2^1(\Omega)}^2 + \|f - \bar{f}\|^2 \right). \quad (6.6)$$

From (6.4) we can conclude (1.10).

References

- [1] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions I, II. *Comm. Pure Appl. Math.*, **12** (1959), 623–727; *ibid* **17** (1964), 35–92.
- [2] R. Farwig, Stationary solutions of compressible Navier-Stokes equations with slip boundary condition. *Comm. Partial Differential Equations*, **14** (1989), 1579–1606.
- [3] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Vol. I, Linearized Steady Problems*, Springer, Heidelberg, New York, (1994).
- [4] S. Goldstein, ed., *Modern Developments in Fluid Dynamics*, Vol. 2, Dover, New York, (1965).
- [5] S. Itoh, N. Tanaka and A. Tani, The initial value problem for the Navier-Stokes equations with general slip boundary condition in Hölder spaces. *J. Math. Fluid Mech.*, **5** (2003), 275–301.
- [6] S. Itoh and A. Tani, The initial value problem for the non-homogeneous Navier-Stokes equations with general slip boundary condition. *Proc. Roy. Soc. Edinburgh Sect. A*, **130** (2000), 827–835.
- [7] O. A. Ladyzhenskaya, *The Mathematical Theory of Viscous Incompressible Flow*. Second English edition, Gordon and Breach, 1969.
- [8] C. L. M. H. Navier, M émoire sur les lois du mouvement des fluides. *M ém. Acad. Sci. Inst. France*, **6** (1823), 389–440.
- [9] V. V. Pukhnachov, Plane steady-state problem with a free boundary for the Navier-Stokes equations. *Zh. Prikl. Mech. Techn. Fiz.*, **13**, (1972), 91–102.
- [10] J. Serrin, Mathematical principles of classical fluid mechanics, in *Handbuch der Physik*, VIII/1, Springer, 1959, 125–263.
- [11] V. A. Solonnikov, General boundary value problems for systems elliptic in the sence of A. Douglis and L. Nirenberg I, II. *Amer. Math. Soc. Transl.*,(2) **56** (1966), 193–232; *Proc. Steklov Inst. Math.*, **92** (1968), 269–339.

- [12] V. A. Solonnikov, Unsteady motion of a finite mass of fluid, bounded by a free surface. *J. Soviet Math.*, **40** (1988), 672–686.
- [13] V. A. Solonnikov and V. E. Ščadilov, On a boundary value problem for a stationary system of Navier-Stokes equations. *Proc. Steklov Inst. Math.*, **125** (1973), 196–210.
- [14] A. Tani, S. Itoh and N. Tanaka, The initial value problem for the Navier-Stokes equations with general slip boundary condition. *Adv. Math. Sci. Appl.*, **4** (1994), 51–69.
- [15] A. Tani, The initial value problem for the equations of motion of general fluid with general slip boundary condition. *Kôkyûroku, RIMS, Kyoto Univ.*, **734** (1990), 123–142.
- [16] L. R. Volevic, Solvability of boundary value problem for general elliptic systems, *Amer. Math. Soc. Transl.*,(2) **67** (1968), 182–225.