



Title	Transformation relations of matrix functions associated to the hypergeometric function of Gauss under modular transformations
Author(s)	Watanabe, Humihiko
Citation	Hokkaido University Preprint Series in Mathematics, 672, 1-11
Issue Date	2004
DOI	10.14943/83823
Doc URL	http://hdl.handle.net/2115/69477
Type	bulletin (article)
File Information	pre672.pdf



[Instructions for use](#)

Transformation relations of matrix functions associated to the hypergeometric function of Gauss under modular transformations

Humihiko Watanabe

Dedicated to Professor Hiroshi Umemura on his sixtieth birthday.

§0 Introduction.

In this paper we consider 2×2 matrix functions analytic on the upper half plane associated to the hypergeometric function of Gauss, and establish transformations of these matrix functions under some modular transformations. The matrix functions studied here are obtained as the lifts of the local solutions of the matrix hypergeometric differential equation of SL type (i.e., whose image of monodromy representation is contained in $SL(2, \mathbf{C})$) at $0, 1, \infty$ to the upper half plane by the lambda function (§2). Each component of the matrix functions is represented by a definite integral with a power product of theta functions as integrand. Such an integral was invented by Wirtinger in order to uniformize the hypergeometric function of Gauss to the upper half plane ([5]). In this paper we call it *Wirtinger integral* (cf. (1.2)). In spite of many possibilities of application of the Wirtinger integral, there seems to be very few examples of application of the Wirtinger integral in literature. One of the advantages of exploiting the matrix functions above in the study of the hypergeometric function is that the monodromy property and the connection relations of the hypergeometric function are all translated as transformations of those matrix functions under modular transformations of the independent variable (§3). Moreover we can derive such transformations by exploiting classical formulas of theta functions without need to use any monodromy property or connection formula of the hypergeometric function. That is to say, this gives another new derivation of the monodromy property and the connection formulas of the hypergeometric function of Gauss.

Acknowledgement. We would like to thank Taro Horie, Naruo Kanou, Hiroshi Sakata, Kenji Iohara, Norio Suzuki for illuminating discussion. This work was supported by JSPS. KAKENHI (15740081).

§1 Wirtinger integral for the hypergeometric function of Gauss.

Following the notation of Chandrasekharan [1], we introduce the four theta functions $\theta(u, \tau), \theta_i(u, \tau)$ ($i = 1, 2, 3$) by

$$\theta(u, \tau) = \frac{1}{i} \sum_{n=-\infty}^{+\infty} (-1)^n e^{(n+\frac{1}{2})^2 \pi i \tau} e^{(2n+1)\pi i u}, \quad \theta_1(u, \tau) = \sum_{n=-\infty}^{+\infty} e^{(n+\frac{1}{2})^2 \pi i \tau} e^{(2n+1)\pi i u}, \quad \theta_2(u, \tau) = \sum_{n=-\infty}^{+\infty} (-1)^n e^{n^2 \pi i \tau} e^{2n\pi i u}, \quad \theta_3(u, \tau) = \sum_{n=-\infty}^{+\infty} e^{n^2 \pi i \tau} e^{2n\pi i u},$$

which are defined for all $(u, \tau) \in \mathbf{C} \times H$, where H denotes the upper half plane. Mumford [2] (see also Umemura [3]) adopts the symbols $\theta_{00}, \theta_{01}, \theta_{10}, \theta_{11}$ to denote the theta functions above. The relations between the two notations are as follows: $\theta(u, \tau) = -\theta_{11}(u, \tau)$, $\theta_1(u, \tau) = \theta_{10}(u, \tau)$, $\theta_2(u, \tau) = \theta_{01}(u, \tau)$, $\theta_3(u, \tau) = \theta_{00}(u, \tau)$.

The lambda function $\lambda(\tau)$ is defined by $\lambda(\tau) = \frac{\theta_1(0, \tau)^4}{\theta_3(0, \tau)^4}$. It defines a mapping of H to the open set $\mathbf{P}^1 - \{0, 1, \infty\}$ of \mathbf{P}^1 the complex projective line, and is

invariant under the action of $\Gamma(2)$ the principal congruence subgroup of level 2: $\lambda\left(\frac{a\tau + b}{c\tau + d}\right) = \lambda(\tau)$ for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(2)$. So the mapping defined by $\lambda(\tau)$ induces an isomorphism of $H/\Gamma(2)$ onto $\mathbf{P}^1 - \{0, 1, \infty\}$. We can choose the set $C = \{\tau \in H \mid -1 \leq \operatorname{Re} \tau < 1, |\tau + \frac{1}{2}| \geq \frac{1}{2}, |\tau - \frac{1}{2}| > \frac{1}{2}\}$ as a fundamental domain of H for the group $\Gamma(2)$. By the behaviour of $\lambda(\tau)$ near the cusps, the points $\tau = 0, \pm 1, \infty$ correspond to the points $x = 1, \infty, 0$ of \mathbf{P}^1 , respectively. Moreover, by the mapping defined by $\lambda(\tau)$, the positive imaginary axis of H maps to the real open interval $(0, 1)$ of \mathbf{P}^1 , each of the upper semi-circles of H centered at $\tau = \pm \frac{1}{2}$ with radius $\frac{1}{2}$ maps to the real ray $(1, +\infty)$ of \mathbf{P}^1 , and each of the rays $(-1, i\infty)$, $(1, i\infty)$ of H parallel to the positive imaginary axis maps to the real ray $(-\infty, 0)$ of \mathbf{P}^1 .

Let $F(\alpha, \beta, \gamma, x)$ denote the hypergeometric series of Gauss or its analytic continuation, and let $E(\alpha, \beta, \gamma)$ denote the hypergeometric differential equation of Gauss: $x(1-x)\frac{d^2y}{dx^2} + \{\gamma - (\alpha + \beta + 1)x\}\frac{dy}{dx} - \alpha\beta y = 0$. The following formula is well-known (e.g. see [4]):

$$(1.1) \quad F(\alpha, \beta, \gamma, x) = \frac{\Gamma(\gamma)}{(1 - e^{2\pi i \alpha})(1 - e^{2\pi i(\gamma - \alpha)})\Gamma(\alpha)\Gamma(\gamma - \alpha)} \int^{(1+, 0+, 1-, 0-)} t^{\alpha-1}(1-t)^{\gamma-\alpha-1}(1-xt)^{-\beta} dt,$$

where $\alpha \neq 1, 2, 3, \dots, \gamma \neq 0, -1, -2, \dots$, and $\gamma - \alpha \neq 1, 2, 3, \dots$. We use the function $\lambda(\tau)$ to lift the function $F(\alpha, \beta, \gamma, x)$ or, strictly speaking, the analytic continuation of $F(\alpha, \beta, \gamma, x)$ to the upper half plane H . Namely, substituting $x = \lambda(\tau)$ and $t = \operatorname{sn}^2 u$ into (1.1), we have, after some calculation,

$$(1.2) \quad F(\alpha, \beta, \gamma, \lambda(\tau)) = \frac{2\pi\Gamma(\gamma)\theta_3(0, \tau)^2}{(1 - e^{2\pi i \alpha})(1 - e^{2\pi i(\gamma - \alpha)})\Gamma(\alpha)\Gamma(\gamma - \alpha)} \lambda(\tau)^{\frac{1-\gamma}{2}} (1 - \lambda(\tau))^{\frac{\gamma-\alpha-\beta}{2}} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau)^{2\alpha-1} \theta_1(u, \tau)^{2\gamma-2\alpha-1} \theta_2(u, \tau)^{2\beta-2\gamma+1} \theta_3(u, \tau)^{-2\beta+1} du.$$

We call the definite integral in (1.2) *Wirtinger integral* for the hypergeometric function of Gauss (see [5]). Note that the function $F(\alpha, \beta, \gamma, \lambda(\tau))$ is single-valued in the variable τ , holomorphic on H .

§2 Hypergeometric functions of matrix form and their lifts to the upper half plane.

Let $Y = Y(x)$ be a 2×2 matrix-valued analytic function of the complex variable x , and let $A(x)$ denote the matrix-valued function given by

$$A(x) = \frac{1}{(\alpha - \beta)x} \begin{bmatrix} \alpha(\beta - \gamma + 1) & \alpha(\gamma - \beta - 1) \\ \beta(\alpha - \gamma + 1) & \beta(\gamma - \alpha - 1) \end{bmatrix} + \frac{1}{(\alpha - \beta)(x - 1)} \begin{bmatrix} \alpha(\gamma - \alpha - 1) & \alpha(\beta - \gamma + 1) \\ \beta(\gamma - \alpha - 1) & \beta(\beta - \gamma + 1) \end{bmatrix},$$

where α, β, γ denote complex parameters. Let us consider the following differential equation of 2×2 matrix form:

$$(2.1) \quad \frac{d}{dx}Y = A(x)Y.$$

This is a hypergeometric differential equation of matrix form. In fact, if we set $Y = \begin{bmatrix} y_{11}(x) & y_{12}(x) \\ y_{21}(x) & y_{22}(x) \end{bmatrix}$, we see that the functions $y_{11}(x)$ and $y_{12}(x)$ satisfy the equation $E(\alpha, \beta + 1, \gamma)$, and the functions $y_{21}(x)$ and $y_{22}(x)$ satisfy the equation $E(\alpha + 1, \beta, \gamma)$. In what follows, we always assume that the parameters α, β, γ satisfy the conditions

$$(2.2) \quad \alpha \notin \mathbf{Z}, \beta \notin \mathbf{Z}, \gamma \notin \mathbf{Z}, \gamma - \alpha \notin \mathbf{Z}, \gamma - \beta \notin \mathbf{Z}, \gamma - \alpha - \beta \notin \mathbf{Z}, \text{ and } \alpha - \beta \notin \mathbf{Z}.$$

Let $Y_0(x), Y_1(x), Y_\infty(x)$ be the local solutions of (2.1) at $x = 0, 1, \infty$, respectively, given by

$$\begin{aligned} Y_0(x) &= \begin{bmatrix} F(\alpha, \beta + 1, \gamma, x) & \alpha(\beta - \gamma + 1)x^{1-\gamma}F(1 + \alpha - \gamma, 2 + \beta - \gamma, 2 - \gamma, x) \\ F(\alpha + 1, \beta, \gamma, x) & \beta(\alpha - \gamma + 1)x^{1-\gamma}F(2 + \alpha - \gamma, 1 + \beta - \gamma, 2 - \gamma, x) \end{bmatrix}, \\ Y_1(x) &= \begin{bmatrix} (\beta - \gamma + 1)F(\alpha, \beta + 1, \alpha + \beta - \gamma + 2, 1 - x) & \alpha(1 - x)^{\gamma - \alpha - \beta - 1}F(\gamma - \alpha, \gamma - \beta - 1, \gamma - \alpha - \beta, 1 - x) \\ (\alpha - \gamma + 1)F(\alpha + 1, \beta, \alpha + \beta - \gamma + 2, 1 - x) & \beta(1 - x)^{\gamma - \alpha - \beta - 1}F(\gamma - \alpha - 1, \gamma - \beta, \gamma - \alpha - \beta, 1 - x) \end{bmatrix}, \\ Y_\infty(x) &= \begin{bmatrix} (\alpha - \beta)(\alpha - \beta + 1)(1 - x)^{-\alpha}F(\alpha, \gamma - \beta - 1, \alpha - \beta, (1 - x)^{-1}) & \alpha(\gamma - \beta - 1)(1 - x)^{-\beta - 1}F(\gamma - \alpha, \beta + 1, \beta - \alpha + 2, (1 - x)^{-1}) \\ \beta(\gamma - \alpha - 1)(1 - x)^{-\alpha - 1}F(\gamma - \beta, \alpha + 1, \alpha - \beta + 2, (1 - x)^{-1}) & (\beta - \alpha)(\beta - \alpha + 1)(1 - x)^{-\beta}F(\beta, \gamma - \alpha - 1, \beta - \alpha, (1 - x)^{-1}) \end{bmatrix}. \end{aligned}$$

The image of the monodromy representation of (2.1) is contained in the general linear group $GL(2, \mathbf{C})$, but not in the special linear group $SL(2, \mathbf{C})$. To obtain from (2.1) a matrix differential equation whose image of the monodromy representation is contained in $SL(2, \mathbf{C})$, we introduce a new 2×2 matrix unknown \tilde{Y} by

$$(2.3) \quad Y = x^{\frac{1-\gamma}{2}}(1-x)^{\frac{\gamma-\alpha-\beta-1}{2}}\tilde{Y}.$$

If we eliminate Y from (2.1) and (2.3), we have a new differential equation

$$(2.4) \quad \frac{d}{dx}\tilde{Y} = \tilde{A}(x)\tilde{Y},$$

where $\tilde{A}(x)$ is given by

$$\tilde{A}(x) = \frac{1}{(\alpha - \beta)x} \begin{bmatrix} \alpha\beta - \frac{(\alpha + \beta)(\gamma - 1)}{2} & \alpha(\gamma - \beta - 1) \\ \beta(\alpha - \gamma + 1) & -\alpha\beta + \frac{(\alpha + \beta)(\gamma - 1)}{2} \end{bmatrix} + \frac{1}{(\alpha - \beta)(x - 1)} \begin{bmatrix} \alpha\beta - \frac{(\alpha + \beta)(\alpha + \beta - \gamma + 1)}{2} & \alpha(\beta - \gamma + 1) \\ \beta(\gamma - \alpha - 1) & -\alpha\beta + \frac{(\alpha + \beta)(\alpha + \beta - \gamma + 1)}{2} \end{bmatrix}.$$

From $Y_0(x), Y_1(x), Y_\infty(x)$, we can obtain via (2.3) the local solutions $\tilde{Y}_0(x), \tilde{Y}_1(x), \tilde{Y}_\infty(x)$ of Equation (2.4) at $x = 0, 1, \infty$, respectively. In fact, we have

$$\begin{aligned}\tilde{Y}_0(x) &= \begin{bmatrix} x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\alpha+\beta-\gamma+1}{2}}F(\alpha, \beta+1, \gamma, x) & \alpha(\beta-\gamma+1)x^{\frac{1-\gamma}{2}}(1-x)^{\frac{\alpha+\beta-\gamma+1}{2}}F(1+\alpha-\gamma, 2+\beta-\gamma, 2-\gamma, x) \\ x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\alpha+\beta-\gamma+1}{2}}F(\alpha+1, \beta, \gamma, x) & \beta(\alpha-\gamma+1)x^{\frac{1-\gamma}{2}}(1-x)^{\frac{\alpha+\beta-\gamma+1}{2}}F(2+\alpha-\gamma, 1+\beta-\gamma, 2-\gamma, x) \end{bmatrix}, \\ \tilde{Y}_1(x) &= \begin{bmatrix} (\beta-\gamma+1)x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\alpha+\beta-\gamma+1}{2}}F(\alpha, \beta+1, \alpha+\beta-\gamma+2, 1-x) & \alpha x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\gamma-\alpha-\beta-1}{2}}F(\gamma-\alpha, \gamma-\beta-1, \gamma-\alpha-\beta, 1-x) \\ (\alpha-\gamma+1)x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\alpha+\beta-\gamma+1}{2}}F(\alpha+1, \beta, \alpha+\beta-\gamma+2, 1-x) & \beta x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\gamma-\alpha-\beta-1}{2}}F(\gamma-\alpha-1, \gamma-\beta, \gamma-\alpha-\beta, 1-x) \end{bmatrix}, \\ \tilde{Y}_\infty(x) &= \begin{bmatrix} (\alpha-\beta)(\alpha-\beta+1)x^{\frac{\gamma-1}{2}}(1-x)^{\frac{-\alpha+\beta-\gamma+1}{2}}F(\alpha, \gamma-\beta-1, \alpha-\beta, (1-x)^{-1}) & \alpha(\gamma-\beta-1)x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\alpha-\beta-\gamma-1}{2}}F(\gamma-\alpha, \beta+1, \beta-\alpha+2, (1-x)^{-1}) \\ \beta(\gamma-\alpha-1)x^{\frac{\gamma-1}{2}}(1-x)^{\frac{-\alpha+\beta-\gamma-1}{2}}F(\gamma-\beta, \alpha+1, \alpha-\beta+2, (1-x)^{-1}) & (\beta-\alpha)(\beta-\alpha+1)x^{\frac{\gamma-1}{2}}(1-x)^{\frac{\alpha-\beta-\gamma+1}{2}}F(\beta, \gamma-\alpha-1, \beta-\alpha, (1-x)^{-1}) \end{bmatrix}.\end{aligned}$$

It is easy to see that the local monodromy matrix of each function $\tilde{Y}_i(x)$ ($i = 0, 1, \infty$) has determinant one.

Let us make the lifts of the functions $\tilde{Y}_0(x), \tilde{Y}_1(x), \tilde{Y}_\infty(x)$ to the upper half plane H , using the Wirtinger integral (1.2). Namely, we set $\tau' = -1/\tau$, $\tau'' = 1/(1-\tau)$, $Z_0(\tau) = \tilde{Y}_0(\lambda(\tau))$, $Z_1(\tau') = \tilde{Y}_1(\lambda(\tau'))$ and $Z_\infty(\tau'') = \tilde{Y}_\infty(\lambda(\tau''))$. Applying (1.2) to each component of $\tilde{Y}_i(\lambda(\tau))$ ($i = 0, 1, \infty$), we have

$$\begin{aligned}Z_0(\tau) &= \begin{bmatrix} \frac{2\pi\Gamma(\gamma)\theta_3(0, \tau)^2}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\gamma-\alpha)})\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau)^{2\alpha-1}\theta_1(u, \tau)^{2\gamma-2\alpha-1}\theta_2(u, \tau)^{2\beta-2\gamma+3}\theta_3(u, \tau)^{-2\beta-1} du \\ \frac{2\pi\Gamma(\gamma)\theta_3(0, \tau)^2}{(1-e^{2\pi i\beta})(1-e^{2\pi i(\gamma-\beta)})\Gamma(\beta)\Gamma(\gamma-\beta)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau)^{2\beta-1}\theta_1(u, \tau)^{2\gamma-2\beta-1}\theta_2(u, \tau)^{2\alpha-2\gamma+3}\theta_3(u, \tau)^{-2\alpha-1} du \\ \frac{2\pi\alpha\Gamma(2-\gamma)\theta_3(0, \tau)^2}{(1-e^{2\pi i(\beta-\gamma)})(1-e^{-2\pi i\beta})\Gamma(-\beta)\Gamma(1+\beta-\gamma)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau)^{2\beta-2\gamma+3}\theta_1(u, \tau)^{-2\beta-1}\theta_2(u, \tau)^{2\alpha-1}\theta_3(u, \tau)^{2\gamma-2\alpha-1} du \\ \frac{2\pi\beta\Gamma(2-\gamma)\theta_3(0, \tau)^2}{(1-e^{2\pi i(\alpha-\gamma)})(1-e^{-2\pi i\alpha})\Gamma(-\alpha)\Gamma(1+\alpha-\gamma)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau)^{2\alpha-2\gamma+3}\theta_1(u, \tau)^{-2\alpha-1}\theta_2(u, \tau)^{2\beta-1}\theta_3(u, \tau)^{2\gamma-2\beta-1} du \end{bmatrix}, \\ Z_1(\tau') &= \begin{bmatrix} \frac{2\pi\Gamma(\alpha+\beta-\gamma+2)\theta_3(0, \tau')^2}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\beta-\gamma)})\Gamma(\alpha)\Gamma(\beta-\gamma+1)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau')^{2\alpha-1}\theta_1(u, \tau')^{2\beta-2\gamma+3}\theta_2(u, \tau')^{2\gamma-2\alpha-1}\theta_3(u, \tau')^{-2\beta-1} du \\ \frac{2\pi\Gamma(\alpha+\beta-\gamma+2)\theta_3(0, \tau')^2}{(1-e^{2\pi i\beta})(1-e^{2\pi i(\alpha-\gamma)})\Gamma(\beta)\Gamma(\alpha-\gamma+1)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau')^{2\beta-1}\theta_1(u, \tau')^{2\alpha-2\gamma+3}\theta_2(u, \tau')^{2\gamma-2\beta-1}\theta_3(u, \tau')^{-2\alpha-1} du \\ \frac{2\pi\alpha\Gamma(\gamma-\alpha-\beta)\theta_3(0, \tau')^2}{(1-e^{2\pi i(\gamma-\alpha)})(1-e^{-2\pi i\beta})\Gamma(-\beta)\Gamma(\gamma-\alpha)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau')^{2\gamma-2\alpha-1}\theta_1(u, \tau')^{-2\beta-1}\theta_2(u, \tau')^{2\alpha-1}\theta_3(u, \tau')^{2\beta-2\gamma+3} du \\ \frac{2\pi\beta\Gamma(\gamma-\alpha-\beta)\theta_3(0, \tau')^2}{(1-e^{2\pi i(\gamma-\beta)})(1-e^{-2\pi i\alpha})\Gamma(-\alpha)\Gamma(\gamma-\beta)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau')^{2\gamma-2\beta-1}\theta_1(u, \tau')^{-2\alpha-1}\theta_2(u, \tau')^{2\beta-1}\theta_3(u, \tau')^{2\alpha-2\gamma+3} du \end{bmatrix},\end{aligned}$$

$$Z_\infty(\tau'') = \left[\begin{array}{l} \frac{2\pi e^{\frac{1}{2}\pi i(\gamma-1)}\Gamma(\alpha-\beta+2)\theta_1(0,\tau'')^2}{(1-e^{2\pi i\alpha})(1-e^{-2\pi i\beta})\Gamma(\alpha)\Gamma(-\beta)} \int^{(\frac{1}{2}+,0+, \frac{1}{2}-,0-)} \theta(u,\tau'')^{2\alpha-1}\theta_1(u,\tau'')^{-2\beta-1}\theta_2(u,\tau'')^{2\gamma-2\alpha-1}\theta_3(u,\tau'')^{2\beta-2\gamma+3} du \\ \frac{-2\pi\beta e^{\frac{1}{2}\pi i(\gamma-1)}\Gamma(\alpha-\beta+2)\theta_1(0,\tau'')^2}{(1-e^{2\pi i(\gamma-\beta)})(1-e^{2\pi i(\alpha-\gamma)})\Gamma(\gamma-\beta)\Gamma(\alpha-\gamma+1)} \int^{(\frac{1}{2}+,0+, \frac{1}{2}-,0-)} \theta(u,\tau'')^{2\gamma-2\beta-1}\theta_1(u,\tau'')^{2\alpha-2\gamma+3}\theta_2(u,\tau'')^{2\beta-1}\theta_3(u,\tau'')^{-2\alpha-1} du \\ \frac{-2\pi\alpha e^{\frac{1}{2}\pi i(\gamma-1)}\Gamma(\beta-\alpha+2)\theta_1(0,\tau'')^2}{(1-e^{2\pi i(\beta-\gamma)})(1-e^{2\pi i(\gamma-\alpha)})\Gamma(\gamma-\alpha)\Gamma(\beta-\gamma+1)} \int^{(\frac{1}{2}+,0+, \frac{1}{2}-,0-)} \theta(u,\tau'')^{2\gamma-2\alpha-1}\theta_1(u,\tau'')^{2\beta-2\gamma+3}\theta_2(u,\tau'')^{2\alpha-1}\theta_3(u,\tau'')^{-2\beta-1} du \\ \frac{2\pi e^{\frac{1}{2}\pi i(\gamma-1)}\Gamma(\beta-\alpha+2)\theta_1(0,\tau'')^2}{(1-e^{2\pi i\beta})(1-e^{-2\pi i\alpha})\Gamma(-\alpha)\Gamma(\beta)} \int^{(\frac{1}{2}+,0+, \frac{1}{2}-,0-)} \theta(u,\tau'')^{2\beta-1}\theta_1(u,\tau'')^{-2\alpha-1}\theta_2(u,\tau'')^{2\gamma-2\beta-1}\theta_3(u,\tau'')^{2\alpha-2\gamma+3} du \end{array} \right].$$

Note that the matrix functions $Z_0(\tau)$, $Z_1(\tau')$, $Z_\infty(\tau'')$ are single-valued in the variable τ , holomorphic on H .

§3 Transformations of $Z_0(\tau)$, $Z_1(\tau')$, $Z_\infty(\tau'')$.

The translation of the local monodromies of the matrix functions $\tilde{Y}_0(x)$, $\tilde{Y}_1(x)$, $\tilde{Y}_\infty(x)$ into $Z_0(\tau)$, $Z_1(\tau')$, $Z_\infty(\tau'')$ is as follows:

$$(3.1) \quad Z_0(\tau+2) = Z_0(\tau) \begin{bmatrix} e^{\pi i(\gamma-1)} & 0 \\ 0 & e^{\pi i(1-\gamma)} \end{bmatrix}; \quad (3.2) \quad Z_1(\tau'+2) = Z_1(\tau') \begin{bmatrix} e^{\pi i(\alpha+\beta-\gamma+1)} & 0 \\ 0 & e^{\pi i(\gamma-\alpha-\beta-1)} \end{bmatrix}; \quad (3.3) \quad Z_\infty(\tau''+2) = Z_\infty(\tau'') \begin{bmatrix} e^{\pi i(\alpha-\beta)} & 0 \\ 0 & e^{\pi i(\beta-\alpha)} \end{bmatrix}.$$

Without need to use any formula for $\tilde{Y}_0(x)$, $\tilde{Y}_1(x)$, $\tilde{Y}_\infty(x)$, we can easily verify these formulas directly by transformation rules of theta functions. The translation of the connection formulas of $\tilde{Y}_0(x)$, $\tilde{Y}_1(x)$, $\tilde{Y}_\infty(x)$ into $Z_0(\tau)$, $Z_1(\tau')$, $Z_\infty(\tau'')$ is as follows:

Theorem. (Gauss-Riemann) Assume the conditions (2.2). Then we have

$$(3.4) \quad Z_0(\tau) = Z_1(\tau') \begin{bmatrix} -\frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta-1)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} & -\frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha-\beta-1)}{\Gamma(-\alpha)\Gamma(-\beta)} \\ \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma+1)}{\Gamma(\alpha+1)\Gamma(\beta+1)} & \frac{\Gamma(2-\gamma)\Gamma(\alpha+\beta-\gamma+1)}{\Gamma(\alpha-\gamma+1)\Gamma(\beta-\gamma+1)} \end{bmatrix}; \quad (3.5) \quad Z_0(\tau) = Z_\infty(\tau'') \begin{bmatrix} \frac{\Gamma(\gamma)\Gamma(\beta-\alpha-1)}{\Gamma(\gamma-\alpha)\Gamma(\beta+1)} & \frac{\Gamma(2-\gamma)\Gamma(\beta-\alpha-1)}{\Gamma(-\alpha)\Gamma(1+\beta-\gamma)} e^{-\pi i\gamma} \\ \frac{\Gamma(\gamma)\Gamma(\alpha-\beta-1)}{\Gamma(\alpha+1)\Gamma(\gamma-\beta)} & \frac{\Gamma(2-\gamma)\Gamma(\alpha-\beta-1)}{\Gamma(1+\alpha-\gamma)\Gamma(-\beta)} e^{-\pi i\gamma} \end{bmatrix}.$$

The proof is given in the next section. Formulas (3.1)-(3.5) determine the monodromy of the hypergeometric function of Gauss completely. For example, combining (3.4) with (3.2), we have immediately the:

Corollary. *We have*

$$(3.6) \quad Z_0\left(\frac{\tau}{-2\tau+1}\right) = Z_0(\tau) \begin{bmatrix} \frac{-\cos\pi(\alpha-\beta) + e^{-\pi i\gamma} \cos\pi(\gamma-\alpha-\beta)}{2\pi i \Gamma(\gamma-1)\Gamma(\gamma)} & \frac{2\pi i \Gamma(1-\gamma)\Gamma(2-\gamma)}{\Gamma(-\alpha)\Gamma(-\beta)\Gamma(1+\alpha-\gamma)\Gamma(1+\beta-\gamma)} \\ \frac{i \sin \pi\gamma}{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} & \frac{\cos\pi(\alpha-\beta) - e^{\pi i\gamma} \cos\pi(\alpha+\beta-\gamma)}{i \sin \pi\gamma} \end{bmatrix}.$$

This is the translation of the monodromy of $\tilde{Y}_0(x)$ along a curve with base point near $x = 0$ turning around $x = 1$ in the anticlockwise direction.

§4 Proof of Theorem.

We set $Z_0(\tau) = \begin{bmatrix} z_{11}(\tau) & z_{12}(\tau) \\ z_{21}(\tau) & z_{22}(\tau) \end{bmatrix}$, $Z_1(\tau') = \begin{bmatrix} \zeta_{11}(\tau') & \zeta_{12}(\tau') \\ \zeta_{21}(\tau') & \zeta_{22}(\tau') \end{bmatrix}$, $Z_\infty(\tau'') = \begin{bmatrix} Z_{11}(\tau'') & Z_{12}(\tau'') \\ Z_{21}(\tau'') & Z_{22}(\tau'') \end{bmatrix}$. First, let us prove Formula (3.4) by exploiting transformation rules of theta functions.

Lemma 1. *We have*

$$(4.1) \quad z_{11}(\tau) = \frac{\Gamma(\gamma)\Gamma(\beta-\gamma+1)e^{-\pi i\alpha}}{\Gamma(\alpha+\beta-\gamma+2)\Gamma(\gamma-\alpha)} \zeta_{11}(\tau') + \frac{\Gamma(\gamma)\Gamma(-\beta)e^{\pi i(\gamma-\alpha)}}{\Gamma(\alpha+1)\Gamma(\gamma-\alpha-\beta)} \zeta_{12}(\tau') - \frac{\Gamma(\gamma)\Gamma(1+\beta-\gamma)\Gamma(-\beta)e^{\pi i(\gamma-\alpha-\beta)}}{\Gamma(\alpha+1)\Gamma(\gamma-\alpha)\Gamma(2-\gamma)} z_{12}(\tau).$$

Proof. Applying Jacobi transformations of theta functions to the expression of $z_{11}(\tau)$, we have

$$(4.2) \quad z_{11}(\tau) = \frac{2\pi\Gamma(\gamma)}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\gamma-\alpha)})\Gamma(\alpha)\Gamma(\gamma-\alpha)} \frac{i}{\tau} \theta_3\left(0, -\frac{1}{\tau}\right)^2 (-i)e^{\pi i\alpha} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta\left(\frac{u}{\tau}, -\frac{1}{\tau}\right)^{2\alpha-1} \theta_1\left(\frac{u}{\tau}, -\frac{1}{\tau}\right)^{2\beta-2\gamma+3} \theta_2\left(\frac{u}{\tau}, -\frac{1}{\tau}\right)^{2\gamma-2\alpha-1} \theta_3\left(\frac{u}{\tau}, -\frac{1}{\tau}\right)^{-2\beta-1} du.$$

Substituting $v = -u/\tau$ into the definite integral of (4.2), we have

$$(4.3) \quad z_{11}(\tau) = \frac{2\pi e^{-\pi i\alpha} \Gamma(\gamma) \theta_3(0, \tau')^2}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\gamma-\alpha)})\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int^{(\frac{\tau'}{2}+, 0+, \frac{\tau'}{2}-, 0-)} \theta(v, \tau')^{2\alpha-1} \theta_1(v, \tau')^{2\beta-2\gamma+3} \theta_2(v, \tau')^{2\gamma-2\alpha-1} \theta_3(v, \tau')^{-2\beta-1} dv,$$

where we chose in the integral of (4.3) the branch of $\theta(v, \tau')^{2\alpha-1}$ satisfying $\theta(-v, \tau')^{2\alpha-1} = -e^{-2\pi i\alpha}\theta(v, \tau')^{2\alpha-1}$. Meanwhile, applying Cauchy's theorem to the integration of the integrand of (4.3) along the parallelogram with vertices $0, \frac{1}{2}\tau', \frac{1}{2}(1+\tau'), \frac{1}{2}$, we have

$$\begin{aligned}
(4.4) \quad & \frac{1}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\gamma-\alpha)})} \int_{(\frac{\tau'}{2}+, 0+, \frac{\tau'}{2}-, 0-)} \theta(v, \tau')^{2\alpha-1} \theta_1(v, \tau')^{2\beta-2\gamma+3} \theta_2(v, \tau')^{2\gamma-2\alpha-1} \theta_3(v, \tau')^{-2\beta-1} dv \\
& = \frac{1}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\beta-\gamma)})} \int_{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau')^{2\alpha-1} \theta_1(v, \tau')^{2\beta-2\gamma+3} \theta_2(v, \tau')^{2\gamma-2\alpha-1} \theta_3(v, \tau')^{-2\beta-1} dv \\
& + \frac{e^{\pi i\gamma}}{(1-e^{-2\pi i\beta})(1-e^{2\pi i(\gamma-\alpha)})} \int_{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau')^{2\gamma-2\alpha-1} \theta_1(v, \tau')^{-2\beta-1} \theta_2(v, \tau')^{2\alpha-1} \theta_3(v, \tau')^{2\beta-2\gamma+3} dv \\
& - \frac{e^{2\pi i(\gamma-\beta)}}{(1-e^{-2\pi i\beta})(1-e^{2\pi i(\beta-\gamma)})} \int_{(\frac{\tau'}{2}+, 0+, \frac{\tau'}{2}-, 0-)} \theta(v, \tau')^{2\beta-2\gamma+3} \theta_1(v, \tau')^{2\alpha-1} \theta_2(v, \tau')^{-2\beta-1} \theta_3(v, \tau')^{2\gamma-2\alpha-1} dv,
\end{aligned}$$

where we chose the branch of $\theta_1(v, \tau')^{2\beta-2\gamma+3}$ satisfying $\theta_1(v + \frac{1}{2}, \tau')^{2\beta-2\gamma+3} = -e^{2\pi i(\gamma-\beta)}\theta(v, \tau')^{2\beta-2\gamma+3}$. Substituting (4.4) into (4.3), we have the desired formula (4.1), which proves Lemma 1.

Lemma 2. *We have*

$$(4.5) \quad z_{12}(\tau) = \frac{\Gamma(2-\gamma)\Gamma(\alpha+1)e^{\pi i(\gamma-\beta)}}{\Gamma(\alpha+\beta-\gamma+2)\Gamma(-\beta)} \zeta_{11}(\tau') + \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha)e^{-\pi i\beta}}{\Gamma(1+\beta-\gamma)\Gamma(\gamma-\alpha-\beta)} \zeta_{12}(\tau') - \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha)\Gamma(\alpha+1)e^{\pi i(\gamma-\alpha-\beta)}}{\Gamma(-\beta)\Gamma(1+\beta-\gamma)\Gamma(\gamma)} z_{11}(\tau).$$

The proof is similar to that of Lemma 1. We omit it.

The system of linear equations (4.1) and (4.5) is unified as the equivalent matrix equation:

$$\begin{aligned}
& (z_{11}(\tau), z_{12}(\tau)) \begin{bmatrix} 1 & \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha)\Gamma(\alpha+1)e^{\pi i(\gamma-\alpha-\beta)}}{\Gamma(1+\beta-\gamma)\Gamma(-\beta)\Gamma(\gamma)} \\ \frac{\Gamma(\gamma)\Gamma(1+\beta-\gamma)\Gamma(-\beta)e^{\pi i(\gamma-\alpha-\beta)}}{\Gamma(\alpha+1)\Gamma(\gamma-\alpha)\Gamma(2-\gamma)} & 1 \end{bmatrix} \\
& = (\zeta_{11}(\tau'), \zeta_{12}(\tau')) \begin{bmatrix} \frac{\Gamma(\gamma)\Gamma(\beta-\gamma+1)e^{-\pi i\alpha}}{\Gamma(\gamma-\alpha)\Gamma(\alpha+\beta-\gamma+2)} & \frac{\Gamma(2-\gamma)\Gamma(\alpha+1)e^{\pi i(\gamma-\beta)}}{\Gamma(\alpha+\beta-\gamma+2)\Gamma(-\beta)} \\ \frac{\Gamma(\gamma)\Gamma(-\beta)e^{\pi i(\gamma-\alpha)}}{\Gamma(\alpha+1)\Gamma(\gamma-\alpha-\beta)} & \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha)e^{-\pi i\beta}}{\Gamma(\gamma-\alpha-\beta)\Gamma(\beta-\gamma+1)} \end{bmatrix},
\end{aligned}$$

from which it follows that

$$(4.6) \quad (z_{11}(\tau), z_{12}(\tau)) = (\zeta_{11}(\tau'), \zeta_{12}(\tau')) \left[\begin{array}{c} \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta-1)}{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma+1)} \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha-\beta-1)}{\Gamma(2-\gamma)\Gamma(\alpha+\beta-\gamma+1)} \\ \frac{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)}{\Gamma(\alpha+1)\Gamma(\beta+1)} \frac{\Gamma(-\alpha)\Gamma(-\beta)}{\Gamma(\alpha-\gamma+1)\Gamma(\beta-\gamma+1)} \end{array} \right].$$

Exchanging the variables α and β in (4.6), we have immediately

$$(4.7) \quad (z_{21}(\tau), z_{22}(\tau)) = (\zeta_{21}(\tau'), \zeta_{22}(\tau')) \left[\begin{array}{c} \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta-1)}{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma+1)} \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha-\beta-1)}{\Gamma(2-\gamma)\Gamma(\alpha+\beta-\gamma+1)} \\ \frac{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)}{\Gamma(\alpha+1)\Gamma(\beta+1)} \frac{\Gamma(-\alpha)\Gamma(-\beta)}{\Gamma(\alpha-\gamma+1)\Gamma(\beta-\gamma+1)} \end{array} \right].$$

The system of equations (4.6) and (4.7) is equivalent to the matrix equality (3.4), which proves the first half of the theorem.

Next, let us prove Formula (3.5).

Lemma 3. *We have*

$$(4.8) \quad z_{11}(\tau) = \frac{\Gamma(\gamma)\Gamma(\beta-\alpha-1)}{\Gamma(1+\beta)\Gamma(\gamma-\alpha)} Z_{11}(\tau'') + \frac{\Gamma(\gamma)\Gamma(\alpha-\beta-1)}{\Gamma(\alpha+1)\Gamma(\gamma-\beta)} Z_{12}(\tau'').$$

Proof. By transformation rules of theta functions, we have

$$(4.9) \quad z_{11}(\tau) = \frac{2\pi e^{\frac{1}{2}\pi i(\gamma-1)}\Gamma(\gamma)\theta_2(0, \tau-1)^2}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\gamma-\alpha)})\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(u, \tau-1)^{2\alpha-1} \theta_1(u, \tau-1)^{2\gamma-2\alpha-1} \theta_2(u, \tau-1)^{-2\beta-1} \theta_3(u, \tau-1)^{2\beta-2\gamma+3} du.$$

Applying Jacobi transformation formulas to (4.9), we have

$$(4.10) \quad z_{11}(\tau) = \frac{2\pi e^{\frac{1}{2}\pi i(\gamma-1)} e^{\frac{1}{2}\pi i(2\alpha-1)} \Gamma(\gamma)}{(1-e^{2\pi i\alpha})(1-e^{2\pi i(\gamma-\alpha)})\Gamma(\alpha)\Gamma(\gamma-\alpha)} \frac{i}{\tau-1} \theta_1\left(0, \frac{1}{1-\tau}\right)^2 \times \\ \times \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta\left(\frac{u}{\tau-1}, \frac{1}{1-\tau}\right)^{2\alpha-1} \theta_1\left(\frac{u}{\tau-1}, \frac{1}{1-\tau}\right)^{-2\beta-1} \theta_2\left(\frac{u}{\tau-1}, \frac{1}{1-\tau}\right)^{2\gamma-2\alpha-1} \theta_3\left(\frac{u}{\tau-1}, \frac{1}{1-\tau}\right)^{2\beta-2\gamma+3} du.$$

Substituting $v = u/(1 - \tau)$ into the integral of (4.10), we have

$$(4.11) \quad z_{11}(\tau) = \frac{2\pi e^{\frac{1}{2}\pi i(\gamma-1)} e^{-\pi i\alpha} \Gamma(\gamma) \theta_1(0, \tau'')^2}{(1 - e^{2\pi i\alpha})(1 - e^{2\pi i(\gamma-\alpha)}) \Gamma(\alpha) \Gamma(\gamma - \alpha)} \int_{(\frac{\tau''}{2}+, 0+, \frac{\tau''}{2}-, 0-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv,$$

where we chose in the integral of (4.11) the branch of $\theta(v, \tau'')^{2\alpha-1}$ satisfying $\theta(-v, \tau'')^{2\alpha-1} = -e^{-2\pi i\alpha} \theta(v, \tau'')^{2\alpha-1}$. Meanwhile, applying Cauchy's theorem to the integration of the integrand of (4.11) along the parallelogram with vertices $0, \frac{1}{2}\tau'', \frac{1}{2}(1 + \tau''), \frac{1}{2}$, we have

$$(4.12) \quad \begin{aligned} & \frac{1}{(1 - e^{2\pi i\alpha})(1 - e^{2\pi i(\gamma-\alpha)})} \int_{(0+, \frac{\tau''}{2}+, 0-, \frac{\tau''}{2}-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv \\ & + \frac{1}{(1 - e^{2\pi i\alpha})(1 - e^{-2\pi i\beta})} \int_{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv \\ & - \frac{e^{2\pi i\beta}}{(1 - e^{-2\pi i\beta})(1 - e^{2\pi i(\beta-\gamma)})} \int_{(\frac{\tau''}{2}+, 0+, \frac{\tau''}{2}-, 0-)} \theta(v, \tau'')^{-2\beta-1} \theta_1(v, \tau'')^{2\alpha-1} \theta_2(v, \tau'')^{2\beta-2\gamma+3} \theta_3(v, \tau'')^{2\gamma-2\alpha-1} dv \\ & - \frac{e^{\pi i\gamma}}{(1 - e^{2\pi i(\gamma-\alpha)})(1 - e^{2\pi i(\beta-\gamma)})} \int_{(0+, \frac{1}{2}+, 0-, \frac{1}{2}-)} \theta(v, \tau'')^{2\gamma-2\alpha-1} \theta_1(v, \tau'')^{2\beta-2\gamma+3} \theta_2(v, \tau'')^{2\alpha-1} \theta_3(v, \tau'')^{-2\beta-1} dv = 0, \end{aligned}$$

where we chose the branch of $\theta_1(v, \tau'')^{-2\beta-1}$ satisfying $\theta_1(v + \frac{1}{2}, \tau'')^{-2\beta-1} = -e^{2\pi i\beta} \theta_1(v, \tau'')^{-2\beta-1}$. Moreover, applying Cauchy's theorem to the integration of the function $\theta(v, \tau'')^{-2\beta-1} \theta_1(v, \tau'')^{2\alpha-1} \theta_2(v, \tau'')^{2\beta-2\gamma+3} \theta_3(v, \tau'')^{2\gamma-2\alpha-1}$ along the same parallelogram, we have

$$(4.13) \quad \begin{aligned} & \frac{1}{(1 - e^{-2\pi i\beta})(1 - e^{2\pi i(\beta-\gamma)})} \int_{(0+, \frac{\tau''}{2}+, 0-, \frac{\tau''}{2}-)} \theta(v, \tau'')^{-2\beta-1} \theta_1(v, \tau'')^{2\alpha-1} \theta_2(v, \tau'')^{2\beta-2\gamma+3} \theta_3(v, \tau'')^{2\gamma-2\alpha-1} dv \\ & + \frac{1}{(1 - e^{2\pi i\alpha})(1 - e^{-2\pi i\beta})} \int_{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau'')^{-2\beta-1} \theta_1(v, \tau'')^{2\alpha-1} \theta_2(v, \tau'')^{2\beta-2\gamma+3} \theta_3(v, \tau'')^{2\gamma-2\alpha-1} dv \\ & - \frac{e^{-2\pi i\alpha}}{(1 - e^{2\pi i\alpha})(1 - e^{2\pi i(\gamma-\alpha)})} \int_{(\frac{\tau''}{2}+, 0+, \frac{\tau''}{2}-, 0-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv \\ & - \frac{e^{-\pi i\gamma}}{(1 - e^{2\pi i(\gamma-\alpha)})(1 - e^{2\pi i(\beta-\gamma)})} \int_{(0+, \frac{1}{2}+, 0-, \frac{1}{2}-)} \theta(v, \tau'')^{2\beta-2\gamma+3} \theta_1(v, \tau'')^{2\gamma-2\alpha-1} \theta_2(v, \tau'')^{-2\beta-1} \theta_3(v, \tau'')^{2\alpha-1} dv = 0, \end{aligned}$$

where we chose the branch of $\theta_1(v, \tau'')^{2\alpha-1}$ satisfying $\theta_1(v + \frac{1}{2}, \tau'')^{2\alpha-1} = -e^{-2\pi i\alpha}\theta(v, \tau'')^{2\alpha-1}$. From (4.12) and (4.13) it follows that

$$\begin{aligned}
(4.14) \quad & \frac{1 - e^{2\pi i(\beta-\alpha)}}{(1 - e^{2\pi i\alpha})(1 - e^{2\pi i(\gamma-\alpha)})} \int^{(\frac{\tau''}{2}+, 0+, \frac{\tau''}{2}-, 0-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv \\
& = \frac{1 - e^{2\pi i\beta}}{(1 - e^{2\pi i\alpha})(1 - e^{-2\pi i\beta})} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv \\
& \quad + \frac{e^{\pi i\gamma} - e^{2\pi i\beta} e^{-\pi i\gamma}}{(1 - e^{2\pi i(\gamma-\alpha)})(1 - e^{2\pi i(\beta-\gamma)})} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau'')^{2\gamma-2\alpha-1} \theta_1(v, \tau'')^{2\beta-2\gamma+3} \theta_2(v, \tau'')^{2\alpha-1} \theta_3(v, \tau'')^{-2\beta-1} dv,
\end{aligned}$$

$$\begin{aligned}
(4.15) \quad & \frac{1 - e^{2\pi i(\beta-\alpha)}}{(1 - e^{-2\pi i\beta})(1 - e^{2\pi i(\beta-\gamma)})} \int^{(\frac{\tau''}{2}+, 0+, \frac{\tau''}{2}-, 0-)} \theta(v, \tau'')^{-2\beta-1} \theta_1(v, \tau'')^{2\alpha-1} \theta_2(v, \tau'')^{2\beta-2\gamma+3} \theta_3(v, \tau'')^{2\gamma-2\alpha-1} dv \\
& = \frac{1 - e^{-2\pi i\alpha}}{(1 - e^{2\pi i\alpha})(1 - e^{-2\pi i\beta})} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau'')^{2\alpha-1} \theta_1(v, \tau'')^{-2\beta-1} \theta_2(v, \tau'')^{2\gamma-2\alpha-1} \theta_3(v, \tau'')^{2\beta-2\gamma+3} dv \\
& \quad - \frac{e^{-2\pi i\alpha} e^{\pi i\gamma} + e^{-\pi i\gamma}}{(1 - e^{2\pi i(\gamma-\alpha)})(1 - e^{2\pi i(\beta-\gamma)})} \int^{(\frac{1}{2}+, 0+, \frac{1}{2}-, 0-)} \theta(v, \tau'')^{2\gamma-2\alpha-1} \theta_1(v, \tau'')^{2\beta-2\gamma+3} \theta_2(v, \tau'')^{2\alpha-1} \theta_3(v, \tau'')^{-2\beta-1} dv.
\end{aligned}$$

Substituting (4.14) into (4.11), we have the desired formula (4.8), which proves Lemma 3.

Lemma 4. *We have*

$$(4.16) \quad z_{12}(\tau) = \frac{\Gamma(2-\gamma)\Gamma(\beta-\alpha-1)e^{-\pi i\gamma}}{\Gamma(1+\beta-\gamma)\Gamma(-\alpha)} Z_{11}(\tau'') + \frac{\Gamma(2-\gamma)\Gamma(\alpha-\beta-1)e^{-\pi i\gamma}}{\Gamma(-\beta)\Gamma(\alpha-\gamma+1)} Z_{12}(\tau'').$$

The proof is similar to that of Lemma 3. We omit it.

Exchanging α and β in (4.8) and (4.16), we have immediately

$$(4.17) \quad z_{21}(\tau) = \frac{\Gamma(\gamma)\Gamma(\beta-\alpha-1)}{\Gamma(1+\beta)\Gamma(\gamma-\alpha)} Z_{21}(\tau'') + \frac{\Gamma(\gamma)\Gamma(\alpha-\beta-1)}{\Gamma(\alpha+1)\Gamma(\gamma-\beta)} Z_{22}(\tau''),$$

$$(4.18) \quad z_{22}(\tau) = \frac{\Gamma(2-\gamma)\Gamma(\beta-\alpha-1)e^{-\pi i\gamma}}{\Gamma(1+\beta-\gamma)\Gamma(-\alpha)} Z_{21}(\tau'') + \frac{\Gamma(2-\gamma)\Gamma(\alpha-\beta-1)e^{-\pi i\gamma}}{\Gamma(-\beta)\Gamma(\alpha-\gamma+1)} Z_{22}(\tau'').$$

Formulas (4.8), (4.16)-(4.18) are unified to the equivalent single matrix equality (3.5), which proves the second half of the theorem. Q.E.D.

References

- [1] K.Chandrasekharan, Elliptic functions, Springer, 1985.
- [2] D.Mumford, Tata lectures on theta I, Birkhäuser, 1983.
- [3] H.Umemura, Theory of elliptic functions: Analysis of elliptic curves, Univ. of Tokyo Press, 2000 (in Japanese).
- [4] H.Watanabe, Sur la dégénérescence de quelques formules de connexion pour les fonctions hypergéométriques de Gauss, preprint, 2003
(<http://coe.math.sci.hokudai.ac.jp/literature/preprint/back/index.html.en>).
- [5] W.Wirtinger, Zur Darstellung der hypergeometrischen Function durch bestimmte Integrale, Akad. Wiss. Wien. S.-B. IIa, III (1902), 894-900.

Humihiko Watanabe,
Kitami Institute of Technology,
165, Koencho, Kitami 0908507, Hokkaido,
Japan,
e-mail: hwatanab@cs.kitami-it.ac.jp