



Title	Weighted L^p Sobolev-Lieb-Thirring inequalities
Author(s)	Tachizawa, Kazuya
Citation	Hokkaido University Preprint Series in Mathematics, 702, 1-6
Issue Date	2005
DOI	10.14943/83853
Doc URL	http://hdl.handle.net/2115/69507
Type	bulletin (article)
File Information	pre702.pdf



[Instructions for use](#)

Weighted L^p Sobolev-Lieb-Thirring inequalities

By Kazuya Tachizawa

Department of Mathematics, Faculty of Science,
Hokkaido University, Sapporo 060-0810

Abstract: We give a weighted L^p version of the Sobolev-Lieb-Thirring inequality for suborthonormal functions.

Key words: Sobolev-Lieb-Thirring inequalities; A_p -weights.

2000 Mathematics Subject Classification. Primary 26D15; Secondary 42B25.

1 Introduction

In 1976 Lieb and Thirring proved the following inequality.

Theorem 1.1 ([4]). *Let $n \in \mathbb{N}$. Then there exists a positive constant c_n such that for every family $\{\phi_i\}_{i=1}^N$ in $H^1(\mathbb{R}^n)$ which is orthonormal in $L^2(\mathbb{R}^n)$, we have*

$$(1) \quad \int_{\mathbb{R}^n} \left(\sum_{i=1}^N |\phi_i(x)|^2 \right)^{1+2/n} dx \leq c_n \sum_{i=1}^N \|\nabla \phi_i\|^2.$$

In this theorem $H^1(\mathbb{R}^n)$ denotes the Sobolev space and $\|\cdot\|$ is the norm of $L^2(\mathbb{R}^n)$. In [4] Lieb and Thirring applied this inequality to the problem of the stability of matter. Ghidaglia, Marion, and Temam proved a generalization of (1) under the suborthonormal condition on $\{\phi_i\}$, where $\{\phi_i\}_{i=1}^N$ in $L^2(\mathbb{R}^n)$ is called suborthonormal if the inequality

$$\sum_{i,j=1}^N \xi_i \bar{\xi}_j (\phi_i, \phi_j) \leq \sum_{i=1}^N |\xi_i|^2$$

holds for all $\xi_i \in \mathbb{C}, i = 1, \dots, N$, where (\cdot, \cdot) means the L^2 inner product ([2]). They applied the inequality (1) to the estimate of the dimension of attractors associated with partial differential equations. In this paper we shall give a weighted L^p version of (1) under the suborthonormal condition on $\{\phi_i\}$.

For the statement of our result we need to recall the definition of A_p -weights (c.f. [3], [5]). By a cube in \mathbb{R}^n we mean a cube which sides are parallel to coordinate axes. Let w be a non-negative, locally integrable function on \mathbb{R}^n . We say that w is an A_p -weight for $1 < p < \infty$ if there exists a positive constant C such that

$$\frac{1}{|Q|} \int_Q w(x) dx \left(\frac{1}{|Q|} \int_Q w(x)^{-1/(p-1)} dx \right)^{p-1} \leq C$$

for all cubes $Q \subset \mathbb{R}^n$. For example, $w(x) = |x|^\alpha$ is an A_p -weight when $-n < \alpha < n(p-1)$.

We say that w is an A_1 -weight if there exists a positive constant C such that

$$\frac{1}{|Q|} \int_Q w(y) dy \leq Cw(x) \quad a.e. x \in Q$$

for all cubes $Q \subset \mathbb{R}^n$. If $-n < \alpha \leq 0$, then $w(x) = |x|^\alpha$ is an A_1 -weight. Let A_p be the class of A_p -weights. The inclusion $A_p \subset A_q$ holds for $p < q$.

A nonnegative, locally integrable function w on \mathbb{R}^n is called a weight function. For a weight function w we define

$$L^p(w) = \left\{ f : \text{measurable on } \mathbb{R}^n, \int_{\mathbb{R}^n} |f(x)|^p w(x) dx < \infty \right\}.$$

The following is a conclusion of [7, Theorem 1.2] and [6, Lemma 3.2].

Theorem 1.2. *Let $n \in \mathbb{N}$, $3 \leq n$, $w \in A_2$, and $w^{-n/2} \in A_{n/2}$. Then there exists a positive constant c such that for every family $\{\phi_i\}_{i=1}^N$ in $L^2(\mathbb{R}^n)$ which is suborthonormal in $L^2(\mathbb{R}^n)$ and $|\nabla \phi_i| \in L^2(w)$, $i = 1, \dots, N$, we have*

$$\int_{\mathbb{R}^n} \left(\sum_{i=1}^N |\phi_i(x)|^2 \right)^{1+2/n} w(x) dx \leq c \sum_{i=1}^N \int_{\mathbb{R}^n} |\nabla \phi_i(x)|^2 w(x) dx,$$

where c depends only on n and w .

By using this theorem we can prove the following weighted L^p version of the Sobolev-Lieb-Thirring inequality.

Theorem 1.3. *Let $n \in \mathbb{N}$ and $3 \leq n$. Let $2n/(n+2) < p < n$, $p \neq 2$, and w be a weight function. When $p > 2$, we assume that $w^{n/(n-p)} \in A_{p(n-2)/(2(n-p))}$. When $p < 2$, we assume that $w^{n/(n-2)} \in A_1$.*

Then there exists a positive constant c such that for every family $\{\phi_i\}_{i=1}^N$ in $L^2(\mathbb{R}^n)$ which is suborthonormal in $L^2(\mathbb{R}^n)$ and $|\nabla \phi_i| \in L^p(w)$, $i = 1, \dots, N$, we have

$$\int_{\mathbb{R}^n} \left(\sum_{i=1}^N |\phi_i(x)|^2 \right)^{(1+2/n)p/2} w(x) dx \leq c \int_{\mathbb{R}^n} \left(\sum_{i=1}^N |\nabla \phi_i(x)|^2 \right)^{p/2} w(x) dx,$$

where c depends only on n, p and w .

This is a new result even in the case $w \equiv 1$. When $2 < p < n$, an example of w is given by $w(x) = |x|^\alpha$, $-n + p < \alpha < n(p-2)/2$. When $2n/(n+2) < p < 2$, an example of w is given by $w(x) = |x|^\alpha$, $-n + 2 < \alpha \leq 0$.

2 Proof of Theorem 1.3

Let M be the Hardy-Littlewood maximal operator, that is,

$$M(f)(x) = \sup_{x \in Q} \frac{1}{|Q|} \int_Q |f(y)| dy,$$

where f is a locally integrable function on \mathbb{R}^n and the supremum is taken over all cubes Q which contain x . The following proposition is proved in [3, Chapter IV] or [5, Chapter V].

Proposition 2.1.

(i) Let $1 < p < \infty$ and w be a weight function on \mathbb{R}^n . Then there exists a positive constant c such that

$$\int_{\mathbb{R}^n} M(f)^p w dx \leq c \int_{\mathbb{R}^n} |f|^p w dx$$

for all $f \in L^p(w)$ if and only if $w \in A_p$.

(ii) Let $1 < p < \infty$ and $w \in A_p$. Then there exists a $q \in (1, p)$ such that $w \in A_q$.

(iii) Let $0 < \tau < 1$ and f be a locally integrable function on \mathbb{R}^n such that $M(f)(x) < \infty$ a.e.. Then $M(f)^\tau \in A_1$.

(iv) Let $1 < p < \infty$. Then $w \in A_p$ if and only if $w^{1-p'} \in A_{p'}$, where $p^{-1} + p'^{-1} = 1$.

(v) Let $1 < p < \infty$ and $w_1, w_2 \in A_1$. Then $w_1 w_2^{1-p} \in A_p$.

Proof of Theorem 1.3

Our proof is very similar to that of the extrapolation theorem in harmonic analysis (c.f. [1, Theorem 7.8]). In our proof the integral means that over \mathbb{R}^n .

Let $2 < p < n$ and $2/p + 1/q = 1$. We remark that the assumption $w^{n/(n-p)} \in A_{p(n-2)/(2(n-p))}$ leads to $w \in A_p$ by an easy calculation. Let $u \in L^q(w)$, $u \geq 0$, and $\|u\|_{L^q(w)} = 1$. Since $w^{n/(n-p)} \in A_{p(n-2)/(2(n-p))}$, we have $w^{-2/(p-2)} \in A_{p(n-2)/(n(p-2))}$ by (iv) of Proposition 2.1. Hence there exists a γ such that $n/(n-2) < \gamma < q$ and $w^{-2/(p-2)} \in A_{p/(\gamma(p-2))}$ by (ii) of Proposition 2.1. Then we have $uw \leq M((uw)^\gamma)^{1/\gamma}$ a.e.. Because

$$w^{-2q/p} = w^{-2/(p-2)} \in A_{p/(\gamma(p-2))} = A_{q/\gamma}$$

and

$$(2) \quad \int M((uw)^\gamma)^{q/\gamma} w^{-2q/p} dx \leq c \int (uw)^q w^{-2q/p} dx = c \int u^q w dx = c$$

by (i) of Proposition 2.1, we get $M((uw)^\gamma)(x) < \infty$ a.e.. Hence $M((uw)^\gamma)^{1/\gamma} \in A_1$ by (iii) of Proposition 2.1. Let $\alpha = \frac{n}{(n-2)\gamma}$. Then $0 < \alpha < 1$ and

$$M((uw)^\gamma)^{-n/(2\gamma)} = \{M((uw)^\gamma)^\alpha\}^{1-n/2} \in A_{n/2},$$

where we used $M((uw)^\gamma)^\alpha \in A_1$ and (v) of Proposition 2.1. Let

$$\rho(x) = \sum_{i=1}^N |\phi_i(x)|^2.$$

Then we have

$$\begin{aligned} \int \rho^{1+2/n} u w dx &\leq \int \rho^{1+2/n} M((uw)^\gamma)^{1/\gamma} dx \leq c \int \left(\sum_{i=1}^N |\nabla \phi_i|^2 \right) M((uw)^\gamma)^{1/\gamma} dx \\ &\leq c \left(\int \left(\sum_{i=1}^N |\nabla \phi_i|^2 \right)^{p/2} w dx \right)^{2/p} \left(\int M((uw)^\gamma)^{q/\gamma} w^{-2q/p} dx \right)^{1/q} \\ &\leq c \left(\int \left(\sum_{i=1}^N |\nabla \phi_i|^2 \right)^{p/2} w dx \right)^{2/p} \end{aligned}$$

where we used Theorem 1.2 and (2). If we take the supremum for all $u \in L^q(w)$, $u \geq 0$, and $\|u\|_{L^q(w)} = 1$, then we get

$$\left(\int \rho^{(1+2/n)p/2} w dx \right)^{2/p} \leq c \left(\int \left(\sum_{i=1}^N |\nabla \phi_i|^2 \right)^{p/2} w dx \right)^{2/p}.$$

Next we consider the case $2n/(n+2) < p < 2$. We remark that $w \in A_1$ by the assumption $w^{n/(n-2)} \in A_1$. Let

$$f = \left(\sum_{i=1}^N |\nabla \phi_i|^2 \right)^{1/2}.$$

We can take γ such that $(2-p)n/2 < \gamma < p$. Then

$$\int M(f^\gamma)^{p/\gamma} w dx \leq c \int f^p w dx < \infty,$$

where we used $w \in A_1 \subset A_{p/\gamma}$ and (i) of Proposition 2.1. Hence we have $M(f^\gamma)(x) < \infty$ a.e. and

$$M(f^\gamma)^{(2-p)n/(2\gamma)} \in A_1$$

by (iii) of Proposition 2.1. Furthermore we have

$$M(f^\gamma)^{-(2-p)/\gamma} w \in A_2,$$

where we used

$$M(f^\gamma)^{(2-p)/\gamma} \in A_1, \quad w \in A_1,$$

and (v) of Proposition 2.1. Moreover

$$\{M(f^\gamma)^{-(2-p)/\gamma} w\}^{-n/2} = M(f^\gamma)^{(2-p)n/(2\gamma)} (w^{n/(n-2)})^{(1-n/2)} \in A_{n/2}$$

because $w^{n/(n-2)} \in A_1$. Therefore

$$\begin{aligned} \int \rho^{(1+2/n)p/2} w \, dx &= \int \rho^{(1+2/n)p/2} w M(f^\gamma)^{-(2-p)p/(2\gamma)} M(f^\gamma)^{(2-p)p/(2\gamma)} \, dx \\ &\leq \left(\int \rho^{1+2/n} M(f^\gamma)^{-(2-p)/\gamma} w \, dx \right)^{p/2} \left(\int M(f^\gamma)^{p/\gamma} w \, dx \right)^{1-p/2} \\ &\leq c \left(\int f^2 M(f^\gamma)^{-(2-p)/\gamma} w \, dx \right)^{p/2} \left(\int f^p w \, dx \right)^{1-p/2} \\ &\leq c \left(\int M(f^\gamma)^{2/\gamma} M(f^\gamma)^{-(2-p)/\gamma} w \, dx \right)^{p/2} \left(\int f^p w \, dx \right)^{1-p/2} \\ &\leq c \left(\int M(f^\gamma)^{p/\gamma} w \, dx \right)^{p/2} \left(\int f^p w \, dx \right)^{1-p/2} \leq c \int f^p w \, dx, \end{aligned}$$

where we used Theorem 1.2 in the second inequality.

Acknowledgment

The author was partly supported by the Grants-in-Aid for formation of COE and for Scientific Research, The Ministry of Education, Science, Sports and Culture, Japan.

References

- [1] Duoandikoetxea, J.: Fourier analysis. Graduate Studies in Mathematics, 29, American Mathematical Society, Providence, RI (2001).
- [2] Ghidaglia, J.-M., Marion, M. and Temam, R.: Generalization of the Sobolev-Lieb-Thirring inequalities and applications to the dimension of attractors. Differential Integral Equations, **1**, 1-21 (1988) .
- [3] García-Cuerva, J. and Rubio de Francia, J.L.: Weighted norm inequalities and related topics. North-Holland Mathematics Studies, 116, North-Holland (1985).

- [4] Lieb, E. and Thirring, W.: Inequalities for the moments of the eigenvalues of the Schrödinger hamiltonian and their relation to Sobolev inequalities. *Studies in Mathematical Physics*, Princeton University Press, 269–303 (1976).
- [5] Stein, E.M.: *Harmonic analysis : real-variable methods, orthogonality, and oscillatory integrals*. Princeton Mathematical Series, **43**, Princeton University Press (1993).
- [6] Tachizawa, K.: On the moments of negative eigenvalues of elliptic operators. *J. Fourier Anal. Appl.*, **8**, 233–244 (2002).
- [7] Tachizawa, K.: Weighted Sobolev-Lieb-Thirring inequalities. *Rev. Mat. Iberoamericana*, **21**, 67–85 (2005).