



Title	HORO-TIGHT IMMERSIONS OF S^1
Author(s)	Buosi, Marcelo; Izumiya, Shyuichi; Soares Ruas, Mria Aoarecida
Citation	Hokkaido University Preprint Series in Mathematics, 711, 1-5
Issue Date	2005
DOI	10.14943/83862
Doc URL	http://hdl.handle.net/2115/69516
Type	bulletin (article)
File Information	pre711.pdf



[Instructions for use](#)

HORO-TIGHT IMMERSIONS OF S^1

MARCELO BUOSI, SHYUICHI IZUMIYA, AND MARIA APARECIDA SOARES RUAS

ABSTRACT. We characterize horo-tight immersions into \mathbb{D}^m in terms of a family of real valued functions parametrized by S^{m-1} . By means of such functions we provide an elementary proof that horo-tightness and tightness are equivalent properties in the class of immersions from S^1 into hyperbolic space.

What are the horo-tight immersions of spheres? This question was proposed by Thomas E. Cecil and Patrick J. Ryan in [5, pg 236]. We address this paper to the simplest case of that question, namely the horo-tight immersions of a simple closed curve.

The concept of horo-tightness was introduced in [4], whose main subjects are tight and taut immersions into hyperbolic space.

One may regard tightness, tautness and horo-tightness as generalizations of Euclidean tightness. An immersion into \mathbb{R}^m is called Euclidean tight if every non-degenerate linear height function has the minimum number of critical points required by the Morse inequalities. The linear height functions in any Euclidean space are parametrized by the sphere of unit vectors. On the other hand, in a Lorentz space there are three types of spheres, each one formed entirely by spacelike, timelike or lightlike vectors. Such trichotomy gives rise to the concepts of tightness, tautness and horo-tightness respectively.

In §2 we observe that, in the Poincaré model, the radii of the hyperhorospheres tangent to its boundary at a fixed point is an interesting function. Because we can characterize horo-tightness by requiring that every non-degenerate such function has the minimum number of critical points.

The main result (Theorem 3.5), states that an immersion into the hyperbolic space of a simple closed curve is horo-tight if and only if it is tight. Then, in codimension greater than 1, tightness, tautness and horo-tightness are all equivalent concepts in the class of such immersions.

1. NOTATION AND DEFINITIONS

We begin with the definition of perfect functions.

A Morse function φ on a manifold M is called a *perfect function* if

- (1) $M_r(\varphi) = \{x \in M \mid \varphi(x) \leq r\}$ is compact for all real r ,
- (2) there exists a field \mathbf{F} such that for all $r \in \mathbb{R}$ and all integers k , the number of critical points of φ of index k which lie in $M_r(\varphi)$ is equal to the k th Betti number of $M_r(\varphi)$ over \mathbf{F} .

As observed in [4] there are manifolds which do not admit perfect functions.

Let $\mathbb{R}^{m+1} = \{(x_0, x_1, \dots, x_m) \mid x_i \in \mathbb{R}\}$ be an $(m+1)$ -dimensional vector space. For any $x = (x_0, x_1, \dots, x_m)$ and $y = (y_0, y_1, \dots, y_m)$ the *Lorentz inner product* of x and y is defined

The first author was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil, grant # 141321/00-8.

by $\mathbf{b}(x, y) = -x_0y_0 + x_1y_1 + \cdots + x_my_m$. The pair $(\mathbb{R}^{m+1}, \mathbf{b})$ is called the $(m+1)$ -Lorentz space and is denoted by \mathbb{R}_1^{m+1} . A non-zero vector $x \in \mathbb{R}_1^{m+1}$ is called spacelike, timelike or lightlike if $\mathbf{b}(x, x) > 0$, $\mathbf{b}(x, x) < 0$ or $\mathbf{b}(x, x) = 0$ respectively.

We use two models of hyperbolic space, \mathbb{H}^m and \mathbb{D}^m . The space \mathbb{H}^m , a sphere of timelike vectors, is given by

$$\mathbb{H}^m = \{x \in \mathbb{R}_1^{m+1} \mid \mathbf{b}(x, x) = -1, x_0 \geq 1\},$$

on which \mathbf{b} restricts to a Riemannian metric of constant sectional curvature -1 . Let \mathbb{D}^m be the open unit ball at the origin of $\mathbb{R}^m = \{\mathbf{0}\} \times \mathbb{R}^m$. The stereographic projection $P : \mathbb{D}^m \rightarrow \mathbb{H}^m$ given by

$$P(x) = -e_0 + 2\frac{x + e_0}{1 - \mathbf{b}(x, x)}$$

induces on \mathbb{D}^m a Riemannian metric \mathbf{g} such that $(\mathbb{D}^m, \mathbf{g})$ is isometric to \mathbb{H}^m . This is the well known Poincaré ball model.

We now define spheres of spacelike and lightlike vectors in \mathbb{R}_1^{m+1} : the *de Sitter m -space* $\mathbb{S}_1^m = \{x \in \mathbb{R}_1^{m+1} \mid \mathbf{b}(x, x) = 1\}$ and the *lightcone $(m-1)$ -sphere* $\mathbb{S}_+^{m-1} = \{x \in \mathbb{R}_1^{m+1} \mid \mathbf{b}(x, x) = 0, x_0 = 1\}$.

An immersion $f : M \rightarrow \mathbb{H}^m$ is called *horo-tight* if every singular function $\mathcal{H}_v(x) = -\mathbf{b}(f(x), v)$, $v \in \mathbb{S}_+^{m-1}$, is either degenerate or a perfect function. The function $L_h(x) = \ln(-\mathbf{b}(x, v))$, $x \in \mathbb{H}^m$ measures the distance from x to the hyperhorosphere $h = \{x \in \mathbb{H}^m \mid \mathbf{b}(x, v) = -1\}$.

An immersion $f : M \rightarrow \mathbb{H}^m$ is called *tight* if, for every $\sigma \in \mathbb{S}_1^m$, the function

$$x \mapsto L_\pi(x) = \sinh^{-1}(-\mathbf{b}(x, \sigma))$$

is either degenerate or a perfect function. Recall that $L_\pi(x)$ is the distance from x to the hyperplane $\pi = \{x \in \mathbb{H}^m \mid \mathbf{b}(x, \sigma) = 0\}$.

An immersion $f : M \rightarrow \mathbb{H}^m$ is called *taut* if, for every $p \in \mathbb{H}^m - f(M)$, the function

$$x \mapsto L_p(x) = (\cosh^{-1}(-\mathbf{b}(x, p)))^2$$

is either degenerate or a perfect function. Recall that $L_p(x)$ is the square of the distance in \mathbb{H}^m from p to x .

The above three concepts are invariant under isometries of \mathbb{H}^m , this follows from the characterization of such isometries, see for instance [6]. Most of the published work on these subjects are contained in the papers [2, 3, 4]. Next we list some results of [4] which will be used latter. First, we observe from Theorem 2.1 that if $f : M \rightarrow \mathbb{D}^m$ is a taut immersion in the Euclidean sense then $P \circ f : M \rightarrow \mathbb{H}^m$ is tight and taut. Next, Corollary 4.2 says that an immersion $f : M \rightarrow \mathbb{D}^m$ is tight if and only if $\Phi \circ f : M \rightarrow \mathbb{D}^m$ is tight in the Euclidean sense for every isometry Φ of (\mathbb{D}^m, g) . Finally, for an immersion $f : M \rightarrow \mathbb{H}^m$ of a compact manifold, Theorem 5.1 says that tightness or tautness implies horo-tightness.

2. HORO-TIGHTNESS IN \mathbb{D}^m

Definition 2.1. An immersion $f : M \rightarrow \mathbb{D}^m$ is *horo-tight* if $P \circ f : M \rightarrow \mathbb{H}^m$ is horo-tight.

Let $f : M^n \rightarrow \mathbb{D}^m$ be an immersion. Let \mathbb{S}^{m-1} denote the sphere of unit vectors at the origin in \mathbb{R}^m . For $w \in \mathbb{S}^{m-1}$, let $\mathcal{R}_w : \mathbb{D}^m \rightarrow \mathbb{R}$ be the function defined by

$$(2.1) \quad R_w(p) = 1 - \frac{1 - \langle p, p \rangle}{2(1 - \langle p, w \rangle)}$$

where $\langle \cdot, \cdot \rangle$ is the Euclidean inner product. Geometrically $\mathcal{R}_w(p)$ is the radius of the unique hyperhorosphere containing p which is tangent to \mathbb{S}^{m-1} at w . The natural restriction of R_w to M is a real-valued function.

We shall compare the singularities of the functions \mathcal{R}_w and \mathcal{H}_v . If (x_1, \dots, x_n) are local coordinates in a neighborhood of $x \in M$, then

$$\frac{\partial}{\partial x_i} \mathcal{R}_w = -\frac{(1 - \langle f, f \rangle) \langle f_i, w \rangle - 2(1 - \langle f, w \rangle) \langle f_i, f \rangle}{2(1 - \langle f, w \rangle)^2}$$

where $f_i = \frac{\partial}{\partial x_i} f$, also, for $v = (1, w) \in \mathbb{S}_+^{m-1}$ and for

$$\mathcal{H}_v(x) = -\mathbf{b}(P \circ f(x), v) = -\frac{2 \langle f(x), w \rangle - \langle f(x), f(x) \rangle - 1}{1 - \langle f(x), f(x) \rangle}$$

we have

$$\frac{\partial}{\partial x_i} \mathcal{H}_v = -2 \frac{(1 - \langle f, f \rangle) \langle f_i, w \rangle - 2(1 - \langle f, w \rangle) \langle f_i, f \rangle}{(1 - \langle f(x), f(x) \rangle)^2}.$$

Thus \mathcal{R}_w and \mathcal{H}_v , $v = (1, w)$, have the same critical points and, at such points, the Hessian of \mathcal{R}_w equals that of \mathcal{H}_v up to a positive scalar factor. Since $M_r(\mathcal{R}_w) = M_{\frac{r}{1-r}}(\mathcal{H}_v)$ for $0 < r < 1$, we have the following:

Proposition 2.2. *An immersion $f : M \rightarrow \mathbb{D}^m$ is horo-tight if and only if every singular \mathcal{R}_w is either degenerate or a perfect function.*

As a direct result we have

Corollary 2.3. *The natural inclusion $i : N \rightarrow \mathbb{D}^m$ of any umbilic submanifold N of $(\mathbb{D}^m, \mathfrak{g})$ is a horo-tight embedding.*

3. HORO-TIGHT IMMERSIONS OF S^1

Let S^1 denote a manifold homeomorphic to the unitary circle. In this section we characterize the horo-tight immersions of S^1 into hyperbolic space. As the next lemma says, we can work with embeddings $S^1 \hookrightarrow \mathbb{H}^m$.

Lemma 3.1. *Every horo-tight immersion $f : S^1 \rightarrow \mathbb{H}^m$ is an embedding.*

Proof. Suppose $f(x_1) = f(x_2) = p$ for two distinct points in S^1 . Let R_w be a Morse function such that the hyperhorosphere $h_s = \{y \in \mathbb{H}^m \mid R_w(y) = s\}$ intersects $f(S^1)$ at p and q , $p \neq q$. It follows from the horo-tightness of f that s is a critical value of R_w and therefore R_w has at least three critical points. A contradiction. \square

In dimension 1 the two-piece-property with respect to hyperhorospheres may be rephrased as follows:

Lemma 3.2. *Let $f : S^1 \rightarrow \mathbb{D}^m$ be a horo-tight immersion. Then for any $w \in \mathbb{S}^{m-1}$ either \mathcal{R}_w is constant or the equation $\mathcal{R}_w(x) = c$ has at most two roots.*

We treat separately the one codimensional case of the main result.

Proposition 3.3. *An immersion $f : S^1 \rightarrow \mathbb{D}^2$ is horo-tight only if it is tight.*

Proof. Let us assume that $\Phi \circ f$ is not tight in the Euclidean sense for some isometry Φ of $(\mathbb{D}^m, \mathfrak{g})$. It follows that $\alpha = \Phi \circ f$ is a non-convex curve. Hence there exists a tangent line r at some point p such that none of its closed half-planes contains $\Phi \circ f(S^1)$ entirely.

Now let h be a tangent horo-circle of $\Phi \circ f$ at p which passes through the interior of $\Phi \circ f(S^1)$. See Figure 1. It follows that h intersects $\Phi \circ f(S^1)$ in at least three points. Since h certainly does not contain $\Phi \circ f(S^1)$ we have a contradiction with Lemma 3.2. Therefore $\Phi \circ f$ is tight in the Euclidean sense for every isometry Φ of $(\mathbb{D}^m, \mathfrak{g})$ and hence the result follows from Theorem 4.2 of [4]. \square




FIGURE 1. Horo-tightness implies convexity.

The second part of the main theorem follows from the lemma below.

Lemma 3.4. *Let $f : S^1 \rightarrow \mathbb{D}^m, m \geq 3$, be an immersion whose image contains the origin of \mathbb{R}^m . Then there exists a hyperhorosphere which intersects $f(S^1)$ in at least three points.*

Proof. We have to find $w \in \mathbb{S}^{m-1}$ and $0 < c < 1$ such that the equation $R_w(x) = c$ has three distinct roots. Since $R_w(\mathbf{0}) = 1/2$ for all $w \in \mathbb{S}^{m-1}$ and $\mathbf{0} \in f(S^1)$, it is sufficient to determine non-zero vectors $f(x), f(y)$ such that the linear system

$$(3.1) \quad \begin{cases} \langle f(x), w \rangle = \langle f(x), f(x) \rangle \\ \langle f(y), w \rangle = \langle f(y), f(y) \rangle \end{cases}$$

has a solution in \mathbb{S}^{m-1} . Let $\pi_u \subset \mathbb{R}^m$ be the hyperplane given by the equation $\langle X, u \rangle = \langle u, u \rangle$. For small enough $f(x)$, the hyperplane $\pi_{f(x)}$ contains a non-zero vector $f(y) \neq f(x)$. Thus any $w \in \pi_{f(x)} \cap \pi_{f(y)}$ is a solution of the system (3.1). \square

We now proof the main result.

Theorem 3.5. *An immersion $f : S^1 \rightarrow \mathbb{D}^m$ is horo-tight if and only if it is tight.*

Proof. We have shown that horo-tightness of $S^1 \hookrightarrow \mathbb{D}^2$ implies tightness, so let us assume $m \geq 3$. We will actually prove that f is taut and tight. In view of Theorem 2.1 of [4], it is enough to show that f is taut in the Euclidean sense.

For any point $p \in \mathbb{H}^m$ one can find an isometry $\Phi : \mathbb{H}^m \rightarrow \mathbb{H}^m$ such that $\Phi(p) = e_0$, see [6]. Hence we may assume that $\mathbf{0} \in f(S^1)$ and thus, by Lemmas 3.4 and 3.2 there exists a hyperhorosphere h which contains $f(S^1)$. For almost all hyperplanes π in \mathbb{R}^m we have a

hyperhorosphere h' such that $\pi \cap h = h' \cap h$. Since f is horo-tight it is also tight in the Euclidean sense. The result now follows from Theorem 2.1 of [1].

The converse is a particular case of Theorem 5.1 of [4]. \square

From the above proof we see that tautness, tightness, and horo-tightness are equivalent concepts in the class of immersions from S^1 into \mathbb{D}^m , $m \geq 3$. Whether the above results hold for higher dimensional spheres is an interesting question which may be studied elsewhere.

Acknowledgement. The first author would like to express his gratitude to his advisers, Maria Aparecida Soares Ruas and Shyuichi Izumiya, for their assistance in this work.

REFERENCES

- [1] S. Carter and A. West, *Tight ant taut immersions*, Proc. London Math. Soc, (3) **25** (1972), 701-720.
- [2] T. E. Cecil, *A characterization of metric spheres in hyperbolic space by Morse theory*, Tohoku Math. J., **26** (1974), 341-351.
- [3] T. E. Cecil and P. J. Ryan, *Distance functions and umbilic submanifolds of hyperbolic space*, Nagoya Math. J., **74** (1979), 67-75.
- [4] T. E. Cecil and P. J. Ryan, *Tight ant taut immersions into hyperbolic space*, J. London Math. Soc., **19** (1979), 561-572.
- [5] T. E. Cecil and P. J. Ryan, *Tight ant taut immersions of manifolds*, Research Notes in Mathematics, **107** (1985).
- [6] R. G. Ratcliffe, *Foundations of hyperbolic manifolds*, Graduate Texts in Math., **149** (1994).

FACULDADES FEDERAIS INTEGRADAS DE DIAMANTINA, RUA DA GLÓRIA 187, 39100-000, DIAMANTINA, MG, BRAZIL.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN.

INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO, DEPARTAMENTO DE MATEMÁTICA, CAIXA POSTAL 668, 13560-970, SÃO CARLOS, SP, BRAZIL.

E-mail address, Marcelo Buosi: pmarpceplo@yahoo.com.br

E-mail address, Shyuichi Izumiya: izumiya@math.sci.hokudai.ac.jp

E-mail address, Maria Aparecida Soares Ruas: maasruas@icmc.usp.br