



Title	Small data scattering for the Klein-Gordon equation with cubic convolution nonlinearity
Author(s)	Sasaki,Hironobu
Citation	Hokkaido University Preprint Series in Mathematics, 745: 1-11
Issue Date	2005
DOI	10.14943/83895
Doc URL	<a href="http://hdl.handle.net/2115/69553">http://hdl.handle.net/2115/69553</a>
Type	bulletin (article)
File Information	pre745.pdf



[Instructions for use](#)

# SMALL DATA SCATTERING FOR THE KLEIN-GORDON EQUATION WITH CUBIC CONVOLUTION NONLINEARITY

HIRONOBU SASAKI\*

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY,  
060-0810, JAPAN.

ABSTRACT. We consider the scattering problem for the Klein-Gordon equation with cubic convolution nonlinearity. We give some estimates for the nonlinearity, and prove the existence of the scattering operator, which improves the known results in some sense. Our proof is based on the Strichartz estimates for the inhomogeneous Klein-Gordon equation.

## 1. INTRODUCTION

This paper is concerned with the scattering problem for the nonlinear Klein-Gordon equation of the form

$$\partial_t^2 u - \Delta u + u = F_\gamma(u) \quad (1.1)$$

in space-time  $\mathbb{R} \times \mathbb{R}^n$ , where  $u$  is a real-valued or a complex-valued unknown function of  $(t, x) \in \mathbb{R} \times \mathbb{R}^n$ ,  $\partial_t = \partial/\partial t$  and  $\Delta$  is the Laplacian in  $\mathbb{R}^n$ . The nonlinearity  $F_\gamma(u)$  is a cubic convolution term  $F_\gamma(u) = -(V_\gamma * |u|^2)u$  with

$$|V_\gamma(x)| \leq C|x|^{-\gamma}. \quad (1.2)$$

Here,  $0 < \gamma < n$  and  $*$  denotes the convolution in the space variables. The term  $F_\gamma(u)$  is an approximative expression of the nonlocal interaction of specific elementary particles. Menzala and Strauss started to study this equation in [5].

In order to treat the scattering problem, we define the scattering operator for (1.1). First, we list some notation to give the definition. Let  $H^s$  be the usual Sobolev space  $(1 - \Delta)^{-s/2} L_2(\mathbb{R}^n)$  and let  $H^{s,\sigma}$  be the weighted Sobolev space  $(1 - \Delta)^{-s/2} \langle x \rangle^{-\sigma} L_2(\mathbb{R}^n)$ . A Hilbert space  $X^{s,\sigma}$

---

2000 *Mathematics Subject Classification.* 35P25, 81Q05, 35B05.

*Key words and phrases.* Global existence, asymptotic behavior, scattering, Klein-Gordon equation, cubic convolution.

\*Research Fellow of the Japan Society for the Promotion of Science.

is denoted by  $H^{s,\sigma} \oplus H^{s-1,\sigma}$ . For a positive number  $\delta$  and a Banach space  $A$ , we denote the set  $\{a \in A; \|a\| \leq \delta\}$  by  $B(\delta; A)$ . Then the scattering operator is defined as the mapping  $S : B(\delta; X^{s,\sigma}) \ni (f_-, g_-) \mapsto (f_+, g_+) \in X^{s,0}$  if the following condition holds for some  $\delta > 0$ :

*For any  $(f_-, g_-) \in B(\delta; X^{s,\sigma})$ , there uniquely exist a time-global solution  $u \in C(\mathbb{R}; H^s)$  of (1.1), and data  $(f_+, g_+) \in X^{s,0}$  such that  $u(t)$  approaches  $u_\pm(t)$  in  $H^s$  as  $t$  tends to  $\pm\infty$ , where  $u_\pm(t)$  are solutions of linear Klein-Gordon equations whose initial data are  $(f_\pm, g_\pm)$ , respectively.*

We call that “ $(S, X^{s,\sigma})$  is well-defined” if we can define the scattering operator  $S : B(\delta; X^{s,\sigma}) \rightarrow X^{s,0}$  for some  $\delta > 0$ .

By Mochizuki [6], it is shown that if  $n \geq 3$ ,  $s \geq 1$ ,  $\gamma < n$  and  $2 \leq \gamma \leq 2s+2$ , then  $(S, X^{s,0})$  is well-defined. By using the methods of Mochizuki and Motai [7] and Strauss [11], we see that if  $n \geq 2$ ,  $s \geq 1$ ,  $4/3 < \gamma < 2$  and  $\sigma > 1/3$ , then  $(S, X^{s,\sigma})$  is well-defined. In view of the condition of  $\sigma$ , there is a gap between the two cases  $\gamma \geq 2$  and  $\gamma < 2$ . Our aim of this paper is to fill the gap. By using the Strichartz estimate for pre-admissible pair and the complex interpolation method for the weighted Sobolev space, we show that  $(S, X^{s,\sigma})$  is well-defined if  $4/3 < \gamma < 2$  and  $\sigma > (2-\gamma)/2$ , which improves the condition above.

In order to state our results, we give notation which will be used in this paper.

For  $s \in \mathbb{R}$  and  $(1/p, 1/q) \in [0, 1] \times [0, 1]$ , let  $H_p^s$  be the Sobolev space  $(1-\Delta)^{-s/2}L_p(\mathbb{R}^n)$  and let  $B_p^s$  be the Besov space  $B_{p,2}^s(\mathbb{R}^n)$  (see, e.g., [1] for the definition of the Besov space). For  $s \in \mathbb{R}$ , we set  $E^s[u](t) = \|(u(t), \partial_t u(t))\|_{X^{s,0}}$ . For  $s_0 \in \mathbb{R}$  and  $Q = (1/q, 1/r) \in [0, 1] \times [0, 1]$ ,  $L(s_0, Q)$  denotes either  $L_q(\mathbb{R}; H_r^{s_0}(\mathbb{R}^n))$  or  $L_q(\mathbb{R}; B_r^{s_0}(\mathbb{R}^n))$ . Put  $J = (0, t)$  or  $(-\infty, t)$  or  $(t, \infty)$ ,  $\omega = \sqrt{1-\Delta}$  and  $U(t) = \exp(\pm it\omega)$ . For a Banach space  $A$ ,  $\mathcal{B}^0(\mathbb{R}; A)$  is the set of all  $A$ -valued, continuous and bounded functions on  $\mathbb{R}$ . Moreover, if  $f$  in  $\mathcal{B}^0(\mathbb{R}; A)$  has its derivative, and if  $\partial_t f \in \mathcal{B}^0(\mathbb{R}; A)$ , then we write  $f \in \mathcal{B}^1(\mathbb{R}; A)$ . For  $s \in \mathbb{R}$ ,  $\mathcal{H}^s$  denotes  $\mathcal{B}^0(\mathbb{R}; H^s) \cap \mathcal{B}^1(\mathbb{R}; H^{s-1})$  with the norm  $\|u\|_{\mathcal{H}^s} = \|u\|_{L(s, (0, 1/2))} + \|\partial_t u\|_{L(s-1, (0, 1/2))}$ . Furthermore, we set

$$\begin{aligned} \underline{\mathcal{H}}^s &= \{u \in \mathcal{H}^s; \text{there exist } f, g \in \dot{\mathcal{S}}(\mathbb{R}^n) \text{ such that} \\ &\quad u(t) = \cos t\omega f + \omega^{-1} \sin t\omega g, \omega^{-1} \partial_t u(t) \in \mathcal{H}^s\}. \end{aligned}$$

We call  $u = u(t, x)$  a free solution if  $u \in \underline{\mathcal{H}}^s$  for some  $s \in \mathbb{R}$ . For a free solution  $u_0$ ,  $u \in \dot{\mathcal{S}}(\mathbb{R}^n)$  is said to be a  $u_0$ -solution if

$$u(t) = u_0(t) + \int_0^t \frac{\sin(t - \tau)\omega}{\omega} F(u(\tau)) d\tau.$$

For  $s, s_0 \in \mathbb{R}$  and  $Q = (1/q, 1/r) \in [0, 1] \times [0, 1]$ , we denote  $L(s_0, Q) \cap \mathcal{H}^s$  and  $L(s_0, Q) \cap \underline{\mathcal{H}}^s$  by  $Z(s_0, s, Q)$  and  $\underline{Z}(s_0, s, Q)$ , respectively. Define  $1/q_\varepsilon = 1/3 - \varepsilon$  and  $1/r_\theta = 1/2 - (1 + \theta)/3n$ . Assume that  $4/3 < \gamma < 2$ . Then we can easily show that there exist sufficiently small  $\varepsilon(\gamma) > 0$  and  $\theta(\gamma) \in (0, 1)$  such that

$$\begin{aligned} \frac{1}{6} &< \frac{n}{2} \left( \frac{1}{2} - \frac{1}{r_{\theta(\gamma)}} \right) < \frac{1}{q_{\varepsilon(\gamma)}} < n \left( \frac{1}{2} - \frac{1}{r_{\theta(\gamma)}} \right), \\ \gamma &= 2 - 2 \left\{ \frac{2}{q_{\varepsilon(\gamma)}} - n \left( \frac{1}{2} - \frac{1}{r_{\theta(\gamma)}} \right) \right\}. \end{aligned}$$

For  $Q_\gamma = (1/q_{\varepsilon(\gamma)}, 1/r_{\theta(\gamma)})$ , we set

$$s(Q_\gamma) = \max \left\{ \frac{n+2}{n} \left( 1 - \frac{3}{q_{\varepsilon(\gamma)}} \right), \frac{2-\gamma}{4} \right\}.$$

We are now ready to state our main results.

**Theorem 1.1.** *Assume that  $n \geq 2$ ,  $4/3 < \gamma < 2$ ,  $s \geq 1$ , and put  $s_\gamma = s(Q_\gamma)$ ,  $Z = Z(s_\gamma + s - 1, s, Q_\gamma)$ . Then there exist some positive numbers  $\delta_0, \delta_+, \delta_-$  satisfying the following properties:*

- (i) *If  $u_0 \in B(\delta_0; \underline{Z})$ , then there uniquely exist  $u \in Z$  and  $u_+, u_- \in \underline{Z}$  such that  $u$  is a  $u_0$ -solution and we have*

$$\lim_{t \rightarrow \pm\infty} E^s[u - u_\pm](t) = 0. \quad (1.3)$$

*Moreover, the operators  $\widetilde{V}_\pm : B(\delta_0; \underline{Z}) \ni u_0 \mapsto u_\pm \in \underline{Z}$  are well defined, injective and continuous.*

- (ii) *If  $u_\pm \in B(\delta_\pm; \underline{Z})$ , then there uniquely exist  $u \in Z$  and  $u_0 \in \underline{Z}$  such that  $u$  is a  $u_0$ -solution and (1.3) holds.*

*Moreover, the operators  $W_\pm : B(\delta_\pm; \underline{Z}) \ni u_\pm \mapsto u_0 \in \underline{Z}$  are well defined, injective and continuous.*

- (iii) *The numbers  $\delta_\pm$  satisfy  $B(\delta_-; \underline{Z}) \subset B(\delta_0; \underline{Z})$ ,  $W_-(B(\delta_-; \underline{Z})) \subset B(\delta_0; \underline{Z})$  and  $B(\delta_+; \underline{Z}) \subset V_+ \circ W_-(B(\delta_-; \underline{Z}))$ . In particular, the operator  $\widetilde{S} = V_+ \circ W_- : B(\delta_-; \underline{Z}) \rightarrow \underline{Z}$  is well defined, injective and continuous.*

The following result follows from Theorem 1.1.

**Corollary 1.2.** *Assume that  $n \geq 2$ ,  $4/3 < \gamma < 2$ ,  $\sigma > (2 - \gamma)/2$ ,  $s \geq 1$  and put  $s_\gamma = s(Q_\gamma)$ ,  $Z = Z(s_\gamma + s - 1, s, Q_\gamma)$ ,  $u_\star(t) = \cos t\omega f_\star +$*

$\omega^{-1} \sin t\omega f_\star$ , where  $\star$  denotes either 0, + or  $-$ . Then there exist some positive numbers  $\eta_0$  and  $\eta_-$  satisfying the following properties:

- (i) If  $(f_0, g_0) \in B(\eta_0; X^{s,\sigma})$ , then there uniquely exist  $u \in Z$  and  $(f_+, g_+), (f_-, g_-) \in X^{s,0}$  such that  $u$  is a  $u_0$ -solution and (1.3) holds.

Moreover, the operators  $V_\pm : B(\eta_0; X^{s,\sigma}) \ni (f_0, g_0) \mapsto (f_\pm, g_\pm) \in X^{s,0}$  are well defined, injective and continuous.

- (ii) If  $(f_-, g_-) \in B(\eta_-; X^{s,\sigma})$ , there uniquely exist  $u \in Z$  and  $(f_+, g_+) \in X^{s,0}$  such that  $u$  satisfies

$$u(t) = u_-(t) + \int_t^\infty \frac{\sin(t-\tau)\omega}{\omega} F(u(\tau)) d\tau \quad (1.4)$$

and (1.3) holds.

Moreover, the scattering operator  $S : B(\eta_-; X^{s,\sigma}) \ni (f_-, g_-) \mapsto (f_+, g_+) \in X^{s,0}$  is well defined, injective and continuous.

*Remark 1.* If  $V$  satisfies  $V(x) = V(-x)$ , and  $0 < \gamma < 3$ , then we can easily show small data global existence for (1.1) using the energy equality (see, e.g., [5]) and an a priori estimate. However, the existence of the scattering operator is not known.

*Remark 2.* For recent results on the wave equation with a cubic convolution, see, e.g., Hidano [3] and Tsutaya [13].

## 2. PRELIMINARIES

In this section, we prove the key lemma for our main results. We first state the generalized Hölder inequality in [10].

**Proposition 2.1.** *If  $s \geq 0$ ,  $1 < p_j < \infty$ ,  $j = 1, \dots, 5$  and  $1/p_1 = 1/p_2 + 1/p_3 = 1/p_4 + 1/p_5$ , then we have*

$$\|fg|H_{p_1}^s\| \lesssim \|f|H_{p_2}^s\| \|g|L_{p_3}\| + \|f|L_{p_4}\| \|g|H_{p_5}^s\|. \quad (2.1)$$

We need the Strichartz estimates proved by Nakamura and Ozawa [9] (see also [8]).

**Proposition 2.2.** *(i) If  $2/q_j = n(1/2 - 1/r_j)$ ,  $2\rho_j = (n+2)(1/2 - 1/r_j)$ ,  $2 \leq q_j, r_j \leq \infty$ ,  $(q_j, r_j) \neq (2, \infty)$ ,  $j = 1, 2$ , then we have*

$$\left\| \int_J U(t-\tau)h(\tau)d\tau \right\|_{L_{q_1} B_{r_1}^{-\rho_1}} \lesssim \|h\|_{L_{q_2} B_{r_2}^{\rho_2}}. \quad (2.2)$$

*(ii) If  $1/r_4 + 2/nq_4 = 1/r_3 + 2/nq_3 + 2/n$ ,  $\max(0, 1/2 - 1/n) < 1/r_j < 1/2$ ,  $0 < 1/q_j < n(1/2 - 1/r_j)$ ,  $1/q_3 < 1/q_4$ ,  $\rho_3 + \rho_4 = (n +$*

2)(1/r'\_4 - 1/r\_3)/2, then we have

$$\left\| \int_J U(t - \tau) h(\tau) d\tau \right\|_{L_{q_3} B_{r_3}^{-\rho_1}} \lesssim \|h\|_{L_{q_4} B_{r_4}^{\rho_4}}. \quad (2.3)$$

We next state the estimates of the nonlinearity.

**Lemma 2.3.** *Assume that  $0 \leq \rho \leq \tilde{\rho}$ ,  $0 < 1/\tilde{r} < 1/2 \leq 1/r < 1$ ,  $0 < \gamma < n$ . If there exist some  $\theta_j \in [0, 1]$ ,  $j = 1, 2$ , satisfying*

$$1 + \frac{1}{r} = \frac{\gamma}{n} + \left(\frac{1}{\tilde{r}} - \theta_1 \frac{\tilde{\rho} - \rho}{n}\right) + 2\left(\frac{1}{\tilde{r}} - \theta_2 \frac{\tilde{\rho}}{n}\right), \quad (2.4)$$

$$\frac{1}{\tilde{r}} - \theta_1 \frac{\tilde{\rho} - \rho}{n}, \frac{1}{\tilde{r}} - \theta_2 \frac{\tilde{\rho}}{n} > 0, \quad (2.5)$$

then we have

$$\|F(u)\|_{H_r^\rho} \lesssim \|u\|_{H_{\tilde{r}}^{\tilde{\rho}}}^3 \quad (2.6)$$

and

$$\|F(u)\|_{B_r^\rho} \lesssim \|u\|_{B_{\tilde{r}}^{\tilde{\rho}}}^3. \quad (2.7)$$

*Proof.* Put

$$\begin{aligned} \frac{1}{p_1} &= -1 + \frac{\gamma}{n} + \left(\frac{1}{\tilde{r}} - \theta_1 \frac{\tilde{\rho} - \rho}{n}\right) + \left(\frac{1}{\tilde{r}} - \theta_2 \frac{\tilde{\rho}}{n}\right), \\ \frac{1}{p_2} &= \frac{1}{\tilde{r}} - \theta_2 \frac{\tilde{\rho}}{n}, \quad \frac{1}{p_3} = -1 + \frac{\gamma}{n} + 2\left(\frac{1}{\tilde{r}} - \theta_2 \frac{\tilde{\rho}}{n}\right), \\ \frac{1}{p_4} &= \frac{1}{\tilde{r}} - \theta_1 \frac{\tilde{\rho} - \rho}{n}, \quad \frac{1}{p_5} = \frac{1}{p_4} + \frac{1}{p_2}. \end{aligned}$$

Then we have  $1/r = 1/p_1 + 1/p_2 = 1/p_3 + 1/p_4$ ,  $1 < r, p_j < \infty$ ,  $j = 1, \dots, 5$ . By Proposition 2.1 and the Hardy-Littlewood-Sobolev inequality, we have

$$\begin{aligned} \|F(f)\|_{H_r^\rho} &= \|(V * |f|^2)f\|_{H_r^\rho} \\ &\lesssim \|V * |f|^2\|_{H_{p_1}^\rho} \|f\|_{L_{p_2}} + \|V * |f|^2\|_{L_{p_3}} \|f\|_{H_{p_4}^\rho}, \\ \|V * |f|^2\|_{H_{p_1}^\rho} &\lesssim \|V * \omega^\rho |f|^2\|_{L_{p_1}} \\ &\lesssim \|\omega^\rho |f|^2\|_{L_{p_5}} \\ &= \| |f|^2 \|_{H_{p_5}^\rho} \\ &\lesssim \|f\|_{H_{p_4}^\rho} \|f\|_{L_{p_2}} \end{aligned}$$

and

$$\|V * |f|^2\|_{L_{p_3}} \lesssim \|f\|_{L_{p_2}}^2.$$

By the embedding  $H_{\tilde{r}}^{\tilde{\rho}} \hookrightarrow L_{p_2}$  and  $H_{\tilde{r}}^{\tilde{\rho}} \hookrightarrow H_{p_4}^\rho$ , we obtain (2.6). From (2.6), we see that (2.7) holds since  $H_r^\rho \hookrightarrow B_r^\rho$  and  $B_{\tilde{r}}^{\tilde{\rho}} \hookrightarrow H_{\tilde{r}}^{\tilde{\rho}}$ .  $\square$

Finally we state the key lemma to prove Theorem 1.1.

**Lemma 2.4.** *Assume that  $n \geq 2$ ,  $4/3 < \gamma < 2$ ,  $s \geq 1$  and put  $L = L(s_\gamma + s - 1, Q_\gamma)$ . Then there exists some  $\delta > 0$  satisfying as follows: If  $u_0 \in B_\delta(L)$ , then there uniquely exists  $u \in L$  such that we have*

$$u(t) = u_0(t) + \int_J \frac{\sin(t - \tau)\omega}{\omega} F(u(\tau)) d\tau, \quad (2.8)$$

$$\|u|L\| \leq \frac{4}{3} \|u_0|L\|, \quad (2.9)$$

$$\left\| \int_J U(t - \tau) F(u(\tau)) d\tau | L(s - 1, (0, 1/2)) \right\| \leq \frac{1}{3} \|u_0|L\|. \quad (2.10)$$

*Proof.* (Step I.) In order to show the existence of a time-global solution, we define the contraction mapping on the suitable complete metric space. Put  $Y = B(\frac{4}{3} \|u_0|L\|; L)$  and  $d(u, v) = \|u - v|L\|$ . Then  $(Y, d)$  is a nonempty complete metric space. We define a mapping  $\Phi$  by

$$\Phi : u \longmapsto u_0 + \int_J \frac{\sin(t - \tau)\omega}{\omega} F(u(\tau)) d\tau.$$

By Proposition 2.2,(2), we have

$$\left\| \int_J \frac{\sin(t - \tau)\omega}{\omega} F(u(\tau)) d\tau | L \right\| \lesssim \|F(u)|L(s_\gamma + s - 2 + \tilde{\rho}, (\frac{3}{q_\varepsilon}, \frac{1}{\tilde{r}_\theta}))\|.$$

Here  $(\varepsilon, \theta)$  denotes  $(\varepsilon(\gamma), \theta(\gamma))$  and  $(\tilde{r}_\theta, \tilde{\rho})$  satisfies

$$\begin{aligned} \frac{1}{\tilde{r}_\theta} + \frac{6}{nq_\varepsilon} &= \frac{1}{r_\theta} + \frac{2}{nq_\varepsilon} + \frac{2}{n}, \\ \tilde{\rho} &= \frac{n+2}{2} \left( \frac{1}{\tilde{r}_\theta} - \frac{1}{r_\theta} \right) = \frac{n+2}{2} \left( \frac{2}{n} - \frac{4}{nq_\varepsilon} \right) = \frac{n+2}{3n} + 2 \frac{n+2}{n} \varepsilon. \end{aligned}$$

Remark that  $\max(0, 1/2 - 1/n) < 1/r_\theta$ ,  $1/\tilde{r}_\theta < 1/2$ ,  $0 < 1/q_\varepsilon < n(1/2 - 1/r_\theta)$  and  $0 < 1 - 3/q_\varepsilon < n(1/2 - 1/\tilde{r}_\theta)$  since

$$\begin{aligned} \frac{1}{2} &< \frac{1}{\tilde{r}_\theta} = \frac{1}{r_\theta} - \frac{4}{nq_\varepsilon} + \frac{2}{n} = \frac{1}{2} + \frac{1-\theta}{3n} + \frac{4\varepsilon}{n} < \frac{1}{2} + \frac{1}{n}, \\ 0 &< 1 - \frac{3}{q_\varepsilon} = 3\varepsilon < \frac{1-\theta}{3} + 4\varepsilon = n \left( \frac{1}{\tilde{r}_\theta} - \frac{1}{2} \right). \end{aligned}$$

Since  $\tilde{\rho} \leq 1$ , we have

$$\|F(u)|L(s_\gamma + s - 2 + \tilde{\rho}, (\frac{3}{q_\varepsilon}, \frac{1}{\tilde{r}_\theta}))\| \lesssim \|F(u)|L(s_\gamma + s - 1, (\frac{3}{q_\varepsilon}, \frac{1}{\tilde{r}_\theta}))\|.$$

It follows from Lemma 2.3 that

$$\|F(u)|L(s_\gamma + s - 1, (\frac{3}{q_\varepsilon}, \frac{1}{\dot{r}_\theta}))\| \lesssim \|u|L\|^3$$

since  $1 + 1/\dot{r}_\theta = \gamma/n + 3/r_\theta$ .

(Step II.) We estimate the left hand side of (2.10). By Proposition 2.2,(2),

$$\| \int_J U(t - \tau) F(u(\tau)) d\tau |L(s - 1, (0, 1/2))\| \lesssim \|F(u)|L(s - 1 + \dot{\rho}, (\frac{3}{q_\varepsilon}, \frac{1}{\dot{r}_\theta}))\|,$$

where

$$\frac{1}{\dot{r}_\theta} = \frac{1}{2} + \frac{2}{n}(1 - \frac{3}{q_\varepsilon}), \dot{\rho} = \frac{n+2}{n}(1 - \frac{3}{q_\varepsilon}).$$

Remark that  $2(1 - 3/q_\varepsilon) = n(1/\dot{r}_\theta - 1/2)$  and  $2\dot{\rho} = (n+2)(1/\dot{r}_\theta - 1/2)$ .

Put

$$\vartheta = \frac{1}{s - 1 + s_\gamma} \left\{ \frac{1}{q_\varepsilon} - \frac{n}{2} \left( \frac{1}{2} - \frac{1}{r_\theta} \right) \right\}.$$

Then we have  $0 \leq \dot{\rho} \leq s_\gamma$ ,  $0 \leq \vartheta \leq 1$ ,

$$\begin{aligned} 1 + \frac{1}{\dot{r}_\theta} &= \frac{\gamma}{n} + \frac{1}{r_\theta} + 2 \left( \frac{1}{r_\theta} - \vartheta \frac{s - 1 + s_\gamma}{n} \right), \\ \frac{1}{r_\theta} - \vartheta \frac{s - 1 + s_\gamma}{n} &> 0. \end{aligned}$$

Thus, by Lemma 2.3, we have

$$\|F(u)|L(s - 1 + \dot{\rho}, (\frac{3}{q_\varepsilon}, \frac{1}{\dot{r}_\theta}))\| \lesssim \|u|L\|^3.$$

(Step III.) We prove (2.9) and (2.10). By Steps I and II, we have

$$\begin{aligned} \|\Phi u|L\| &\leq \|u_0|L\| + C\|u|L\|^3, \\ \|\Phi u - \Phi v|L\| &\leq C(\|u|L\|^2 + \|v|L\|^2)\|u - v|L\| \end{aligned}$$

and

$$\begin{aligned} \| \int_J U(t - \tau) F(u(\tau)) d\tau |L(s - 1, (0, 1/2))\| \\ \leq C\|u|L\|^3 \end{aligned}$$

for all  $u, v \in Y$ . Thus, if we put  $\|u_0|L\| \leq \delta$  and  $\delta \leq 3/(8\sqrt{C})$ , then we obtain

$$\|\Phi u|L\| \leq (1 + \frac{64}{27}C\delta^2)\|u_0|L\| \leq \frac{4}{3}\|u_0|L\|, \quad (2.11)$$

$$d(\Phi u, \Phi v) \leq \frac{32}{9}C\delta^2\|u - v|L\| \leq \frac{1}{2}d(u, v) \quad (2.12)$$



and

$$\left\| \int_J U(t-\tau)F(u(\tau))d\tau \middle| L(s-1, (0, 1/2)) \right\| \leq \frac{64}{27}C\delta^2\|u_0\|L \leq \frac{1}{3}\|u_0\|L. \quad (2.13)$$

By (2.11) and (2.12),  $\Phi$  is a contraction mapping from  $(Y, d)$  into itself. Thus, there exists a unique fixed point  $u \in Y$ . By (2.13), we obtain (2.10).  $\square$

### 3. PROOF OF THEOREM 1.1

In this section, we give a proof of Theorem 1.1 dividing into five steps.

(Step I.) For all  $u_0 \in B_\delta(\underline{Z}(s_\gamma + s - 1, s, Q_\gamma))$ , by Lemma 2.4, there uniquely exists  $u_0$ -solution satisfying (2.9) and (2.10). We have

$$\begin{aligned} \|u\|\mathcal{H}^s &\leq \|u_0\|\mathcal{H}^s + \left\| \int_0^t U(t-\tau)F(u(\tau))d\tau \middle| L(s-1, (0, 1/2)) \right\| \\ &\leq \|u_0\|\mathcal{H}^s + \|u_0\|L/3 \\ &\leq \|u_0\|Z. \end{aligned}$$

Thus,  $\|u\|Z \leq 3\|u_0\|Z$ . Put

$$u_\pm = u_0 + \int_0^{\pm\infty} \frac{\sin(t-\tau)\omega}{\omega} F(u(\tau))d\tau.$$

Then we have

$$\begin{aligned} \|u_\pm\|Z &\leq \|u_0\|Z + \left\| \left( \int_0^t + \int_t^{\pm\infty} \right) \frac{\sin(t-\tau)\omega}{\omega} F(u(\tau))d\tau \middle| Z \right\| \\ &\leq 5/3\|u_0\|Z \end{aligned}$$

and  $u_\pm \in \underline{\mathcal{H}}^s$ .

(Step II.) It follows from Lemma 2.4 that

$$E[u - u_\pm](t) \lesssim \|\chi_{(t, \pm\infty)}u\|L \rightarrow 0$$

as  $t \rightarrow \pm\infty$ .

Assume that there exist some  $\tilde{u}_\pm \in \mathcal{H}^s$  such that  $E[u - \tilde{u}_\pm](t) \rightarrow 0$  as  $t \rightarrow 0$ . By the energy equality, we have

$$E[u_\pm - \tilde{u}_\pm](0) \simeq E[u_\pm - \tilde{u}_\pm](t) \rightarrow 0$$

as  $t \rightarrow 0$ . Hence  $u_\pm = \tilde{u}_\pm$ . Thus,  $V_\pm : B_\delta(\underline{Z}) \ni u_0 \mapsto u_\pm \in \underline{Z}$  is well defined.

(Step III.) It is clear to see  $u_0 \mapsto u$  is injective. If we put  $u \mapsto u_\pm$  and  $\tilde{u} \mapsto u_\pm$ , then we have

$$u_0 + \int_0^{\pm\infty} \frac{\sin(t-\tau)\omega}{\omega} F(u(\tau)) d\tau = \tilde{u}_0 + \int_0^{\pm\infty} \frac{\sin(t-\tau)\omega}{\omega} F(\tilde{u}(\tau)) d\tau.$$

Since  $u$  is  $u_0$ -solution, we have

$$u + \int_t^{\pm\infty} \frac{\sin(t-\tau)\omega}{\omega} F(u(\tau)) d\tau = \tilde{u} + \int_t^{\pm\infty} \frac{\sin(t-\tau)\omega}{\omega} F(\tilde{u}(\tau)) d\tau.$$

Thus,

$$\begin{aligned} \|u - \tilde{u}|Z\| &= \left\| \int_t^{\pm\infty} \frac{\sin(t-\tau)\omega}{\omega} \{F(u) - F(\tilde{u})\} d\tau \right\| \\ &\leq \frac{1}{2} \|u - \tilde{u}|Z\| \end{aligned}$$

if  $\delta$  is sufficiently small. Hence  $V_\pm$  is injective.

(Step IV.) For sufficiently small  $\delta$ , we have

$$\begin{aligned} \|u - v|Z\| &\leq \|u_0 - v_0|Z\| + \frac{1}{2} \|u - v|Z\|, \\ \|u_\pm - v_\pm|Z\| &\leq \|u_0 - v_0|Z\| + \frac{1}{2} \|u - v|Z\|. \end{aligned}$$

Thus,  $u_\pm \rightarrow v_\pm$  in  $Z$  if  $u_0 \rightarrow v_0$  in  $Z$ . Hence Theorem 1.1,(i) holds. Theorem 1.1,(ii) can be proved analogously.

(Step V.) We put  $\delta_- = \delta_0/3$ . Then we have  $B_{\delta_-}(\underline{Z}) \hookrightarrow B_{\delta_0}(\underline{Z})$  and  $W_-(B_{\delta_-}(\underline{Z})) \hookrightarrow B_{\delta_0}(\underline{Z})$  since  $\|V_\pm(u_0)|Z\| \leq 3\|u_0|Z\|$  and  $\|W_\pm(u_\pm)|Z\| \leq 3\|u_\pm|Z\|$ . Put  $\delta_+ = \delta_-/9 = \delta_0/27$ . Then  $V_- \circ W_+(B_{\delta_+}(\underline{Z})) \subset B_{\delta_+}(\underline{Z})$ . Thus  $B_{\delta_+}(\underline{Z}) \hookrightarrow V_+ \circ W_-(B_{\delta_-}(\underline{Z}))$ . Hence we have completed the proof of Theorem 1.1.

#### 4. PROOF OF COROLLARY

It is sufficient to show the following lemma:

**Lemma 4.1.** *Assume that  $n \geq 2$ ,  $\max(0, 1/2 - 1/n) < 1/r < 1/2$  and  $(n/2 - n/r)/2 < 1/q < (n/2 - n/r)$ . Then we have*

$$\|U(\cdot)f|L_q L_r\| \lesssim \|f|H^{s,\sigma}\| \quad (4.1)$$

if

$$s > \frac{n+2}{2} \left( \frac{1}{2} - \frac{1}{r} \right) + \frac{n+2}{2n} \left\{ \frac{2}{q} - n \left( \frac{1}{2} - \frac{1}{r} \right) \right\}$$

and

$$\sigma > \frac{2}{q} - n \left( \frac{1}{2} - \frac{1}{r} \right).$$

*Proof.* We can choose  $\vartheta \in (0, 1)$  and sufficiently small  $\delta > 0$  which satisfy

$$\frac{1}{q} = (1 - \vartheta) \frac{n}{2} \left( \frac{1}{2} - \frac{1}{r} \right) + \vartheta \left\{ n \left( \frac{1}{2} - \frac{1}{r} \right) - \delta \right\}.$$

By using the Strichartz estimate for the homogeneous Klein-Gordon equation prove by [4], we have

$$\|U(\cdot)f|L_{q_0}L_r\| \lesssim \|f|H^{(n+2)(1/2-1/r)/2}\|,$$

where  $1/q_0 = (n/2 - n/r)/2$ . On the other hand, by [2], we see that

$$\begin{aligned} \|U(t)f|L_r\| &\lesssim \langle t \rangle^{-(n/2-n/r)} \max\{\|f|H^{n(1/2-1/r)}\|, \|f|H_r^{(n+2)(1/2-1/r)}\|\} \\ &\lesssim \langle t \rangle^{-(n/2-n/r)} \|f|H^{s_1, \sigma_1}\|, \end{aligned}$$

where  $s_1 = (n+2)(1/2 - 1/r)$ ,  $\sigma_1 = n(1/2 - 1/r) + \delta$ . Thus, it holds

$$\|U(\cdot)f|L_{q_1}L_r\| \lesssim \|f|H^{s_1, \sigma_1}\|,$$

where  $1/q_1 = n(1/2 - 1/r) - \delta$ . By (24) in [12], we have

$$[H^{s_1/2, 0}, H^{s_1, \sigma_1}]_{\vartheta} = H^{(1+\vartheta)s_1/2, \vartheta\sigma_1}.$$

Here,  $[\cdot, \cdot]$  denotes the complex interpolation method and

$$\vartheta = \left\{ \frac{n}{2} \left( \frac{1}{2} - \frac{1}{r} \right) - \delta \right\}^{-1} \left\{ \frac{1}{q} - \frac{n}{2} \left( \frac{1}{2} - \frac{1}{r} \right) \right\}.$$

Hence, we obtain

$$\|U(\cdot)f|L_qL_r\| \lesssim \|f|H^{(1+\vartheta)s_1/2, \vartheta\sigma_1}\|.$$

We can put  $s > (1 + \vartheta)s_1/2$  and  $\sigma > \vartheta\sigma_1$  with sufficiently small  $\delta$ . Thus, (4.1) holds.  $\square$

By substituting  $q = q_{\varepsilon(\gamma)}$  and  $r = r_{\theta(\gamma)}$  for (4.1), we can easily see that Corollary 1.2 holds.

**Acknowledgement** The author would like to acknowledge the helpful guidance and encouragement of Professors T. Ozawa and K. Tsutaya. The author is grateful to the referee for pointing out some gaps in the proof and mistakes in the manuscript.

## REFERENCES

- [1] J. Bergh and J. Löfström, *Interpolation spaces*, Grundlehren der Mathematischen Wissenschaften, Springer, Berlin, 1976.
- [2] P. Brenner, *On scattering and everywhere defined scattering operators for nonlinear Klein-Gordon equations*, J. Differential Equations 56 (1985), 310-344.
- [3] K. Hidano, *Small data scattering and blow-up for a wave equation with a cubic convolution*, Funkcialaj Ekvacioj 43 (2000), 559-588.

- [4] S. Machihara, K. Nakanishi and T. Ozawa, *Small global solutions and the non-relativistic limit for the nonlinear Dirac equation*, Rev. Mat. Iberoamericana 19 (2002), 179-194.
- [5] G. Menzala and W. Strauss, *On a wave equation with a cubic convolution*, J. Differential Equations 43 (1982), 93-105.
- [6] K. Mochizuki, *On small data scattering with cubic convolution nonlinearity*, J. Math. Soc. Japan 41 (1989), 143-160.
- [7] K. Mochizuki and T. Motai, *On small data scattering for some nonlinear wave equations*, in "Patterns and Waves -qualitative analysis of nonlinear differential equations -," pp.543-560, Stud. Math. Appl., 18, North-Holland, Amsterdam, 1986.
- [8] M. Nakamura, *A miniature scattering theory for nonlinear Klein-Gordon Equations*, Kyushu J. Math. 57 (2003), 255-263.
- [9] M. Nakamura and T. Ozawa, *The Cauchy problem for nonlinear Klein-Gordon equations in the Sobolev spaces*, Publ. Res. Inst. Math. Sci. 37 (2001), 255-293.
- [10] G. Ponce, *On the global well-posedness of the Benjamin-Ono equation*, Differential and Integral Equations 4 (1991), 527-542.
- [11] W. Strauss, *Nonlinear scattering theory at low energy: sequel*, J. Funct. Anal. 43 (1981), 281-294.
- [12] H. Triebel, *Spaces of distributions with weights. Multipliers in  $L_p$ -spaces with weights*, Math. Nachr. 78 (1977), 339-355.
- [13] K. Tsutaya, *Global existence and blow-up for a wave equation with a potential and a cubic convolution*, Nonlinear analysis and applications: to V. Lakshmikantham on his 80th birthday. vol. 1, 2, 913-937, Kluwer Acad. Publ., Dordrecht, 2003.
- [14] K. Tsutaya and Y. Tsutsumi, private communication.

*E-mail address:* hisasaki@math.sci.hokudai.ac.jp