



Title	High regularity of solutions of compressible Navier-Stokes equations
Author(s)	Cho, Yonggeun
Citation	Hokkaido University Preprint Series in Mathematics, 776, 1-64
Issue Date	2006
DOI	10.14943/83926
Doc URL	<a href="http://hdl.handle.net/2115/69584">http://hdl.handle.net/2115/69584</a>
Type	bulletin (article)
File Information	pre776.pdf



[Instructions for use](#)

# HIGH REGULARITY OF SOLUTIONS OF COMPRESSIBLE NAVIER-STOKES EQUATIONS

YONGGEUN CHO

ABSTRACT. We study the Navier-Stokes equations for compressible *barotropic* fluids in a bounded or unbounded domain  $\Omega$  of  $\mathbf{R}^3$ . The initial density may vanish in an open subset of  $\Omega$  or to be positive but vanish at space infinity. We first prove the local existence of solutions  $(\rho^{(j)}, u^{(j)})$  in  $C([0, T_*]; H^{2(k-j)+3} \times D_0^1 \cap D^{2(k-j)+3}(\Omega))$ ,  $0 \leq j \leq k, k \geq 1$  under the assumptions that the data satisfy compatibility conditions and that the initial density is sufficiently small. To control the nonnegativity or decay at infinity of density, we need to establish a boundary value problem of  $(k+1)$ -coupled elliptic system which may not be in general solvable. The smallness condition of initial density is necessary for the solvability, which is not necessary in case that the initial density has positive lower bound. Secondly, we prove the global existence of smooth radial solutions of *isentropic* compressible Navier-Stokes equations on a bounded annulus or a domain which is the exterior of a ball under a smallness condition of initial density.

## 1. INTRODUCTION

We consider the following compressible Navier-Stokes equations describing the motion of a viscous compressible barotropic fluid in a domain  $\Omega$  of  $\mathbf{R}^3$

$$(1.1) \quad \rho_t + \operatorname{div}(\rho u) = 0 \quad \text{in } (0, T) \times \Omega,$$

$$(1.2) \quad (\rho u)_t + \operatorname{div}(\rho u \otimes u) + L u + \nabla p = \rho f \quad \text{in } (0, T) \times \Omega,$$

$$(1.3) \quad L u = -\mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u,$$

and the initial and boundary conditions

$$(1.4) \quad (\rho, u)|_{t=0} = (\rho_0, u_0) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } (0, T) \times \partial\Omega,$$

$$(1.5) \quad (\rho - \rho^\infty, u)(t, x) \rightarrow (0, 0) \quad \text{as } |x| \rightarrow \infty, \quad (t, x) \in (0, T) \times \Omega.$$

Here we denote by  $\rho$  and  $u$  the unknown density and velocity fields of the fluid, respectively.  $p$  is the pressure which is a smooth function of  $\rho$ .  $f$  denotes a given external force and the constants  $\mu, \lambda$  are the viscosity coefficients. We assume that the viscosity coefficients  $\mu$  and  $\lambda$  satisfy the natural physical restrictions  $\mu > 0$  and  $3\lambda + 2\mu \geq 0$  so that  $L = -\mu \Delta - (\lambda + \mu) \nabla \operatorname{div}$  is a strongly elliptic operator. We assume that  $T$  is a finite positive number and  $\Omega$  is either a bounded domain in  $\mathbf{R}^3$  with smooth boundary or a usual unbounded domain such as the whole space  $\mathbf{R}^3$ ,

---

2000 *Mathematics Subject Classification.* 35Q30, 76N10.

*Key words and phrases.* viscous compressible fluids, compressible Navier-Stokes equations, vacuum.

The author is JSPS Research Fellow.

the half space  $\mathbf{R}^2 \times \mathbf{R}_+$  and an exterior domain with smooth boundary. If  $\Omega$  is a bounded domain (or the whole space), then the condition (1.5) at infinity (or the boundary condition in (1.4) respectively) is unnecessary and should be neglected.  $\rho^\infty$  is a nonnegative constant. For the simplicity of presentation, we always assume that  $\rho^\infty = 0$ . When the density has positive lower bound on a domain, we mean it by that  $\rho^\infty > 0$ . In particular,  $\rho(x) \rightarrow \rho^\infty$  as  $|x| \rightarrow \infty$  for unbounded domain. For the mathematical derivation of the equations (1.1) and (1.2), see [19] for instance.

Throughout this paper, we use the following simplified notations for the homogeneous and inhomogeneous Sobolev spaces.

$$\begin{aligned} L^r &= L^r(\Omega), \quad D^{k,r} = \{v \in L^1_{loc}(\Omega) : |v|_{D^{k,r}} < \infty\}, \\ W^{k,r} &= L^r \cap D^{k,r}, \quad H^k = W^{k,2}, \quad D^k = D^{k,2}, \\ D_0^1 &= \{v \in L^6(\Omega) : |v|_{D_0^1} < \infty \text{ and } v = 0 \text{ on } \partial\Omega\}, \\ H_0^1 &= L^2 \cap D_0^1, \quad |v|_{D^{k,r}} = |\nabla^k v|_{L^r} \quad \text{and} \quad |v|_{D_0^1} = |\nabla v|_{L^2}. \end{aligned}$$

Hereafter we use the obvious notation

$$|\cdot|_{X \cap Y} = |\cdot|_X + |\cdot|_Y \quad \text{for (semi-)normed spaces } X, Y$$

and  $C$  denotes a generic positive constant depending only on the constants  $k$  (Sobolev space index),  $\mu$ ,  $\lambda$ ,  $T$  and the norms of  $p = p(\cdot)$  and  $f$ . For a detailed study of homogeneous Sobolev spaces, we refer the readers to the Galdi's book [13].

In this paper, we study the local and global existence and the regularity for the initial boundary value problem (simply IBVP) (1.1)-(1.5). Our result is a sequel of those in [5] and [4], in which the existence and uniqueness of strong and classical solutions are considered under a general assumption on initial density that  $\rho_0 \geq 0$ . In the present case, we consider a high regularity of local and global solutions in Sobolev space. There has been a large amount of literature on the existence and regularity related to IBVP (1.1)–(1.16). See [18, 19] and [9, 10, 11, 12, 16, 17] for the weak solutions, and [22, 15, 26, 20, 21, 8, 14, 25, 27, 28, 29, 23] and the recent works [3, 4, 5, 6] for strong solutions. For the finite time blowup of smooth solutions, see [2] and [30].

To treat the high regularity of solutions, we need to control the high order time derivatives of solutions. For this purpose it is necessary to assume the compatibility conditions for the time derivatives of solutions at the initial time zero. More specifically, we denote  $(\partial/\partial t)^k v$ , the  $k$ -th order time derivative of  $v$ , by  $v^{(k)}$  and differentiate the equation (1.2)  $k$ -times with respect to  $t$ . Then we have

$$\begin{aligned} (1.6) \quad & \rho u^{(k+1)} + Lu^{(k)} + \nabla p^{(k)} \\ &= \sum_{0 \leq j \leq k} \binom{k}{j} \rho^{(k-j)} (f - u \cdot \nabla u)^{(j)} \\ & \quad - k \rho^{(1)} u^{(k)} - \sum_{1 \leq j \leq k-1} \binom{k}{j-1} \rho^{(k+1-j)} u^{(j)}. \end{aligned}$$

For the regularity of  $u^{(j)}$  at time zero we then define a  $(k+1)$ -coupled compatibility condition as follows: for some functions  $g_j \in D_0^1 \cap D^{2(k-j)+3}$ ,  $1 \leq j \leq k+1$ , the data satisfy that

$$(1.7) \quad Lu_0 + \nabla p(\rho_0) = \rho_0 (f(0) - u_0 \cdot \nabla u_0 - g_1) \quad \text{in } \Omega$$

and

$$(1.8) \quad \begin{aligned} & L(u^{(j)}(0)) + \nabla p^{(j)}(0) \\ &= -j\rho^{(1)}(0)u^{(j)}(0) - \sum_{1 \leq l \leq j-1} \binom{j}{l-1} \rho^{(j+1-l)}(0)u^{(l)}(0) \\ &+ \sum_{0 \leq l \leq j} \binom{j}{l} \rho^{(j-l)}(0) \left( f^{(l)} - \sum_{0 \leq m \leq l} \binom{l}{m} u^{(l-m)}(0) \cdot \nabla u^{(m)}(0) \right) \\ &- \rho_0 g_{j+1} \quad \text{in } \Omega \end{aligned}$$

for all  $1 \leq j \leq k$ . Here we define  $p^{(k)}(0)$  and  $\rho^{(k)}(0)$  recursively as follows:

$$\begin{aligned} p^{(j)}(0) &= \left( \frac{\partial}{\partial t} \right)^j (p(\rho)) \Big|_{t=0}, \\ \rho^{(j+1)}(0) &= -\operatorname{div} \left( \sum_{0 \leq l \leq j} \binom{j}{l} \rho^{(j-l)}(0)u^{(l)}(0) \right), \quad 0 \leq j \leq k-1. \end{aligned}$$

We call the condition (1.7) zeroth compatibility condition and (1.8)  $j$ -th one. Under the smallness of condition on  $\rho_0$ , the solution  $u^{(j)}(0)$  to the mapping  $(g_1, \dots, g_{k+1}) \mapsto (u_0, u^{(1)}(0), \dots, u^{(k)}(0))$  is unique. See Section 4 and 5 for the existence and uniqueness.

In [5, 4], the authors used the zeroth compatibility condition (1.7) to prove the local existence of strong and classical solutions. See also [3]. We follow the similar strategy in [4] summarized as follows:

- (1) linearization on bounded domain with initial density having positive lower bound,
- (2) a priori estimates independent of domain size and lower bound of initial density, and domain expansion,
- (3) construction of a sequence of approximate solutions to the linearized problem,
- (4) convergence of the sequence in a strong sense.

If we consider only the zeroth compatibility condition (1.7), then in the course of linearization (1) and domain expansion (2), it is usually necessary to reconstruct  $u_0$  satisfying (1.7) for a given  $g_1$  (see Lemma 9 in [4]). Since (1.7) is a single elliptic equation, the existence of  $u_0$  is well-known by the classical elliptic theory. However, in the present case, the situation is quite different. We encounter the case that we should find  $(u_0, u^{(1)}(0), \dots, u^{(k)}(0))$  satisfying (1.8) for a given pair  $(g_1, \dots, g_{k+1})$ . In other words, we must find solutions of  $(k+1)$ -coupled elliptic system. In general, it is difficult to solve such a system. However, if we assume

that the initial density is sufficiently small, then we can find solution successfully (see Lemma 4.2 below). From this solvability of  $(k + 1)$ -coupled system, we can construct approximate solutions to the linearized problem, which is convergent to a solution of the original equations. For this purpose, we also have to construct a sequence of system of compatibility conditions and show the strong convergence of the sequence to the original system of compatibility conditions (1.7) and (1.8) which are possible under the smallness condition of the initial density. See Section 5 below.

Now we state the first main result.

**Theorem 1.1.** *Assume that*

$$(1.9) \quad \begin{aligned} & \rho_0 \in L^1 \cap H^{2k+3}, \quad \rho_0 \geq 0 \quad \text{in } \Omega, \quad u_0 \in D_0^1 \cap D^{2k+3}, \\ & f^{(j)} \in C(0, T; H^{2(k-j)+1}) \cap L^2(0, T; H^{2(k-j)+2}) \quad \text{for } 0 \leq j \leq k \\ & f^{(k+1)} \in L^2(0, T; L^2) \end{aligned}$$

and that the data  $\rho_0, u_0$  and  $f$  satisfy the compatibility conditions (1.7) and (1.8). Assume further that  $|\rho_0|_{L^1 \cap H^{2k+3}}$  is sufficiently small. Then there exist a time  $T_* \in (0, T)$  and a unique solution  $(\rho, u)$  to the IBVP (1.1)–(1.5) such that for all  $0 \leq j \leq k$ ,  $\rho \in C([0, T_*]; L^1)$

$$(1.10) \quad \begin{aligned} & \rho^{(j)} \in C([0, T_*]; H^{2(k-j)+3}) \cap L^2(0, T_*; H^{2(k-j)+4}), \\ & u^{(j)} \in C([0, T_*]; D_0^1 \cap D^{2(k-j)+3}) \cap L^2(0, T_*; D^{2(k-j)+4}) \\ & u^{(k+1)} \in L^\infty(0, T_*; D_0^1) \cap L^2(0, T_*; D^2), \\ & \sqrt{\rho} u^{(k+1)} \in L^\infty(0, T_*; L^2). \end{aligned}$$

*Remark 1.2.* If the initial density has positive lower bound, then the smallness condition of initial density can be removed. See Remark 3.3 and Section 4 below

*Remark 1.3.* For each  $k \geq 1$ , one can observe a smoothing effect and apply the methods used to get a classical solution in [4] to the improvement of regularity slightly.

Now we study the global existence of radial solutions. The global existence of strong solution ( $k = 0$ ) is known by H. Kim and H.J. Choe [7]. They showed the density is bounded on any finite time interval by using the radial symmetry and the effective viscous effect. The second main result of this paper is to show that every Sobolev norm of the solution is controlled by  $L^\infty$  norm of density. See Theorem 6.2 below.

To do this, we need to estimate the  $L^\infty$  norm of pressure  $p$  (see (6.9) below). This is possible in the case of the isentropic compressible Navier-Stokes equations with  $p = A\rho^\gamma$ ,  $A > 0$  and  $\gamma > 1$ . Since from (1.1), the pressure  $p$  satisfies the equation  $p_t + \nabla p \cdot u + \gamma p \operatorname{div} u = 0$ . Hence for a high regularity for any  $\gamma > 1$ , as in [4] we consider the following slightly general IBVP of isentropic compressible

Navier-Stokes equations:

$$(1.11) \quad \rho_t + \operatorname{div}(\rho u) = 0 \quad \text{in } (0, T) \times \Omega,$$

$$(1.12) \quad p_t + \nabla p \cdot u + \gamma p \operatorname{div} u = 0 \quad \text{in } (0, T) \times \Omega,$$

$$(1.13) \quad (\rho u)_t + \operatorname{div}(\rho u \otimes u) + L u + \nabla p = \rho f \quad \text{in } (0, T) \times \Omega,$$

$$(1.14) \quad L u = -\mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u,$$

and the initial and boundary conditions

$$(1.15) \quad (\rho, p, u)|_{t=0} = (\rho_0, p_0, u_0) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } (0, T) \times \partial\Omega,$$

$$(1.16) \quad (\rho, p, u)(t, x) \rightarrow (0, 0, 0) \quad \text{as } |x| \rightarrow \infty, \quad (t, x) \in (0, T) \times \Omega.$$

In fact, the barotropic fluids satisfy the equation  $p_t + \nabla p \cdot u + \rho p'(\rho) \operatorname{div} u = 0$  and the initial pressure  $p(0) = p(\rho_0) \in H^{2k+3}$ . The isentropic case above is the one that  $\rho p'(\rho) = \gamma p$ . Hence the local existence of IBVP (1.11)-(1.16) follows from Theorem 1.1, provided the initial pressure  $p_0$  of (1.15) is in  $H^{2k+3}$ . Moreover, if  $p_0 = A\rho_0^\gamma$ , then the solution  $p$  of (1.12) satisfies the equation of state  $p = A\rho^\gamma$ . For this, see the proof of Theorem 4 of [4].

Now let us introduce the second result. Let  $B(0, r)$ ,  $r > 0$  be the ball with radius  $r$  centered at the origin.

**Theorem 1.4.** *If  $(\rho_0, u_0, f)$  is a radially symmetric data satisfying the conditions in Theorem 1.1 with  $p_0 = A\rho_0^\gamma \in H^{2k+3}$  on an annular domain  $\Omega = B(0, b) \setminus B(0, a)$  with  $0 < a < b \leq \infty$ , then there exists a unique global radially symmetric solution  $(\rho, p, u)$  of IBVP (1.11)-(1.16) satisfying the regularity (1.10) and the state equation  $p = A\rho^\gamma$  with  $\Omega = B(0, b) \setminus B(0, a)$ .*

*Remark 1.5.* In this isentropic case, dividing the density and pressure by a large constant, we can remove the smallness condition of initial density. But instead, we need a large viscosity coefficients.

This paper is organized as follows. In Section 2, we reduce the original problem to a linearized problem on a bounded domain under a positive lower bound assumption on the initial density. In Section 3, we prove uniform a priori estimates independent of domain size and lower bound of density. In Section 4, to prove Theorem 1.1 expanding domain size, we remove the lower bound restriction on the density. At this point, we solve the  $(k+1)$ -coupled elliptic system under a smallness assumption on the initial density. Then we prove Theorem 1.1 by constructing a sequence of approximate solutions and showing a strong convergence of it. Finally, Section 6 is devoted to proving Theorem 1.4.

## 2. LINEARIZATION

In this section, we introduce some existence and regularity results on solutions of a linear parabolic system and a linear transport equation. Then assuming the

positivity of the initial density, we linearize the original equations on a bounded domain.

Let  $\Omega$  be a bounded domain in  $\mathbf{R}^3$  with smooth boundary, and we consider the following linear parabolic problem

$$(2.1) \quad \begin{cases} \rho w_t + Lw = F & \text{in } (0, T) \times \Omega, \\ w(0) = w_0 & \text{in } \Omega, \quad u = 0 & \text{on } (0, T) \times \partial\Omega, \end{cases}$$

where  $\rho$  is a known scalar field in  $(0, T) \times \Omega$  such that

$$(2.2) \quad \rho^{(j)} \in C([0, T]; H^{2(k-j)+3}) \quad \text{and} \quad \rho \geq \delta \quad \text{on } [0, T] \times \bar{\Omega}$$

for any  $0 \leq j \leq k$ ,  $k \geq 1$  and for some constant  $\delta > 0$ . Then applying the arguments in the proof of Lemma 2 and Lemma 3 in [4] and induction, one can show the following existence and regularity results on solutions to the linear parabolic problem (2.1). See also [27, 28, 29].

**Lemma 2.1.** *If  $w_0 \in H_0^1 \cap H^{2k+3}$ ,  $F^{(j)} \in C([0, T]; H^{2(k-j)+1}) \cap L^2(0, T; H^{2(k-j)+2})$ ,  $F^{(k+1)} \in L^2(0, T; L^2)$  and*

$$(2.3) \quad \begin{aligned} & \rho(0)^{-1} \left( (F^{(j)}(0) - Lw^{(j)}(0) - \sum_{0 \leq l \leq j-1} \binom{j}{l} \rho^{(j-l)}(0) w^{(l+1)}(0) \right) \\ & \in H_0^1 \cap H^{2(k-j)+1} \end{aligned}$$

for all  $0 \leq j \leq k$ ,  $k \geq 1$ , then the solution  $w$  also satisfies

$$\begin{aligned} & w^{(j)} \in C([0, T]; H^{2(k-j)+3}) \quad \text{for all } 0 \leq j \leq k, \\ & w^{(k+1)} \in C([0, T]; H_0^1) \cap L^2(0, T; H^2), \quad \text{and} \quad w^{(k+2)} \in L^2(0, T; L^2). \end{aligned}$$

*Remark 2.2.* From the elliptic regularity theory [1, 5], we see that the condition (2.3) means  $w^{(j)}(0) \in H_0^1 \cap H^{2(k-j)+3}$ .

Now we consider the following linear hyperbolic equation

$$(2.4) \quad \rho_t + \operatorname{div}(\rho v) = 0 \quad \text{in } (0, T) \times \Omega, \quad \rho(0) = \rho_0 \quad \text{in } \Omega,$$

where  $v$  is a known vector field in  $(0, T) \times \Omega$  such that for some integer  $k \geq 1$

$$v^{(j)} \in C([0, T]; D_0^1 \cap D^{2(k-j)+3}) \cap L^2(0, T; D^{2(k-j)+4})$$

for any  $0 \leq j \leq k$ . Let  $\rho^{(j+1)}(0)$  be defined recursively by

$$\rho^{(j+1)}(0) = -\operatorname{div} \left( \sum_{0 \leq l \leq j} \binom{j}{l} \rho^{(j-l)}(0) v^{(l)}(0) \right), \quad 0 \leq j \leq k-1.$$

Then we have

**Lemma 2.3.** *Assume that  $\rho^{(j)}(0) \in H^{2(k-j)+3}$  and  $\rho_0 \geq 0$  in  $\Omega$ . Then*

- (i) there exists a unique solution  $\rho$  to the problem (2.4) such that for all  $0 \leq j \leq k$

$$\rho^{(j)} \in C([0, T]; H^{2(k-j)+3}),$$

- (ii) the solutions  $\rho$  and  $p$  satisfies the following estimate

$$|\rho(t)|_{H^{2k+3}} \leq |\rho_0|_{H^{2k+3}} \exp \left( C \int_0^t |v(s)|_{D_0^1 \cap D^{2k+4}} ds \right)$$

for  $0 \leq t \leq T$  and finally,

- (iii) the solution  $\rho$  is represented by the formula

$$(2.5) \quad \rho(t, x) = \rho_0(U(0, t, x)) \exp \left[ - \int_0^t \operatorname{div} v(s, U(s, t, x)) ds \right],$$

where  $U \in C([0, T] \times [0, T] \times \bar{\Omega})$  is the solution to the initial value problem

$$(2.6) \quad \begin{cases} \frac{\partial}{\partial t} U(t, s, x) = v(t, U(t, s, x)), & 0 \leq t \leq T, \\ U(s, s, x) = x, & 0 \leq s \leq T, \quad x \in \bar{\Omega}. \end{cases}$$

*Proof.* (i) with  $j = 0$ , (ii) and (iii) follow from Lemma 2.1 in [4]. And an induction on  $j$  yields the case (i) with  $j \geq 1$ .  $\square$

To prove Theorem 1.1, we consider the following linearized problem

$$(2.7) \quad \rho_t + \operatorname{div}(\rho v) = 0 \quad \text{in } (0, T) \times \Omega,$$

$$(2.8) \quad \rho u_t + Lu + \nabla p = \rho(f - v \cdot \nabla v) \quad \text{in } (0, T) \times \Omega,$$

$$(2.9) \quad (\rho, u)|_{t=0} = (\rho_0, u_0) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } (0, T) \times \partial\Omega,$$

where  $v$  is a known vector field in  $(0, T) \times \Omega$  such that for  $0 \leq j \leq k$

$$(2.10) \quad \begin{aligned} v^{(j)} &\in C([0, T]; D_0^1 \cap D^{2(k-j)+3}) \cap L^2(0, T; D^{2(k-j)+4}) \\ \text{and } v^{(k+1)} &\in L^\infty(0, T; D_0^1) \cap L^2(0, T; D^2). \end{aligned}$$

Recall again that  $Lu = -\mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u$ .

First, from Lemma 2.1 and Lemma 2.2, we obtain an existence result for positive initial densities.

**Lemma 2.4.** *Let  $\Omega$  be a bounded domain in  $\mathbf{R}^3$  with smooth boundary. In addition to (1.9) and (2.10), we assume that  $\rho_0 \geq \delta$  in  $\Omega$  for some constant  $\delta > 0$  and*

$$(2.11) \quad \begin{aligned} &(f - v \cdot \nabla v)^{(j)}(0) + \rho_0^{-1} \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)}(0) (f - v \cdot \nabla v)^{(m)}(0) \\ &- \rho_0^{-1} \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)}(0) u^{(m)}(0) - \rho_0^{-1} (Lu^{(j)}(0) + \nabla p^{(j)}(0)) \\ &\in H_0^1 \cap H^{2(k-j)+1} \end{aligned}$$



for  $0 \leq j \leq k$ . Then there exists a unique solution  $(\rho, u)$  to the linearized problem (2.7), (2.8) and (2.9) such that for all  $0 \leq j \leq k+1$

$$(2.12) \quad \begin{aligned} & \rho \in C(0, T]; H^{2k+3}), \\ & \rho^{(j)} \in C([0, T]; H^{2(k-j)+2}) \cap L^2(0, T; H^{2(k-j)+3}), \\ & u^{(j)} \in C([0, T]; H_0^1 \cap H^{2(k-j)+3}) \cap L^2(0, T; H^{2(k-j)+4}), \\ & \text{and } \rho \geq \underline{\delta} \text{ on } [0, T] \times \bar{\Omega} \end{aligned}$$

for some constant  $\underline{\delta} > 0$ .

*Proof.* The existence and regularity of a unique solution  $\rho$  to the linear hyperbolic problem (2.7) and (2.9) were already proved in Lemma 2.3. To prove the remaining part of the lemma, let us define  $F$  by  $\rho(f - v \cdot \nabla v) - \nabla p$ . Then

$$\begin{aligned} F^{(j)} &= \rho(f - v \cdot \nabla v)^{(j)} + \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \\ &\quad - \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)} u^{(m)} - \nabla p^{(j)} \end{aligned}$$

for  $1 \leq j \leq k$ . Then by virtue of (1.9), (2.10) and the regularity of  $\rho$ , we can easily show that  $F^{(j)} \in C([0, T]; H^{2(k-j)+1}) \cap L^2(0, T; H^{2(k-j)+2})$ ,  $F^{(k+1)} \in L^2(0, T; L^2)$ . Moreover since  $F$  and  $u$  satisfies the condition (2.3) by (2.11), Lemma 2.1 allows us to deduce the existence and regularity of a unique solution  $u$  to the linear parabolic problem (2.8) and (2.9). This completes the proof of Lemma 2.4.  $\square$

### 3. A PRIORI ESTIMATES FOR POSITIVE DENSITY

Assume that  $\rho_0, u_0, v, f$  and  $\Omega$  satisfy the hypotheses of Lemma 2.4. Then it follows from Lemma 2.4 that there exists a unique smooth solution  $(\rho, u)$  to the linear problem (2.7), (2.8) and (2.9) satisfying the regularity (2.12). We derive some *local (in time) a priori estimates* for  $(\rho, u)$  which are independent of the lower bound  $\delta$  of  $\rho_0$  and size of the domain  $\Omega$ . Let us choose constants  $\varepsilon > 0$  and  $c_0 > 1$  such that  $\varepsilon \leq c_0$

$$(3.1) \quad |\rho_0|_{L^1 \cap H^{2k+3}} < \varepsilon, \quad 1 + \sum_{1 \leq j \leq k+1} |g_j|_{D_0^1} < c_0,$$

where  $g_1 = f(0) - v(0) \cdot \nabla v(0) - \rho_0^{-1} (Lu_0 + \nabla p_0) = u_t(0) \in D_0^1 \cap D^{2k+1}$  and for  $1 \leq j \leq k$

$$\begin{aligned} g_{j+1} &= (f - v \cdot \nabla v)^{(j)}(0) + \rho_0^{-1} \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)}(0) (f - v \cdot \nabla v)^{(m)}(0) \\ &\quad - \rho_0^{-1} \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)}(0) g_{m+1} - \rho_0^{-1} (Lg_j + \nabla p^{(j)}(0)) \\ &= u^{(j+1)}(0) \in D_0^1 \cap D^{2(k-j)+1}. \end{aligned}$$

Assume for all  $0 \leq j \leq k+1$  that

$$(3.2) \quad |v^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} \leq 1 + c_{0,1,j},$$

and for all  $0 \leq l \leq k$  and  $0 \leq j \leq k-l$  that

$$(3.3) \quad \begin{aligned} & \sup_{0 \leq t \leq T_*} \left( |v^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} \right) + \int_0^{T_*} |v^{(j)}(t)|_{D^2 \cap D^{2l+2}}^2 dt \leq 1 + c_{l,2,j}, \\ & \sup_{0 \leq t \leq T_*} \left( |v^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} \right) \\ & + \int_0^{T_*} \left( |v^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}}^2 + |v^{(j)}(t)|_{D^{2l+3}}^2 \right) dt \leq 1 + c_{l,3,j}, \\ & \text{ess sup}_{0 < t < T_*} \left( |v^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}} + |v^{(j)}(t)|_{D^{2l+3}} \right) \\ & + \int_0^{T_*} \left( |v^{(j+1)}(t)|_{D^{2l+2}}^2 + |v^{(j)}(t)|_{D^{2l+4}}^2 \right) dt \leq 1 + c_{l,4,j} \end{aligned}$$

for some time  $T_* \in (0, T)$  and constants  $c_{l,i,j}$  such that

$$1 \leq c_{l,i,j} \leq c_{l+1,i,j} \quad \text{for any } i, j, l$$

$$c_{l,i,j} \leq c_{l,i+1,j} \quad \text{for each fixed } l, j,$$

$$\text{and } c_{l,i,j} \leq c_{l,i,l+1} \quad \text{for each fixed } l \text{ and for any } i.$$

The constants  $c_{l,i,j}$ 's and  $T_*$  will be determined later and depend only on  $c_0$  and the parameters of  $C$ .

**Lemma 3.1.** *For each  $1 \leq j \leq k+1$*

$$|\rho^{(j)}(0)|_{L^{\frac{3}{2}} \cap H^{2(k-j)+3}} \leq C \varepsilon c_{0,1,j-1}^j.$$

*Proof.* Using the equation

$$(3.4) \quad \rho^{(j+1)} = - \sum_{0 \leq m \leq j} \binom{j}{m} \text{div}(\rho^{(j-m)} v^{(m)}),$$

for any multi-index  $\alpha = (\alpha_1, \alpha_2, \alpha_3)$  with  $|\alpha| = \alpha_1 + \alpha_2 + \alpha_3 \leq 2(k - (j+1)) + 3$  we have

$$\begin{aligned} & |D^\alpha \rho^{(j+1)}(0)|_{L^2} \\ & \leq C \sum_{\substack{0 \leq m \leq j \\ \alpha_1 + \alpha_2 = \alpha}} \left( |\nabla D^{\alpha_1} \rho^{(j-m)}(0) \cdot D^{\alpha_2} v^{(m)}(0)|_{L^2} \right. \\ & \quad \left. + |D^{\alpha_1} \rho^{(j-m)}(0) \text{div} D^{\alpha_2} v^{(m)}(0)|_{L^2} \right) \\ & \leq C \sum_{\substack{0 \leq m \leq j \\ \alpha_1 + \alpha_2 = \alpha}} |\rho^{(j-m)}(0)|_{H^{|\alpha_1|+2}} |v^{(m)}(0)|_{D_0^1 \cap D^{|\alpha_2|+1}} \\ & \leq C \sum_{\substack{0 \leq m \leq j \\ \alpha_1 + \alpha_2 = \alpha}} |\rho^{(j-m)}(0)|_{H^{2(k-j)+3}} |v^{(m)}(0)|_{D_0^1 \cap D^{2(k-j)+2}}, \end{aligned}$$

where  $D^\alpha = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \partial x_3^{\alpha_3}}$ . Since  $0 \leq m \leq j$ ,  $2(k-j)+3 = 2(k-(j-m))+3-2m \leq 2(k-(j-m))+3$  and  $2(k-j)+2 \leq 2(k-m)+2-2(j-m) \leq 2(k-m)+2$ . Thus by the initial bound (3.1), we obtain

$$\begin{aligned} & |\rho^{(j+1)}(0)|_{H^{2(k-(j+1))+3}} \\ & \leq C \sum_{0 \leq m \leq j} |\rho^{(j-m)}(0)|_{H^{2(k-(j-m))+3}} |v^{(m)}(0)|_{D_0^1 \cap D^{2(k-m)+2}} \\ & \leq C c_{0,1,j} \sum_{0 \leq m \leq j} |\rho^{(j-m)}(0)|_{H^{2(k-(j-m))+3}}. \end{aligned}$$

Therefore the induction on  $j$  yields that

$$|\rho^{(j)}(0)|_{H^{2(k-j)+3}} \leq C \varepsilon c_{0,1,j-1}^j$$

for all  $1 \leq j \leq k+1$ . Similarly, we can treat  $L^{\frac{3}{2}}$  norm.  $\square$

**Lemma 3.2.** *If  $\varepsilon$  is sufficiently small, then we have*

$$\sum_{0 \leq j \leq k+1} |u^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} \leq C c_0.$$

*Proof.* We first consider the case  $1 \leq j \leq k$ . Since  $g_j \in D_0^1$  satisfies the following boundary value problem:

$$\begin{aligned} L(u^{(j)}(0)) &= \rho_0(f - v \cdot \nabla v)^{(j)}(0) + \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)}(0) (f - v \cdot \nabla v)^{(m)}(0) \\ &+ \sum_{0 \leq m \leq j-1} \binom{j}{m} \rho^{(j-m)}(0) u^{(m+1)}(0) - \nabla p^{(j)}(0) - \rho_0 g_{j+1}, \end{aligned}$$

we have

$$\begin{aligned} |u^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} &\leq C |u^{(j)}(0)|_{D_0^1} + C |\rho_0(f - v \cdot \nabla v)^{(j)}(0)|_{H^{2(k-j)+1}} \\ &+ C \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}(0) (f - v \cdot \nabla v)^{(m)}(0)|_{H^{2(k-j)+1}} \\ &+ C \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}(0) u^{(m+1)}(0)|_{H^{2(k-j)+1}} \\ &+ C |p^{(j)}(0)|_{H^{2(k-j)+2}} + C |\rho_0 g_{j+1}|_{H^{2(k-j)+1}} \end{aligned}$$

from the elliptic regularity results [1, 5], where  $C$  does not depend on the size of domain. We easily show that

$$\begin{aligned} & |\rho_0(f - v \cdot \nabla v)^{(j)}(0)|_{H^{2(k-j)+1}} \\ & \leq C |\rho_0|_{H^{2(k-j)+3}} |f^{(j)}(0)|_{H^{2(k-j)+1}} \\ & \quad + C \sum_{0 \leq m \leq j} |\rho_0|_{H^{2(k-j)+3}} |v^{(j-m)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} |v^{(m)}(0)|_{D_0^1 \cap D^{2(k-j)+2}} \\ & \leq C \varepsilon c_{0,1,j}^2. \end{aligned}$$

Similarly, we have

$$\begin{aligned} & \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}(0)(f - v \cdot \nabla v)^{(m)}(0)|_{H^{2(k-j)+1}} \\ & \leq C \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}(0)|_{H^{2(k-j)+3}} |(f - v \cdot \nabla v)^{(m)}(0)|_{H^{2(k-j)+1}} \leq C\varepsilon c_{0,1,j-1}^{j+2}, \end{aligned}$$

$$|p^{(j)}(0)|_{H^{2(k-j)+2}} \leq C\varepsilon c_{0,1,j-1}^j,$$

$$|\rho_0 g_{j+1}|_{H^{2(k-j)+1}} \leq C|\rho_0|_{H^{2(k-j)+3}} |g_{j+1}|_{H^{2(k-j)+1}} \leq C\varepsilon |g_{j+1}|_{D_0^1 \cap D^{2(k-(j+1))+3}},$$

and

$$\begin{aligned} & \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}(0)u^{(m+1)}(0)|_{H^{2(k-j)+1}} \\ & \leq C \sum_{0 \leq m \leq j} |\rho^{(j-m)}(0)|_{H^{2(k-j)+1}} |u^{(m+1)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} \\ & \leq C\varepsilon c_{0,1,j-1}^j A_{j-1} + C\varepsilon c_{0,1,0} |u^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}}, \end{aligned}$$

where

$$A_j = \sum_{1 \leq m \leq j} |u^{(m)}(0)|_{D_0^1 \cap D^{2(k-m)+3}} \quad \text{and} \quad A_0 = 0.$$

Thus if we choose  $\varepsilon < \frac{1}{2Mc_{0,1,k}^{k+2}}$  for some large  $M \geq C$ , then since  $g_j \in D_0^1$ , we obtain

$$\begin{aligned} A_j & \leq C \sum_{1 \leq j \leq k+1} |g_j|_{D_0^1} + C\varepsilon |g_{j+1}|_{D_0^1 \cap D^{2(k-(j+1))+3}} \\ & \leq Cc_0 + C\varepsilon |g_{j+1}|_{D_0^1 \cap D^{2(k-(j+1))+3}} \end{aligned}$$

for all  $1 \leq j \leq k$ . Since  $|g_{k+1}|_{D_0^1} \leq c_0$ , we have  $G_k \leq Cc_0$ .

Now using the energy estimate for  $u_0$ , we easily get  $|u_0|_{D_0^1} \leq Cc_0$  for small  $\varepsilon$ . By the another use of elliptic regularity result, we finally have for  $j = 0$  that

$$\begin{aligned} |u_0|_{D_0^1 \cap D^{2k+3}} & \leq C|u_0|_{D_0^1} + C|\rho_0(f(0) - v(0) \cdot \nabla v(0))|_{H^{2k+1}} \\ & \quad + C|p_0|_{H^{2k+2}} + C|\rho_0 g_1|_{H^{2k+1}} \\ & \leq Cc_0 + C\varepsilon c_{0,1,0}^2 + C\varepsilon c_0 \leq Cc_0. \end{aligned}$$

This completes the proof of the lemma.  $\square$

*Remark 3.3.* Since the estimate of Lemma 3.2 is stable under a small perturbation of  $g_j$ , this lemma will be used for the reconstruction of initial data in Lemma 4.2 below. If the initial data has positive lower bound, then this lemma is not necessary.

**Lemma 3.4.** *For any fixed  $\delta > 0$  we have*

$$\inf_{\Omega} \rho(t) \geq C^{-1}\delta,$$

and for each  $1 \leq j \leq k+1$

$$|\rho^{(j)}(t)|_{H^{2(k-j)+3}} \leq C\varepsilon c_{0,1,j-1}^j, \quad \int_0^t |\rho^{(k+2)}(s)|_{L^2}^2 ds \leq C\varepsilon^2,$$

if  $0 \leq t \leq \min(T_*, T_1)$  where  $T_1 = (1 + c_{k,4,0})^{-(2k+6)}$ .

*Proof.* From Lemma 2.3, we recall that

$$|\rho(t)|_{H^{2k+3}} \leq |\rho_0|_{H^{2k+3}} \exp \left( C \int_0^t |v(s)|_{D_0^1 \cap D^{2k+4}} ds \right)$$

and

$$\inf_{\Omega} \rho(t) \geq \left( \inf_{\Omega} \rho_0 \right) \exp \left( -C \int_0^t |v(s)|_{D_0^1 \cap D^{2k+4}} ds \right)$$

for  $0 \leq t \leq T$ . Hence from the observation that

$$\begin{aligned} \int_0^t |v(s)|_{D_0^1 \cap D^{2k+4}} ds &\leq t^{\frac{1}{2}} \left( \int_0^t |v(s)|_{D_0^1 \cap D^{2k+4}}^2 ds \right)^{\frac{1}{2}} \\ &\leq C(1 + c_{k,4,0})t + C((1 + c_{k,4,0})t)^{\frac{1}{2}}, \end{aligned}$$

we obtain the desired estimate for  $j = 0$ . For the estimate of the case  $j \geq 1$ , multiplying (3.4) by  $D^\alpha \rho^{(j)}$  for any multi-index  $\alpha$  with  $|\alpha| \leq 2(k-j) + 3$ , we have

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} |D^\alpha \rho^{(j)}|_{L^2}^2 \\ &= - \int \operatorname{div} (D^\alpha \rho^{(j)} v) D^\alpha \rho^{(j)} dx \\ &\quad - \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \neq 0}} C_{\alpha_1, \alpha_2} \int \operatorname{div} (D^{\alpha_1} \rho^{(j)} D^{\alpha_2} v^{(m)}) D^\alpha \rho^{(j)} dx \\ &\quad - \sum_{\substack{1 \leq m \leq j \\ \alpha_1 + \alpha_2 = \alpha}} C_{j, \alpha} \int \operatorname{div} (D^{\alpha_1} \rho^{(j-m)} D^{\alpha_2} v^{(m)}) D^\alpha \rho^{(j)} dx \\ &= \sum_{1 \leq i \leq 3} J_i \end{aligned} \tag{3.5}$$

where

$$C_{\alpha_1, \alpha_2} = \frac{\alpha!}{\alpha_1! \alpha_2!}, \quad C_{j, \alpha} = \binom{j}{m} \frac{\alpha!}{\alpha_1! \alpha_2!}.$$

Each term can be estimated as follows:

$$\begin{aligned} J_1 &\leq C |\operatorname{div} v|_{D_0^1 \cap D^2} |D^\alpha \rho^{(j)}|_{L^2}^2, \\ J_2 &\leq C \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \neq 0}} \int \left( |\nabla D^{\alpha_1} \rho^{(j)}| |D^{\alpha_2} v| + |D^{\alpha_1} \rho^{(j)}| |D^{\alpha_2} \operatorname{div} v| \right) |D^\alpha \rho^{(j)}| dx \\ &\leq C \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \neq 0}} |v|_{D_0^1 \cap D^{|\alpha_2|+2}} |\rho^{(j)}|_{H^{|\alpha_1|+1}} |D^\alpha \rho^{(j)}|_{L^2}, \\ J_3 &\leq C \sum_{\substack{1 \leq m \leq j \\ \alpha_1 + \alpha_2 = \alpha}} |\rho^{(j-m)}|_{H^{|\alpha_1|+1}} |v^{(m)}|_{D_0^1 \cap D^{|\alpha_2|+2}} |D^\alpha \rho^{(j)}|_{L^2}. \end{aligned}$$

Substituting all these estimate into (3.5) and integrating over  $(0, t)$ , we have

$$\begin{aligned} |\rho^{(j)}(t)|_{H^{2(k-j)+3}} &\leq |\rho^{(j)}(0)|_{H^{2(k-j)+3}} + C \int_0^t |\operatorname{div} v|_{D_0^1 \cap D^2} |D^\alpha \rho^{(j)}|_{L^2} ds \\ &\quad + C \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \neq 0}} \int_0^t |v(s)|_{D_0^1 \cap D^{|\alpha_2|+2}} |\rho^{(j)}(s)|_{H^{|\alpha_1|+1}} ds \\ &\quad + C \sum_{\substack{1 \leq m \leq j \\ \alpha_1 + \alpha_2 = \alpha}} \int_0^t |\rho^{(j-m)}|_{H^{|\alpha_1|+2}} |v^{(m)}|_{D_0^1 \cap D^{|\alpha_2|+1}} ds . \end{aligned}$$

Since  $2(k-j) + 5 \leq 2k + 3$  for  $j \geq 1$ ,

$$2(k-j) + 5 \leq 2(k-(j-m)) + 5 - 2m \leq 2(k-(j-m)) + 3 \text{ for } m \geq 1$$

$$\text{and } 2(k-j) + 4 \leq 2(k-m) + 4 - 2(j-m) \leq 2(k-m) + 4 \text{ for } m \leq j,$$

we obtain

$$\begin{aligned} |\rho^{(j)}(t)|_{H^{2(k-j)+3}} &\leq C \varepsilon c_{0,1,j-1}^j + C c_{1,2,0} \int_0^t |\rho^{(j)}(s)|_{H^{2(k-j)+3}} ds \\ &\quad + C \max_{0 \leq m \leq j-1} \left( \sup_{0 \leq s \leq t} |\rho^{(m)}(s)|_{H^{2(k-m)+3}} \right) (1 + c_{k,4,0})^{\frac{1}{2}} t^{\frac{1}{2}} . \end{aligned}$$

Hence the Gronwall's inequality yields that if  $t \leq (1 + c_{k,4,0})^{-1}$

$$|\rho^{(j)}(t)|_{H^{2(k-j)+3}} \leq C \left( \varepsilon c_{0,1,j-1}^j + \max_{0 \leq m \leq j-1} \left( \sup_{0 \leq s \leq t} |\rho^{(m)}(s)|_{H^{2(k-m)+3}} \right) \right) .$$

Using the induction on  $j$  again, we have

$$|\rho^{(j)}(t)|_{H^{2(k-j)+3}} \leq C \varepsilon c_{0,1,j-1}^j$$

for all  $t \in [0, \min(T_*, T_1)]$ . Similarly (more easily), we can estimate  $|\rho^{(j)}|_{L^1}$ .

Finally we have

$$\begin{aligned} |\rho^{(k+2)}|_{L^2} &\leq C \sum_{0 \leq m \leq k+1} \left( |\nabla \rho^{(k+1-m)} \cdot v^{(m)}|_{L^2} + |\rho^{(k+1-m)} v|_{L^2} \right) \\ &\leq C \sum_{1 \leq m \leq k+1} |\rho^{(k+1-m)}|_{H^2} |\nabla v^{(m)}|_{L^2} + C |\rho^{(k+1)}|_{H^1} |v|_{D_0^1 \cap D^2} \\ &\leq C \varepsilon c_{0,1,k-1}^k c_{0,4,k} + C c_{0,1,k}^{k+1} c_{0,3,0} \end{aligned}$$

and hence

$$\int_0^t |\rho^{(k+2)}(s)|_{L^2}^2 ds \leq C \varepsilon^2 .$$

□

Using the equation  $p_t + \nabla p \cdot u + \rho p'(\rho) \operatorname{div} u = 0$ , by the similar estimates as in the proof of Lemma 3.1 and Lemma 3.4 we have

**Lemma 3.5.** *For any  $0 \leq j \leq k+1$ , we have*

$$|(p - p(0))^{(j)}(t)|_{L^1 \cap H^{2(k-j)+3}} \leq C \varepsilon c_{0,1,j-1}^j$$

for  $0 \leq t \leq \min(T_*, T_1)$  where  $T_1 = (1 + c_{k,4,0})^{-(2k+6)}$ , where  $p(0)$  is the value of  $p$  at point zero.

The three lemmas below, Lemma 3.6, 3.7 and 3.8, are similar to those in [4]. But since they are needed for a priori estimates for high regularity, we revisit the proof of them.

**Lemma 3.6.**

$$\int_0^t |\sqrt{\rho}u_t(s)|_{L^2}^2 ds + |u(t)|_{D_0^1}^2 \leq Cc_0^2, \quad \int_0^t |u(s)|_{D^2}^2 ds \leq Cc_0^3$$

for  $0 \leq t \leq \min(T_*, T_1)$ .

*Proof.* Multiplying the equation (2.8) by  $u_t$  and integrating over  $\Omega$ , we obtain

$$\begin{aligned} & \int \rho |u_t|^2 dx + \frac{1}{2} \frac{d}{dt} \int \mu |\nabla u|^2 + (\lambda + \mu) (\operatorname{div} u)^2 dx \\ (3.6) \quad & = - \int \nabla p \cdot u_t dx + \int \rho (f - v \cdot \nabla v) \cdot u_t dx. \end{aligned}$$

Using the condition  $\varepsilon \leq c_0$  and Lemma 3.4 together with (3.3), we can estimate the second term of the right hand side in (3.14) as follows:

$$\begin{aligned} & \int \rho (f - v \cdot \nabla v) \cdot u_t dx \leq |\rho|_{L^\infty}^{\frac{1}{2}} |f - v \cdot \nabla v|_{L^2} |\sqrt{\rho}u_t|_{L^2} \\ & \leq C |\rho|_{L^\infty} \left( |f|_{L^2}^2 + |v|_{D_0^1 \cap D^2}^4 \right) + \frac{1}{2} |\sqrt{\rho}u_t|_{L^2}^2 \\ & \leq Cc_0c_{0,3,0}^4 + \frac{1}{2} |\sqrt{\rho}u_t|_{L^2}^2. \end{aligned}$$

To estimate the first term, we observe from Lemma 3.5 that

$$\begin{aligned} & - \int \nabla p \cdot u_t dx = \int (p - p(0)) \operatorname{div} u_t dx \\ & = \frac{d}{dt} \int (p - p(0)) \operatorname{div} u dx - \int p_t \operatorname{div} u dx, \\ & \int (p - p(0)) \operatorname{div} u dx \leq C |p(\rho) - p(0)|_{L^2}^2 + \frac{\mu}{4} |\nabla u|_{L^2}^2 \leq Cc_0^2 + \frac{\mu}{4} |\nabla u|_{L^2}^2 \end{aligned}$$

and

$$- \int p_t \operatorname{div} u dx \leq |p_t|_{L^2}^2 + |\nabla u|_{L^2}^2 \leq Cc_0^2c_{0,1,0}^2 + |\nabla u|_{L^2}^2.$$

Hence integrating (3.14) in time over  $(0, t)$ , we have

$$\begin{aligned} & \int_0^t |\sqrt{\rho}u_t(s)|_{L^2}^2 ds + |\nabla u(t)|_{L^2}^2 \\ & \leq Cc_0^2 (1 + |\nabla u_0|_{L^2}^2) + Cc_0^2c_{0,3,0}^4 t + C \int_0^t |\nabla u(s)|_{L^2}^2 ds \end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_1)$ . Therefore, in view of Gronwall's inequality, we conclude that

$$\int_0^t |\sqrt{\rho}u_t(s)|_{L^2}^2 ds + |\nabla u(t)|_{L^2}^2 \leq Cc_0^2 \quad \text{for } 0 \leq t \leq \min(T_*, T_1).$$

Moreover, since for each  $t \in (0, T)$ ,  $u = u(t) \in D_0^1 \cap D^2$  is a solution of the elliptic system

$$Lu = -\nabla p + \rho(f - v \cdot \nabla v) - \rho u_t \quad \text{in } \Omega,$$

it follows from the elliptic regularity result in [1, 5] that

$$\begin{aligned} |u|_{D^2} &\leq C \left( |-\nabla p + \rho(f - v \cdot \nabla v) - \rho u_t|_{L^2} + |u|_{D_0^1} \right) \\ &\leq C \left( c_{0,3,0}^2 + c_0^{\frac{1}{2}} |\sqrt{\rho} u_t|_{L^2} \right) \end{aligned}$$

and thus

$$\int_0^t |u(s)|_{D^2}^2 ds \leq C c_0^3 \quad \text{for } 0 \leq t \leq \min(T_*, T_1).$$

This completes the proof of Lemma 3.6.  $\square$

**Lemma 3.7.**

$$|\sqrt{\rho} u_t(t)|_{L^2} + |u(t)|_{D^2} + \int_0^t \left( |u_t(s)|_{D_0^1}^2 + |u(s)|_{D^3}^2 \right) ds \leq C c_0^3 c_{0,2,0}^{\frac{3}{2}} c_{0,3,0}^{\frac{1}{2}}$$

for  $0 \leq t \leq \min(T_*, T_1)$ .

*Proof.* We differentiate (2.8) with respect to  $t$  and have

$$(3.7) \quad \rho u_{tt} + Lu_t + \nabla p_t = \rho(f - v \cdot \nabla v)_t + \rho_t(f - v \cdot \nabla v - u_t).$$

Multiplying this by  $u_t$  and integrating over  $\Omega$ , we obtain

$$(3.8) \quad \begin{aligned} &\frac{1}{2} \frac{d}{dt} \int \rho |u_t|^2 dx + \int \mu |\nabla u_t|^2 + (\lambda + \mu) (\operatorname{div} u_t)^2 dx \\ &= \int \left( -\nabla p_t + \rho(f - v \cdot \nabla v)_t + \rho_t(f - v \cdot \nabla v - \frac{1}{2} u_t) \right) \cdot u_t dx. \end{aligned}$$

To estimate each term in the right hand side of (3.8), we follow the arguments in [3, 5, 6]; we first apply the standard inequalities such as Hölder, Sobolev and Young's inequalities and Lemma 3.4.

$$-\int \nabla p_t \cdot u_t dx = \int p_t \operatorname{div} u_t dx \leq C |p_t|_{L^2}^2 + \frac{\mu}{8} |\nabla u_t|_{L^2}^2 \leq C c_0^2 c_{0,1,0}^2 + \frac{\mu}{8} |\nabla u_t|_{L^2}^2,$$

$$\int \rho f_t \cdot u_t dx \leq |f_t|_{L^2} |\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho} u_t|_{L^2} \leq |f_t|_{L^2}^2 + C c_0 |\sqrt{\rho} u_t|_{L^2}^2,$$

$$\begin{aligned} -\int \rho(v \cdot \nabla v)_t \cdot u_t dx &\leq C |\rho|_{L^\infty}^{\frac{1}{2}} |v_t|_{D_0^1} |v|_{D_0^1} |\sqrt{\rho} u_t|_{L^3} \\ &\leq C |\rho|_{L^\infty}^{\frac{3}{4}} |v_t|_{D_0^1} |v|_{D_0^1} |\sqrt{\rho} u_t|_{L^2}^{\frac{1}{2}} |\nabla u_t|_{L^2}^{\frac{1}{2}} \\ &\leq \eta^{-2} C |\rho|_{L^\infty}^3 |v|_{D_0^1}^4 |\sqrt{\rho} u_t|_{L^2}^2 + \eta |v_t|_{D_0^1}^2 + \frac{\mu}{8} |\nabla u_t|_{L^2}^2 \\ &\leq \eta^{-2} C c_{0,2,0}^7 |\sqrt{\rho} u_t|_{L^2}^2 + \eta |v_t|_{D_0^1}^2 + \frac{\mu}{8} |\nabla u_t|_{L^2}^2, \end{aligned}$$



$$\begin{aligned}
\int \rho_t (f - v \cdot \nabla v) \cdot u_t \, dx &\leq C |\rho_t|_{H^1} \left( |f|_{L^2} + |v|_{D_0^1 \cap D^2}^2 \right) |\nabla u_t|_{L^2} \\
&\leq C c_0^2 c_{0,1,0}^2 \left( |f|_{L^2}^2 + c_{0,3,0}^4 \right) + \frac{\mu}{8} |\nabla u_t|_{L^2}^2 \\
&\leq C c_{0,3,0}^8 + \frac{\mu}{8} |\nabla u_t|_{L^2}^2
\end{aligned}$$

and finally

$$\begin{aligned}
-\int \rho_t \left( \frac{1}{2} |u_t|^2 \right) \, dx &= \int \operatorname{div}(\rho v) \left( \frac{1}{2} |u_t|^2 \right) \, dx \\
&\leq \int \rho |v| |u_t| |\nabla u_t| \, dx \\
&\leq C |\rho|_{L^\infty}^{\frac{3}{4}} |v|_{D_0^1} |\sqrt{\rho} u_t|_{L^2}^{\frac{1}{2}} |\nabla u_t|_{L^2}^{\frac{3}{2}} \\
&\leq C c_{0,2,0}^7 |\sqrt{\rho} u_t|_{L^2}^2 + \frac{\mu}{8} |\nabla u_t|_{L^2}^2.
\end{aligned}$$

Here  $\eta \in (0, 1)$  is a small number. Substituting these estimates into (3.8) and taking  $\eta = (1 + c_{0,3,0})^{-1}$ , we have

$$\begin{aligned}
&\frac{d}{dt} \int \rho |u_t|^2 \, dx + \mu \int |\nabla u_t|^2 \, dx \\
(3.9) \quad &\leq C \left( |f_t|_{L^2}^2 + c_{0,3,0}^8 \right) + C c_{0,3,0}^9 |\sqrt{\rho} u_t|_{L^2}^2 + (1 + c_{0,3,0})^{-1} |v_t|_{D_0^1}^2
\end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_1)$ . On the other hand, since  $u_t(0) = g_2$ , it follows that

$$(3.10) \quad |\sqrt{\rho} u_t(0)|_{L^2} + |u_t(0)|_{D_0^1} \leq C c_0^3.$$

Hence integrating (3.9) over  $(0, t)$ , we also have

$$|\sqrt{\rho} u_t(t)|_{L^2}^2 + \int_0^t |\nabla u_t(s)|_{L^2}^2 \, ds \leq C \left( c_0^3 + c_{0,3,0}^8 t \right) + C c_{0,3,0}^9 \int_0^t |\sqrt{\rho} u_t(s)|_{L^2}^2 \, ds.$$

Therefore, in view of Gronwall's inequality, we conclude that

$$|\sqrt{\rho} u_t(t)|_{L^2}^2 + \int_0^t |u_t(s)|_{D_0^1}^2 \, ds \leq C c_0^3 \quad \text{for } 0 \leq t \leq \min(T_*, T_1).$$

Moreover, since for each  $t \in (0, T)$ ,  $u = u(t) \in D_0^1 \cap D^3$  is a solution of the elliptic system

$$Lu = -\nabla p + \rho(f - v \cdot \nabla v) - \rho u_t \quad \text{in } \Omega,$$

it follows from the elliptic regularity result in [1, 5] that

$$\begin{aligned}
|u(t)|_{D^2} &\leq C c_0^2 (1 + |v \cdot \nabla v|_{L^2}) \\
&\leq C c_0^2 \left( 1 + |v|_{D_0^1}^{\frac{3}{2}} |v|_{D_0^1 \cap D^2}^{\frac{1}{2}} \right) \leq C c_0^2 c_{0,2,0}^{\frac{3}{2}} c_{0,3,0}^{\frac{1}{2}}
\end{aligned}$$

and

$$\int_0^t |u(s)|_{D^3}^2 \, ds \leq C c_0^2 \int_0^t \left( 1 + |v(s)|_{D_0^1 \cap D^2}^4 + |u_t(s)|_{D_0^1}^2 \right) \, ds \leq C c_0^5$$

for  $0 \leq t \leq \min(T_*, T_1)$ . This completes the proof of Lemma 3.7.  $\square$

**Lemma 3.8.**

$$|u_t(t)|_{D_0^1}^2 + |u(t)|_{D^3}^2 + \int_0^t (|\sqrt{\rho}u_{tt}(s)|_{L^2}^2 + |u_t(s)|_{D^2}^2 + |u(s)|_{D^4}^2) ds \leq Cc_0^7c_{0,3,0}^{12}$$

for  $0 \leq t \leq \min(T_*, T_1)$ .

*Proof.* Multiplying (3.7) by  $u_{tt}$  and integrating over  $\Omega$ , we have

$$(3.11) \quad \begin{aligned} & \int \rho |u_{tt}|^2 dx + \frac{1}{2} \frac{d}{dt} \int \mu |\nabla u_t|^2 + (\lambda + \mu) (\operatorname{div} u_t)^2 dx \\ &= \int (-\nabla p_t + \rho(f - v \cdot \nabla v)_t + \rho_t(f - v \cdot \nabla v - u_t)) \cdot u_{tt} dx \end{aligned}$$

We can estimate the first two terms in the right hand side of (3.11) as follows:

$$\begin{aligned} - \int \nabla p_t \cdot u_{tt} dx &= \int p_t \operatorname{div} u_{tt} dx = \frac{d}{dt} \int p_t \operatorname{div} u_t dx - \int p_{tt} \operatorname{div} u_t dx \\ &\leq \frac{d}{dt} \int p_t \operatorname{div} u_t dx + |p_{tt}|_{L^2}^2 + |\nabla u_t|_{L^2}^2 \end{aligned}$$

and

$$\begin{aligned} \int \rho(f - v \cdot \nabla v)_t \cdot u_{tt} dx &\leq C|\rho|_{L^\infty}^{\frac{1}{2}} \left( |f_t|_{L^2} + |v|_{D_0^1 \cap D^2} |v_t|_{D_0^1} \right) |\sqrt{\rho}u_{tt}|_{L^2} \\ &\leq Cc_0 \left( |f_t|_{L^2}^2 + c_{0,3,0}^2 |v_t|_{D_0^1}^2 \right) + \frac{1}{2} |\sqrt{\rho}u_{tt}|_{L^2}^2. \end{aligned}$$

To estimate the last term, we observe that

$$\begin{aligned} & \int \rho_t (f - v \cdot \nabla v) \cdot u_{tt} dx \\ &= \frac{d}{dt} \int \rho_t (f - v \cdot \nabla v) \cdot u_t dx - \int \rho_{tt} (f - v \cdot \nabla v) \cdot u_t dx \\ &\quad - \int \rho_t (f - v \cdot \nabla v)_t \cdot u_t dx \end{aligned}$$

and

$$- \int \rho_t u_t \cdot u_{tt} dx = - \frac{d}{dt} \int \rho_t \left( \frac{1}{2} |u_t|^2 \right) dx + \int \rho_{tt} \left( \frac{1}{2} |u_t|^2 \right) dx.$$

Then by virtue of Lemma 3.4, we obtain

$$\begin{aligned} - \int \rho_{tt} (f - v \cdot \nabla v) \cdot u_t dx &\leq C|\rho_{tt}|_{L^2} \left( |f|_{H^1} + |v|_{D_0^1 \cap D^2}^2 \right) |\nabla u_t|_{L^2} \\ &\leq Cc_{0,3,0}^4 |\rho_{tt}|_{L^2}^2 + |\nabla u_t|_{L^2}^2, \\ - \int \rho_t (f - v \cdot \nabla v)_t \cdot u_t dx &\leq C|\rho_t|_{L^3} \left( |f_t|_{L^2} + |v|_{D_0^1 \cap D^2} |v_t|_{D_0^1} \right) |\nabla u_t|_{L^2} \\ &\leq Cc_{0,3,0}^4 \left( |f_t|_{L^2}^2 + c_3^2 |v_t|_{D_0^1}^2 \right) + |\nabla u_t|_{L^2}^2 \end{aligned}$$

and

$$\begin{aligned}
\int \rho_{tt} \left( \frac{1}{2} |u_t|^2 \right) dx &= - \int \operatorname{div}(\rho_t v + \rho v_t) \left( \frac{1}{2} |u_t|^2 \right) dx \\
&\leq \int (|\rho_t| |v| + \rho |v_t|) |u_t| |\nabla u_t| dx \\
&\leq C c_{0,3,0}^3 |\nabla u_t|_{L^2}^2 + C c_0^{\frac{3}{4}} |v_t|_{D_0^1} |\sqrt{\rho} u_t|_{L^2}^{\frac{1}{2}} |\nabla u_t|_{L^2}^{\frac{3}{2}} \\
&\leq C c_{0,3,0}^3 |\nabla u_t|_{L^2}^2 + (1 + c_{0,3,0})^{-1} |v_t|_{D_0^1}^2 |\sqrt{\rho} u_t|_{L^2} |\nabla u_t|_{L^2} \\
&\leq C c_{0,3,0}^3 |u_t|_{D_0^1}^2 + (1 + c_{0,3,0})^{-1} |v_t|_{D_0^1}^2 (|\sqrt{\rho} u_t|_{L^2}^2 + |\nabla u_t|_{L^2}^2).
\end{aligned}$$

Substituting all the above estimates into (3.11), we have

$$\begin{aligned}
&\int \rho |u_{tt}|^2 dx + \frac{d}{dt} \int \mu |\nabla u_t|^2 + (\lambda + \mu) (\operatorname{div} u_t)^2 dx \\
&\leq \frac{d}{dt} \int (2p_t \operatorname{div} u_t + 2\rho_t (f - v \cdot \nabla v) \cdot u_t - \rho_t |u_t|^2) dx \\
(3.12) \quad &+ C \left( |p_{tt}|_{L^2}^2 + c_{0,3,0}^4 |\rho_{tt}|_{L^2}^2 + c_{0,3,0}^4 |f_t|_{L^2}^2 + c_{0,3,0}^6 |v_t|_{D_0^1}^2 + c_{0,3,0}^3 |u_t|_{D_0^1}^2 \right) \\
&\quad + |v_t|_{D_0^1}^2 |\sqrt{\rho} u_t|_{L^2}^2 + (1 + c_{0,3,0})^{-1} |v_t|_{D_0^1}^2 |\nabla u_t|_{L^2}^2
\end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_1)$ . Now let us define a function  $\Lambda$  by

$$\begin{aligned}
\Lambda(t) &= \int (\mu |\nabla u_t|^2 + (\lambda + \mu) (\operatorname{div} u_t)^2) (t) dx \\
&\quad - \int (2p_t \operatorname{div} u_t + 2\rho_t (f - v \cdot \nabla v) \cdot u_t - \rho_t |u_t|^2) (t) dx.
\end{aligned}$$

Then it follows from Lemma 3.4, Lemma 3.7 and (3.10) that

$$\begin{aligned}
|\Lambda| &\leq C \left( |\nabla u_t|_{L^2}^2 + |p_t|_{L^2}^2 + |\rho_t|_{L^3}^2 |f - v \cdot \nabla v|_{L^2}^2 + |\rho|_{L^\infty}^3 |v|_{D_0^1}^4 |\sqrt{\rho} u_t|_{L^2}^2 \right) \\
&\leq C |\nabla u_t|_{L^2}^2 + C c_0^7 c_{0,3,0}^8, \\
\Lambda &\geq C^{-1} |\nabla u_t|_{L^2}^2 - C c_0^7 c_{0,3,0}^8 \quad \text{and} \quad |\Lambda(0)| \leq C c_0^7 c_{0,3,0}^8.
\end{aligned}$$

Hence integrating (3.12) over  $(0, t)$  and using Lemma 3.4 and Lemma 3.7, we deduce that

$$\begin{aligned}
&\int_0^t |\sqrt{\rho} u_{tt}(s)|_{L^2}^2 ds + |\nabla u_t(t)|_{L^2}^2 \\
&\leq C c_0^7 c_{0,3,0}^{12} + \int_0^t C (1 + c_{0,3,0})^{-1} |v_t|_{D_0^1}^2 |\nabla u_t(s)|_{L^2}^2 ds
\end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_1)$ . Therefore, in view of Gronwall's inequality, we conclude that

$$\int_0^t |\sqrt{\rho} u_{tt}(s)|_{L^2}^2 ds + |u_t(t)|_{D_0^1}^2 \leq C c_0^7 c_{0,3,0}^{12}$$

for  $0 \leq t \leq \min(T_*, T_1)$ . Moreover, since  $Lu = -\nabla p + \rho(f - v \cdot \nabla v - u_t)$  in  $\Omega$ , it follows from the elliptic regularity result that

$$\int_0^t |u_t(s)|_{D^2}^2 ds + |u(t)|_{D^3}^2 \leq C c_0^7 c_{0,3,0}^{12} \quad \text{for } 0 \leq t \leq \min(T_*, T_1).$$

This completes the proof of Lemma 3.8.  $\square$

**Lemma 3.9.** For  $1 \leq j \leq k+1$  we have

$$\begin{aligned} & \int_0^t |\sqrt{\rho}u^{(j+1)}|_{L^2}^2 ds + \max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |u_{(m)}(s)|_{D_0^1}^2 \right) \\ & \leq C c_{0,3,j-1}^{j(2j+6)} \left( 1 + \max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |\sqrt{\rho}u^{(m)}(s)|_{L^2}^2 \right) \right). \end{aligned}$$

*Proof.* From (1.6) we deduce that for all  $1 \leq j \leq k+1$

$$\begin{aligned} & \rho u^{(j+1)} + Lu^{(j)} + \nabla p^{(j)} \\ (3.13) \quad & = \sum_{0 \leq m \leq j} \binom{j}{m} \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \\ & \quad - j \rho^{(1)} u^{(j)} - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \rho^{(j+1-m)} u^{(m)}. \end{aligned}$$

Multiplying (3.13) by  $u^{(j+1)}$  and integrating over  $\Omega$ , we have

$$\begin{aligned} & \int \rho |u^{(j+1)}|^2 dx + \frac{1}{2} \frac{d}{dt} \int \left( \mu |\nabla u^{(j)}|^2 + (\lambda + \mu) (\operatorname{div} u^{(j)})^2 \right) dx \\ & = - \int \nabla p^{(j)} \cdot u^{(j+1)} dx \\ (3.14) \quad & + \sum_{0 \leq m \leq j} \binom{j}{m} \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j+1)} dx \\ & - j \int \rho^{(1)} u^{(j)} \cdot u^{(j+1)} dx \\ & - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \rho^{(j+1-m)} u^{(m)} \cdot u^{(j+1)} dx. \end{aligned}$$

To estimate the first term, we observe from Lemma 3.5 that

$$\begin{aligned} - \int \nabla p^{(j)} \cdot u^{(j+1)} dx & = \int p^{(j)} \operatorname{div} u^{(j+1)} dx \\ & = \frac{d}{dt} \int p^{(j)} \operatorname{div} u^{(j)} dx - \int p^{(j+1)} \operatorname{div} u^{(j)} dx \end{aligned}$$

$$\int p^{(j)} \operatorname{div} u^{(j)} dx \leq C |p^{(j)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2 \leq C c_{0,1,j-1}^{2j+2} + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2$$

and

$$- \int p^{(j+1)} \operatorname{div} u^{(j)} dx \leq C |p^{(j+1)}|_{L^2}^2 + |\nabla u^{(j)}|_{L^2}^2.$$

To estimate the second term of the right hand side in (3.14), we rewrite it as follows:

$$\begin{aligned} & \sum_{0 \leq m \leq j} \binom{j}{m} \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j+1)} dx \\ &= \sum_{0 \leq m \leq j-1} \binom{j}{m} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j+1)} dx \\ & \quad + \int \rho (f - v \cdot \nabla v)^{(j)} \cdot u^{(j+1)} dx. \end{aligned}$$

Using Lemma 3.4 together with (3.3), we have

$$\begin{aligned} \int \rho (f - v \cdot \nabla v)^{(j)} \cdot u^{(j+1)} dx &\leq |\rho|_{L^\infty}^{\frac{1}{2}} |(f - v \cdot \nabla v)^{(j)}|_{L^2} |\sqrt{\rho} u^{(j+1)}|_{L^2} \\ &\leq C |\rho|_{L^\infty} |(f - v \cdot \nabla v)^{(j)}|_{L^2}^2 + \frac{1}{2} |\sqrt{\rho} u^{(j+1)}|_{L^2}^2 \\ &\leq C c_{0,3,j}^5 + \frac{1}{2} |\sqrt{\rho} u^{(j+1)}|_{L^2}^2, \end{aligned}$$

$$\begin{aligned} & \sum_{0 \leq m \leq j-1} \binom{j}{m} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j+1)} dx \\ &= \sum_{0 \leq m \leq j-1} \binom{j}{m} \frac{d}{dt} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j)} dx \\ & \quad - \sum_{0 \leq m \leq j-1} \binom{j}{m} \int \left( \rho^{(j+1-m)} (f - v \cdot \nabla v)^{(m)} \right. \\ & \quad \left. + \rho^{(j-m)} (f - v \cdot \nabla v)^{(m+1)} \right) \cdot u^{(j)} dx. \end{aligned}$$

Since for all  $0 \leq m \leq j-1$ ,

$$|\rho^{(j-m)}|_{H^1} \leq C c_{0,1,j-1}^{j+1} \quad \text{and} \quad |(v \cdot \nabla v)^{(m)}|_{L^2 \cap L^3} \leq C c_{0,3,j-1}^2,$$

we have

$$\begin{aligned} & \sum_{0 \leq m \leq j-1} \binom{j}{m} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j)} dx \\ &\leq \sum_{0 \leq m \leq j-1} \binom{j}{m} |\rho^{(j-m)}|_{L^3} |(f - v \cdot \nabla v)^{(m)}|_{L^3} |\nabla u^{(j)}|_{L^2} \\ &\leq C \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}|_{H^1}^2 |(f - v \cdot \nabla v)^{(m)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2 \\ &\leq C c_{0,3,j-1}^{2j+6} + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2. \end{aligned}$$

On the other hand, we obtain

$$\begin{aligned}
& \sum_{0 \leq m \leq j-1} \binom{j}{m} \int \left( \rho^{(j+1-m)} (f - v \cdot \nabla v)^{(m)} \right. \\
& \quad \left. + \rho^{(j-m)} (f - v \cdot \nabla v)^{(m+1)} \right) \cdot u^{(j)} dx \\
& \leq C c_{0,3,j-1}^{2j+6} + c_{0,3,j-1}^4 |\rho^{(j+1)}|_{L^2}^2 \\
& \quad + C \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}|_{H^1}^2 |(f - v \cdot \nabla v)^{(m)}|_{L^2}^2 + |\nabla u^{(j)}|_{L^2}^2 \\
& \leq C c_{0,3,j-1}^{2j+6} + C c_{0,3,j-1}^4 |\rho^{(j+1)}|_{L^2}^2 + |\nabla u^{(j)}|_{L^2}^2.
\end{aligned}$$

We estimate the third term of (3.14) as follows:

$$\begin{aligned}
-j \int \rho^{(1)} u^{(j)} \cdot u^{(j+1)} dx &= -\frac{j}{2} \frac{d}{dt} \int \rho^{(1)} |u^{(j)}|^2 dx + \frac{j}{2} \int \rho^{(2)} |u^{(j)}|^2 dx, \\
-\frac{j}{2} \int \rho^{(1)} |u^{(j)}|^2 dx &\leq C \int \rho |v| |u^{(j)}| |\nabla u^{(j)}| dx \\
&\leq C |\rho|_{L^\infty}^{\frac{3}{4}} |v|_{D_0^1} |\sqrt{\rho} u^{(j)}|_{L^2}^{\frac{1}{2}} |\nabla u^{(j)}|_{L^2}^{\frac{3}{2}} \\
&\leq C c_{0,2,0}^7 |\sqrt{\rho} u^{(j)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2,
\end{aligned}$$

and

$$\begin{aligned}
\frac{j}{2} \int \rho^{(2)} |u^{(j)}|^2 dx &= -\frac{j}{2} \int (\rho^{(1)} v + \rho v^{(1)}) \cdot (u^{(j)} \cdot \nabla u^{(j)}) dx \\
&\leq C (|\rho^{(1)}|_{H^1} |v|_{D_0^1} + |\rho|_{L^3} |v^{(1)}|_{D_0^1}) |\nabla u^{(j)}|_{L^2}^2 \\
&\leq C c_{0,3,0}^3 (1 + |v^{(1)}|_{D_0^1}) |\nabla u^{(j)}|_{L^2}^2.
\end{aligned}$$

Finally, we have for the last term

$$\begin{aligned}
& - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int \rho^{(j+1-m)} u^{(m)} \cdot u^{(j+1)} dx \\
&= - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \frac{d}{dt} \int \rho^{(j+1-m)} u^{(m)} \cdot u^{(j)} dx \\
& \quad + \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho^{(j+2-m)} u^{(m)} + \rho^{(j+1-m)} u^{(m+1)}) \cdot u^{(j)} dx, \\
& - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int \rho^{(j+1-m)} u^{(m)} \cdot u^{(j)} dx \\
&= \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho v)^{(j-m)} (u^{(j)} \cdot \nabla u^{(m)} + u^{(m)} \cdot \nabla u^{(j)}) dx \\
&\leq C \sum_{1 \leq m \leq j-1} |(\rho v)^{(j-m)}|_{L^3}^2 |\nabla u^{(m)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2 \\
&\leq C c_{0,2,j-1}^{2j+4} \sum_{1 \leq m \leq j-1} |\nabla u^{(m)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2,
\end{aligned}$$

$$\begin{aligned}
& \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho^{(j+2-m)} u^{(m)} + \rho^{(j+1-m)} u^{(m+1)}) \cdot u^{(j)} dx \\
&= - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho v)^{(j+1-m)} (u^{(j)} \cdot \nabla u^{(m)} + u^{(m)} \cdot \nabla u^{(j)}) dx \\
&\quad + \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int \rho^{(j+1-m)} u^{(m+1)} \cdot u^{(j)} dx \\
&\leq C \sum_{1 \leq m \leq j-1} |(\rho v)^{(j+1-m)}|_{L^3}^2 |\nabla u^{(m)}|_{L^2}^2 + |\nabla u^{(j)}|_{L^2}^2 \\
&\quad + C \sum_{2 \leq m \leq j-1} |\rho^{(j+2-m)}|_{L^{\frac{3}{2}}}^2 |\nabla u^{(m)}|_{L^2}^2 + |\nabla u^{(j)}|_{L^2}^2 - \int \rho_{(2)} |u^{(j)}|^2 dx \\
&\leq C \left( c_{0,3,j-1}^{2j+4} + c_{0,1,j-1}^{2j+4} \right) \sum_{1 \leq m \leq j-1} |\nabla u^{(m)}|_{L^2}^2 \\
&\quad + \int (|\rho^{(1)}| |v| + \rho |v^{(1)}|) |\nabla u^{(j)}| |u^{(j)}| dx \\
&\leq C c_{0,3,j-1}^{2j+4} \sum_{1 \leq m \leq j-1} |\nabla u^{(m)}|_{L^2}^2 + C c_{0,3,0}^3 (1 + |\nabla v^{(1)}|_{L^2}) |\nabla u^{(j)}|_{L^2}^2.
\end{aligned}$$

Now integrating (3.14) over  $(0, t) \subset (0, T_1)$ , we have

$$\begin{aligned}
& \int_0^t |\sqrt{\rho} u^{(j)}(s)|^2 ds + |\nabla u^{(j)}(t)|_{L^2}^2 \\
&\leq C c_{0,3,j-1}^{2j+6} + C c_{0,2,0}^7 |\sqrt{\rho} u^{(j)}(t)|_{L^2}^2 + C c_{0,3,j-1}^{2j+4} \sum_{1 \leq m \leq j-1} |\nabla u^{(m)}|_{L^2}^2 \\
&\quad + C \int_0^t |p^{(j+1)}(s)|_{L^2}^2 ds + C c_{0,3,j-1}^4 \int_0^t |\rho^{(j+1)}(s)|_{L^2}^2 ds \\
&\quad + C c_{0,3,j-1}^{2j+4} \sum_{1 \leq m \leq j-1} \int_0^t |\nabla u^{(m)}|_{L^2}^2 ds \\
&\quad + C c_{0,3,0}^3 \int_0^t (1 + |v^{(1)}(s)|_{D_0^1}) |\nabla u^{(j)}(s)|_{L^2}^2 ds.
\end{aligned}$$

Let us define functions  $I_j$  and  $II_j(t)$  by

$$\begin{aligned}
I_j(t) &= \max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |\nabla u^{(m)}|_{L^2}^2 \right), \\
II_j(t) &= \max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |\sqrt{\rho} u^{(j)}(s)|_{L^2}^2 \right).
\end{aligned}$$

Then

$$\begin{aligned}
|\nabla u^{(j)}|_{L^2}^2 &\leq C c_{0,3,j-1}^{2j+6} + C c_{0,2,0}^7 |\sqrt{\rho} u^{(j)}(t)|_{L^2}^2 \\
&\quad + C (c_{0,3,j-1}^{2j+4} + c_{0,3,j-1}^{2j+4} t) I_{j-1}(t) \\
&\quad + C c_{0,3,0}^3 \int_0^t (1 + |v^{(1)}(s)|_{D_0^1}) |\nabla u^{(j)}(s)|_{L^2}^2 ds.
\end{aligned}$$

Therefore Gronwall's inequality implies that

$$(3.15) \quad I_j(t) \leq Cc_{0,3,j-1}^{2j+6} + Cc_{0,2,0}^7 I_j(t) + C(j-1)c_{0,3,j-1}^{2j+4} I_{j-1}(t)$$

for  $0 \leq t \leq T_1$ . Then by induction on  $j$  we have

$$\begin{aligned} I_j(t) &\leq C \left( c_{0,3,j-1}^{2j+6} + c_{0,2,0}^7 I_j(t) \right) \\ &\quad \times \left( 1 + c_{0,3,j-1}^{2j+4} + c_{0,3,j-1}^{2j+4} c_{0,3,j-2}^{2(j-1)+4} + \cdots \right. \\ &\quad \left. + c_{0,3,j-1}^{2j+4} c_{0,3,j-2}^{2(j-1)+4} \cdots c_{0,3,1}^{2 \cdot 2+4} \right) \end{aligned}$$

and hence

$$I_j(t) \leq Cc_{0,3,j-1}^{(j-1)(j+6)} \left( c_{0,3,j-1}^{2j+6} + c_{0,2,0}^7 I_j(t) \right). \quad \square$$

**Lemma 3.10.** *For each  $1 \leq j \leq k+1$  we have*

$$\begin{aligned} &\max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |\sqrt{\rho} u^{(m)}(s)|_{L^2}^2 \right) \leq Cc_0^2, \\ &\int_0^t \left( |\sqrt{\rho} u^{(j+1)}(s)|_{L^2}^2 + |u^{(j)}(s)|_{D^2}^2 \right) ds \\ &\quad + \max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |\nabla u^{(m)}(s)|_{L^2}^2 \right) \leq Cc_{0,3,j-1}^{(j+1)(2j+6)}. \end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_2)$ , where  $T_2 = \min \left( ((k+1)c_{0,3,k}^{(k+2)(2k+8)})^{-1}, T_1 \right)$ .

*Proof.* Multiplying (3.13) by  $u^{(j)}$  and integrating over  $\Omega$ , we have

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int \rho |u^{(j)}|^2 dx + \mu |\nabla u^{(j)}|_{L^2}^2 + (\lambda + \mu) \int (\operatorname{div} u^{(j)})^2 dx \\ &= - \int \nabla p^{(j)} \cdot u^{(j)} dx \\ (3.16) \quad &+ \sum_{0 \leq m \leq j} \binom{j}{m} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j)} dx \\ &- j \int \rho^{(1)} |u^{(j)}|^2 dx - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \rho^{(j+1-m)} u^{(m)} \cdot u^{(j)} dx. \end{aligned}$$

We estimate each term of right hand side of (3.16) as follows:

$$- \int \nabla p^{(j)} \cdot u^{(j)} dx = \int p^{(j)} \operatorname{div} u^{(j)} dx \leq C|p^{(j)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2,$$



$$\begin{aligned}
& \sum_{0 \leq m \leq j} \binom{j}{m} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j)} dx \\
&= \sum_{0 \leq m \leq j-1} \binom{j}{m} \int \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \cdot u^{(j)} dx \\
&\quad + \int \rho (f - v \cdot \nabla v)^{(j)} \cdot u^{(j)} dx \\
&\leq C \sum_{0 \leq m \leq j-1} |\rho^{(j-m)}|_{L^2}^2 |(f - v \cdot \nabla v)^m|_{L^3}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2 \\
&\quad + |\rho|_{L^\infty} |(f - v \cdot \nabla v)^{(j)}|_{L^2}^2 + |\sqrt{\rho} u^{(j)}|_{L^2}^2 \\
&\leq C(c_{0,3,j-1}^{2j+6} + c_{0,3,j}^3) + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2 + |\sqrt{\rho} u^{(j)}|_{L^2}^2, \\
&\quad -j \int \rho^{(1)} |u^{(j)}|^2 dx = 2j \int \rho v \cdot (u^{(j)} \cdot \nabla u^{(j)}) dx \\
&\quad \leq Cc_{0,3,0}^3 |\sqrt{\rho} u^{(j)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2,
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int \rho^{(j+1-m)} u^{(m)} \cdot u^{(j)} dx \\
&= \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho v)^{(j-m)} (u^{(m)} \cdot \nabla u^{(j)} + u^{(j)} \cdot \nabla u^{(m)}) dx \\
&\leq C \sum_{1 \leq m \leq j-1} |(\rho v)^{(j-m)}|_{L^3} |\nabla u^{(m)}| |\nabla u^{(j)}|_{L^2} \\
&\leq Cc_{0,3,j-1}^{2j+4} \sum_{1 \leq m \leq j-1} |\nabla u^{(m)}|_{L^2}^2 + \frac{\mu}{8} |\nabla u^{(j)}|_{L^2}^2.
\end{aligned}$$

Substituting all these estimate into (3.16), we have

$$\begin{aligned}
(3.17) \quad & \frac{d}{dt} \int \rho |u^{(j)}|^2 dx + \mu |\nabla u^{(j)}|_{L^2}^2 \\
& \leq C(c_{0,1,j-1}^{2j+2} + c_{0,3,j-1}^{2j+6} + c_{0,3,j}^3) + Cc_{0,3,0}^3 |\sqrt{\rho} u^{(j)}|_{L^2}^2 \\
& \quad + Cc_{0,3,j-1}^{2j+4} \sum_{1 \leq m \leq j-1} |\nabla u^{(m)}|_{L^2}^2.
\end{aligned}$$

Recall that  $I_j(t) = \max_{1 \leq m \leq j} (\sup_{0 \leq s \leq t} |\nabla u^{(m)}(s)|_{L^2}^2)$ . Then integrating (3.17) over  $(0, t)$ , we have for  $t \in [0, T_1]$

$$\begin{aligned}
& \int \rho |u^{(j)}|^2 dx + \mu \int_0^t |\nabla u^{(j)}|_{L^2}^2 ds \\
& \leq |\sqrt{\rho} u^{(j)}(0)|_{L^2}^2 + C + Cc_{0,3,j-1}^{2j+4} jt I_{j-1}(t) + Cc_{0,3,0}^3 \int_0^t |\sqrt{\rho} u^{(j)}|_{L^2}^2 ds.
\end{aligned}$$

From the observation that

$$|\sqrt{\rho} u^{(j)}(0)|_{L^2} = |\sqrt{\rho_0} g_j|_{L^2} \leq C |\rho_0|_{L^{\frac{3}{2}}} |g_j|_{D_0^1} \leq Cc_0^2,$$

we deduce that

$$|\sqrt{\rho}u^{(j)}(t)|_{L^2}^2 \leq Cc_0^2 + Cc_{0,3,j-1}^{2j+4}jtI_{j-1}(t)$$

and hence

$$(3.18) \quad \max_{1 \leq m \leq j} \left( \sup_{0 \leq s \leq t} |\sqrt{\rho}u^{(m)}|_{L^2}^2 \right) \leq Cc_0^2 + Cc_{0,3,j-1}^{2j+4}jtI_{j-1}(t)$$

for  $0 \leq t \leq T_1$ . Therefore from Lemma 3.9, we conclude that

$$(3.19) \quad I_j(t) \leq Cc_{0,3,j-1}^{(j+1)(2j+6)} + Cjc_{0,3,j-1}^{(j+1)(2j+6)}tI_{j-1}(t).$$

Thus if we choose  $t \leq ((k+1)c_{0,3,k}^{(k+2)(2k+8)})^{-1} \leq (jc_{0,3,j-1}^{(j+1)(2j+6)})^{-1}$ , then using (3.18), (3.19) and Lemma 3.6, we can prove the lemma by induction.  $\square$

**Lemma 3.11.** *For each  $1 \leq j \leq k$ , we have*

$$\begin{aligned} |u^{(j)}(t)|_{D^2} + \int_0^t \left( |u^{(j+1)}(s)|_{D_0^1}^2 + |u^{(j)}(s)|_{D^3}^2 \right) ds &\leq Cc_{0,2,j}^{(j+1)(2j+7)}c_{0,3,j}^{\frac{1}{2}}, \\ |u^{(j+1)}(t)|_{D_0^1} + |u^{(j)}(t)|_{D^3} \\ &+ \int_0^t \left( |u^{(j+1)}(s)|_{D^2}^2 + |u^{(j)}(s)|_{D^4}^2 \right) ds \leq Cc_{0,3,j}^{(j+3)(2j+8)} \end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_2)$ .

*Proof.* Since for each  $t$ ,  $u_{(j)}(t)$  is a solution to the elliptic boundary value problem:

$$\begin{aligned} Lu^{(j)} &= \sum_{0 \leq m \leq j} \binom{j}{m} \rho^{(j-m)} (f - v \cdot \nabla v)^{(m)} \\ &- \sum_{1 \leq m \leq j} \binom{j}{m} \rho^{(j+1-m)} u^{(m)} - \nabla p^{(j)} - \rho u^{(j+1)}, \end{aligned}$$

from the elliptic regularity (see [5]) we have

$$\begin{aligned} |u^{(j)}|_{D_0^1 \cap D^2} &\leq C|u^{(j)}|_{D_0^1} + C \sum_{0 \leq m \leq j} |\rho^{(j-m)} (f - v \cdot \nabla v)^{(m)}|_{L^2} \\ &+ C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)} u^{(m)}|_{L^2} + |\nabla p^{(j)}|_{L^2} + |\rho u^{(j+1)}|_{L^2} \\ &\leq C|u^{(j)}|_{D_0^1} + C \sum_{0 \leq m \leq j} |\rho^{(j-m)}|_{H^2} |(f - v \cdot \nabla v)^{(m)}|_{L^2} \\ &+ C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)}|_{H^1} |\nabla u^{(m)}|_{L^2} + C|p^{(j)}|_{H^1} \\ &+ C|\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho}u^{(j+1)}|_{L^2}. \end{aligned}$$

Using the bound (3.3), Lemma 3.4 and Lemma 3.10, we can easily show that for  $1 \leq j \leq k$

$$|u^{(j)}(t)|_{D_0^1 \cap D^2} \leq Cc_{0,3,j}^{(j+1)(2j+7)} + Cc_{0,2,j}^{\frac{3}{2}}c_{0,3,j}^{\frac{1}{2}}$$

for  $0 \leq t \leq \min(T_*, T_2)$ . Further, we have

$$\begin{aligned} |u^{(j+1)}|_{D^2} &\leq C|u^{(j+1)}|_{D_0^1} + C \sum_{0 \leq m \leq j+1} |\rho^{(j+1-m)}|_{H^2} |(f - v \cdot \nabla v)^{(m)}|_{L^2} \\ &\quad + C \sum_{1 \leq m \leq j} |\rho^{(j+2-m)}|_{H^1} |\nabla u^{(m)}|_{L^2} + C|p^{(j+1)}|_{H^1} \\ &\quad + C|\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho}u^{(j+2)}|_{L^2} \end{aligned}$$

and hence

$$\begin{aligned} (3.20) \quad &\int_0^t |u^{(j+1)}(s)|_{D^2}^2 ds \\ &\leq C \sum_{0 \leq m \leq j+1} \int_0^t |\rho^{(j+1-m)}(s)|_{H^2}^2 |(f - v \cdot \nabla v)^{(m)}|_{L^2}^2 ds \\ &\quad + C \sum_{1 \leq m \leq j+1} \int_0^t \left( |\rho^{(j+2-m)}|_{H^1}^2 |\nabla u^{(m)}|_{L^2}^2 + |p^{(j+1)}|_{H^1}^2 \right) ds \\ &\quad + C \int_0^t |\rho u^{(j+2)}|_{L^2}^2 ds \leq Cc_{0,3,j}^{(j+3)(2j+8)}. \end{aligned}$$

On the other hand, using the elliptic regularity [1, 5], we obtain

$$\begin{aligned} |u^{(j)}|_{D^3} &\leq C|u^{(j)}|_{D_0^1} + C \sum_{0 \leq m \leq j} |\rho^{(j-m)}(f - v \cdot \nabla v)^{(m)}|_{H^1} \\ &\quad + C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)}u^{(m)}|_{H^1} + C|p^{(j)}|_{H^2} + C|\rho u^{(j+1)}|_{H^1} \\ &\leq Cc_{0,3,j-1}^{(j+1)(2j+6)} + \sum_{0 \leq m \leq j} |\rho^{(j-m)}|_{H^2} |(f - v \cdot \nabla v)^{(m)}|_{H^1} \\ &\quad + C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)}|_{H^2} |\nabla u^{(m)}|_{L^2} \\ &\quad + Cc_{0,1,j-1}^{j+1} + |\rho|_{H^2} |\nabla u^{(j+1)}|_{L^2} \\ &\leq C \left( c_{0,3,j-1}^{(j+1)(2j+6)} + c_{0,1,j-1}^{j+1} (1 + c_{0,3,j}^2) \right. \\ &\quad \left. + c_{0,1,j-1}^{j+1} c_{0,3,j-1}^{(j+1)(2j+6)} + c_{0,3,j}^{(j+2)(2j+8)} \right) \\ &\leq Cc_{0,3,j}^{(j+3)(2j+8)}. \end{aligned}$$

And finally we have

$$\begin{aligned}
|u^{(j)}|_{D^4} &\leq C|u^{(j)}|_{D_0^1} + C \sum_{0 \leq m \leq j} |\rho^{(j-m)}(f - v \cdot \nabla v)^{(m)}|_{H^2} \\
&\quad + C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)}u^{(m)}|_{H^2} + C|p^{(j)}|_{H^3} + C|\rho u^{(j+1)}|_{H^2} \\
&\leq Cc_{0,3,j-1}^{(j+1)(2j+6)} + \sum_{0 \leq m \leq j} |\rho^{(j-m)}|_{H^2} |(f - v \cdot \nabla v)^{(m)}|_{H^2} \\
&\quad + C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)}|_{H^2} |u^{(m)}|_{D_0^1 \cap D^2} \\
&\quad + Cc_{0,1,j-1}^{j+1} + C|\rho|_{H^2} |u^{(j+1)}|_{D_0^1 \cap D^2}.
\end{aligned}$$

and therefore by (3.20)

$$\int_0^t |u_{(j)}(s)|_{D^4}^2 ds \leq Cc_{0,3,j}^{(j+3)(2j+8)}$$

for  $0 \leq t \leq \min(T_*, T_2)$ . This completes the proof of lemma.  $\square$

**Lemma 3.12.** *For all  $l$  and  $j$  with  $1 \leq l \leq k$  and  $0 \leq j \leq k - l$ , we have*

$$\begin{aligned}
|u^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} &\leq Cc_{l-1,3,j}^{(j+2l+1)(2j+2l+6)} \\
|u^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} &\leq Cc_{l,2,j}^{(j+2l+2)(2j+2l+7)} \\
|u^{(j)}(t)|_{D_0^1 \cap D^{2l+3}} &\leq Cc_{l,3,j}^{(j+2l+3)(2j+2l+8)}
\end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_3)$ , where  $T_3 = \min(c_{k,4,0}^{-(2k+9)^2}, T_2)$ .

*Proof.* For the proof in case when  $j = 0$ , we regard  $c_{l,i,-1}$  as  $c_{l,i,0}$ . From the elliptic regularity, we have

$$\begin{aligned}
&|u^{(j)}|_{D_0^1 \cap D^{2l+1}} \\
&\leq C|u^{(j)}|_{D_0^1} + C \sum_{0 \leq m \leq j} |\rho^{(j-m)}(f - v \cdot \nabla v)^{(m)}|_{H^{2l-1}} \\
&\quad + C \sum_{1 \leq m \leq j} |\rho^{(j+1-m)}u^{(m)}|_{H^{2l-1}} + C|p^{(j)}|_{H^{2l}} + C|\rho u^{(j+1)}|_{L^{2l-1}} \\
&\leq Cc_{0,3,j-1}^{(j+1)(2j+6)} + C \sum_{0 \leq m \leq j} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ |\alpha| \leq 2l-1}} |D^{\alpha_1} \rho^{(j-m)}|_{H^2} |D^{\alpha_2} (f - v \cdot \nabla v)^{(m)}|_{L^2} \\
&\quad + C \sum_{1 \leq m \leq j} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ |\alpha| \leq 2l-1}} |D^{\alpha_1} \rho^{(j+1-m)} D^{\alpha_2} u^{(m)}|_{L^2} \\
&\quad + Cc_{0,1,j-1}^{j+1} + C|\rho|_{H^{2l}} |u^{(j+1)}|_{D_0^1 \cap D^{2l-1}} \\
&\leq Cc_{0,3,j-1}^{(j+1)(2j+6)} + Cc_{l-1,3,j}^3 + Cc_{0,1,j-1}^{j+1} \max_{1 \leq m \leq j+1} |u^{(m)}|_{D_0^1 \cap D^{2(l-1)+1}} \\
&\leq Cc_{l-1,3,j}^{(j+1)(2j+6)} + Cc_{0,1,j-1}^{j+1} M_{l-1,j+1},
\end{aligned}$$

where  $M_{l,n}(t) = \max_{0 \leq m \leq n} |u^{(m)}|_{D_0^1 \cap D^{2l+1}}$  for  $n \geq 0$ . Thus we have

$$M_{l,j}(t) \leq Cc_{l-1,3,j}^{(j+1)(2j+6)} + Cc_{0,1,j-1}^{j+1} M_{l-1,j+1}$$

for all  $0 \leq t \leq \min(T_*, T_2)$ . Since by Lemma 3.10

$$M_{0,j+1} = \max_{0 \leq m \leq j+1} |u^{(m)}|_{D_0^1} \leq Cc_{0,3,j}^{(j+2)(2j+8)},$$

we obtain

$$M_{1,j} \leq Cc_{0,3,j}^{(j+3)(2j+8)}.$$

By induction, we deduce that

$$M_{l,j}(t) \leq Cc_{l-1,3,j}^{(j+2l+1)(2j+2l+6)}.$$

Now let us define a function  $N_{l,n}$  by

$$N_{l,n}(t) = \max_{1 \leq m \leq n} |u^{(m)}(t)|_{D_0^1 \cap D^{2l+2}}.$$

Then we similarly have

$$\begin{aligned} |u^{(j)}|_{D_0^1 \cap D^{2l+2}} &\leq Cc_{0,3,j-1}^{(j+1)(2j+6)} + Cc_{l,2,j}^3 + Cc_{0,1,j-1}^{j+1} N_{l-1,j+1} \\ &\leq Cc_{l,2,j}^{(j+1)(2j+6)} + Cc_{0,1,j-1}^{j+1} N_{l-1,j+1} \end{aligned}$$

and hence

$$N_{l,j}(t) \leq Cc_{l,2,j}^{(j+1)(2j+6)} + Cc_{0,1,j-1}^{j+1} N_{l-1,j+1}(t)$$

for all  $0 \leq t \leq \min(T_*, T_2)$ . Since by Lemma 3.11

$$N_{0,j+1} = \max_{0 \leq m \leq j+1} |u^{(m)}|_{D_0^1 \cap D^2} \leq Cc_{0,2,j+1}^{(j+2)(2j+9)} c_{0,3,j+1}^{\frac{1}{2}} \leq Cc_{0,3,j+1}^{(j+3)(2j+9)},$$

we obtain

$$N_{1,j} \leq Cc_{1,2,j}^{(j+1)(2j+6)} + Cc_{0,3,j+1}^{(j+4)(2j+9)} \leq Cc_{1,2,j}^{(j+4)(2j+9)}.$$

Hence by induction we have

$$N_{l,j}(t) \leq Cc_{l,2,j}^{(j+2l+2)(2j+2l+7)}$$

for all  $0 \leq t \leq \min(T_*, T_2)$ .

From the similar argument as above results, we can easily deduce that

$$\begin{aligned} |u^{(j)}|_{D_0^1 \cap D^{2l+3}} &\leq Cc_{l,3,j}^{(j+2l+3)(2j+2l+8)}, \\ |u^{(j+1)}|_{D_0^1 \cap D^{2l+1}} &\leq Cc_{l-1,3,j+1}^{(j+2l+2)(2j+2l+8)}, \\ \int_0^t \left( |u^{(j+1)}|_{D_0^1 \cap D^{2l+2}}^2 + |u^{(j)}|_{D_0^1 \cap D^{2l+4}}^2 \right) &\leq Cc_{l,3,j}^{(j+2l+3)(2j+2l+8)} \end{aligned}$$

for all  $0 \leq t \leq \min(T_*, T_3)$ .  $\square$

Summarizing the estimates up to now, we have the following.

**Proposition 3.13.** *If  $(\rho, u)$  is a solution to the linearized problem (2.7)-(2.10) with the initial density  $\rho_0 > \delta$  the known vector field  $v$  satisfying (3.1), (3.2) and (3.3), then for sufficiently small  $\varepsilon$ , there exists a time interval  $[0, T_*]$  such that the velocity  $u$  satisfies the following estimates that for all  $0 \leq j \leq k+1$*

$$(3.21) \quad |u^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} \leq c_{0,1,j},$$

and for all  $0 \leq l \leq k$  and  $0 \leq j \leq k - l$  that  $\inf_{(0, T_*) \times \Omega} \rho(t, x) \geq C^{-1} \delta$ .

$$\begin{aligned}
(3.22) \quad & \sup_{0 < t < T_*} \left( |\rho(t)|_{L^1} + |(\rho^{(j)}(t), (p - p(0))^{(j)}(t))|_{H^{2(k-j)+3}} \right) \\
& \quad + \int_0^{T_*} |\rho^{(k+2)}(s)|_{L^2}^2 dt \leq \varepsilon c_{0,1,j} \\
& \sup_{0 \leq t \leq T_*} \left( |u^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} \right) + \int_0^{T_*} |u^{(j)}(t)|_{D^2 \cap D^{2l+2}}^2 dt \leq c_{l,2,j}, \\
& \sup_{0 \leq t \leq T_*} \left( |u^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} \right) \\
& \quad + \int_0^{T_*} \left( |u^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}}^2 + |u^{(j)}(t)|_{D^{2l+3}}^2 \right) dt \leq c_{l,3,j}, \\
& \text{ess sup}_{0 < t < T_*} \left( |u^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}} + |u^{(j)}(t)|_{D^{2l+3}} \right) \\
& \quad + \int_0^{T_*} \left( |u^{(j+1)}(t)|_{D^{2l+2}}^2 + |u^{(j)}(t)|_{D^{2l+4}}^2 \right) dt \leq c_{l,4,j}, \\
& \text{ess sup}_{0 \leq t \leq T_*} |\sqrt{\rho} u^{(j+1)}(t)|_{L^2} + \int_0^{T_*} |\sqrt{\rho} u^{(j+2)}(t)|_{L^2}^2 dt \leq c_{0,4,j}.
\end{aligned}$$

*Proof.* From Lemma 3.4, we see the lower bound of  $\rho$  by choosing  $c_{0,1,j} \geq C c_{0,1,j-1}^j$ . If we let the constant  $c_{0,1,j}$  be  $C c_0$  for all  $j$ , by Lemma 3.2 the estimate (3.21) follows immediately.

For the estimate (3.22), we consider the following three cases: (1)  $l = 0$  and  $j = 0$ , (2)  $l = 0$  and  $1 \leq j \leq k$ , (3)  $l \geq 1$  and  $0 \leq j \leq k - l$ .

Case (1):  $l = 0$  and  $j = 0$ .

From Lemma 3.6 to Lemma 3.8, it follows that

$$\begin{aligned}
& |u(t)|_{D_0^1} + \int_0^t |u(s)|_{D^2}^2 ds \leq C c_0^3, \\
& |u(t)|_{D^2} + \int_0^t \left( |u_t(s)|_{D_0^1}^2 + |u(s)|_{D^3}^2 \right) ds \leq C c_0^3 c_{0,2,0}^{\frac{3}{2}} c_{0,3,0}^{\frac{1}{2}}, \\
& |u_t(t)|_{D_0^1} + |u(t)|_{D^3} + \int_0^t \left( |u_t(s)|_{D^2}^2 + |u(s)|_{D^4}^2 \right) ds \leq C c_0^7 c_{0,3,0}^{12}, \\
& |\sqrt{\rho} u_t(t)|_{L^2} + \int_0^t |\sqrt{\rho} u_{tt}(s)|_{L^2}^2 ds \leq C c_0^7 c_{0,3,0}^{12}
\end{aligned}$$

for  $0 \leq t \leq \min(T_*, T_3)$ . Therefore, defining the constants  $c_{0,i,0}$ 's and  $T_*$  by

$$(3.23) \quad c_{0,1,0} = C c_0, \quad c_{0,2,0} = C c_0^3, \quad c_{0,3,0} = C^2 c_0^6 c_{0,2,0}^3, \quad c_{0,4,0} = C c_0^7 c_{0,3,0}^{12}$$

and

$$(3.24) \quad T_* = \min(T, T_3),$$

we conclude that

$$\begin{aligned}
(3.25) \quad & \sup_{0 \leq t \leq T_*} |u(t)|_{D_0^1} + \int_0^{T_*} |u(t)|_{D^2}^2 dt \leq c_{0,2,0}, \\
& \sup_{0 \leq t \leq T_*} |u(t)|_{D^2} + \int_0^{T_*} \left( |u_t(t)|_{D_0^1}^2 + |u(t)|_{D^3}^2 \right) dt \leq c_{0,3,0}, \\
& \text{ess sup}_{0 \leq t \leq T_*} \left( |u_t(t)|_{D_0^1} + |u(t)|_{D^3} \right) + \int_0^{T_*} \left( |u_t(t)|_{D^2}^2 + |u(t)|_{D^4}^2 \right) dt \leq c_{0,4,0}, \\
& \text{ess sup}_{0 \leq t \leq T_*} \left( |\sqrt{\rho}u_t(t)|_{L^2} \right) + \int_0^{T_*} |\sqrt{\rho}u_{tt}(t)|_{L^2}^2 dt \leq c_{0,4,0}.
\end{aligned}$$

Case (2):  $l = 0$  and  $1 \leq j \leq k$ .

From Lemma 3.9 to Lemma 3.11, we obtain that

$$\begin{aligned}
(3.26) \quad & |u^{(j)}(t)|_{D_0^1} + \int_0^t |u^{(j)}(s)|_{D^2}^2 ds \leq C c_{0,3,j-1}^{(j+1)(2j+6)}, \\
& |u_{(j)}(t)|_{D^2} + \int_0^t \left( |u_{(j+1)}|_{D_0^1}^2 + |u_{(j)}|_{D^3}^2 \right) ds \leq C c_{0,2,j}^{(j+1)(2j+7)} c_{0,3,j}^{\frac{1}{2}}, \\
& |u_{(j+1)}(t)|_{D_0^1} + |u_{(j)}(t)|_{D^3} \\
& \quad + \int_0^t \left( |u_{(j+1)}(s)|_{D^2}^2 + |u_{(j)}(s)|_{D^4}^2 \right) ds \leq C c_{0,3,j}^{(j+3)(2j+8)}.
\end{aligned}$$

Thus if we define the constants  $c_{0,i,j}$  by

$$c_{0,2,j} = C c_{0,3,j-1}^{(j+1)(2j+6)}, \quad c_{0,3,j} = C^2 c_{0,2,j}^{(2j+2)(2j+7)}, \quad c_{0,4,j} = C c_{0,3,j}^{(j+3)(2j+8)},$$

then we have

$$\begin{aligned}
(3.27) \quad & \sup_{0 \leq t \leq T_*} |u^{(j)}(t)|_{D_0^1} + \int_0^{T_*} |u^{(j)}(t)|_{D^2}^2 dt \leq c_{0,2,j}, \\
& \sup_{0 \leq t \leq T_*} |u^{(j)}(t)|_{D^2} + \int_0^{T_*} \left( |u^{(j+1)}(t)|_{D_0^1}^2 + |u^{(j)}(t)|_{D^3}^2 \right) dt \leq c_{0,3,j}, \\
& \text{ess sup}_{0 \leq t \leq T_*} \left( |u^{(j+1)}(t)|_{D_0^1} + |u^{(j)}(t)|_{D^3} \right) \\
& \quad + \int_0^{T_*} \left( |u^{(j+1)}(t)|_{D^2}^2 + |u^{(j)}(t)|_{D^4}^2 \right) dt \leq c_{0,4,j}, \\
& \text{ess sup}_{0 \leq t \leq T_*} \left( |\sqrt{\rho}u^{(j+1)}(t)|_{L^2} \right) + \int_0^{T_*} |\sqrt{\rho}u^{(j+2)}(t)|_{L^2}^2 dt \leq c_{0,4,j}.
\end{aligned}$$

Case (3):  $l \geq 1$  and  $0 \leq j \leq k - l$ .

From Lemma 3.12, we easily show that

$$\begin{aligned}
& |u^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} + \int_0^t |u^{(j)}(s)|_{D_0^1 \cap D^{2l+2}}^2 ds \\
& \leq C c_{l-1,3,j}^{(j+2l+1)(2j+2l+6)}, \\
(3.28) \quad & |u^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} + \int_0^t \left( |u^{(j+1)}|^2_{D_0^1 \cap D^{2l+1}} + |u^{(j)}|^2_{D_0^1 \cap D^{2l+3}} \right) ds \\
& \leq C c_{l,2,j}^{(j+2l+2)(2j+2l+7)} \\
& |u^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}} + |u^{(j)}(t)|_{D_0^1 \cap D^{2l+3}} \\
& \quad + \int_0^t \left( |u^{(j+1)}(s)|_{D_0^1 \cap D^{2l+2}}^2 + |u^{(j)}(s)|_{D_0^1 \cap D^{2l+4}}^2 \right) ds \\
& \leq C c_{l,3,j}^{(j+2l+3)(2j+2l+8)}.
\end{aligned}$$

and hence if we choose the constants  $c_{l,i,j}$  such that

$$\begin{aligned}
c_{l,2,j} &= C c_{l-1,3,j}^{(j+2l+1)(2j+2l+6)}, \\
c_{l,3,j} &= C c_{l,2,j}^{(j+2l+2)(2j+2l+7)}, \\
c_{l,4,j} &= C c_{l,3,j}^{(j+2l+3)(2j+2l+8)},
\end{aligned}$$

then we finally have

$$\begin{aligned}
(3.29) \quad & \sup_{0 \leq t \leq T_*} |u^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} + \int_0^{T_*} |u^{(j)}(t)|_{D_0^1 \cap D^{2l+2}}^2 dt \leq c_{l,2,j}, \\
& \sup_{0 \leq t \leq T_*} |u^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} \\
& \quad + \int_0^{T_*} \left( |u^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}}^2 + |u^{(j)}(t)|_{D_0^1 \cap D^{2l+3}}^2 \right) dt \leq c_{l,3,j}, \\
& \text{ess sup}_{0 \leq t \leq T_*} \left( |u_{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}} + |u_{(j)}(t)|_{D_0^1 \cap D^{2l+3}} \right) \\
& \quad + \int_0^{T_*} \left( |u_{(j+1)}(t)|_{D_0^1 \cap D^{2l+2}}^2 + |u_{(j)}(t)|_{D_0^1 \cap D^{2l+4}}^2 \right) dt \leq c_{l,4,j}.
\end{aligned}$$

Combining all the results above and Lemma 3.4, we complete the proof of proposition.  $\square$

#### 4. LINEAR PROBLEM WITH NONNEGATIVE DENSITY

In this section, we establish the existence and local time boundedness of solution to the linearized problem with nonnegative initial density on a general domain.

The following is the concerning linearized problem.

$$(4.1) \quad \rho_t + \text{div}(\rho v) = 0 \quad \text{in } (0, T) \times \Omega,$$

$$(4.2) \quad \rho u_t + Lu + \nabla p = \rho(f - v \cdot \nabla v) \quad \text{in } (0, T) \times \Omega,$$

$$(4.3) \quad (\rho, u)|_{t=0} = (\rho_0, u_0) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } (0, T) \times \partial\Omega,$$

$$(4.4) \quad (\rho, u)(t, x) \rightarrow (0, 0) \quad \text{as } |x| \rightarrow \infty, \quad (t, x) \in (0, T) \times \Omega,$$



We assume that the initial data satisfy the bound and compatibility conditions that

$$(4.5) \quad \rho_0 \geq 0, \quad |\rho_0|_{L^1 \cap H^{2k+3}} \leq \frac{\varepsilon}{2}, \quad 2 + \sum_{1 \leq j \leq k+1} |g_j|_{D_0^1} \leq c_0,$$

and

$$(4.6) \quad L(u^{(j)}(0)) = j \operatorname{div}(\rho_0 v(0)) u^{(j)}(0) + G_{j-1} + H_j - \nabla p^{(j)}(0) - \rho_0 g_{j+1}$$

for some  $g_j \in D_0^1 \cap D^{2(k-j)+3}$  and all  $0 \leq j \leq k$ , where

$$\begin{aligned} G_{j-1} &= 0 \quad \text{for } j = 0, 1, \\ G_{j-1} &= - \sum_{1 \leq m \leq j-1} \binom{j}{m} \rho^{(j+1-m)}(0) u^{(m)}(0) \quad \text{for } j \geq 2, \\ H_j &= \sum_{0 \leq m \leq j} \binom{j}{m} \rho^{(j-m)}(0) (f - v \cdot \nabla v)^{(m)}(0). \end{aligned}$$

We also assume that the known  $v$  satisfies the bound conditions that for all  $0 \leq j \leq k+1$ ,

$$(4.7) \quad |v^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} \leq c_{0,1,j},$$

and for all  $0 \leq l \leq k$  and  $0 \leq j \leq k-l$ ,

$$(4.8) \quad \begin{aligned} & \sup_{0 \leq t \leq T_*} \left( |v^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} \right) + \int_0^{T_*} |v^{(j)}(t)|_{D^2 \cap D^{2l+2}}^2 dt \leq c_{l,2,j}, \\ & \sup_{0 \leq t \leq T_*} \left( |v^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} \right) \\ & + \int_0^{T_*} \left( |v^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}}^2 + |v^{(j)}(t)|_{D^{2l+3}}^2 \right) dt \leq c_{l,3,j}, \\ & \operatorname{ess\,sup}_{0 < t < T_*} \left( |v^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}} + |v^{(j)}(t)|_{D^{2l+3}} \right) \\ & + \int_0^{T_*} \left( |v^{(j+1)}(t)|_{D^{2l+2}}^2 + |v^{(j)}(t)|_{D^{2l+4}}^2 \right) dt \leq c_{l,4,j} \end{aligned}$$

for some time  $T_* \in (0, T)$  and constants  $c_{l,i,j}$  such that

$$\begin{aligned} 1 &\leq c_0 \leq c_{l,i,j} \leq c_{l+1,i,j} \quad \text{for any } i, j, l, \\ c_{l,i,j} &\leq c_{l,i+1,j} \quad \text{for each fixed } l, j, \\ \text{and } c_{l,i,j} &\leq c_{l,i,l+1} \quad \text{for each fixed } l \text{ and for any } i. \end{aligned}$$

**Proposition 4.1.** *If the data  $\rho_0, u_0, f$  and the vector field  $v$  satisfy the conditions (4.5), (4.6), (4.7) and (4.8) for some sufficiently small constant  $\varepsilon$ , some fixed  $c_0$  and  $c_{l,i,j}$  appearing in Proposition 3.13, then there exists a time  $T_* > 0$  and a unique solution  $(\rho, u)$  to the linearized problem (4.1) – (4.4) satisfying the estimate (3.21) and (3.22).*

The above proposition will be proved at the end of this section.

For the proof of Proposition 4.1, we will use a standard domain expansion technique. To do so, we choose a cut-off function  $\varphi \in C_c^\infty(B_1)$  such that  $\varphi = 1$  in  $B_{\frac{1}{2}}$ .

Let  $R_0 > 1$  be a sufficiently large number so that

$$\Omega \subset B_{R_0/2} \quad \text{if} \quad \Omega \subset\subset \mathbf{R}^3, \quad \text{and} \quad \Omega \subset \mathbf{R}^3 \setminus B_{R_0/2} \quad \text{in} \quad \mathbf{R}^3 \setminus \subset\subset \mathbf{R}^3.$$

We define

$$\begin{aligned} \varphi_R(x) &= \varphi(x/R), \quad g_{R,j}(x) = \varphi_R(x)g_j(x), \quad 1 \leq j \leq k+1, \\ v_R(t, x) &= \varphi_R(x)v(t, x) \quad \text{and} \quad f_R(t, x) = \varphi_R(x)f(t, x) \end{aligned}$$

for  $(t, x) \in [0, T_*] \times \Omega$ . Note that if  $\Omega \subset\subset \mathbf{R}^3$ , then  $g_{R,j} = g_j$ ,  $v_R = v$  and  $f_R = f$  for each  $R > R_0$  and otherwise, they are supported in  $\Omega_R$  or  $[0, T_*] \times \Omega_R$ , where  $\Omega_R = \Omega \cap B_R$ <sup>1</sup>.

For the domain expansion, we lift the initial density to make use of the estimates independent of lower bound of initial density in Section 3. More specifically, let us define  $\rho_{R,0} = \rho_0 + R^{-3}$  on  $\Omega_R$  and  $\rho_R$  be the solution of the linear transport equation

$$(4.9) \quad (\rho_R)_t + \operatorname{div}(\rho_R v_R) = 0, \quad \text{on} \quad [0, T] \times \Omega_R; \quad \rho_R(0) = \rho_{R,0} \quad \text{in} \quad \Omega_R.$$

Then we can define  $\rho_R^{(j)}(0)$  as in Section 3. Let  $p_R = p(\rho_R)$ . Then we have

**Lemma 4.2.** *If  $\varepsilon$  is sufficiently small and  $R$  is sufficiently large, then there exists a unique solution*

$$(w_{R,0}, w_{R,1}, \dots, w_{R,k}) \in D_0^1 \cap D^{2k+3}(\Omega_R) \times D_0^1 \cap D^{2k+1}(\Omega_R) \times \dots \times D_0^1 \cap D^3(\Omega_R)$$

to the coupled elliptic system: for  $0 \leq j \leq k$ ,  $g_{R,j} = 0$  on  $\partial\Omega_R$  and

$$(4.10) \quad L(w_{R,j}) = j \operatorname{div}(\rho_{R,0} v_R(0))w_{R,j} + G_{R,j-1} + H_{R,j} - \nabla p_R^{(j)}(0) - \rho_{R,0} g_{R,j+1}$$

in  $\Omega_R$ , where

$$\begin{aligned} G_{R,j-1} &= 0 \quad \text{for} \quad j = 0, 1, \\ G_{R,j-1} &= - \sum_{1 \leq m \leq j-1} \binom{j}{m} \rho_R^{(j+1-m)}(0) w_{R,m} \quad \text{for} \quad j \geq 2, \\ H_{R,j} &= \sum_{0 \leq m \leq j} \binom{j}{m} \rho_R^{(j-m)}(0) (f_R - v_R \cdot \nabla v_R)^{(j)}(0). \end{aligned}$$

Furthermore, for such  $\varepsilon$  and  $R$ , there holds

$$|w_{R,j}|_{D_0^1 \cap D^{2(k-j)+3}} \leq Cc_0$$

for all  $0 \leq j \leq k$ .

<sup>1</sup>If  $\Omega$  is the half space  $\mathbf{R}^2 \times \mathbf{R}_+$ , then the non-smooth domain  $\Omega_R$  should be replaced by a smooth domain  $\tilde{\Omega}_R$  such that  $\Omega_R \subset \tilde{\Omega}_R \subset \Omega_{2R}$ .

*Proof.* We first observe that

$$(4.11) \quad \begin{aligned} & \rho_{R,0} \rightarrow \rho_0 \quad \text{in } L^1 \cap H^{2k+3}, \\ & v_R^{(j)} \rightarrow v^{(j)} \quad \text{in } C([0, T_*]; D_0^1 \cap D^{2(k-j)+3}) \quad \text{for } 1 \leq j \leq k, \\ & \text{and } v_R^{(k+1)} \rightarrow v^{(k+1)} \quad \text{in } L^\infty(0, T_*; D_0^1) \end{aligned}$$

as  $R \rightarrow \infty$ . Hence we deduce from Lemma 3.1, 3.4 and 3.5 that for any given  $\varepsilon_1$  there exists a small  $\varepsilon_0$  and a large  $R_1 > R_0$  such that if  $\varepsilon < \varepsilon_0$  and  $R > R_1$ , then

$$\begin{aligned} & |p_R^{(j)}(0)|_{L^2} \leq \varepsilon_1, \quad \sum_{1 \leq j \leq k} |H_{R,j}|_{L^{\frac{6}{5}} \cap D_0^1}^2 \leq \varepsilon_1, \\ \text{and } & \sum_{2 \leq j \leq k} \left( \sum_{1 \leq m \leq j-1} \binom{j}{m-1} |\rho_R^{(j+1-m)}(0)|_{L^{\frac{3}{2}}} \right)^2 \leq \varepsilon_1. \end{aligned}$$

Now we use an iteration method to prove the lemma. Consider the following decoupled elliptic system of infinitely iterated equations:

$$(4.12) \quad \begin{aligned} L(w_{R,j}^n) &= j \operatorname{div}(\rho_{R,0} v_R(0)) w_{R,j}^n \\ &+ G_{R,j-1}^{n-1} + H_{R,j} - \nabla p_R^{(j)}(0) - \rho_{R,0} g_{R,j+1} \quad \text{in } \Omega_R, \end{aligned}$$

where

$$\begin{aligned} G_{R,j-1}^{n-1} &= G_{j-1}(w_{R,1}^{n-1}, \dots, w_{R,j-1}^{n-1}) \quad \text{for } j \geq 2 \quad \text{and } n \geq 1, \\ G_{R,j-1}^{n-1} &= 0 \quad \text{for } n \geq 1 \quad \text{and } j = 0, 1, \\ \text{and } w_{R,j}^0 &= 0 \quad \text{for } 1 \leq j \leq k-1. \end{aligned}$$

The solution  $w_{R,j}^n$  is constructed by the following process:

(i) first find  $(w_{R,0}^1, w_{R,1}^1, \dots, w_{R,k}^1)$  satisfying the system

$$\left\{ \begin{aligned} L(w_{R,k}^1) &= k \operatorname{div}(\rho_{R,0} v_R(0)) w_{R,k}^1 + H_{R,k} - \nabla p_R^{(k)}(0) - \rho_{R,0} g_{R,k+1}, \\ L(w_{R,k-1}^1) &= (k-1) \operatorname{div}(\rho_{R,0} v_R(0)) w_{R,k-1}^1 + H_{R,k-1} - \nabla p_R^{(k-1)}(0) - \rho_{R,0} g_{R,k}, \\ &\vdots \\ L(w_{R,0}^1) &= H_{R,0} - \nabla p_0 - \rho_{R,0} g_{R,1}, \end{aligned} \right.$$

(ii) then find  $(w_{R,0}^2, \dots, w_{R,k}^2)$  such that

$$\left\{ \begin{aligned} L(w_{R,k}^2) &= k \operatorname{div}(\rho_{R,0} v_R(0)) w_{R,k}^2 + G_{R,k-1}^1 + H_{R,k} - \nabla p_R^{(k)}(0) - \rho_{R,0} g_{R,k+1}, \\ L(w_{R,k-1}^2) &= (k-1) \operatorname{div}(\rho_{R,0} v_R(0)) w_{R,k-1}^2 \\ &\quad + G_{R,k-2}^1 + H_{R,k-1} - \nabla p_R^{(k-1)}(0) - \rho_{R,0} g_{R,k}, \\ &\vdots \\ L(w_{R,0}^2) &= H_{R,0} - \nabla p_0 - \rho_{R,0} g_{R,1}, \end{aligned} \right.$$

(iii) and in this way we find  $(w_{R,0}^n, \dots, w_{R,k}^n)$  such that

$$\begin{cases} L(w_{R,k}^n) = k \operatorname{div}(\rho_{R,0} v_R(0)) w_{R,k}^n + G_{R,k-1}^{n-1} + H_{R,k} - \nabla p_R^{(k)}(0) - \rho_{R,0} g_{R,k+1}, \\ L(w_{R,k-1}^n) = (k-1) \operatorname{div}(\rho_{r,0} v_R(0)) w_{R,k-1}^n \\ \quad + G_{R,k-2}^{n-1} + H_{R,k-1} - \nabla p_R^{(k-1)}(0) - \rho_{R,0} g_{R,k}, \\ \quad \vdots \\ L(w_{R,0}^n) = H_{R,0} - \nabla p_0 - \rho_{R,0} g_{R,1}. \end{cases}$$

Since every system is decoupled, the existence of the solution  $(g_{R,0}^n, \dots, g_{R,k}^n)$  in  $D_0^1(\Omega_R) \times \dots \times D_0^1(\Omega_R)$  is always guaranteed by the classical elliptic theory (see for instance [1]), provided  $\varepsilon$  is sufficiently small.

We now prove that  $\sum_{0 \leq j \leq k} |w_{R,j}^n|_{D_0^1}$  is uniformly bounded on  $n$ . Multiplying (4.12) by  $w_{R,j}^n$  and integrating over  $\Omega_R$ , by integration by parts, we have for  $j \geq 2$

$$\begin{aligned} & \int L w_{R,j}^n \cdot w_{R,j}^n dx \\ &= j \int \operatorname{div}(\rho_{R,0} v_R(0)) |w_{R,j}^n|^2 dx + \int G_{R,j-1}^{n-1} \cdot w_{R,j}^n dx \\ & \quad + \int H_{R,j} \cdot w_{R,j}^n dx - \int p_R^{(j)}(0) \operatorname{div} w_{R,j}^n dx + \int \rho_{R,0} g_{R,j+1} \cdot w_{R,j}^n dx \\ & \leq 2jC |\rho_{R,0}|_{L^6} |\nabla v_R(0)|_{L^2} |\nabla w_{R,j}^n|_{L^2}^2 + C(|G_{R,j-1}^{n-1}|_{L^{\frac{6}{5}}}^2 + |H_{R,j}|_{L^{\frac{6}{5}}} + |p_R^{(j)}(0)|_{L^2}^2) \\ & \quad + C|\rho_{R,0}|_{L^{\frac{3}{2}}}^2 |\nabla g_{R,j+1}|_{L^2}^2 + \frac{\mu}{4} |\nabla w_{R,j}^n|_{L^2}^2. \end{aligned}$$

Thus if  $\varepsilon_1$  and  $R_1$  are small and large enough, respectively, then

$$\begin{aligned} |\nabla w_{R,j}^n|_{L^2}^2 & \leq C(|G_{R,j-1}^{n-1}|_{L^{\frac{6}{5}}}^2 + |H_{R,j}|_{L^{\frac{6}{5}}}^2 + |p_R^{(j)}(0)|_{L^2}^2) + C\varepsilon_1^2 |\nabla g_{R,j+1}|_{L^2}^2 \\ & \leq C \left( \left( \sum_{1 \leq m \leq j-1} \binom{j}{m} |\rho_R^{(j+1-m)}(0)|_{L^{\frac{3}{2}}} \right)^2 \max_{1 \leq m \leq j-1} |\nabla w_{R,m}^{n-1}|_{L^2}^2 \right) \\ & \quad + C(|H_{R,j}|_{L^{\frac{6}{5}}}^2 + |p_R^{(j)}(0)|_{L^2}^2) + C\varepsilon_1^2 |\nabla g_{R,j+1}|_{L^2}^2 \\ & \leq \varepsilon_2 \left( \max_{1 \leq m \leq j-1} |\nabla w_{R,m}^n|_{L^2}^2 \right) + \varepsilon_2 + \varepsilon_2 |\nabla g_{j+1}|_{L^2}^2 \end{aligned}$$

for  $j \geq 2$ , where  $\varepsilon_2 = C\varepsilon_1$ . For the case  $j = 0, 1$ , we have

$$\begin{aligned} |\nabla w_{R,1}^n|_{L^2}^2 & \leq \varepsilon_2 + \varepsilon_2 |\nabla g_2|_{L^2}^2, \\ |\nabla w_{R,0}^n|_{L^2}^2 & \leq \varepsilon_2 + \varepsilon_2 |\nabla g_1|_{L^2}^2. \end{aligned}$$

Fixing  $R$ , let us define  $|\nabla w_{R,j}^n|_{L^2}^2$  and  $|\nabla g_{R,j}|_{L^2}^2$  by  $a_j^n$  and  $A_j$ , respectively. Then we have

$$(4.13) \quad a_j^n \leq \varepsilon_2 \left( \max_{1 \leq m \leq j-1} a_m^{n-1} \right) + \varepsilon_2 + \varepsilon_2 A_{j+1} \quad \text{for } j \geq 0, n \geq 1,$$

where  $a_{-1}^{n-1} = a_0^{n-1} = a_j^0 = 0$  for  $0 \leq j \leq k, n \geq 1$ . Now we claim that if  $\varepsilon_2 < \frac{1}{2}$ , then there holds

$$(4.14) \quad a_j^n \leq \left( \frac{\varepsilon_2}{1 - \varepsilon_2} + A\varepsilon_2^k \right) \left( \frac{1 - \varepsilon_2}{1 - 2\varepsilon_2} \right)$$

for  $1 \leq j \leq k$  and  $n \geq 1$ , where  $A = \max_{1 \leq j \leq k+1} A_j$ .

By induction we first observe that

$$\begin{aligned} a_k^1 &\leq \varepsilon_2 + \varepsilon_2 A, \\ a_{k-1}^1 &\leq \varepsilon_2 + \varepsilon_2(\varepsilon_2 + \varepsilon_2 A) = \varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^2 A, \\ a_{k-2}^1 &\leq \varepsilon_2 + \varepsilon_2(\varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^2 A) = \varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^3 + \varepsilon_2^3 A, \\ &\vdots \\ a_1^1 &\leq \varepsilon_2 + \varepsilon_2^2 + \cdots + \varepsilon_2^k + \varepsilon_2^k A \leq \frac{\varepsilon_2}{1 - \varepsilon_2} + \varepsilon_2^k A. \end{aligned}$$

Let  $B = \frac{\varepsilon_2}{1 - \varepsilon_2} + \varepsilon_2^k A$ . Then

$$\begin{aligned} a_k^2 &\leq \varepsilon_2 B + \varepsilon_2 + \varepsilon_2 A, \\ a_{k-1}^2 &\leq \varepsilon_2 B + \varepsilon_2 + \varepsilon_2(\varepsilon_2 B + \varepsilon_2 + \varepsilon_2 A) = \varepsilon_2 + \varepsilon_2^2 + \varepsilon_2 B + \varepsilon_2^2 B + \varepsilon_2^2 A, \\ a_{k-2}^2 &\leq \varepsilon_2 B + \varepsilon_2 + \varepsilon_2(\varepsilon_2 + \varepsilon_2^2 + \varepsilon_2 B + \varepsilon_2^2 B + \varepsilon_2^2 A) \\ &= \varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^3 + (\varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^3)B + \varepsilon_2^3 A, \\ &\vdots \\ a_1^2 &\leq \varepsilon_2 + \varepsilon_2^2 + \cdots + \varepsilon_2^k + (\varepsilon_2 + \varepsilon_2^2 + \cdots + \varepsilon_2^k)B + \varepsilon_2^k A \leq B + \frac{\varepsilon_2}{1 - \varepsilon_2} B. \end{aligned}$$

Suppose that

$$a_1^n \leq \left( 1 + \frac{\varepsilon_2}{1 - \varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1 - \varepsilon_2} \right)^{n-1} \right) B$$

for some  $n \geq 2$ . We then have

$$\begin{aligned}
a_k^{n+1} &\leq \varepsilon_2 \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B + \varepsilon_2 + \varepsilon_2 A, \\
a_{k-1}^{n+1} &\leq \varepsilon_2 \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B \\
&\quad + \varepsilon_2 + \varepsilon_2^2 \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B + \varepsilon_2^2 + A\varepsilon_2^2 \\
&= (\varepsilon_2 + \varepsilon_2^2) \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B + \varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^2 A, \\
a_{k-2}^{n+1} &\leq (\varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^3) \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B \\
&\quad + \varepsilon_2 + \varepsilon_2^2 + \varepsilon_2^3 + \varepsilon_2^3 A, \\
&\quad \vdots \\
a_1^{n+1} &\leq (\varepsilon_2 + \varepsilon_2^2 + \cdots + \varepsilon_2^k) \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B \\
&\quad + \varepsilon_2 + \cdots + \varepsilon_2^k + \varepsilon_2^k A \\
&\leq \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^n \right) B.
\end{aligned}$$

Thus by induction we have for all  $n \geq 1$  and  $j \geq 1$

$$a_j^n \leq \left( 1 + \frac{\varepsilon_2}{1-\varepsilon_2} + \cdots + \left( \frac{\varepsilon_2}{1-\varepsilon_2} \right)^{n-1} \right) B.$$

Since  $\varepsilon_2 < \frac{1}{2}$  and hence  $\frac{\varepsilon_2}{1-\varepsilon_2} < 1$ , we the above inequality implies the estimate (4.14).

If  $\varepsilon_2 < \frac{1}{4}$  and  $R$  is large, then we therefore have

$$(4.15) \quad |\nabla w_{R,j}^n|_{L^2}^2 \leq \frac{1}{2} + |\nabla g_{R,j}|_{L^2}^2 \leq 1 + |g_j|_{D_0^1}^2 \quad \text{for all } n \geq 1, 0 \leq j \leq k.$$

Following the elliptic regularity estimate in the proof of Lemma 3.2, one can prove from (4.15) that

$$(4.16) \quad |w_{R,j}^n|_{D_0^1 \cap D^{2(k-j)+3}} \leq C(1 + |g_j|_{D_0^1}) \leq Cc_0$$

for all large  $R$ ,  $n \geq 1$  and  $0 \leq j \leq k$ . We leave the details of proof to the readers.

Now using the weak compactness of Sobolev space, we find a solution to the problem (4.10)

$$(w_{R,0}, w_{R,1}, \cdots, w_{R,k}) \in D_0^1 \cap D^{2k+3} \times D_0^1 \cap D^{2k+1} \times \cdots \times D_0^1 \cap D^3$$

such that<sup>2</sup>

$$w_{R,j}^n \rightharpoonup w_{R,j} \quad \text{in } D_0^1 \cap D^{2(k-j)+3}, \quad |w_{R,j}|_{D_0^1 \cap D^{2(k-j)+3}} \leq Cc_0.$$

Finally, we prove the uniqueness. Let  $(w_{R,0}^1, \dots, w_{R,k}^1)$  and  $(w_{R,0}^2, \dots, w_{R,k}^2)$  be the solutions to the elliptic system (4.10). Then we have

$$\begin{aligned} & L(w_{R,j}^1 - w_{R,j}^2) \\ &= j \operatorname{div}(\rho_{R,0} v_R(0))(w_{R,j}^1 - w_{R,j}^2) \\ & - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \rho_R^{(j+1-m)}(0)(w_{R,m}^1 - w_{R,m}^2). \end{aligned}$$

Thus by integration by parts we have

$$\begin{aligned} |\nabla(w_{R,k}^1 - w_{R,k}^2)|_{L^2}^2 &\leq \varepsilon_2 \max_{1 \leq m \leq k-1} |\nabla(w_{R,j}^1 - w_{R,j}^2)|_{L^2}^2, \\ |\nabla(w_{R,k-1}^1 - w_{R,k-1}^2)|_{L^2}^2 &\leq \varepsilon_2 \max_{1 \leq m \leq k-2} |\nabla(w_{R,j}^1 - w_{R,j}^2)|_{L^2}^2, \\ &\vdots \\ |\nabla(w_{R,1}^1 - w_{R,1}^2)|_{L^2}^2 &\leq \varepsilon_2 |\nabla(w_{R,2}^1 - w_{R,2}^2)|_{L^2}^2, \\ |\nabla(w_{R,0}^1 - w_{R,0}^2)|_{L^2}^2 &\leq \varepsilon_2 |\nabla(w_{R,1}^1 - w_{R,1}^2)|_{L^2}^2. \end{aligned}$$

Therefore,

$$\sum_{0 \leq j \leq k} |\nabla(w_{R,j}^1 - w_{R,j}^2)|_{L^2}^2 \leq k \frac{\varepsilon_2}{1 - \varepsilon_2} \sum_{0 \leq j \leq k} |\nabla(w_{R,j}^1 - w_{R,j}^2)|_{L^2}^2$$

and hence choosing  $\varepsilon_2$  such that  $\frac{k\varepsilon_2}{1-\varepsilon_2} < 1$ , the uniqueness is proved. This completes the proof of lemma.  $\square$

**Lemma 4.3.** *If  $(u_0, u^{(1)}(0), \dots, u^{(k)}(0))$  is the given pair satisfying the compatibility condition (4.6) and  $(w_{R,0}, w_{R,1}, \dots, w_{R,k})$  is the pair of solution to the elliptic system (4.10), then extending  $(w_{R,0}, \dots, w_{R,k})$  in  $\Omega$  by defining 0 outside  $\Omega_R$ , the extended pair converges to  $(u_0, u^{(1)}(0), \dots, u^{(k)}(0))$  in  $D_0^1(\Omega)$ .*

*Proof.* Let us denote the pair of extensions by  $(w_{R,0}, \dots, w_{R,k})$  again. Let  $\varphi_R$  be the cut-off function as above and  $w_j = u^{(j)}(0)$ . Then we have for  $0 \leq j \leq k$

$$\begin{aligned} L(\varphi_R w_j) &= j \operatorname{div}(\rho_0 v(0)) \varphi_R w_j - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \rho^{(j+1-m)}(0) \varphi_R w_m \\ &+ \varphi_R H_j - \nabla(\varphi_R p^{(j)}(0)) - \rho \varphi_R g_{R,j+1} + \Phi_{R,j}, \end{aligned}$$

<sup>2</sup>Actually using the argument of uniqueness proof, we can prove that  $w_{R,j}^n$  converges strongly to  $w_{R,j}$  in  $D_0^1$ .

where  $\Phi_{R,j} = L(\varphi_R w_j) - \varphi_R L w_j + \nabla \varphi_R p^{(j)}(0)$ . Thus the difference  $w_{R,j} - \varphi_R w_j$  satisfies the system in  $\Omega_R$

$$\begin{aligned}
& L(w_{R,j} - \varphi_R w_j) \\
&= j(\operatorname{div}(\rho_{R,0} v_R(0)) w_{R,j} - \operatorname{div}(\rho_0 v(0)) \varphi_R w_j) \\
(4.17) \quad & - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \left( \rho_R^{(j+1-m)}(0) w_{R,m} - \rho^{(j+1-m)}(0) \varphi_R w_m \right) \\
& + (H_{R,j} - \varphi_R H_j) - \nabla(p_R^{(j)}(0) - \varphi_R p^{(j)}(0)) \\
& - R^{-3} g_{R,j+1} - \Phi_{R,j}.
\end{aligned}$$

Multiplying (4.17) by  $\Gamma_{R,j} = w_{R,j} - \varphi_R w_j$  and integrating over  $\Omega_R$ , we have

$$\begin{aligned}
& \mu |\nabla \Gamma_{R,j}|_{L^2}^2 + (\lambda + \mu) \int (\operatorname{div} \Gamma_{R,j})^2 dx \\
&= \int j \operatorname{div}(\rho_{R,0} v_R(0)) |\Gamma_{R,j}|^2 dx \\
& + \int j (\operatorname{div}(\rho_{R,0} v_R(0) - \rho_0 v(0)) \varphi_R g_j \cdot \Gamma_{R,j} dx \\
& - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho_R^{(j+1-m)}(0) \Gamma_{R,m}) \cdot \Gamma_{R,j} dx \\
(4.18) \quad & - \sum_{1 \leq m \leq j-1} \binom{j}{m-1} \int (\rho_R^{(j+1-m)}(0) - \rho^{(j+1-m)}(0)) \varphi_R g_m \cdot \Gamma_{R,j} dx \\
& + \int (H_{R,j} - \varphi_R H_j) \cdot \Gamma_{R,j} dx \\
& + \int (p_R^{(j)}(0) - \varphi_R p^{(j)}(0)) \operatorname{div} \Gamma_{R,j} dx \\
& - R^{-3} \int g_{R,j+1} \cdot \Gamma_{R,j} dx \\
& - \int \Phi_{R,j} \cdot \Gamma_{R,j} dx \\
& \equiv \sum_{1 \leq i \leq 8} I_j^i.
\end{aligned}$$

We estimate each  $I_j^i$  as follows:

$$\begin{aligned}
I_j^1 &= -2j \int \rho_{R,0} v_R(0) \cdot (\Gamma_{R,j} \cdot \nabla \Gamma_{R,j}) dx \\
&\leq C |\rho_{R,0}|_{L^6} |\nabla v_R(0)|_{L^2} |\nabla \Gamma_{R,j}|_{L^2}^2 \leq \varepsilon_2 |\nabla \Gamma_{R,j}|_{L^2}^2,
\end{aligned}$$

where  $\varepsilon_2$  is the small constant appearing in Lemma 4.2.

$$\begin{aligned}
I_j^2 &\leq C |\operatorname{div}(\rho_{R,0} v_R(0) - \rho_0 v(0))|_{L^{\frac{3}{2}}} |g_j|_{L^6} |\nabla \Gamma_{R,j}|_{L^2} \\
&\leq C c_0^2 |\operatorname{div}(\rho_{R,0} v_R(0) - \rho_0 v(0))|_{L^{\frac{3}{2}}}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2,
\end{aligned}$$



$$I_j^3 \leq \varepsilon_2 \max_{1 \leq m \leq j-1} |\nabla \Gamma_{R,m}|_{L^2}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2,$$

$$\begin{aligned} I_j^4 &\leq C \sum_{1 \leq m \leq j-1} |\rho_R^{(j+1-m)}(0) - \rho^{(j+1-m)}(0)|_{L^{\frac{3}{2}}}^2 |g_m|_{L^6}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2 \\ &\leq C c_0^2 \sum_{2 \leq m \leq j} |\rho_R^{(m)}(0) - \rho^{(m)}(0)|_{L^{\frac{3}{2}}}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2, \end{aligned}$$

$$I_j^5 \leq C |H_{R,j} - \varphi_R H_j|_{L^{\frac{6}{5}}}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2,$$

$$I_j^6 \leq C |p_R^{(j)}(0) - \varphi_R p^{(j)}(0)|_{L^2}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2,$$

$$I_j^7 \leq C R^{-6} |g_{R,j+1}|_{L^{\frac{6}{5}}}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2,$$

and finally

$$I_j^8 \leq C |\Phi_{R,j}|_{L^{\frac{6}{5}}}^2 + \frac{\mu}{32} |\nabla \Gamma_{R,j}|_{L^2}^2.$$

Substituting all these estimate into (4.18) and then choose  $\varepsilon_2$  to be sufficiently small, we have for all  $0 \leq j \leq k$

$$(4.19) \quad |\nabla \Gamma_{R,j}|_{L^2}^2 \leq \varepsilon_2 \max_{1 \leq m \leq j-1} |\nabla \Gamma_{R,m}|_{L^2}^2 + \Lambda_{R,j},$$

where

$$\begin{aligned} \Lambda_{R,j} &= C c_0^2 |\operatorname{div}(\rho_{R,0} v_R(0) - \rho_0 v(0))|_{L^{\frac{3}{2}}}^2 + C c_0^2 \sum_{2 \leq m \leq j} |\rho_R^{(m)}(0) - \rho^{(m)}(0)|_{L^{\frac{3}{2}}}^2 \\ &\quad + C |p_R^{(j)}(0) - \varphi_R p^{(j)}(0)|_{L^2}^2 + C R^{-6} |g_{R,j+1}|_{L^{\frac{6}{5}}}^2 + C |\Phi_{R,j}|_{L^{\frac{6}{5}}}^2. \end{aligned}$$

Here we interpret  $\max_{1 \leq m \leq j-1} |\nabla \Gamma_{R,m}|_{L^2}^2$  as 0 in case when  $j = 0, 1$ .

One can easily show that  $\Lambda_{R,j} = o(1)$ , where  $o(1)$  denotes a function of  $R$ , which tends to zero as  $R \rightarrow \infty$ . We leave the details of its proof to the readers.

Now let  $b_{R,j} = |\nabla \Gamma_{R,j}|_{L^2}^2$ . Then we have for  $j \geq 2$

$$b_{R,j} \leq \varepsilon_2 \left( \max_{1 \leq m \leq j-1} b_{R,m} \right) + o(1)$$

and

$$b_{R,1} = o(1), \quad b_{R,0} = o(1).$$

Thus letting  $\Gamma_R = \max_{0 \leq m \leq k} b_{R,m}$ , we have  $\Gamma_R = o(1)$ . Therefore we have just proven that  $|w_{R,j} - \varphi_R w_j|_{D_0^1(\Omega_R)} \rightarrow 0$  as  $R \rightarrow \infty$ . Since  $\varphi_R w_j \rightarrow w_j$  in  $D_0^1$ , this implies that each extension  $w_{R,j}$  converges to  $w_j$  in  $D_0^1$ .  $\square$

*Proof of Proposition 4.1.* To prove the existence, we consider the following initial boundary value problem together with (4.9)

$$(4.20) \quad \rho_R(u_R)_t + Lu_R + \nabla p_R = \rho_R(f_R - v_R \cdot \nabla v_R) \quad \text{in } (0, T_*) \times \Omega_R,$$

$$(4.21) \quad u_R^{(j)}(0) = w_{R,j} \quad (0 \leq j \leq k+1) \quad \text{in } \Omega_R,$$

where  $w_{R,j}$  are the functions constructed in Lemma 4.2 and  $w_{R,0} = u_{R,0} = u_R(0)$ .

From Lemma 4.3 and the observation (4.11), we deduce that

$$(4.22) \quad |\rho_{R,0}|_{L^1 \cap H^{2k+3}} \leq \varepsilon, \quad 1 + |u_{R,0}|_{D_0^1} + \sum_{1 \leq j \leq k+1} |w_{R,j}|_{D_0^1} \leq c_0,$$

and for all  $0 \leq j \leq k+1$ ,

$$(4.23) \quad |v_R^{(j)}(0)|_{D_0^1 \cap D^{2(k-j)+3}} \leq 1 + c_{0,1,j},$$

and for all  $0 \leq l \leq k$  and  $0 \leq j \leq k-l$ ,

$$(4.24) \quad \begin{aligned} & \sup_{0 \leq t \leq T_*} \left( |v_R^{(j)}(t)|_{D_0^1 \cap D^{2l+1}} \right) + \int_0^{T_*} |v_R^{(j)}(t)|_{D^2 \cap D^{2l+2}}^2 dt \leq 1 + c_{l,2,j}, \\ & \sup_{0 \leq t \leq T_*} \left( |v_R^{(j)}(t)|_{D_0^1 \cap D^{2l+2}} \right) \\ & + \int_0^{T_*} \left( |v_R^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}}^2 + |v_R^{(j)}(t)|_{D^{2l+3}}^2 \right) dt \leq 1 + c_{l,3,j}, \\ & \text{ess sup}_{0 < t < T_*} \left( |v_R^{(j+1)}(t)|_{D_0^1 \cap D^{2l+1}} + |v_R^{(j)}(t)|_{D^{2l+3}} \right) \\ & + \int_0^{T_*} \left( |v_R^{(j+1)}(t)|_{D^{2l+2}}^2 + |v_R^{(j)}(t)|_{D^{2l+4}}^2 \right) dt \leq 1 + c_{l,4,j} \end{aligned}$$

Therefore, applying all the estimates in Section 3 to  $(\rho_R, u_R)$  with  $(\rho_{R,0}, w_R, v_R)$  satisfying (4.23) and (4.24) instead of  $(\rho_0, g_j, v)$  in (3.1), we conclude that if  $\varepsilon$  is sufficiently small, then for each  $R > R_1$ , the solution  $(\rho_R, u_R)$  satisfies the estimate (3.21) and (3.22) with the domain being  $\Omega_R$ . We extend  $(\rho_R, u_R)$  by defining to be zero outside  $\Omega_R$ . Then by the uniform estimate (3.22) on  $R$ , we deduce that there exists a sequence  $\{R_j\}$ ,  $R_j \rightarrow \infty$  as  $j \rightarrow \infty$  such that  $(\rho_{R_j}, u_{R_j})$  converges in a weak or weak-\* sense to a limit  $(\rho, u)$ . Moreover, since  $(\rho, u)$  also satisfies (3.22) with the domain being  $\Omega_R$  for each  $R > R_1$ , it follows that for  $0 \leq j \leq k+1$

$$(4.25) \quad \begin{aligned} & \rho \in L^\infty(0, T_*), \quad (\rho^{(j)}, p^{(j)}) \in L^\infty(0, T_*; H^{2(k-j)+3}), \\ & u^{(j)} \in L^\infty(0, T_*; D_0^1 \cap D^{2(k-j)+3}), \quad \sqrt{\rho} u^{(k+1)} \in L^\infty(0, T_*; L^2). \end{aligned}$$

We will show that  $(\rho, u)$  is a solution to the original problem (4.1)-(4.4). It is obvious that  $(\rho, u)$  satisfies the boundary conditions in (4.3) and (4.4). Let  $R > R_1$  be a fixed large number. Then since for all sufficiently large  $l$ ,  $(\rho_{R_l}, u_{R_l})$  satisfies the uniform estimate (3.21) and (3.22) with the domain being  $\Omega_R$ , it follows from a standard compactness result (see [24]) that a subsequence of  $(\rho_{R_l}^{(j)}, u_{R_l}^{(j)})$  converges to  $(\rho^{(j)}, u^{(j)})$  in  $C([0, T_*]; H^1(\Omega_R))$  for all  $0 \leq j \leq k$ . Using this result together with Lemma 4.3, we can show that  $(\rho, u)$  satisfies the equations (2.7)-(2.8) in  $(0, T_*) \times \Omega_R$  and  $(\rho(0), u(0)) = (\rho_0, u_0)$  in  $\Omega_R$ . Let  $w_j$  be the limit of  $w_{R,j}$

constructed in Lemma 4.2 as in Lemma 4.3. Then  $u^{(j)}(0) = w_j$  in  $\Omega_R$ . Since  $R$  can be arbitrarily large, we have proved the existence of a solution  $(\rho, u)$  to the original problem (4.1)-(4.2) satisfying the compatibility condition 4.6 and the regularity (4.25) as in [5]. The uniqueness of solutions with this regularity can be proved by a simple energy estimate. We deduce from a classical embedding result that  $u^{(j)} \in C([0, T_*]; D_0^1 \cap D^{2(k-j)+3})$  for all  $0 \leq j \leq k$ . Then the time-continuity of  $\rho$  follows from Lemma 2.3. This completes the proof of Proposition 4.1.  $\square$

## 5. PROOF OF THEOREM 1.1

In this section, we prove Theorem 1.1. We construct a sequence of approximate solutions to the linearized problem which converges to a solution to the original nonlinear problem in a strong sense.

Let us invoke the system of compatibility conditions for linearized problem such that

$$(5.1) \quad L(w_j) = j \operatorname{div}(\rho_0 v(0)) w_j + G_{j-1} + H_j - \nabla p^{(j)}(0) - \rho_0 g_{j+1}$$

for some given  $\rho_0 \in L^1 \cap H^{2k+3}$ ,  $g_j \in D_0^1 \cap D^{2(k-j)+3}$  and all  $1 \leq j \leq k+1$ , where

$$\begin{aligned} G_{j-1} &= 0 \quad \text{for } j = 0, 1, \\ G_{j-1} &= - \sum_{1 \leq m \leq j-1} \binom{j}{m} \rho^{(j+1-m)}(0) w_m \quad \text{for } j \geq 2, \\ H_j &= \sum_{0 \leq m \leq j} \binom{j}{m} \rho^{(j-m)}(0) (f - v \cdot \nabla v)^{(j)}(0). \end{aligned}$$

Now let  $w_{0,0}, w_{0,1}, \dots, w_{0,k}$  be the unique solutions of (5.1) with  $v = 0$ . This is possible for small initial density  $\rho_0$ . Then we find a unique solution  $(\underline{\rho}, \underline{u})$  to the linearized problem (4.1)–(4.4) with initial data  $(\underline{u}(0), \underline{u}^{(1)}(0), \dots, \underline{u}^{(k)}(0))$  satisfying

$$(\underline{u}(0), \underline{u}^{(1)}(0), \dots, \underline{u}^{(k)}(0)) = (w_{0,0}, w_{0,1}, \dots, w_{0,k}).$$

In this turn, we find the unique solutions  $w_{1,0}, w_{1,1}, \dots, w_{1,k}$  of the system (5.1) and also the unique solution  $(\rho_1, u_1)$  to the linear problem (4.1)–(4.4) with  $v = \underline{u}$  and initial data satisfying

$$(u_1(0), u_1^{(1)}(0), \dots, u_1^{(k)}(0)) = (w_{1,0}, w_{1,1}, \dots, w_{1,k}).$$

In this way, we find  $w_{n,j}$  and  $u_n$  for  $n \geq 1, 0 \leq j \leq k$  with  $v = u_{n-1}$  of the equations (5.1), and (4.1)–(4.4), respectively satisfying

$$(u_n(0), u_n^{(1)}(0), \dots, u_n^{(k)}(0)) = (w_{n,0}, w_{n,1}, \dots, w_{n,k}).$$

If  $\sum_{1 \leq j \leq k+1} |g_j|_{D_0^1} \leq c_0$  for some fixed  $c_0$  and  $\varepsilon$  is sufficiently small, then from Proposition 3.13, Lemma 4.2 and Lemma 4.3 we deduce that for all  $n \geq 1$  and for all  $0 \leq j \leq k$

$$\sum_{0 \leq j \leq k} |w_{n,j}|_{D_0^1 \cap D^{2(k-j)+3}} \leq C c_0,$$

$$\begin{aligned}
(5.2) \quad & \sup_{0 < t < T_*} \left( |\rho_n(t)|_{L^1} + |(\rho_n^{(j)}(t), (p_n - p_n(0))^{(j)}(t))|_{H^{2(k-j)+3}} \right) \leq \tilde{C}\varepsilon \\
& \sup_{0 \leq t \leq T_*} \left( |u_n^{(j)}(t)|_{D_0^1 \cap D^{2(k-j)+3}} \right) + \int_0^{T_*} |u_n^{(j)}(t)|_{D^{2(k-j)+4}}^2 dt \\
& \quad + \operatorname{ess\,sup}_{0 < t < T_*} |u_n^{(k+1)}(t)|_{D_0^1} + \int_0^{T_*} |u_n^{(k+1)}(t)|_{D^2}^2 dt \\
& \quad + \operatorname{ess\,sup}_{0 \leq t \leq T_*} |\sqrt{\rho_n} u_n^{(k+1)}(t)|_{L^2} \leq \tilde{C}.
\end{aligned}$$

Throughout the proof, we denote by  $\tilde{C} \geq 1$  a generic positive constant depending only on  $c_0$  and the parameters of  $C$ , but independent of  $n$ .

We first show the full sequence  $\{(w_{n,0}, \dots, w_{n,k})\}_{n \geq 1}$  converges to  $(w_0, \dots, w_k)$  in  $D_0^1 \cap D^{2k+3} \times \dots \times D_0^1 \cap D^3$ .

Let us denote  $\rho_{n-1}^{(j)}(0)$  and  $p_{n-1}^{(j)}(0)$  by  $\rho^{(j)}(0)$  and  $p^{(j)}(0)$  with  $v^{(l)}(0) = w_{n-1,l}$  for  $0 \leq l \leq j$ . Then  $(w_{n,0}, \dots, w_{n,k})$  satisfies the following equations: for all  $0 \leq j \leq k$

$$L(w_{n,j}) = j \operatorname{div}(\rho_0 w_{n-1,0}) w_{n,j} + G_{n,j-1} + H_{n-1,j} - \nabla p_{n-1}^{(j)}(0) - \rho_0 g_{j+1},$$

where

$$\begin{aligned}
G_{n,j-1} &= 0 \quad \text{for } j = 0, 1, \\
G_{n,j-1} &= - \sum_{1 \leq m \leq j-1} \binom{j}{m} \rho_{n-1}^{(j+1-m)}(0) w_{n,m} \quad \text{for } j \geq 2
\end{aligned}$$

and

$$H_{n-1,j} = \sum_{0 \leq l \leq j} \binom{j}{l} \rho_{n-1}^{(j-l)}(0) \left( f^{(l)}(0) - \sum_{0 \leq m \leq l} \binom{l}{m} w_{n-1,l-m} \cdot \nabla w_{n-1,m} \right).$$

Now let  $\bar{w}_{n+1,j} = w_{n+1,j} - w_{n,j}$ . Then

$$\begin{aligned}
(5.3) \quad L(\bar{w}_{n+1,j}) &= j \operatorname{div}(\rho_0 \bar{w}_{n,0}) w_{n+1,j} + j \operatorname{div}(\rho_0 w_{n-1,0}) \bar{w}_{n+1,j} \\
& \quad + \bar{G}_{n+1,j-1} + \bar{H}_{n,j} - \nabla(p_{n+1}^{(j)}(0) - p_n^{(j)}(0)),
\end{aligned}$$

where

$$\begin{aligned}
\bar{G}_{n+1,j-1} &= - \sum_{1 \leq l \leq j-1} \binom{j}{l} (\rho_n^{(j+1-l)}(0) - \rho_{n-1}^{(j+1-l)}(0)) w_{n+1,l} \\
& \quad - \sum_{1 \leq l \leq j-1} \binom{j}{l} \rho_{n-1}^{(j+1-l)}(0) \bar{w}_{n+1,l}, \\
\bar{H}_{n,j} &= \sum_{0 \leq m \leq j} \binom{j}{l} (\rho_n^{(j-l)}(0) - \rho_{n-1}^{(j-l)}(0)) (f^{(l)}(0) \\
& \quad - \sum_{0 \leq m \leq l} \binom{l}{m} w_{n,l-m} \cdot \nabla w_{n,l}) \\
& \quad - \sum_{0 \leq l \leq j} \sum_{0 \leq m \leq l} \binom{j}{l} \binom{l}{m} \rho_{n-1}^{(j-l)}(0) (w_{n,l-m} \cdot \nabla w_{n,l} \\
& \quad - w_{n-1,l-m} \cdot \nabla w_{n-1,m}).
\end{aligned}$$

Then multiplying (5.3) by  $\bar{w}_{n,j}$  and integrating it over  $\Omega$ , we have

$$\begin{aligned} & \mu |\nabla \bar{w}_{n+1,j}|_{L^2}^2 + (\lambda + \mu) |\operatorname{div} \bar{w}_{n+1,j}|_{L^2}^2 \\ &= \int (j \operatorname{div}(\rho_0 w_{n-1,0}) |\bar{w}_{n+1,j}|^2 + j \operatorname{div}(\rho_0 \bar{w}_{n,0}) w_{n+1,j} \cdot \bar{w}_{n+1,j} \\ &\quad + \bar{G}_{n+1,j-1} \cdot \bar{w}_{n+1,j} + \bar{H}_{n,j} \cdot \bar{w}_{n+1,j} \\ &\quad + (p_n^{(j)} - p_{n-1}^{(j)}) \operatorname{div} \bar{w}_{n+1,j}) dx \\ &\equiv \sum_{1 \leq i \leq 5} I_i. \end{aligned}$$

Using Hölder and Sobolev inequalities, we get the following estimates.

$$\begin{aligned} I_1 &\leq C |\rho_0 w_{n-1,0}|_{L^3} |\nabla \bar{w}_{n+1,j}|_{L^2}^2 \leq C c_0 \varepsilon |\nabla \bar{w}_{n+1,j}|_{L^2}^2, \\ I_2 &\leq C |\rho_0|_{H^1} |\nabla \bar{w}_{n,0}|_{L^2} |\nabla w_{n+1,j}|_{L^2} |\nabla \bar{w}_{n+1,j}|_{L^2} \\ &\leq C c_0^2 \frac{\varepsilon^2}{\eta} |\nabla \bar{w}_{n,0}|_{L^2}^2 + \eta |\nabla \bar{w}_{n+1,j}|_{L^2}^2, \end{aligned}$$

where  $\eta \in (0, 1)$  is a small number. By the definition of  $\rho_n^{(j)}(0)$  and  $p_n^{(j)}(0)$ , one can easily show that

$$(5.4) \quad \begin{aligned} & |\rho_n^{(j)}(0) - \rho_{n-1}^{(j)}(0)|_{L^{\frac{3}{2}} \cap H^{2(k-j)+3}} \leq C c_0^j \varepsilon \sum_{0 \leq l \leq j-1} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}} \\ & |p_n^{(j)}(0) - p_{n-1}^{(j)}(0)|_{H^{2(k-j)+3}} \leq C c_0^j \varepsilon \sum_{0 \leq l \leq j-1} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}}. \end{aligned}$$

Hence

$$\begin{aligned} I_3 &\leq C c_0^{2j} \frac{\varepsilon^2}{\eta} \sum_{0 \leq l \leq j-1} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}}^2 + \eta |\nabla \bar{w}_{n+1,j}|_{L^2}^2 + C c_0^j \varepsilon \sum_{0 \leq l \leq j} |\nabla \bar{w}_{n+1,l}|_{L^2}^2 \\ I_4 &\leq C c_0^{2j} \frac{\varepsilon^2}{\eta} \sum_{0 \leq l \leq j-1} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}}^2 + C c_0^{2j} \frac{\varepsilon^2}{\eta} \sum_{0 \leq l \leq j} |\nabla \bar{w}_{n,l}|_{L^2}^2 + \eta |\nabla \tilde{w}_{n+1,j}|_{L^2}^2 \\ I_5 &\leq C c_0^{2j} \frac{\varepsilon^2}{\eta} \sum_{0 \leq l \leq j-1} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}}^2 + \eta |\nabla \bar{w}_{n+1,j}|_{L^2}^2. \end{aligned}$$

For sufficiently small  $\eta$  and  $\varepsilon$ , we have

$$(5.5) \quad \sum_{0 \leq l \leq j} |\nabla \bar{w}_{n+1,l}|_{L^2}^2 \leq C c_0^{2j} \varepsilon \sum_{0 \leq l \leq j} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}}^2.$$

Now we use the elliptic regularity for  $\bar{w}_{n+1,j}$  of (5.3). Then by (5.4) and (5.5) we have

$$\begin{aligned} & |\bar{w}_{n+1,j}|_{D_0^1 \cap D^{2(k-j)+3}} \\ & \leq C |\bar{w}_{n+1,j}|_{D_0^1} + C |\operatorname{div}(\rho_0 \bar{w}_{n,0}) w_{n+1,j}|_{D^{2(k-j)+1}} \\ & \quad + |\operatorname{div}(\rho_0 w_{n-1,0}) \bar{w}_{n+1,j}|_{D^{2(k-j)+1}} + |\bar{G}_{n+1,j-1}|_{D^{2k-j+1}} \\ & \quad + |\bar{H}_{n,j}|_{D^{2(k-j)+1}} + |p_n^{(j)}(0) - p_{n-1}^{(j)}(0)|_{D^{2(k-j)+2}} \\ & \leq C c_0^{2j} \varepsilon^{\frac{1}{2}} \sum_{0 \leq l \leq j} |\bar{w}_{n,l}|_{D_0^1 \cap D^{2(k-l)+3}} + C c_0^{2j} \varepsilon |\bar{w}_{n+1,j}|_{D_0^1 \cap D^{2(k-j)+3}}. \end{aligned}$$

Hence for small  $\varepsilon$  we conclude that

$$\sum_{0 \leq j \leq k} |\bar{w}_{n+1,j}|_{D_0^1 \cap D^{2(k-j)+3}} \leq \frac{1}{2} \sum_{0 \leq j \leq k} |\bar{w}_{n,j}|_{D_0^1 \cap D^{2(k-j)+3}}.$$

This means that

$$\sum_{n \geq 0} \sum_{0 \leq j \leq k} |\bar{w}_{n+1,j}|_{D_0^1 \cap D^{2(k-j)+3}} \leq C \sum_{0 \leq j \leq k} |w_{0,j}|_{D_0^1 \cap D^{2(k-j)+3}}$$

Therefore  $(w_{n,0}, \dots, w_{n,k})$  converges to a pair of functions  $(w_0, \dots, w_k)$  in  $D_0^1 \cap D^{2k+3} \times \dots \times D_0^1 \cap D^3$ . Apparently, these functions  $w_n$  satisfies by their regularity the equations

$$(5.6) \quad L(w_j) = j \operatorname{div}(\rho_0 w_0) w_j + G_{j-1} + H_j - \nabla p^{(j)}(0) - \rho_0 g_{j+1},$$

where

$$\begin{aligned} G_{j-1} &= 0 \quad \text{for } j = 0, 1, \\ G_{j-1} &= - \sum_{1 \leq l \leq j-1} \binom{j}{l} \rho^{(j+1-l)}(0) w_l \quad \text{for } j \geq 2, \\ H_j &= \sum_{0 \leq l \leq j} \binom{j}{l} \rho^{(j-l)}(0) \left( f^{(l)}(0) - \sum_{0 \leq m \leq l} \binom{l}{m} w_{l-m} \cdot \nabla w_m \right) \\ \rho^{(j+1)}(0) &= -\operatorname{div} \left( \sum_{0 \leq l \leq j} \binom{j}{l} \rho^{(j-l)}(0) w_l \right), \quad 0 \leq j \leq k. \end{aligned}$$

It can be easily shown by the argument of convergence that  $w_n$  are unique functions.

From now on, we show that the full sequence  $\{(\rho_n, u_n)\}$  of approximate solutions converges to a solution to the original problem (1.1)-(1.5) in a strong sense. To do this, let us define

$$\bar{\rho}_{n+1} = \rho_{n+1} - \rho_n, \quad \bar{u}_{n+1} = u_{n+1} - u_n \quad \text{and} \quad p_n = p(\rho_n).$$

Then from the equation (4.1), we derive

$$(5.7) \quad (\bar{\rho}_{n+1})_t + \operatorname{div}(\bar{\rho}_{n+1} u_n) + \operatorname{div}(\rho_n \bar{u}_n) = 0.$$

Multiplying this by  $\bar{\rho}_{n+1}$  and integrating over  $\Omega$ , we obtain

$$\begin{aligned} & \frac{d}{dt} \int |\bar{\rho}_{n+1}|^2 dx \\ & \leq C \int |\nabla u_n| |\bar{\rho}_{n+1}|^2 + (|\nabla \rho_n| |\bar{u}_n| + \rho_n |\nabla \bar{u}_n|) |\bar{\rho}_{n+1}| dx \\ & \leq C |\nabla u_n|_{L^\infty} |\bar{\rho}_{n+1}|_{L^2}^2 + C (|\nabla \rho_n|_{H^1} + |\rho_n|_{L^\infty}) |\nabla \bar{u}_n|_{L^2} |\bar{\rho}_{n+1}|_{L^2}. \end{aligned}$$

Hence it follows from the uniform bound (5.2) that

$$(5.8) \quad \frac{d}{dt} |\bar{\rho}_{n+1}|_{L^2}^2 \leq \eta^{-1} \tilde{C} |\bar{\rho}_{n+1}|_{L^2}^2 + \eta |\nabla \bar{u}_n|_{L^2}$$

for  $0 \leq t \leq T_*$ .

We need an estimate for  $|\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}$  in addition to (5.8). Multiplying (5.7) by  $\text{sgn}(\bar{\rho}_{n+1})|\bar{\rho}_{n+1}|^{\frac{1}{2}}$  and integrating over  $\Omega$ , we get

$$\begin{aligned} & \frac{d}{dt} \int |\bar{\rho}_{n+1}|^{\frac{3}{2}} dx \\ & \leq C \int |\nabla u_n| |\bar{\rho}_{n+1}|^{\frac{3}{2}} + (|\nabla \rho_n| |\bar{u}_n| + \rho_n |\nabla \bar{u}_n|) |\bar{\rho}_{n+1}|^{\frac{1}{2}} dx \\ & \leq C |\nabla u_n|_{L^\infty} |\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}^{\frac{3}{2}} + C |\rho_n|_{H^1} |\nabla \bar{u}_n|_{L^2} |\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}^{\frac{1}{2}}. \end{aligned}$$

Hence multiplying this by  $|\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}^{\frac{1}{2}}$  and using (5.2), we have

$$(5.9) \quad \frac{d}{dt} |\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}^2 \leq \eta^{-1} \tilde{C} |\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}^2 + \eta |\nabla \bar{u}_n|_{L^2}^2$$

for  $0 \leq t \leq T_*$ .

Next from the equation (4.2), we derive

$$\begin{aligned} & \rho_{n+1} (\bar{u}_{n+1})_t + \rho_{n+1} u_n \cdot \nabla \bar{u}_{n+1} + L \bar{u}_{n+1} + \nabla (p_{n+1} - p_n) \\ & = \bar{\rho}_{n+1} (f - (u_n)_t - u_{n-1} \cdot \nabla u_{n-1}) \\ & \quad + \rho_{n+1} (u_n \cdot \nabla \bar{u}_{n+1} - \bar{u}_n \cdot \nabla u_n - u_{n-1} \cdot \nabla \bar{u}_n). \end{aligned}$$

Multiplying this by  $\bar{u}_{n+1}$ , integrating over  $\Omega$  and using the equation (4.1) with  $(\rho, v) = (\rho_{n+1}, u_n)$ , we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int \rho_{n+1} |\bar{u}_{n+1}|^2 dx + \mu \int |\nabla \bar{u}_{n+1}|^2 dx \\ & \leq C \int |\bar{\rho}_{n+1}| |(u_n)_t| |\bar{u}_{n+1}| dx + C \int |p_{n+1} - p_n| |\nabla \bar{u}_{n+1}| dx \\ (5.10) \quad & + C \int |\bar{\rho}_{n+1}| |f - u_{n-1} \cdot \nabla u_{n-1}| |\bar{u}_{n+1}| dx \\ & + C \int \rho_{n+1} (|u_n| |\nabla \bar{u}_{n+1}| + |\bar{u}_n| |\nabla u_n| + |u_{n-1}| |\nabla \bar{u}_n|) |\bar{u}_{n+1}| dx. \end{aligned}$$

Using the uniform bound (5.2), we can estimate the last three integrals of the right hand side in (5.10) as follows:

$$C \int |p_{n+1} - p_n| |\nabla \bar{u}_{n+1}| dx \leq \tilde{C} |\bar{\rho}_{n+1}|_{L^2}^2 + \frac{\mu}{12} |\nabla \bar{u}_{n+1}|_{L^2}^2,$$

$$\begin{aligned} & C \int |\bar{\rho}_{n+1}| |f - u_{n-1} \cdot \nabla u_{n-1}| |\bar{u}_{n+1}| dx \\ & \leq C |\bar{\rho}_{n+1}|_{L^2} |f - u_{n-1} \cdot \nabla u_{n-1}|_{H^1} |\nabla \bar{u}_{n+1}|_{L^2} \leq \tilde{C} |\bar{\rho}_{n+1}|_{L^2}^2 + \frac{\mu}{12} |\nabla \bar{u}_{n+1}|_{L^2}^2, \end{aligned}$$

$$C \int \rho_{n+1} |u_n| |\nabla \bar{u}_{n+1}| |\bar{u}_{n+1}| dx \leq \tilde{C} |\sqrt{\rho_{n+1}} \bar{u}_{n+1}|_{L^2}^2 + \frac{\mu}{12} |\nabla \bar{u}_{n+1}|_{L^2}^2$$

and

$$\begin{aligned} & C \int \rho_{n+1} (|\bar{u}_n| |\nabla u_n| + |u_{n-1}| |\nabla \bar{u}_n|) |\bar{u}_{n+1}| dx \\ & \leq C |\rho_{n+1}|_{L^\infty}^{\frac{1}{2}} \left( |u_n|_{D_0^1 \cap D^2} + |u_{n-1}|_{D_0^1 \cap D^2} \right) |\sqrt{\rho_{n+1}} \bar{u}_{n+1}|_{L^2} |\nabla \bar{u}_n|_{L^2} \\ & \leq \eta^{-1} \tilde{C} |\sqrt{\rho_{n+1}} \bar{u}_{n+1}|_{L^2}^2 + \eta |\nabla \bar{u}_n|_{L^2}^2. \end{aligned}$$

The first integral is readily bounded by

$$C |\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}} |(u_n)_t|_{D_0^1} |\nabla \bar{u}_{n+1}|_{L^2} \leq \tilde{C} |\bar{\rho}_{n+1}|_{L^{\frac{3}{2}}}^2 + \frac{\mu}{12} |\nabla \bar{u}_{n+1}|_{L^2}^2.$$

Therefore, substituting all the estimates into (5.10), we deduce that

$$(5.11) \quad \begin{aligned} & \frac{d}{dt} |\sqrt{\rho_{n+1}} \bar{u}_{n+1}(t)|_{L^2}^2 + \mu |\nabla \bar{u}_{n+1}(t)|_{L^2}^2 \\ & \leq \eta^{-1} \tilde{C} \varphi_{n+1}(t) + 2\eta |\nabla \bar{u}_n(t)|_{L^2}^2 \end{aligned}$$

for  $0 \leq t \leq T_1$ , where

$$\varphi_{n+1}(t) = |\sqrt{\rho_{n+1}} \bar{u}_{n+1}(t)|_{L^2}^2 + |\bar{\rho}_{n+1}(t)|_{L^{\frac{3}{2}} \cap L^2}^2.$$

By virtue of (5.8), (5.9) and (5.11), we deduce that

$$(5.12) \quad \frac{d}{dt} \varphi_{n+1}(t) + \mu \psi_{n+1}(t) \leq \eta^{-1} \tilde{C} \varphi_{n+1}(t) + 4\eta \psi_n(t)$$

for  $0 \leq t \leq T_1$ , where  $\psi_{n+1}(t) = |\nabla \bar{u}_{n+1}(t)|_{L^2}^2$ . Note that  $\varphi_{n+1}(0) = 0$ . Hence integrating (5.12) over  $(0, t)$ , we have

$$\varphi_{n+1}(t) + \mu \int_0^t \psi_{n+1}(s) ds \leq 4\eta \int_0^t \psi_n(s) ds + \eta^{-1} \tilde{C} \int_0^t \varphi_{n+1}(s) ds,$$

which implies, in view of Gronwall's inequality, that

$$(5.13) \quad \varphi_{n+1}(t) + \int_0^t \psi_{n+1}(s) ds \leq \eta \tilde{C} \exp(\eta^{-1} \tilde{C} t) \left( \int_0^t \psi_n(s) ds \right).$$

Choosing  $\eta > 0$  and then  $T_2 > 0$  so small that

$$\eta \tilde{C} \leq \frac{1}{4}, \quad T_2 < T_1 \quad \text{and} \quad \exp(\eta^{-1} \tilde{C} T_2) < 2,$$

we deduce from (5.13) that

$$\sum_{n=1}^{\infty} \left( \sup_{0 \leq t \leq T_2} \varphi_{n+1}(t) + \int_0^{T_2} \psi_{n+1}(t) dt \right) \leq \tilde{C} \int_0^{T_2} \psi^1(t) dt < \infty.$$

Therefore, we conclude that  $\{(\rho_n, u_n)\}$  converges in a strong sense to a limit  $(\rho, u)$  satisfying the regularity estimate (5.2) by weak compactness of  $\{u_n^{(j)}\}$  with  $T_*$  replaced by  $T_2$ . Adapting the proof of Proposition 4.1, we can show that  $(\rho, u)$  is a solution to the original IBV problem (1.1)-(1.5) with  $T$  replaced by  $T_2$ . From the weak formulation and the uniqueness of solution of the system (5.6), we conclude that  $u^{(j)}(0) = w_j$  (and hence  $u^{(j)}(0)$  satisfies the compatibility condition (5.6)). This completes the proof of the existence. The proof of the uniqueness is similar



to (indeed easier than) the proof of the convergence and so omitted. We have completed the proof of Theorem 1.1.

## 6. PROOF OF THEOREM 1.4

In this section we consider the global existence of radially symmetric smooth solution  $(\rho(t, x), u(t, x)) = (\rho(t, |x|, \bar{u}(t, |x|)))$  with  $p(t, x) = A\rho(t, x)^\gamma$  for the initial boundary value problem (1.1)-(1.5) with

$$(6.1) \quad \Omega = \{x \in \mathbf{R}^3 : a < |x| < b\}, \quad f(t, x) = \bar{f}(t, |x|) \frac{x}{|x|},$$

$$(6.2) \quad (\rho_0(x), p_0(x)) = (\rho_0(|x|), p_0(|x|)), \quad u_0(x) = \bar{u}_0(|x|) \frac{x}{|x|}$$

for some constant  $a, b$  with  $0 < a < b \leq \infty$ , where  $\bar{u}, \bar{u}_0, \bar{f}$  are scalar functions.

We first introduce a result of H. Kim and H. J. Choe [7] which are crucial for the proof of global existence.

**Lemma 6.1** (Lemma 3.2 of [7]). *Whenever a strong solution  $(\rho, u)$  exists on a time interval  $[0, T_1]$ , we have*

$$\sup_{0 \leq t \leq T_1} |\rho(t)|_{L^\infty} \leq C_1$$

for some  $C_1$  depending only on  $a, T_1$  and the parameter of  $C$  not on  $b$ .

For the proof of Theorem 1.4, we will use a contradiction argument. Let  $T^*$  be the maximal existence time for the smooth solution  $(\rho, p, u)$  to exist with regularity in Theorem 1.1. Suppose that  $T^* < \infty$ . Then from Lemma 6.1 we deduce that

$$(6.3) \quad \sup_{0 \leq t < T^*} |\rho(t)|_{L^\infty} \leq C_1,$$

where the constant  $C_1$  now depends on  $T^*$ .

Let us define a function  $J_k(t)$  representing all regularity of smooth solution  $(\rho, u)$  as stated in Theorem 1.4 by

$$\begin{aligned} J_k(t) \equiv & \sum_{0 \leq j \leq k+1} |(\rho(t), p(t))|_{L^1} + |(\rho^{(j)}(t), p^{(j)}(t))|_{L^1 \cap H^{2(k-j)+3}} \\ & + \int_\tau^t |(\rho^{(k+2)}(s), p^{(k+2)}(s))|_{L^2}^2 ds \\ & + \sum_{0 \leq j \leq k+1} |u^{(j)}(t)|_{D_0^1 \cap D^{2(k-j)+3}} \\ & + \int_\tau^t |\sqrt{\rho} u^{(k+2)}(s)|_{L^2}^2 ds \end{aligned}$$

for some fixed  $\tau \in (0, T^*)$ , where  $p = A\rho^\gamma$  and hence  $p$  satisfies  $p_t + \nabla p \cdot u + \gamma p \operatorname{div} u = 0$ . Then from the hypothesis that  $T^* < \infty$ , we deduce that

$$\lim_{t \rightarrow T^*} J_k(t) = \infty.$$

But we will prove that this does not occur in the radially symmetric case. For the proof of Theorem 1.4 we have only to prove the following.

**Theorem 6.2.** Fix  $\tau \in (0, T^*)$  and define a function  $\Phi(t)$  by

$$\Phi(t) = 1 + \sup_{\tau < s < t} |\rho(s)|_{L^\infty} \quad \text{for all } t \in (\tau, T^*).$$

Then there exists an increasing continuous function  $M(\cdot)$  such that

$$J_k(t) \leq M_k(\Phi(t)) \quad \text{for all } t \in (\tau, T^*).$$

*Proof.* Since  $u$  is radially symmetric,  $\Delta u = \nabla \operatorname{div} u$ . Thus  $(\rho, p, u)$  satisfies the modified equation

$$(6.4) \quad \rho u_t + \rho u \cdot \nabla u + \nu \nabla \operatorname{div} u + \nabla p = \rho f,$$

where  $\nu = \lambda + 2\mu$ .

Multiplying (6.4) by  $u$  and integrating over  $\Omega$ , we have

$$(6.5) \quad \frac{1}{2} \frac{d}{dt} \int \rho |u|^2 dx + \nu \int (\operatorname{div} u)^2 dx - \int p \operatorname{div} u dx \leq \int \rho |f| |u| dx.$$

From (1.12), we also have

$$\frac{1}{\gamma - 1} \frac{d}{dt} |p|_{L^1} = - \int p \operatorname{div} u dx.$$

Substituting this into (6.5), we have

$$\frac{d}{dt} \left( \frac{1}{2} |\sqrt{\rho} u|_{L^2}^2 + \frac{1}{\gamma - 1} |p|_{L^1} \right) + \nu \int (\operatorname{div} u)^2 dx \leq |\rho|_{L^1}^{\frac{1}{2}} |\sqrt{\rho} u|_{L^2} |f|_{L^\infty}.$$

Integrating over  $(\tau, t)$ , by Gronwall's inequality we obtain for all  $\tau < t < T^*$

$$(6.6) \quad |\sqrt{\rho} u(t)|_{L^2}^2 + |p(t)|_{L^1} + \int_\tau^t |\nabla u(s)|_{L^2}^2 ds \leq \tilde{C}.$$

Throughout this section, we denote  $\tilde{C}$  by a generic constant depending only on  $\tau$ ,  $T^*$ ,  $c_0$  and the parameter of  $C$ .

Now we multiply (6.4) by  $u_t$  and integrate over  $\Omega$ . Then we have

$$(6.7) \quad \begin{aligned} & \frac{1}{2} \int \rho |u_t|^2 dx + \frac{d}{dt} \int \frac{\nu}{2} (\operatorname{div} u)^2 dx \\ & \leq \int \rho |f|^2 dx + \int \rho |u|^2 |\nabla u|^2 dx + \int p \operatorname{div} u_t dx \end{aligned}$$

and also from (1.12)

$$\begin{aligned} \int p \operatorname{div} u_t dx &= \frac{d}{dt} \int p \operatorname{div} u dx - \frac{d}{dt} \int \frac{4\gamma - 3}{2\nu(2\gamma - 1)} p^2 dx \\ & \quad + \frac{\gamma - 1}{\nu^2} \int p(G^2 - p^2) dx - \frac{1}{\nu} \int p u \cdot \nabla G dx, \end{aligned}$$

where  $G$  is the effective viscous flux  $\nu \operatorname{div} u - p$ . Since

$$p \operatorname{div} u - \frac{4\gamma - 3}{2\nu(2\gamma - 1)} p^2 \leq \frac{\nu}{2} (\operatorname{div} u)^2 - \nu \frac{2\gamma - 1}{2(4\gamma - 3)} (\operatorname{div} u)^2,$$

integrating (6.7) over  $(\tau, t)$ , we have

$$\begin{aligned}
& \frac{1}{2} \int_{\tau}^t |\sqrt{\rho}u_t|_{L^2}^2 ds + \frac{\nu}{2} |\nabla u(t)|_{L^2}^2 \\
& \leq \frac{\nu}{2} |\nabla u(\tau)|_{L^2}^2 + \int_{\tau}^t |\rho|_{L^1} |f|_{L^\infty}^2 ds + \frac{\nu}{2} \left(1 - \frac{2\gamma-1}{4\gamma-3}\right) |\nabla u(t)|_{L^2}^2 \\
& \quad - \frac{\nu}{2} \left(1 - \frac{2\gamma-1}{4\gamma-3}\right) |\nabla u(\tau)|_{L^2}^2 + \int_{\tau}^t \int \rho |u|^2 |\nabla u|^2 dx ds \\
& \quad + \frac{\gamma-1}{\nu^2} \int_{\tau}^t \int p(G^2 - p^2) dx ds - \frac{1}{\nu} \int_{\tau}^t \int pu \cdot \nabla G dx ds.
\end{aligned}$$

Thus

$$\begin{aligned}
(6.8) \quad & \int_{\tau}^t |\sqrt{\rho}u_t|_{L^2}^2 ds + |\nabla u(t)|_{L^2}^2 \\
& \leq \tilde{C} + C \int_{\tau}^t \int pG^2 dx ds + C \int_{\tau}^t \int p|u| |\nabla G| dx ds \\
& \quad + C \int_{\tau}^t \int \rho |u|^2 |\nabla u|^2 dx ds
\end{aligned}$$

Now if we use the identity  $\nabla G = \rho u_t + \rho u \cdot \nabla u - \rho f$ ,

$$\begin{aligned}
\int_{\tau}^t \int p|u| |\nabla G| dx ds & \leq \int_{\tau}^t \int p|u| (\rho |u_t| + \rho |u| |\nabla u| + \rho |f|) dx ds \\
& \leq \int_{\tau}^t |p(s)|_{L^\infty} |\sqrt{\rho}u(s)|_{L^2} |\sqrt{\rho}u_t(s)|_{L^2} ds \\
& \quad + \int_{\tau}^t |p(s)|_{L^\infty} |u|_{L^\infty} |\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho}u|_{L^2} |\nabla u|_{L^2} ds \\
& \quad + \int_{\tau}^t |\rho|_{L^\infty}^{\frac{1}{2}} |p|_{L^\infty} |\sqrt{\rho}u|_{L^2} |f|_{L^2} ds \\
& \leq \tilde{C} \int_{\tau}^t |p(s)|_{L^\infty}^2 ds + \frac{1}{2C} \int_{\tau}^t |\sqrt{\rho}u_t|_{L^2}^2 ds \\
& \quad + \tilde{C} \int_{\tau}^t |p(s)|_{L^\infty} \Phi(s)^{\frac{1}{2}} |\nabla u|_{L^2}^2 ds \\
& \quad + \tilde{C} \int_{\tau}^t \Phi(s)^{\frac{1}{2}} |p(s)|_{L^\infty} ds.
\end{aligned}$$

Here we used the embedding  $|u|_{L^\infty} \leq C|\nabla u|_{L^2}$  of radial function  $u$ . Substituting this into (6.8), we have

$$\begin{aligned}
& \int_{\tau}^t |\sqrt{\rho}u_t|_{L^2}^2 ds + |\nabla u(t)|_{L^2}^2 \\
& \leq \tilde{C} + C \int_{\tau}^t \int p(|\nabla u|^2 + p^2) dx ds + \tilde{C} \int_{\tau}^t |p(s)|_{L^\infty}^2 ds \\
& \quad + \tilde{C} \int_{\tau}^t |p(s)|_{L^\infty} \Phi(s)^{\frac{1}{2}} |\nabla u|_{L^2}^2 ds + \tilde{C} \int_{\tau}^t \Phi(s)^{\frac{1}{2}} |p(s)|_{L^\infty} ds \\
& \quad + \int_{\tau}^t \int \rho |u|^2 |\nabla u|^2 dx ds \\
& \leq \tilde{C} + \tilde{C} \int_{\tau}^t (|p(s)|_{L^3}^3 + |p(s)|_{L^\infty}^2 + \Phi(s)) ds \\
& \quad + \tilde{C} \int_{\tau}^t (|p(s)|_{L^\infty} \Phi(s)^{\frac{1}{2}} + |p(s)|_{L^\infty}) |\nabla u(s)|_{L^2}^2 ds \\
& \quad + C \int_{\tau}^t |\rho(s)|_{L^\infty} |u(s)|_{L^\infty}^2 |\nabla u|_{L^2}^2 ds \\
& \leq \tilde{C} + \tilde{C} \int_{\tau}^t (|p(s)|_{L^3}^3 + |p(s)|_{L^\infty}^2 + \Phi(s)) ds \\
& \quad + \tilde{C} \int_{\tau}^t (|p(s)|_{L^\infty} \Phi(s)^{\frac{1}{2}} + |p(s)|_{L^\infty} + \Phi(s) |\nabla u(s)|_{L^2}^2) |\nabla u(s)|_{L^2}^2 ds.
\end{aligned}$$

Using Gronwall's inequality, (6.6) and (6.8), we have

$$\begin{aligned}
& \int_{\tau}^t |\sqrt{\rho}u_t|_{L^2}^2 ds + |\nabla u(t)|_{L^2}^2 \\
& \leq \tilde{C} \left( 1 + \int_{\tau}^t (|p(s)|_{L^3}^3 + |p(s)|_{L^\infty}^2 + \Phi(s)) ds \right) \\
& \quad \times \exp \left( \tilde{C} \int_{\tau}^t (|p(s)|_{L^\infty}^2 + |p(s)|_{L^\infty} + \Phi(s) + \Phi(s) |\nabla u(s)|_{L^2}^2) ds \right) \\
& \leq \tilde{C} \left( 1 + \tilde{C} \int_{\tau}^t (|p(s)|_{L^\infty}^2 + \Phi(s)) ds \right) \\
& \quad \times \exp \left( \tilde{C} \int_{\tau}^t (|p(s)|_{L^\infty}^2 + |p(s)|_{L^\infty} + \Phi(s)) ds \right).
\end{aligned}$$

By the estimate  $|p(s)|_{L^\infty} \leq A\Phi(s)^\gamma$ , we conclude that

$$\begin{aligned}
& \int_{\tau}^t |\sqrt{\rho}u_t|_{L^2}^2 ds + |\nabla u(t)|_{L^2}^2 \\
(6.9) \quad & \leq \tilde{C} \left( 1 + \int_{\tau}^t \Phi(s)^{2\gamma} ds \right) \exp \left( \tilde{C} \int_{\tau}^t \Phi(s)^{2\gamma} ds \right) \\
& \leq \tilde{C} \exp \left( \tilde{C} \Phi(t)^{2\gamma} \right).
\end{aligned}$$

From the definition of effective viscous flux  $G = \nu \operatorname{div} u - p$ , we have

$$\begin{aligned}
\int_{\tau}^t |G|_{L^\infty}^2 ds &\leq C \int_{\tau}^t |G|_{H^1}^2 ds \leq C \int_{\tau}^t (|G|_{L^2}^2 + |\nabla G|_{L^2}^2) ds \\
&\leq C \int_{\tau}^t (|\nabla u|_{L^2}^2 + |p|_{L^2}^2) ds \\
&\quad + C \int_{\tau}^t (|\rho u_t|_{L^2}^2 + |\rho u \cdot \nabla u|_{L^2}^2 + |\rho f|_{L^2}^2) ds \\
&\leq C \int_{\tau}^t (|\nabla u|_{L^2}^2 + \Phi^\gamma |p|_{L^1}) ds \\
&\quad + C \int_{\tau}^t (\Phi + |\sqrt{\rho} u_t|_{L^2}^2 + \Phi^2 |\nabla u|_{L^2}^4 + \Phi^2 |f|_{L^2}^2) ds \\
&\leq \tilde{C} \Phi(t)^{6\gamma} \exp(\tilde{C} \Phi(t)^{2\gamma})
\end{aligned}$$

and also

$$\begin{aligned}
\int_{\tau}^t |\nabla u|_{L^\infty}^2 ds &\leq C \int_{\tau}^t (|G|_{L^\infty}^2 + |p|_{L^\infty}^2 + |u|_{L^\infty}^2) ds \\
&\leq \tilde{C} \Phi(t)^{6\gamma} \exp(\tilde{C} \Phi(t)^{2\gamma}).
\end{aligned}$$

For future reference, we need to show a higher regularity boundedness of  $\rho$  and  $p$ . Here we claim that for  $1 \leq m \leq 2k + 3$

$$\begin{aligned}
&|\rho(t)|_{H^m} + |p(t)|_{H^m} \\
(6.10) \quad &\leq \tilde{C} \left( 1 + \Phi(t)^2 \int_{\tau}^t |G(s)|_{H^m}^2 ds \right) \\
&\quad \times \exp \left( C \int_{\tau}^t (1 + \Phi^2 + |\nabla u|_{L^\infty} + |\nabla u|_{H^{m-1}}) ds \right)
\end{aligned}$$

From (3.5), we have

$$\begin{aligned}
\frac{d}{dt} |D^\alpha \rho|_{L^2}^2 &\leq C |\nabla u|_{L^\infty} |D^\alpha \rho|_{L^2}^2 + C \sum_{1 \leq l \leq |\alpha| - 1} \int |\nabla^{|\alpha| + 1 - l} u| |\nabla^l \rho| |D^\alpha \rho| dx \\
&\quad + C \sum_{0 \leq l \leq |\alpha| - 1} \int |\nabla^{|\alpha| - l} (\operatorname{div})| |\nabla^l \rho| |D^\alpha \rho| dx \\
&\leq C |\nabla u|_{L^\infty} |D^\alpha \rho|_{L^2}^2 + C |\rho|_{L^\infty} |\nabla^{|\alpha|} (G + p)|_{L^2} |D^\alpha \rho|_{L^2} \\
&\quad + C \sum_{1 \leq l \leq |\alpha| - 1} |\nabla^{|\alpha| + 1 - l} u|_{L^2} |\nabla^l \rho|_{L^2} |D^\alpha \rho|_{L^2}
\end{aligned}$$

and get

$$\frac{d}{dt} |\rho|_{H^m}^2 \leq C(1 + |\nabla u|_{L^\infty} + |\nabla u|_{H^{m-1}}) |\rho|_{H^m}^2 + C\Phi^2 (|G|_{H^m}^2 + |p|_{H^m}^2).$$

Similarly, from the equation (1.12), it follows that

$$\frac{d}{dt} |p|_{H^m}^2 \leq C(1 + \Phi^2 + |\nabla u|_{L^\infty} + |\nabla u|_{H^{m-1}}) |p|_{H^m}^2 + C\Phi^2 |G|_{H^m}^2.$$

Summing the estimates above for  $\rho$  and  $p$ , and integrating them over  $(\tau, t)$ , we readily obtain

$$(6.11) \quad \begin{aligned} & |\rho(t)|_{H^m}^2 + |p(t)|_{H^m}^2 \\ & \leq \tilde{C} + C \int_{\tau}^t (1 + \Phi^2 + |\nabla u|_{L^\infty} + |\nabla u|_{H^{m-1}})(|\rho(s)|_{H^m}^2 + |p(s)|_{H^m}^2) ds \\ & \quad + C\Phi^2(t) \int_{\tau}^t |G|_{H^m}^2 ds \end{aligned}$$

and hence by Gronwall's inequality we obtain (6.10).

In particular, if  $m = 1$ , then

$$\begin{aligned} & |\rho(t)|_{H^1}^2 + |p(t)|_{H^1}^2 \\ & \leq \tilde{C} \left( 1 + \Phi^2 \int_{\tau}^t |G|_{H^1}^2 ds \right) \\ & \quad \times \exp \left( C \int_{\tau}^t (1 + \Phi^2 + |\nabla u|_{L^\infty} + |\nabla u|_{L^2}) ds \right) \\ & \leq \tilde{C}\Phi(t)^{6\gamma} \exp \left( \tilde{C}\Phi(t)^{2\gamma} \right). \end{aligned}$$

And hence

$$\begin{aligned} \int_{\tau}^t |\nabla^2 u|_{L^2}^2 ds & \leq C \int_{\tau}^t (|\Delta u|_{L^2}^2 + |\nabla u|_{L^2}^2) ds \leq C \int_{\tau}^t (|\nabla \operatorname{div} u|_{L^2}^2 + |\nabla u|_{L^2}^2) ds \\ & \leq C \int_{\tau}^t (|\nabla G|_{L^2}^2 + |\nabla p|_{L^2}^2 + |\nabla u|_{L^2}^2) ds \leq \tilde{C}\Phi(t)^{6\gamma} \exp \left( \tilde{C}\Phi(t)^{2\gamma} \right). \end{aligned}$$

Now we estimate  $|u_t|_{D_0^1}$ . We differentiate (6.4) with respect to time, multiply  $u_t$  and then integrate over  $(\tau, t)$ . Using the identity  $\rho_t = -\operatorname{div}(\rho u)$  and

$$\begin{aligned} - \int p_t \operatorname{div} u_t dx & = \int (\nabla p \cdot u + \gamma p \operatorname{div} u_t) \\ & = \frac{d}{dt} \int \frac{\gamma}{2} p (\operatorname{div} u)^2 dx + \int \nabla p \cdot (u \operatorname{div} u_t) dx \\ & \quad + \frac{\gamma}{2} \int (-pu \cdot \nabla (\operatorname{div} u)^2 + (\gamma - 1)p (\operatorname{div} u)^3), \end{aligned}$$

integration by parts yields

$$\begin{aligned} & \frac{d}{dt} \int \left( \frac{1}{2} \rho |u_t|^2 + \frac{\gamma}{2} p (\operatorname{div} u)^2 \right) dx + \nu \int |\nabla u_t|^2 dx \\ & \leq \int (2\rho |u| |u_t| |\nabla u_t| + \rho |u| |u_t| |\nabla u|^2 + \rho |u|^2 |u_t| |\nabla^2 u| \\ & \quad + \rho |u|^2 |\nabla u| |\nabla u_t| + \rho |u_t|^2 |\nabla u| + |\nabla p| |u| |\nabla u_t| \\ & \quad + \gamma p |u| |\nabla u| |\nabla^2 u| + \gamma^2 p |\nabla u|^3 + \rho |u| |u_t| |\nabla f| \\ & \quad + \rho |u| |f| |\nabla u_t| + \rho |u_t| |f_t|) dx \\ & \equiv \sum_{1 \leq j \leq 11} I_j. \end{aligned}$$

We estimate each  $I_j$  as follows:

$$I_1 \leq C|\rho|_{L^\infty}|u|_{L^\infty}^2|\sqrt{\rho}u_t|_{L^2}^2 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2 \leq \tilde{C}\Phi^{2\gamma+1}|\sqrt{\rho}u_t|_{L^2}^2 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2,$$

$$\begin{aligned} I_2 &\leq |\rho|_{L^\infty}|u|_{L^\infty}|u_t|_{L^\infty}|\nabla u|_{L^2}^2 \leq C\Phi^2|\nabla u|_{L^2}^6 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2 \\ &\leq \tilde{C}\exp\left(\tilde{C}\Phi^{2\gamma}\right) + \frac{\nu}{20}|\nabla u_t|_{L^2}^2, \end{aligned}$$

$$I_3 \leq C|\rho|_{L^\infty}|u|_{L^\infty}^4|\nabla^2 u|_{L^2}^2 + |\sqrt{\rho}u_t|_{L^2}^2 \leq \tilde{C}\exp\left(\tilde{C}\Phi^{2\gamma}\right)|\nabla^2 u|_{L^2}^2 + |\sqrt{\rho}u_t|_{L^2}^2,$$

$$I_4 \leq C|\rho|_{L^\infty}^2|u|_{L^\infty}^4|\nabla u|_{L^2}^2 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2 \leq \tilde{C}\exp\left(\tilde{C}\Phi^{2\gamma}\right) + \frac{\nu}{20}|\nabla u_t|_{L^2}^2,$$

$$I_5 \leq |\nabla u|_{L^\infty}|\sqrt{\rho}u_t|_{L^2}^2,$$

$$\begin{aligned} I_6 &\leq C|\rho|_{L^\infty}^{\gamma-1}|u|_{L^\infty}|\nabla\rho|_{L^2}|\nabla u_t|_{L^2} \leq C\Phi^{2(\gamma-1)}|\nabla u|_{L^2}^2|\nabla\rho|_{L^2}^2 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2 \\ &\leq \tilde{C}\Phi^{8\gamma-2}\exp\left(\tilde{C}\Phi^{2\gamma}\right) + \frac{\nu}{20}|\nabla u_t|_{L^2}^2, \end{aligned}$$

$$I_7 \leq C|\rho|_{L^\infty}^\gamma|u|_{L^\infty}|\nabla u|_{L^2}|\nabla^2 u|_{L^2} \leq \tilde{C}\exp\left(\tilde{C}\Phi^{2\gamma}\right)|\nabla^2 u|_{L^2}^2,$$

$$I_8 \leq C|\rho|_{L^\infty}^\gamma|\nabla u|_{L^\infty}|\nabla u|_{L^2}^2 \leq \tilde{C}\exp\left(\tilde{C}\Phi^{2\gamma}\right)|\nabla u|_{L^\infty},$$

$$I_9 \leq C|\rho|_{L^\infty}^{\frac{1}{2}}|\sqrt{\rho}u|_{L^2}|\nabla u_t|_{L^2}|\nabla f|_{H^1} \leq \tilde{C}\Phi|f|_{H^2}^2 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2,$$

$$I_{10} \leq |\rho|_{L^\infty}^{\frac{1}{2}}|\sqrt{\rho}u_t|_{L^2}|f|_{H^1}|\nabla u_t|_{L^2} \leq \tilde{C}\Phi|f|_{H^1}^2 + \frac{\nu}{20}|\nabla u_t|_{L^2}^2,$$

$$I_{11} \leq |\rho|_{L^\infty}|f_t|_{L^2}^2 + |\sqrt{\rho}u_t|_{L^2}^2.$$

Then we have

$$\begin{aligned} &|\sqrt{\rho}u_t(t)|_{L^2}^2 + \int p(\operatorname{div}u)^2(t) dx + \int_\tau^t |\nabla u_t|_{L^2}^2 ds \\ &\leq \tilde{C} + \tilde{C}\Phi(t)^{8\gamma-2}\exp\left(\tilde{C}\Phi(t)^{2\gamma}\right) + \tilde{C}\Phi(t)^{2\gamma+1} \int_\tau^t |\sqrt{\rho}u_t(s)|_{L^2}^2 ds. \end{aligned}$$

Thus from the Gronwall's inequality we deduce that for all  $\tau < t < T^*$

$$(6.12) \quad |\sqrt{\rho}u_t(t)|_{L^2}^2 + \int p(\operatorname{div}u)^2(t) dx + \int_\tau^t |\nabla u_t|_{L^2}^2 ds \leq \tilde{C}\exp\left(\tilde{C}\Phi^{2\gamma}\right)$$

and that

$$\begin{aligned}
(6.13) \quad |\nabla u|_{H^1} &\leq C(|\nabla \operatorname{div} u|_{L^2} + |\nabla u|_{L^2}) \\
&\leq C(|\nabla G|_{L^2} + |\nabla p|_{L^2} + |\nabla u|_{L^2}) \\
&\leq C(|\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho} u_t|_{L^2} + |\rho|_{L^\infty} |\nabla u|_{L^2}^2 \\
&\quad + |\rho|_{L^\infty} |f|_{L^2} + |\nabla p|_{L^2} + |\nabla u|_{L^2}) \\
&\leq \tilde{C} \exp\left(\tilde{C} \Phi^{2\gamma}\right).
\end{aligned}$$

From the identity  $\nabla G = \rho u_t + \rho u \cdot \nabla u - \rho f$ , we have

$$\begin{aligned}
|\nabla^2 G|_{L^2} &\leq C(|\nabla \rho|_{L^\infty} |\nabla u_t|_{L^2} + |\rho|_{L^\infty} |\nabla u_t|_{L^2} + |\nabla \rho|_{H^1} |\nabla u|_{L^2} |\nabla u|_{H^1} \\
&\quad + |\rho|_{L^\infty} |\nabla u|_{L^\infty} |\nabla u|_{L^2} + |\rho|_{L^3} |\nabla u|_{L^2} |\nabla^2 u|_{L^2} \\
&\quad + |\nabla \rho|_{L^2} |f|_{H^1} + |\rho|_{L^\infty} |\nabla f|_{L^2}) \\
&\leq C(|\rho|_{H^2} |\nabla u_t|_{L^2} + \Phi |\nabla u_t|_{L^2} + |\rho|_{H^2} |\nabla u|_{L^2} |\nabla u|_{H^1} \\
&\quad + \Phi |\nabla u|_{L^\infty} |\nabla u|_{L^2} + \Phi |\nabla u|_{L^2} |\nabla u|_{H^1} \\
&\quad + |\rho|_{H^1} |\nabla u|_{L^2} |\nabla u|_{H^1} + |\nabla \rho|_{L^2} |f|_{H^1} + \Phi |\nabla f|_{L^2})
\end{aligned}$$

and get

$$\begin{aligned}
&\int_\tau^t |G(s)|_{H^2}^2 ds \\
&\leq \int_\tau^t |G|_{H^1}^2 ds + C\Phi^2(t) \int_\tau^t (|\nabla u_t|_{L^2}^2 + |\nabla u|_{L^2}^2 + |\nabla u|_{L^\infty}^2 + |\nabla f|_{L^2}^2) ds \\
&\quad + C \int_\tau^t (|\nabla u_t|_{L^2}^2 + |\nabla u|_{L^2}^2 |\nabla u|_{H^1}^2 + |f|_{H^1}^2) |\rho|_{H^2}^2 ds
\end{aligned}$$

We then have from (6.11) with  $m = 2$  that

$$\begin{aligned}
&|\rho(t)|_{H^2}^2 + |p(t)|_{H^2}^2 \\
&\leq \tilde{C} + C\Phi^4 \int_\tau^t (|\nabla u_t|_{L^2}^2 + |\nabla u|_{L^2}^2 |\nabla u|_{L^\infty}^2 + |\nabla f|_{L^2}^2) ds \\
&\quad + C\Phi^2 \int_\tau^t |G|_{H^1}^2 ds + C\Phi^4(t) \int_\tau^t (1 + |\nabla u|_{L^\infty} + |\nabla u|_{H^1} + |\nabla u_t|_{L^2}^2 \\
&\quad + |\nabla u|_{L^2} |\nabla u|_{H^1}^2 + |f|_{H^1}^2) (|\rho|_{H^2}^2 + |p|_{H^2}^2) ds.
\end{aligned}$$

Thus by Gronwall's inequality, we deduce that

$$(6.14) \quad \int_\tau^t |G(s)|_{H^2}^2 ds + |\rho(t)|_{H^2}^2 + |p(t)|_{H^2}^2 \leq \tilde{C} \exp\left(\exp\left(\tilde{C} \Phi^{2\gamma}\right)\right).$$

Now we claim that for  $\tau < t < T^*$

$$(6.15) \quad \int_\tau^t |\sqrt{\rho} u_{tt}|_{L^2}^2 ds + |\nabla u_t(t)|_{L^2}^2 \leq \tilde{C} \exp\left(\tilde{C} \exp\left(\tilde{C} \exp\left(\tilde{C} \Phi^{2\gamma}\right)\right)\right)$$



and

$$(6.16) \quad \begin{aligned} & |\nabla u(t)|_{H^2}^2 + \int_{\tau}^t (|u_t(s)|_{D^2}^2 + |u(s)|_{D^4}^2) ds \\ & \leq \tilde{C} \exp \left( \tilde{C} \exp \left( \tilde{C} \exp \left( \tilde{C} \Phi^{2\gamma} \right) \right) \right). \end{aligned}$$

To show the estimates (6.15) and (6.16), we first differentiate (6.4) with respect to  $t$ , multiply  $u_{tt}$  and integrate over  $\Omega$ . Then we obtain

$$(6.17) \quad \begin{aligned} & \int \rho |u_{tt}|^2 dx + \frac{\nu}{2} \frac{d}{dt} \int (\operatorname{div} u_t)^2 dx \\ & = \int p_t \operatorname{div} u_{tt} dx + \int ((\rho f)_t - \rho_t (u_t + u \cdot \nabla u) - \rho u_t \cdot \nabla u_t) \cdot u_{tt} dx. \end{aligned}$$

Since  $p$  satisfies the pressure equation (1.12), we have

$$\begin{aligned} \int p_t \operatorname{div} u_{tt} dx &= \frac{d}{dt} \int p_t \operatorname{div} u_t dx - \int p_{tt} \operatorname{div} u_t dx \\ &\leq \frac{d}{dt} \int p_t \operatorname{div} u_t dx + \int \nabla p_t u \cdot \operatorname{div} u_t dx + \int \nabla p \cdot u_t \operatorname{div} u_t dx \\ &\quad + \gamma \int p_t \operatorname{div} u \operatorname{div} u_t dx + \gamma \int p (\operatorname{div} u_t)^2 dx \\ &\leq \frac{d}{dt} \int p_t \operatorname{div} u_t dx + C |\nabla p_t|_{L^2} |\nabla u|_{L^2} |\nabla u_t|_{L^2} \\ &\quad + C |\nabla p|_{L^2} |\nabla u_t|_{L^2}^2 + C |p_t|_{H^1} |\nabla u|_{L^2} |\nabla u_t|_{L^2} + \Phi^\gamma |\nabla u_t|_{L^2}^2. \end{aligned}$$

Observing from the pressure equation (1.12) that

$$\begin{aligned} |p_t|_{L^2} &\leq C |u|_{L^\infty} |\nabla p|_{L^2} + C |p|_{L^\infty} |\nabla u|_{L^\infty} \leq C |\nabla u|_{L^2} |\nabla p|_{L^2} + C \Phi^\gamma |\nabla u|_{L^2} \\ &\leq \tilde{C} \exp \left( \tilde{C} \Phi^{2\gamma} \right) \end{aligned}$$

and from the identity

$$\nabla p_t = -\nabla^2 p u - \nabla p \nabla u - \gamma \nabla p \operatorname{div} u - p \nabla \operatorname{div} u,$$

that

$$\begin{aligned} |\nabla p_t|_{L^2} &\leq C |\nabla u|_{L^2} |\nabla^2 p|_{L^2} + C |\nabla p|_{H^1} |\nabla u|_{H^1} + C |\nabla p|_{L^2} |G|_{L^\infty} \\ &\quad + |\nabla p|_{L^2} |p|_{L^\infty} + |p|_{L^\infty} (|\nabla G|_{L^2} + |\nabla p|_{L^2}) \\ &\leq \tilde{C} \exp \left( \tilde{C} \Phi^{2\gamma} \right), \end{aligned}$$

we deduce that

$$(6.18) \quad \int p_t \operatorname{div} u_{tt} dx \leq \frac{d}{dt} \int p_t \operatorname{div} u_t dx + \tilde{C} \exp \left( \tilde{C} \Phi^{2\gamma} \right) (1 + |\nabla u_t|_{L^2}^2).$$

We estimate the second integration in (6.17) as follows:

$$\begin{aligned}
& \int \rho(f - u \cdot \nabla u)_t \cdot u_{tt} \, dx \\
(6.19) \quad & \leq C |\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho} u_{tt}|_{L^2} (|f_t|_{L^2} + |\nabla u_t|_{L^2} |\nabla u|_{H^1} + |\nabla u|_{L^2} |\nabla u_t|_{L^2}) \\
& \leq C |\rho|_{L^\infty} (|f_t|_{L^2}^2 + |\nabla u|_{H^1}^2 |\nabla u_t|_{L^2}^2 + |\nabla u|_{L^2}^2 |\nabla u_t|_{L^2}^2) + \frac{1}{2} |\sqrt{\rho} u_{tt}|_{L^2}^2 \\
& \leq C \exp\left(\tilde{C} \exp\left(\tilde{C} \Phi^{2\gamma}\right)\right) (|f_t|_{L^2}^2 + |\nabla u_t|_{L^2}^2) + \frac{1}{2} |\sqrt{\rho} u_{tt}|_{L^2}^2.
\end{aligned}$$

$$\begin{aligned}
& \int \rho_t(f - u \cdot \nabla u) \cdot u_{tt} \, dx \\
(6.20) \quad & = \frac{d}{dt} \int \rho_t(f - u \cdot \nabla u) \cdot u_{tt} \, dx - \int \rho_{tt}(f - u \cdot \nabla u) \cdot u_t \, dx \\
& \quad - \int \rho_t(f_t - u_t \cdot \nabla u - u \cdot \nabla u_t) \cdot u_t \, dx \\
& \leq \frac{d}{dt} \int \rho_t(f - u \cdot \nabla u) \cdot u_t \, dx + |\rho_{tt}|_{L^2} (|f|_{L^2} + |\nabla u|_{L^2}^2) |\nabla u_t|_{L^2} \\
& \quad + |\rho_t|_{L^2} (|f_t|_{L^2} + |\nabla u|_{L^2} |\nabla u_t|_{L^2} + |\nabla u|_{L^2} |\nabla u_t|_{L^2}) |\nabla u_t|_{L^2} \\
& \leq \frac{d}{dt} \int \rho_t(f - u \cdot \nabla u) \cdot u_t \, dx + C |\rho_{tt}|_{L^2} (1 + |\nabla u|_{L^2}^4) \\
& \quad + C (1 + |\rho_{tt}|_{L^2}^2 |f_t|_{L^2}^2 + |\rho_t|_{L^2} |\nabla u|_{L^2}) |\nabla u_t|_{L^2}^2 \\
& \leq \frac{d}{dt} \int \rho_t(f - u \cdot \nabla u) \cdot u_t \, dx \\
& \quad + \tilde{C} \exp\left(\tilde{C} \Phi^{2\gamma}\right) (|\rho_{tt}|_{L^2}^2 + |f_t|_{L^2}^2 + |\nabla u_t|_{L^2}^2).
\end{aligned}$$

$$\begin{aligned}
& - \int \rho_t u_t \cdot u_{tt} \, dx \\
(6.21) \quad & = - \frac{d}{dt} \int \rho_t \left(\frac{1}{2} |u_t|^2\right) \, dx + \int \rho_{tt} \left(\frac{1}{2} |u_t|^2\right) \, dx \\
& = - \frac{d}{dt} \int \rho_t \left(\frac{1}{2} |u_t|^2\right) \, dx - \int \operatorname{div}(\rho_t u + \rho u_t) \left(\frac{1}{2} |u_t|^2\right) \, dx \\
& = - \frac{d}{dt} \int \rho_t \left(\frac{1}{2} |u_t|^2\right) \, dx + \int (\rho_t u + \rho u_t) u_t \cdot \nabla u_t \, dx \\
& \leq - \frac{d}{dt} \int \rho_t \left(\frac{1}{2} |u_t|^2\right) \, dx + C (|\rho_t|_{L^2} |\nabla u|_{L^2} + |\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho} u_t|_{L^2}) |\nabla u_t|_{L^2}^2 \\
& \leq - \frac{d}{dt} \int \rho_t \left(\frac{1}{2} |u_t|^2\right) \, dx + \tilde{C} \exp\left(\tilde{C} \Phi^{2\gamma}\right) (1 + |\sqrt{\rho} u_t|_{L^2}) |\nabla u_t|_{L^2}^2.
\end{aligned}$$

Substituting the estimates (6.18)–(6.21) into (6.17), we have

$$\begin{aligned}
& \frac{1}{2} \int \rho |u_{tt}|_{L^2}^2 \, dx + \frac{d}{dt} \int \left( \frac{\nu}{2} (\operatorname{div} u_t)^2 - \rho_t(f - u \cdot \nabla u) \cdot u_t + \rho_t \left(\frac{1}{2} |u_t|^2\right) \right) \, dx \\
& \leq \tilde{C} \exp\left(\tilde{C} \exp\left(\tilde{C} \Phi^{2\gamma}\right)\right) (1 + |f_t|_{L^2}^2 + |\rho_{tt}|_{L^2}^2 + |\sqrt{\rho} u_t|_{L^2}^2) |\nabla u_t|_{L^2}^2.
\end{aligned}$$

Integrating over  $(\tau, t)$ , we obtain that

$$\begin{aligned} & \int_{\tau}^t |\sqrt{\rho}u_{tt}|_{L^2}^2 ds + \Lambda_1(t) \\ & \leq \tilde{C} + \tilde{C} \exp\left(\tilde{C} \exp\left(\tilde{C}\Phi^{2\gamma}\right)\right) \\ & \quad \times \int_{\tau}^t (1 + |f_t|_{L^2}^2 + |\rho_{tt}|_{L^2}^2 + |\sqrt{\rho}u_t|_{L^2}^2) |\nabla u_t|_{L^2}^2 ds, \end{aligned}$$

where

$$\Lambda_1(t) = \int \left( \frac{\nu}{2} (\operatorname{div} u_t)^2 - \rho_t (f - u \cdot \nabla u) \cdot u_t + \rho_t \left( \frac{1}{2} |u_t|^2 \right) \right) dx.$$

From the following two estimates

$$\begin{aligned} & \int \rho_t (f - u \cdot \nabla u) \cdot u_t dx \\ & \leq C |\rho_t|_{H^1} (|f|_{L^2} + |\nabla u|_{L^2}^2) |\nabla u_t|_{L^2} \\ & \leq C |\rho_t|_{H^1}^2 (|f|_{L^2}^2 + |\nabla u|_{L^2}^4) + \frac{\nu}{8} |\nabla u_t|_{L^2}^2 \end{aligned}$$

and

$$\begin{aligned} & \frac{1}{2} \int \rho_t |u_t|^2 dx \\ & = - \int \rho u \cdot u_t \cdot \nabla u_t dx \leq C |\rho|_{L^\infty}^{\frac{1}{2}} |\sqrt{\rho}u_t|_{L^2} |\nabla u|_{L^2} |\nabla u_t|_{L^2} \\ & \leq C |\rho_t|_{H^1}^2 (|f|_{L^2}^2 + |\nabla u|_{L^2}^4) + |\rho|_{L^\infty} |\sqrt{\rho}u_t|_{L^2}^2 |\nabla u|_{L^2}^2 + \frac{\nu}{8} |\nabla u_t|_{L^2}^2, \end{aligned}$$

we have

$$\begin{aligned} (6.22) \quad & \int_{\tau}^t |\sqrt{\rho}u_{tt}|_{L^2}^2 ds + C^{-1} |\nabla u_t|_{L^2}^2 \\ & \leq \tilde{C} \exp\left(\tilde{C} \exp\left(\tilde{C}\Phi^{2\gamma}\right)\right) \\ & \quad \times \left( 1 + \int_{\tau}^t (1 + |f_t|_{L^2}^2 + |\rho_{tt}|_{L^2}^2) |\nabla u_t|_{L^2}^2 ds \right). \end{aligned}$$

Now we observe from the mass equation (1.1) and Sobolev embedding of radial function that

$$\rho_{tt} = -\nabla \rho_t \cdot u - \nabla \rho \cdot u_t - \rho_t \operatorname{div} u - \rho \operatorname{div} u_t$$

and

$$|\rho_{tt}|_{L^2} \leq C (|\nabla \rho_t|_{L^2} |\nabla u|_{L^2} + |\nabla \rho|_{H^1} |\nabla u_t|_{L^2} + |\nabla u|_{H^1} |\rho_t|_{L^2} + |\rho|_{L^\infty} |\nabla u_t|_{L^2})$$

and

$$(6.23) \quad \int_{\tau}^t |\rho_{tt}|_{L^2}^2 ds \leq \tilde{C} \exp\left(\tilde{C} \exp\left(\tilde{C}\Phi^{2\gamma}\right)\right).$$

Thus substituting (6.23) into (6.22) and using Gronwall's inequality, we obtain the required (6.15). By using the elliptic regularity of the operator  $L$ , we get the estimate (6.16).

Let us define a function  $M_0$  by

$$M_0(\Phi) = \tilde{C} \exp \left( \tilde{C} \exp \left( \tilde{C} \exp \left( \tilde{C} \Phi^{2\gamma} \right) \right) \right),$$

we have just proven the theorem for  $k = 0$ .

Suppose that the proposition holds for all  $j$  with  $0 \leq j \leq k-1, k \geq 1$ . Then by induction we have only to prove that  $J_k \leq M_k(\Phi)$  for some  $M_k$ .

Differentiate (6.4)  $(k+1)$ -times with respect to time. Then we have

$$(6.24) \quad \begin{aligned} & \rho u^{(k+2)} - \nu \nabla \operatorname{div} u^{(k+1)} + \nabla p^{(k+1)} \\ &= (\rho f)^{(k+1)} - (\rho u \cdot \nabla u)^{(k+1)} - \sum_{0 \leq j \leq k} \binom{k+1}{j} \rho^{(k+1-j)} u^{(j+1)}. \end{aligned}$$

Multiplying (6.24) by  $u^{(k+2)}$  and integrating over  $\Omega$ , we have

$$(6.25) \quad \begin{aligned} & \int \rho |u^{(k+2)}|^2 dx + \frac{\nu}{2} \frac{d}{dt} \int (\operatorname{div} u^{(k+1)})^2 dx - \int p^{(k+1)} \operatorname{div} u^{(k+2)} dx \\ &= \int (\rho f)^{(k+1)} \cdot u^{(k+2)} dx - \int (\rho u \cdot \nabla u)^{(k+1)} \cdot u^{(k+2)} dx \\ &\quad - \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+2)} dx \\ &= \int \rho (f^{(k+1)} - (u \cdot \nabla u)^{(k+1)}) \cdot u^{(k+2)} dx \\ &\quad + \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} (f^{(j)} - (u \cdot \nabla u)^{(j)}) \cdot u^{(k+2)} dx \\ &\quad - \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+2)} dx. \end{aligned}$$

For the third integration in the left-hand side of (6.25) we have

$$\begin{aligned} & \int p^{(k+1)} \operatorname{div} u^{(k+2)} dx \\ &= \frac{d}{dt} \int p^{(k+1)} \operatorname{div} u^{(k+1)} dx - \int p^{(k+2)} \operatorname{div} u^{(k+1)} dx \\ &\leq \frac{d}{dt} \int p^{(k+1)} \operatorname{div} u^{(k+1)} dx + C |p^{(k+2)}|_{L^2}^2 + |\nabla u^{(k+1)}|_{L^2}^2. \end{aligned}$$

Now we estimate three integrations in the right-hand side of (6.25) as follows. First we have

$$\begin{aligned} & \int \rho (f^{(k+1)} - (u \cdot \nabla u)^{(k+1)}) \cdot u^{(k+2)} dx \\ &\leq C |\rho|_{L^\infty} |f^{(k+1)} - (u \cdot \nabla u)^{(k+1)}|_{L^2}^2 + \frac{1}{2} |\sqrt{\rho} u^{(k+2)}|_{L^2}^2. \end{aligned}$$

For the second integration we have

$$\begin{aligned}
& \int \rho^{(k+1-j)} (f^{(j)} - (u \cdot \nabla u)^{(j)}) \cdot u^{(k+2)} dx \\
&= \frac{d}{dt} \int \rho^{(k+1-j)} (f^{(j)} - (u \cdot \nabla u)^{(j)}) \cdot u^{(k+1)} dx \\
&\quad - \int (\rho^{(k-j+2)} f^{(j)} + \rho^{(k+1-j)} f^{(j+1)}) \cdot u^{(k+1)} dx \\
&\quad + \int (\rho^{(k+2-j)} (u \cdot \nabla u)^{(j)} + \rho^{(k+1-j)} (u \cdot \nabla u)^{(j+1)}) \cdot u^{(k+1)} dx.
\end{aligned}$$

Using the identity  $\rho_t = -\operatorname{div}(\rho u)$ , by integration by parts, we have

$$\begin{aligned}
& \int \rho (f^{(k+1)} - (u \cdot \nabla u)^{(k+1)}) \cdot u^{(k+2)} dx \\
&= \frac{d}{dt} \int \rho^{(k+1-j)} (f^{(j)} - (u \cdot \nabla u)^{(j)}) \cdot u^{(k+1)} dx \\
&\quad + \int (\rho u)^{(k+1-j)} \cdot \left( \nabla (f^{(j)} \cdot u^{(k+1)}) - \nabla ((u \cdot \nabla u)^{(j)} \cdot u^{(k+1)}) \right) dx \\
&\quad - \int \rho^{(k+1-j)} (f^{(j+1)} - (u \cdot \nabla u)^{(j+1)}) \cdot u^{(k+1)} dx \\
&\leq \frac{d}{dt} \int (\rho^{(k+1-j)} f^{(j)} - \rho^{(k+1-j)} (u \cdot \nabla u)^{(j)}) \cdot u^{(k+1)} dx \\
&\quad + C \left( |(\rho u)^{(k+1-j)}|_{H^1}^2 |f^{(j)}|_{H^1}^2 + |(\rho u)^{(j)}|_{H^1}^2 |\nabla (u \cdot \nabla u)^{(j)}|_{L^2}^2 \right) \\
&\quad + C \left( |\nabla u^{(k+1)}|_{L^2}^2 + |\rho^{(k+1-j)}|_{H^1}^2 (|f^{(j+1)}|_{L^2}^2 + |(u \cdot \nabla u)^{(j+1)}|_{L^2}^2) \right) \\
&\quad + C |\nabla u^{(k+1)}|_{L^2}^2
\end{aligned}$$

and hence

$$\begin{aligned}
& \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} (f^{(j)} - (u \cdot \nabla u)^{(k+1-j)}) \cdot u_{(k+2)} dx \\
&\leq \sum_{0 \leq j \leq k} \binom{k+1}{j} \left( \frac{d}{dt} \int (\rho^{(k+1-j)} f^{(j)} - \rho^{(k+1-j)} (u \cdot \nabla u)^{(j)}) \cdot u^{(k+1)} dx \right) \\
&\quad + C \sum_{0 \leq j \leq k} \left( |(\rho u)^{(k+1-j)}|_{H^1}^2 |f^{(j)}|_{H^1}^2 + |(\rho u)^{(j)}|_{H^1}^2 |\nabla (u \cdot \nabla u)^{(j)}|_{L^2}^2 \right) \\
&\quad + C \sum_{0 \leq j \leq k} \left( |\nabla u^{(k+1)}|_{L^2}^2 + |\rho^{(k+1-j)}|_{H^1}^2 (|f^{(j+1)}|_{L^2}^2 + |(u \cdot \nabla u)^{(j+1)}|_{L^2}^2) \right) \\
&\quad + C \sum_{0 \leq j \leq k} |\nabla u^{(k+1)}|_{L^2}^2.
\end{aligned}$$

And finally,

$$\begin{aligned}
& - \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+2)} dx \\
&= - \sum_{0 \leq j \leq k} \binom{k+1}{j} \left( \frac{d}{dt} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+1)} dx \right. \\
&\quad \left. - \int (\rho^{(k-j+2)} u^{(j+1)} + \rho^{(k+1-j)} u^{(j+2)}) \cdot u^{(k+1)} dx \right) \\
&= - \sum_{0 \leq j \leq k} \binom{k+1}{j} \frac{d}{dt} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+1)} dx \\
&\quad + \sum_{0 \leq j \leq k} \binom{k+1}{j} \int (\rho u)^{(k+1-j)} \cdot (\nabla u^{(j+1)} \cdot u^{(k+1)} \\
&\quad + u^{(j+1)} \cdot \nabla u^{(k+1)}) dx \\
&\quad + \sum_{0 \leq j \leq k-1} \binom{k+1}{j} \int \rho^{(k+1-j)} u^{(j+2)} \cdot u^{(k+1)} dx \\
&\quad + \frac{k+1}{2} \int \rho^{(2)} |u^{(k+1)}|^2 dx \\
&\quad + \frac{k+1}{2} \frac{d}{dt} \int \rho^{(1)} |u^{(k+1)}|^2 dx.
\end{aligned}$$

Thus we have

$$\begin{aligned}
& - \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+2)} dx \\
&\leq - \sum_{0 \leq j \leq k} \binom{k+1}{j} \frac{d}{dt} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+1)} dx \\
&\quad + C \sum_{0 \leq j \leq k-1} |(\rho u)^{(k+1-j)}|_{H^1}^2 + |\nabla u^{(k+1)}|_{L^2}^2 \\
&\quad + (k+1) \int (\rho u)^{(1)} (\nabla u^{(k+1)} \cdot u^{(k+1)} + u^{(k+1)} \cdot \nabla u^{(k+1)}) dx \\
&\quad + C \sum_{0 \leq j \leq k-2} |\rho^{(k+1-j)}|_{H^1}^2 |\nabla u^{(j+2)}|_{L^2}^2 \\
&\quad + C |(\rho u)^{(1)}|_{H^1} |\nabla u^{(k+1)}|_{L^2}^2 \\
&\quad + \frac{k+1}{2} \frac{d}{dt} \int \rho^{(1)} |u^{(k+1)}|^2 dx.
\end{aligned}$$

Substituting all these estimates into (6.25), we have

$$\begin{aligned}
& \int \rho |u^{(k+2)}|^2 dx + \frac{\nu}{2} \frac{d}{dt} |\nabla u^{(k+1)}|_{L^2}^2 \\
& \leq \frac{d}{dt} \left( \int p^{(k+1)} \operatorname{div} u^{(k+1)} dx \right. \\
& \quad + \sum_{0 \leq j \leq k+1} \binom{k+1}{j} \int (\rho^{(k+1-j)} f^{(j)} - \rho^{(k+1-j)} (u \cdot \nabla u)^{(j)}) \cdot u^{(k+1)} dx \\
& \quad - \sum_{0 \leq j \leq k} \binom{k+1}{j} \int \rho^{(k+1-j)} u^{(j+1)} \cdot u^{(k+1)} dx \\
& \quad \left. + \frac{k+1}{2} \int \rho^{(1)} |u^{(k+1)}|^2 dx \right) \\
(6.26) \quad & + \Phi |f^{(k+1)} - (u \cdot \nabla u)^{(k+1)}|_{L^2}^2 + \frac{1}{2} |\sqrt{\rho} u^{(k+2)}|_{L^2}^2 \\
& + C \sum_{0 \leq j \leq k} |(\rho u)^{(k+1-j)}|_{H^1}^2 \left( |f^{(j)}|_{H^1}^2 + |\nabla (u \cdot \nabla u)^{(j)}|_{L^2}^2 \right. \\
& \quad \left. + |f^{(j+1)}|_{H^1}^2 + |(u \cdot \nabla u)^{(j+1)}|_{L^2}^2 \right) \\
& + C \sum_{0 \leq j \leq k-1} |(\rho u)^{(k+1-j)}|_{H^1}^2 |\nabla u^{(j+1)}|_{L^2}^2 \\
& + C \sum_{0 \leq j \leq k-2} |(\rho u)^{(k-j+2)}|_{H^1}^2 |\nabla u^{(j+2)}|_{L^2}^2 \\
& + C(1 + |(\rho u)^{(1)}|_{H^1}) |\nabla u^{(k+1)}|_{L^2}^2.
\end{aligned}$$

Now to end up the estimate (6.26) we need the followings.

$$\begin{aligned}
|(u \cdot \nabla u)^{(k+1)}|_{L^2} & \leq C \sum_{0 \leq j \leq k+1} |u^{(k+1-j)} \cdot \nabla u^{(j)}|_{L^2} \\
& \leq C \sum_{0 \leq j \leq k+1} |\nabla u^{(k+1-j)}|_{L^2} |\nabla u^{(j)}|_{L^2} \\
& \leq C \sum_{1 \leq j \leq k} |\nabla u^{(k+1-j)}|_{L^2} |\nabla u^{(j)}|_{L^2} + C |\nabla u^{(k+1)}|_{L^2} |\nabla u|_{L^2},
\end{aligned}$$

$$|\nabla (u \cdot \nabla u)^{(j)}|_{L^2} \leq C \sum_{0 \leq m \leq j} |\nabla u^{(j-m)}|_{H^1} |\nabla u^{(m)}|_{H^1} + |\nabla u^{(j-m)}|_{L^2} |\nabla u^{(m)}|_{L^2}$$

for  $0 \leq j \leq k$ ,

$$|(u \cdot \nabla u)^{(j)}|_{L^2} \leq C \sum_{0 \leq m \leq j} |\nabla u^{(j-m)}|_{L^2} |\nabla u^{(m)}|_{L^2}$$

for  $1 \leq j \leq k$  and

$$|(\rho u)^{(j)}|_{H^1} \leq C \sum_{0 \leq m \leq j} |\rho^{(j-m)} u^{(m)}|_{H^1} \leq C \sum_{0 \leq m \leq j} |\rho^{(j-m)}|_{H^1} |\nabla u^{(m)}|_{L^2}$$

for all  $1 \leq j \leq k+1$ . Since  $J_j \leq M_{k-1}$  for all  $0 \leq j \leq k-1$  from induction hypothesis, integrating (6.26) over  $(\tau, t)$ , we finally have

$$\begin{aligned} & \int_{\tau}^t |\sqrt{\rho}u^{(k+2)}|_{L^2}^2 ds + |\nabla u^{(k+1)}|_{L^2}^2 + \int_{\tau}^t |u^{(k+1)}|_{D^2}^2 ds \\ & + |\rho^{(k+1)}|_{H^1} + \int_{\tau}^t |\rho^{(k+2)}|_{L^2}^2 ds \leq \tilde{C} (M_{k-1}(\Phi))^N \end{aligned}$$

for some large number  $N$ . Furthermore, if we use the elliptic regularity results, we can also obtain

$$\sum_{0 \leq j \leq k} |u^{(j)}(t)|_{D_0^1 \cap D^{2(k-j)+3}} + \int_{\tau}^t |u^{(k)}|_{D^4}^2 ds \leq \tilde{C} (M_{k-1}(\Phi))^N$$

and hence

$$|(\rho^{(j)}, p^{(j)})(t)|_{H^{2(k-j)+2}} + \int_{\tau}^t |p^{(k+2)}|_{L^2}^2 ds \leq \tilde{C} (M_{k-1}(\Phi))^N.$$

Therefore let  $M_k(\Phi) = \tilde{C} (M_{k-1}(\Phi))^N$ . Then the proof of theorem has been completed.  $\square$

#### REFERENCES

- [1] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II. *Comm. Pure Appl. Math.* 17 (1964), 35-92.
- [2] Y. Cho and B. J. Jin, Blow-up of viscous heat-conducting compressible flows. *J. Math. Anal. Appl.*, in press.
- [3] Y. Cho and H. Kim, Existence results for viscous polytropic fluids with vacuum, Hokkaido University Preprint Series in Math. #675.
- [4] Y. Cho and H. Kim, On classical solutions of the compressible Navier-Stokes equations with nonnegative initial densities, *manuscripta math.*, in press; Hokkaido University Preprint Series in Math. #676.
- [5] Y. Cho, H.J. Choe and H. Kim, Unique solvability of the initial boundary value problems for compressible viscous fluids, *J. Math. Pures Appl.* 83 (2004), 243-275.
- [6] H.J. Choe and H. Kim, Strong solutions of the Navier-Stokes equations for isentropic compressible fluids, *J. Differential Equations* 190 (2003), 504-523.
- [7] H.J. Choe and H. Kim, Global existence of the radially symmetric solutions of the Navier-Stokes equations for the isentropic compressible fluids, *Math. Methods Appl. Sci.* 28 (2005), 1-28.
- [8] R. Danchin, Global existence in critical spaces for compressible Navier-Stokes equations, *Invent. math.* 141 (2000), 579-614.
- [9] B. Desjardins, Regularity of weak solutions of the compressible isentropic Navier-Stokes equations, *Comm. Partial Differential Equations* 22 (1997), 977-1008.
- [10] E. Feireisl, Compressible Navier-Stokes equations with a non-monotone pressure law. *J. Differential Equations* 184 (2002), 97-108.
- [11] E. Feireisl, A. Novotný and H. Petzeltová, On the existence of globally defined weak solutions to the Navier-Stokes equations, *J. Math. fluid Mech.* 3 (2001), 358-392.
- [12] E. Feireisl, A. Novotný and H. Petzeltová, On the domain dependence of solutions to the compressible Navier-Stokes equations of a barotropic fluid. *Math. Methods Appl. Sci.* 25 (2002), 1045-1073.
- [13] G. P. Galdi, *An introduction to the mathematical theory of the Navier-Stokes equations*, Springer-Verlag, New York, 1994.



- [14] D. Hoff, Global solutions of the Navier-Stokes equations for multidimensional compressible flow with discontinuous initial data, *J. Differential Equations* 120 (1995), 215-254.
- [15] N. Itaya, On the Cauchy problem for the system of fundamental equations describing the movement of compressible viscous fluids, *Kodai Math. Sem. Rep.* 23 (1971), 60-120.
- [16] S. Jiang and P. Zhang, On spherically symmetric solutions of the compressible isentropic Navier-Stokes equations, *Comm. Math. Phys.* 215 (2001), 559-581.
- [17] S. Jiang and P. Zhang, Axisymmetric solutions of the 3D Navier-Stokes equations for compressible isentropic fluids. *J. Math. Pures Appl.* 82 (2003), 949-973.
- [18] P. L. Lions, Existence globale de solutions pour les équations de Navier-Stokes compressibles isentropiques. *C.R. Acad. Sci.* 316 (1993), 1335-1340.
- [19] P. L. Lions, *Mathematical Topics in Fluid Mechanics Vol. 1, 2.* Clarendon Press, Oxford, 1996.
- [20] A. Matsumura and T. Nishida, The initial value problem for the equations of motion of viscous and heat-conductive gases. *J. Math. Kyoto Univ.* 20 (1980), 67-104.
- [21] A. Matsumura and T. Nishida, The initial boundary value problems for the equations of motion of compressible and heat-conductive fluids. *Comm. Math. Phys.* 89 (1983), 445-464.
- [22] J. Nash, Le problème de Cauchy pour les équations différentielles d'un fluide général. *Bull. Soc. Math. France* 90 (1962), 487-497.
- [23] R. Salvi and I. Straškraba, Global existence for viscous compressible fluids and their behavior as  $t \rightarrow \infty$ . *J. Fac. Sci. Univ. Tokyo Sect. IA, Math.* 40 (1993), 17-51.
- [24] J. Simon, Compact sets in the space  $L^p(0, T; B)$ , *Ann. Mat. Pura Appl.* 146 (1987), 65-96.
- [25] V. A. Solonnikov, Solvability of the initial boundary value problem for the equation of a viscous compressible fluid. *J. Sov. Math.* 14 (1980), 1120-1133.
- [26] A. Tani, On the first initial-boundary value problem of compressible viscous fluid motion. *Publ. Res. Inst. Math. Sci. Kyoto Univ.* 13 (1971), 193-253.
- [27] A. Valli, An existence theorem for compressible viscous fluids, *Ann. Mat. Pura Appl. (IV)* 130 (1982), 197-213; (IV) 132 (1982), 399-400.
- [28] A. Valli, Periodic and stationary solutions for compressible Navier-Stokes equations via a stability method, *Ann. Sc. Norm. Super. Pisa, Cl. Sci.* 10 (1983), 607-647.
- [29] A. Valli and W. M. Zajączkowski, Navier-Stokes equations for compressible fluids: global existence and qualitative properties of the solutions in the general case, *Commun. Math. Phys.* 103 (1986), 259-296.
- [30] Z. Xin, Blowup of smooth solutions to the compressible Navier-Stokes equation with compact density, *Comm. Pure Appl. Math.* 51 (1998), 229-240.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN  
*E-mail address:* ygcho@math.sci.hokudai.ac.jp