Title	Lightlike surfaces of spacelike curves in de Sitter 3-space
Author(s)	FUSHO, Takesi; IZUMIYA, Shyuichi
Citation	Hokkaido University Preprint Series in Mathematics, 784, 1-10
Issue Date	2006-04-14
DOI	10.14943/83934
Doc URL	http://hdl.handle.net/2115/69592
Type	bulletin (article)
File Information	pre784.pdf

[Instructions for use](#)

Lightlike surfaces of spacelike curves in de Sitter 3-space

Takesi FUSHO and Shyuichi IZUMIYA*

April 14, 2006

April 14, 2006

Abstract

The Lorentzian space form with the positive curvature is called de Sitter space which is an important subject in the theory of relativity. In this paper we consider spacelike curves in de Sitter 3-space. We define the notion of lightlike surfaces of spacelike curves in de Sitter 3-space. We investigate the geometric meanings of the singularities of such surfaces.

1 Introduction

The Lorentzian space form with the positive curvature is called de Sitter space. It is an important subject in the theory of relativity and the astrophysics. De Sitter 4-space is known as a vacuum solution of the Einstein equation. In this paper, we consider, however, spacelike curves in de Sitter 3-space as the most elementary case for the study of higher codimensional spacelike submanifolds in non-flat Lorentzian space forms. For spacelike curves in Minkowski 3-space (i.e., the flat Lorentzian space form) and curves in hyperbolic 3-space have been studied in [4, 5]. In this paper we consider a spacelike curve in de Sitter 3-space and the lightlike surface associated to the curve. The lightlike surface is a line bundle along a spacelike curve whose fibres are lightlike lines which is also characterized that each tangent plane at a regular point is lightlike. The singular point of the lightlike surface corresponds to the point of the spacelike curve which has degenerate contact with a lightcone. We can also consider the contact of spacelike curves with a hyperboloid or an ellipsoid in de Sitter space. However the most interesting case is the study on the contact of spacelike curves with lightcones. Moreover, from the point of view of physics, lightlike surfaces are of importance because they are models of different types of horizons studied in relativity theory [1, 3, 7]. Therefore we stick to lightlike surfaces in this paper. The main results are Theorems 2.1 and 6.3. These results give a classification

*Work partially supported by Grant-in-Aid for formation of COE. ‘Mathematics of Nonlinear Structure via Singularities’

2000 Mathematics Subject classification. 53A35, 58C27, 58C28

Key Words and Phrases. de Sitter space, lightlike surface, spacelike curves, lightcone

of the singularities of lightlike surfaces for generic spacelike curves in de Sitter 3-space and a geometric characterization of the singularities. Eventually we find a new Lorentzian invariant σ for spacelike curves which describe the contact with lightcones.

All maps considered here are of class C^∞ unless otherwise stated.

2 Basic concepts and the main result

We now introduce some basic notions on Minkowski 4-space and spacelike curves. For basic results, see [8]. Let $\mathbb{R}^4 = \{(x_1, x_2, x_3, x_4) | x_1, x_2, x_3, x_4 \in \mathbb{R}\}$ be a 4-dimensional vector space. For any vectors $\mathbf{x} = (x_1, x_2, x_3, x_4)$, $\mathbf{y} = (y_1, y_2, y_3, y_4)$ in $\mathbb{R}^4$, the *pseudo scalar product* of $\mathbf{x}$ and $\mathbf{y}$ is defined to be $\langle \mathbf{x}, \mathbf{y} \rangle = -x_1y_1 + x_2y_2 + x_3y_3 + x_4y_4$. We call $(\mathbb{R}^4, \langle \cdot, \cdot \rangle)$ a *Minkowski 4-space*. We write $\mathbb{R}_1^4$ instead of $(\mathbb{R}^4, \langle \cdot, \cdot \rangle)$. We say that a non-zero vector $\mathbf{x} \in \mathbb{R}_1^4$ is *spacelike*, *lightlike* or *timelike* if $\langle \mathbf{x}, \mathbf{x} \rangle > 0$, $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ or $\langle \mathbf{x}, \mathbf{x} \rangle < 0$ respectively. We define the signature of a vector $\mathbf{x}$:

$$\text{sign}(\mathbf{x}) = \begin{cases} 1 & \mathbf{x} \text{ is spacelike} \\ 0 & \mathbf{x} \text{ is lightlike} \\ -1 & \mathbf{x} \text{ is timelike.} \end{cases}$$

The norm of the vector $\mathbf{x} \in \mathbb{R}_1^4$ is defined by $\|\mathbf{x}\| = \sqrt{|\langle \mathbf{x}, \mathbf{x} \rangle|}$. For a vector $\mathbf{v} \in \mathbb{R}_1^4$ and a real number c , we define a *hyperplane with pseudo normal* $\mathbf{v}$ by $HP(\mathbf{v}, c) = \{\mathbf{x} \in \mathbb{R}_1^4 | \langle \mathbf{x}, \mathbf{v} \rangle = c\}$. We call $HP(\mathbf{v}, c)$ a *spacelike hyperplane*, a *timelike hyperplane* or a *lightlike hyperplane* if $\mathbf{v}$ is timelike, spacelike or lightlike respectively. We also define *de Sitter 3-space* by

$$S_1^3 = \{\mathbf{x} \in \mathbb{R}_1^4 | \langle \mathbf{x}, \mathbf{x} \rangle = 1\}.$$

For any $\mathbf{x} = (x_1, x_2, x_3, x_4)$, $\mathbf{y} = (y_1, y_2, y_3, y_4)$, $\mathbf{z} = (z_1, z_2, z_3, z_4) \in \mathbb{R}^4$, we define a vector

$$\mathbf{x} \wedge \mathbf{y} \wedge \mathbf{z} = \begin{vmatrix} -\mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 & \mathbf{e}_4 \\ x_1 & x_2 & x_3 & x_4 \\ y_1 & y_2 & y_3 & y_4 \\ z_1 & z_2 & z_3 & z_4 \end{vmatrix},$$

where $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4)$ is the canonical basis of $\mathbb{R}_1^4$ and $\mathbf{x}_i = (x_1^i, x_2^i, x_3^i, x_4^i)$. We can easily show that $\langle \mathbf{a}, (\mathbf{x} \wedge \mathbf{y} \wedge \mathbf{z}) \rangle = \det(\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{z})$.

We now describe the explicit differential geometry on spacelike curves in S_1^3 . Let $\gamma : I \longrightarrow S_1^3$ be a regular curve. We say that a regular curve γ is *spacelike* if $\dot{\gamma}(t)$ is spacelike at any $t \in I$, where $\dot{\gamma} = d\gamma/dt$. Since γ is a spacelike regular curve, we can reparametrise γ by the arc-length $s = s(t)$. Hence, we may assume that $\gamma(s)$ is a unit speed curve. So we have the tangent vector $\mathbf{t}(s) = \gamma'(s)$ with $\|\mathbf{t}(s)\| = 1$. In the case when $\langle \mathbf{t}'(s), \mathbf{t}'(s) \rangle \neq 1$, we have a unit vector $\mathbf{n}(s) = \frac{\mathbf{t}'(s) - \gamma(s)}{\|\mathbf{t}'(s) - \gamma(s)\|}$. Moreover, define $\mathbf{e}(s) = \gamma(s) \wedge \mathbf{t}(s) \wedge \mathbf{n}(s)$, then we have a pseudo orthonormal frame $\{\gamma(s), \mathbf{t}(s), \mathbf{n}(s), \mathbf{e}(s)\}$ of $\mathbb{R}_1^4$ along γ . By the standard arguments, we can

show the following *Frenet-Serret type formulae*: Under the assumption that $\langle \mathbf{t}'(s), \mathbf{t}'(s) \rangle \neq 1$,

$$\begin{cases} \gamma'(s) = \mathbf{t}(s) \\ \mathbf{t}'(s) = -\gamma(s) + \kappa_g(s)\mathbf{n}(s) \\ \mathbf{n}'(s) = \kappa_g(s)\delta(\gamma(s))\mathbf{t}(s) + \tau_g(s)\mathbf{e}(s) \\ \mathbf{e}'(s) = \tau_g(s)\mathbf{n}(s) \end{cases},$$

where $\delta(\gamma(s)) = -\text{sign}(\mathbf{n}(s))$, $\kappa_g(s) = \|\mathbf{t}'(s) + \gamma(s)\|$ and

$$\tau_g(s) = -\frac{\delta(\gamma(s))}{\kappa_g^2(s)} \det(\gamma(s), \gamma'(s), \gamma''(s), \gamma'''(s)).$$

We now define a surface in de Sitter 3-space associated to a spacelike curve. Let $\gamma : I \longrightarrow S_1^3$ be a unit speed spacelike curve. We define $LS_\gamma : I \times \mathbb{R} \longrightarrow S_1^3$ by

$$LS_\gamma^\pm(s, u) = \gamma(s) + u(\mathbf{n}(s) \pm \mathbf{e}(s)).$$

We call $LS_\gamma^\pm(s, u)$ the *lightlike surface* of γ . We also define the following model surfaces: For any $\mathbf{v}_0 \in S_1^3$, $LC^2(\mathbf{v}_0) = \{\mathbf{u} \in S_1^3 \mid \langle \mathbf{u}, \mathbf{v}_0 \rangle - 1 = 0\}$ is called a *lightcone*. The main purpose in this paper is to study the Lorentzian geometric meanings of the singularities of the lightlike surface. We define a new invariant σ of a spacelike curve in S_1^3 by

$$\sigma(s) = \kappa_g'(s) \mp \kappa_g(s)\tau_g(s).$$

On the other hand, Let $F : S_1^3 \longrightarrow \mathbb{R}$ be a submersion and $\gamma : I \longrightarrow S_1^3$ be a spacelike curve. We say that γ and $F^{-1}(0)$ have *k-point contact* for $t = t_0$ if the function $g(t) = F \circ \gamma(t)$ satisfies $g(t_0) = g'(t_0) = \dots = g^{(k-1)}(t_0) = 0$, $g^{(k)}(t_0) \neq 0$. We also say that γ and $F^{-1}(0)$ have *at least k-point contact* for $t = t_0$ if the function $g(t) = F \circ \gamma(t)$ satisfies $g(t_0) = g'(t_0) = \dots = g^{(k-1)}(t_0) = 0$. The main result in this paper is as follows.

Theorem 2.1 *Let $\gamma : I \longrightarrow S_1^3$ be a unit speed spacelike curve with $\langle \mathbf{t}'(s), \mathbf{t}'(s) \rangle \neq 1$. For $\mathbf{v}_0 = LS_\gamma^\pm(s_0, u_0)$ and the lightcone $LC^2(\mathbf{v}_0) = \{\mathbf{u} \in S_1^3 \mid \langle \mathbf{u}, \mathbf{v}_0 \rangle - 1 = 0\}$, we have the followings:*

- (1) γ and $LC^2(\mathbf{v}_0)$ have at least 2-point contact for s_0 .
- (2) γ and $LC^2(\mathbf{v}_0)$ have 3-point contact for s_0 if and only if

$$\mathbf{v}_0 = \gamma(s_0) - \frac{1}{\delta(\gamma(s_0))\kappa_g(s_0)}(\mathbf{n}(s_0) \pm \mathbf{e}(s_0)) \text{ and } \sigma(s_0) \neq 0.$$

Under this condition, the germ of Image $LS_\gamma^\pm$ at $LS_\gamma^\pm(s_0, u_0)$ is diffeomorphic to the cuspidal edge $C \times \mathbb{R}$.

- (3) γ and $LC^2(\mathbf{v}_0)$ have 4-point contact for s_0 if and only if

$$\mathbf{v}_0 = \gamma(s_0) - \frac{1}{\delta(\gamma(s_0))\kappa_g(s_0)}(\mathbf{n}(s_0) \pm \mathbf{e}(s_0)), \sigma(s_0) = 0 \text{ and } \sigma'(s_0) \neq 0$$

Under this condition, the germ of Image $LS_\gamma^\pm$ at $LS_\gamma^\pm(s_0, u_0)$ is diffeomorphic to the swallowtail SW.

Here, $C \times \mathbb{R} = \{(x_1, x_2) | x_1^2 = x_2^3\} \times \mathbb{R}$ is the cuspidal edge and

$$SW = \{(x_1, x_2, x_3) | x_1 = 3u^4 + u^2v, x_2 = 4u^3 + 2uv, x_3 = v\}$$

is the swallowtail (cf., Fig. 1).

cuspidal edge

swallowtail

Fig.1.

We remark that the above theorem gives a classification of the singularities of the lightlike surface of a generic spacelike curve in de Sitter 3-space (cf., Theorem 6.3).

3 Extended spacelike height functions

In this section we consider a family of Lorentzian invariant functions on a spacelike curve in de Sitter 3-space in order to describe the lightlike surface of the spacelike curve. Let $\gamma : I \longrightarrow S_1^3$ be a unit speed spacelike curve with $\langle \mathbf{t}'(s), \mathbf{t}'(s) \rangle \neq 1$. We now define a family of functions

$$\overline{H} : I \times S_1^3 \longrightarrow \mathbb{R}$$

by $\overline{H}(s, \mathbf{v}) = \langle \gamma(s), \mathbf{v} \rangle - 1$. We call $\overline{H}$ a *extended spacelike height function* of γ . We denote that $\overline{h}_v(s) = \overline{H}(s, \mathbf{v})$.

Proposition 3.1 *For a unit speed spacelike curve $\gamma : I \longrightarrow S_1^3$ with $\langle \mathbf{t}'(s), \mathbf{t}'(s) \rangle \neq 1$, we have the followings:*

(1) $\overline{h}_v(s) = 0$ if and only if there exist $\lambda, \mu, \nu \in \mathbb{R}$ with $\lambda^2 - \delta(\gamma(s))\mu^2 + \delta(\gamma(s))\nu^2 = 0$ such that $\mathbf{v} = \gamma(s) + \lambda\mathbf{t}(s) + \mu\mathbf{n}(s) + \nu\mathbf{e}(s)$.

(2) $\overline{h}_v(s) = \overline{h}'_v(s) = 0$ if and only if there exists $\mu \in \mathbb{R}$ such that $\mathbf{v} = \gamma(s) + \mu\mathbf{n}(s) \pm \mu\mathbf{e}(s)$.

(3) $\overline{h}_v(s) = \overline{h}'_v(s) = \overline{h}''_v(s) = 0$ if and only if

$$\mathbf{v} = \gamma(s) - \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{n}(s) \mp \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{e}(s).$$

(4) $\overline{h}_v(s) = \overline{h}'_v(s) = \overline{h}''_v(s) = \overline{h}'''_v(s) = 0$ if and only if

$$\mathbf{v} = \gamma(s) - \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{n}(s) \mp \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{e}(s) \text{ and } \sigma(s) = 0.$$

(5) $\overline{h}_v(s) = \overline{h}'_v(s) = \overline{h}''_v(s) = \overline{h}'''_v(s) = \overline{h}^{(4)}_v(s) = 0$ if and only if

$$\mathbf{v} = \gamma(s) - \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{n}(s) \mp \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{e}(s), \sigma(s) = 0 \text{ and } \sigma'(s) = 0.$$

Proof. (1) Since $\mathbf{v} \in S_1^3$, there exist $\omega, \lambda, \mu, \nu \in \mathbb{R}$ with $\omega^2 + \lambda^2 - \delta(\gamma(s))\mu^2 + \delta(\gamma(s))\nu^2 = 1$ such that $\mathbf{v} = \lambda\mathbf{t}(s) + \mu\mathbf{n}(s) + \nu\mathbf{e}(s)$. It follows $\bar{h}_v(s) = \langle \gamma(s), \mathbf{v} \rangle - 1 = 0$ that we have $\omega = 1$, so that $\mathbf{v} = \gamma(s) + \lambda\mathbf{t}(s) + \mu\mathbf{n}(s) + \nu\mathbf{e}(s)$. The converse direction also holds.

(2) Since $\bar{h}'_v(s) = \langle \mathbf{t}(s), \mathbf{v} \rangle = 0$, we have $\lambda = 0$. It follows from the fact $-\delta(\gamma(s))\mu^2 + \delta(\gamma(s))\nu^2 = 0$ that $\nu = \pm\mu$. Therefore, we have $\mathbf{v} = \gamma(s) + \mu\mathbf{n}(s) \pm \mu\mathbf{e}(s)$.

(3) Since $\bar{h}''_v(s) = \langle -\gamma(s) + \kappa_g(s)\mathbf{n}(s), \mathbf{v} \rangle = 0$, we have $\mu = -\frac{1}{\delta(\gamma(s))\kappa_g(s)}$, so that

$$\mathbf{v} = \gamma(s) - \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{n}(s) \mp \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{e}(s).$$

(4) Since $\bar{h}'''_v(s) = \langle (-1 + \kappa_g^2(s))\mathbf{t}(s) + \kappa'_g(s)\mathbf{n}(s) + \kappa_g(s)\tau_g(s)\mathbf{e}(s), \mathbf{v} \rangle, \frac{\kappa'_g(s)}{\kappa_g(s)} \mp \tau_g(s) = 0$. It follows that $\sigma(s) = 0$.

(5) We have $\bar{h}^{(4)}_v(s) = \langle (1 - \kappa_g^2(s))\gamma(s) + 3\kappa_g(s)\kappa'_g(s)\mathbf{t} + (-\kappa_g(s) + \kappa''_g(s) + \kappa_g^3(s) + \kappa_g(s)\tau_g^2(s))\mathbf{n}(s) + (\kappa'_g(s)\tau_g(s) + \kappa_g(s)\tau'_g(s))\mathbf{e}(s), \mathbf{v} \rangle = 0$, so that

$$\frac{1}{\kappa_g(s)}(\kappa''_g(s) \mp \kappa'_g(s)\tau_g(s) \mp \kappa_g(s)\tau'_g(s)) = 0.$$

This is equivalent to the condition that $\sigma'(s) = 0$. □

4 Singularities of lightlike surfaces

In this section we study the geometric properties of the lightlike surface of a spacelike curve in de Sitter 3-space. By the propositions in the last section, we can recognize the functions $\kappa_g(s)$ and $\sigma(s)$ have special meanings. Here, we consider the case when the lightlike surface has the most degenerate singularities. We have the following proposition.

Proposition 4.1 *For a unit speed spacelike curve $\gamma : I \longrightarrow S_1^3$ with $\langle t'(s), t'(s) \rangle \neq 1$, we have the followings:*

- (1) *The singularities of $LS_\gamma^\pm$ is the set $\{(s, u) \mid u = -\frac{1}{\kappa_g(s)\delta(\gamma(s))} \ s \in I\}$.*
- (2) *If $LS_\gamma^\pm(s, -\frac{1}{\kappa_g(s)\delta(\gamma(s))}) = \mathbf{v}_0^\pm$ is a contact vector, then $\gamma(s) \in LC^2(\mathbf{u}_0^\pm)$ for any $s \in I$ and $\sigma(s) \equiv 0$. Moreover, $\text{Image } LS^\pm \subset LC^2(\mathbf{u}_0)$.*

Proof. By straightforward calculations, we have

$$\begin{aligned} \frac{\partial LS_\gamma^\pm}{\partial s} &= (1 + u\kappa_g(s)\delta(\gamma(s)))\mathbf{t}(s) + u\tau_g(s)(\mathbf{e}(s) \pm \mathbf{n}(s)) \\ \frac{\partial LS_\gamma^\pm}{\partial u} &= \mathbf{n}(s) \pm \mathbf{e}(s). \end{aligned}$$

The above two vectors are linearly dependent if and only if $1 + u\kappa_g(s)\delta(\gamma(s)) = 0$. This completes the proof of the assertion (1).

For a smooth function $\mu : I \longrightarrow \mathbb{R}$, we define a mapping $f_\mu : I \longrightarrow \mathbb{R}_1^4$ by

$$f_\mu(s) = \gamma(s) + \mu(s)\mathbf{n}(s) \pm \mu(s)\mathbf{e}(s).$$

Suppose that $f_\mu(s) = \mathbf{v}_0^\pm$ is a constant. Then we have

$$\begin{aligned} \frac{df_\mu(s)}{ds} &= (-1 + \mu(s)\kappa_g(s)\delta(\gamma(s)))\mathbf{t}(s) \\ &\quad + (\mu'(s) \pm \mu(s)\tau_g(s))\mathbf{n}(s) + (\mu(s)\tau_g(s) \pm \mu'(s))\mathbf{e}(s) \\ &= 0 \end{aligned}$$

Since the singularities are $\mu(s) = -\frac{1}{\kappa_g(s)\delta(\gamma(s))}$, $\mu'(s) = \frac{\kappa_g'(s)\delta(\gamma(s))}{\kappa_g^2(s)}$. Substituting this relation into the previous equality, we have $\sigma(s) = 0$. Under this condition $LS_\gamma^\pm(s, -\frac{1}{\kappa_g(s)\delta(\gamma(s))}) = \gamma(s) - \frac{1}{\kappa_g(s)\delta(\gamma(s))}\mathbf{n}(s) \mp \frac{1}{\kappa_g(s)\delta(\gamma(s))}\mathbf{e}(s) = \mathbf{v}_0^\pm$ is constant, so that the relation

$$\langle \gamma(s), \mathbf{v}_0^\pm \rangle = \langle \gamma(s), \gamma(s) - \frac{1}{\kappa_g(s)\delta(\gamma(s))}(\mathbf{n}(s) \pm \mathbf{e}(s)) \rangle = 1$$

holds. Moreover, we also have

$$\langle LS_\gamma^\pm(s, u), \mathbf{v}_0^\pm(s) \rangle = \langle \gamma(s) + u(\mathbf{n}(s) \pm \mathbf{e}(s)), \gamma(s) - \frac{1}{\kappa_g(s)\delta(\gamma(s))}(\mathbf{n}(s) \pm \mathbf{e}(s)) \rangle = 1$$

□

5 Unfoldings of functions of one-variables

In this section we use some general results on the singularity theory for families of function germs. Detailed descriptions are found in the book[2]. Let $F : (\mathbb{R} \times \mathbb{R}^r, (s_0, x_0)) \rightarrow \mathbb{R}$ be a function germ. We call F an r -parameter unfolding of f , where $f(s) = F_{x_0}(s, x_0)$. We say that f has A_k -singularity at s_0 if $f^{(p)}(s_0) = 0$ for all $1 \leq p \leq k$, and $f^{(k+1)}(s_0) \neq 0$. We also say that f has $A_{\geq k}$ -singularity at s_0 if $f^{(p)}(s_0) = 0$ for all $1 \leq p \leq k$. Let F be an unfolding of f and $f(s)$ has A_k -singularity ($k \geq 1$) at s_0 . We denote the $(k-1)$ -jet of the partial derivative $\frac{\partial F}{\partial x_i}$ at s_0 by $j^{(k-1)}(\frac{\partial F}{\partial x_i}(s, x_0))(s_0) = \sum_{j=1}^{k-1} \alpha_{ji}s^j$ for $i = 1, \dots, r$. Then F is called a *versal unfolding* if the $k \times r$ matrix of coefficients $(\alpha_{0i}, \alpha_{ji})$ has rank k ($k \leq r$), where $\alpha_{0i} = \frac{\partial F}{\partial x_i}(s_0, x_0)$.

We now introduce an important set concerning the unfolding. The *discriminant set* of F is the set

$$\mathfrak{D}_F = \{x \in \mathbb{R}^r \mid \text{there exists } s \text{ with } F = \frac{\partial F}{\partial s} = 0 \text{ at } (s, x)\}.$$

Then we have the following well-known result (cf., [4]).

Theorem 5.1 *Let $F : (\mathbb{R} \times \mathbb{R}^r, (s_0, x_0)) \rightarrow \mathbb{R}$ be an r -parameter unfolding of $f(s)$ which has the A_k singularity at s_0 . Suppose that F is an versal unfolding.*

- (a) *If $k = 1$, then $\mathfrak{D}_F$ is locally diffeomorphic to $\{0\} \times \mathbb{R}^{r-1}$.*
- (b) *If $k = 2$, then $\mathfrak{D}_F$ is locally diffeomorphic to $C \times \mathbb{R}^{r-2}$.*
- (c) *If $k = 3$, then $\mathfrak{D}_F$ is locally diffeomorphic to $SW \times \mathbb{R}^{r-3}$. We also say that a point $x_0 \in \mathbb{R}^r$ is a fold point of a map germ $f : (\mathbb{R}^r, x_0) \rightarrow (\mathbb{R}^r, f(x_0))$ if there exist diffeomorphism germs $\phi : (\mathbb{R}^r, x_0) \rightarrow (\mathbb{R}^r, 0)$ and $\psi : (\mathbb{R}^r, f(x_0)) \rightarrow (\mathbb{R}^r, 0)$ such that $\psi \circ \phi(x_1, \dots, x_r) = (x_1, \dots, x_{r-1}, x_r^2)$.*

By Proposition 2.1, the discriminant set of the extended height function $\overline{H}(s, \mathbf{v})$ is given as follows:

$$\mathfrak{D}_{\overline{H}} = \{\gamma(s) + u(\mathbf{n}(s) \pm \mathbf{e}(s)) \mid s \in I, u \in \mathbb{R}\}$$

We consider a unit speed spacelike curve $\gamma : I \longrightarrow S_1^3$ with $\langle \mathbf{t}'(s), \mathbf{t}'(s) \rangle \neq 1$ and the extended height function $\overline{H}$ of γ . Then we have the following proposition.

Proposition 5.2 *Suppose that $\mathbf{v} \in \mathfrak{D}_{\overline{H}}$ and $\overline{h}_v(s) = \overline{H}(s, \mathbf{v})$. If $\overline{h}_v$ has A_k -singularity at s ($k = 1, 2, 3$), then $\overline{H}$ is a versal unfolding of $\overline{h}_v$.*

Proof. We denote that

$$\gamma(s) = (x_1(s), x_2(s), x_3(s), x_4(s)) \text{ and } \mathbf{v} = (v_1, v_2, v_3, \pm\sqrt{1-\nu}),$$

where $\nu = v_1^2 + v_2^2 + v_3^2$. Under this notation we have

$$\overline{H}(s, \mathbf{v}) = -x_1(s)v_1 + x_2(s)v_2 + x_3(s)v_3 \pm x_4(s)\sqrt{1-\nu} - 1.$$

Thus we have

$$\begin{aligned} \frac{\partial \overline{H}}{\partial v_1}(s, \mathbf{v}) &= -x_1(s) \pm \frac{v_1}{\sqrt{1-\nu}}x_4(s) \\ \frac{\partial \overline{H}}{\partial v_i}(s, \mathbf{v}) &= x_i(s) \mp \frac{v_i}{\sqrt{1-\nu}}x_4(s) (i = 2, 3). \end{aligned}$$

We also have

$$\begin{aligned} \frac{\partial}{\partial s} \frac{\partial \overline{H}}{\partial v_1}(s, \mathbf{v}) &= -x_1'(s) \mp \frac{v_1}{\sqrt{1-\nu}}x_4'(s) \\ \frac{\partial}{\partial s} \frac{\partial \overline{H}}{\partial v_i}(s, \mathbf{v}) &= x_i'(s) \pm \frac{v_i}{\sqrt{1-\nu}}x_4'(s) (i = 2, 3) \end{aligned}$$

and

$$\begin{aligned} \frac{\partial^2}{\partial s^2} \frac{\partial \overline{H}}{\partial v_1}(s, \mathbf{v}) &= -x_1''(s) \pm \frac{v_1}{\sqrt{1-\nu}}x_4''(s) \\ \frac{\partial^2}{\partial s^2} \frac{\partial \overline{H}}{\partial v_i}(s, \mathbf{v}) &= x_i''(s) \mp \frac{v_i}{\sqrt{1-\nu}}x_4''(s) (i = 2, 3). \end{aligned}$$

Therefore the 2-jet of $\frac{\partial \overline{H}}{\partial v_i}(s, \mathbf{v})$ ($i = 1, 2, 3$) at s_0 is given by

$$\begin{aligned} \frac{\partial \overline{H}}{\partial v_i}(s, \mathbf{v}) &= \frac{\partial \overline{H}}{\partial v_i}(s_0, \mathbf{v}) + \frac{1}{2} \frac{\partial}{\partial s} \frac{\partial \overline{H}}{\partial v_i}(s_0, \mathbf{v})(s - s_0) + \frac{1}{6} \frac{\partial^2}{\partial s^2} \frac{\partial \overline{H}}{\partial v_i}(s_0, \mathbf{v})(s - s_0)^2 \\ &= \alpha_{0,i} + \frac{1}{2} \alpha_{1,i}(s - s_0) + \frac{1}{6} \alpha_{2,i}(s - s_0)^2. \end{aligned}$$

It is enough to show that the rank of the matrix A is 3, where

$$A = \begin{pmatrix} \alpha_{0,1} & \alpha_{0,2} & \alpha_{0,3} \\ \alpha_{1,1} & \alpha_{1,2} & \alpha_{1,3} \\ \alpha_{2,1} & \alpha_{2,2} & \alpha_{2,3} \end{pmatrix}$$

We denote that

$$A(i, j, k) = \det \begin{pmatrix} x_i(s) & x_j(s) & x_k(s) \\ x_i'(s) & x_j'(s) & x_k'(s) \\ x_i''(s) & x_j''(s) & x_k''(s) \end{pmatrix}$$

then we have

$$\begin{aligned} \det A &= -(A(1, 2, 3) \mp \frac{v_1}{\sqrt{1-\nu}} A(4, 2, 3) \mp \frac{v_2}{\sqrt{1-\nu}} A(1, 4, 3) \mp \frac{v_3}{\sqrt{1-\nu}} A(1, 2, 4)) \\ &= \pm \langle (v_1, v_2, v_3, \pm\sqrt{1-\nu}), (-A(2, 3, 4), -A(1, 3, 4), A(1, 2, 4), -A(1, 2, 3)) \rangle \\ &= \pm \left\langle (v_1, v_2, v_3, \pm\sqrt{1-\nu}), \begin{vmatrix} -\mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 & \mathbf{e}_4 \\ x_1(s) & x_2(s) & x_3(s) & x_4(s) \\ x_1'(s) & x_2'(s) & x_3'(s) & x_4'(s) \\ x_1''(s) & x_2''(s) & x_3''(s) & x_4''(s) \end{vmatrix} \right\rangle \\ &= \pm \frac{1}{\sqrt{1-\nu}} \langle \mathbf{v}, \gamma(s) \wedge \gamma'(s) \wedge \gamma''(s) \rangle. \end{aligned}$$

Since $\mathbf{v} \in \mathfrak{D}_{\overline{H}}$ is a singular point,

$$\mathbf{v} = \gamma(s) - \frac{1}{\delta(\gamma(s))\kappa_g(s)}(\mathbf{n}(s) \pm \mathbf{e}(s)).$$

Moreover, we have

$$\begin{aligned} \gamma(s) \wedge \gamma'(s) \wedge \gamma''(s) &= \gamma(s) \wedge \mathbf{t}(s) \wedge (-\gamma(s) + \kappa_g(s)\mathbf{n}(s)) \\ &= \kappa_g(s)\gamma(s) \wedge \mathbf{t}(s) \wedge \mathbf{n}(s) \\ &= \kappa_g(s)\mathbf{e}(s). \end{aligned}$$

Therefore we have

$$\begin{aligned} \det A &= \pm \frac{1}{\sqrt{1-\nu}} \langle \gamma(s) - \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{n}(s) \mp \frac{1}{\delta(\gamma(s))\kappa_g(s)}\mathbf{e}(s), \kappa_g(s)\mathbf{e}(s) \rangle \\ &= -\frac{\delta(\gamma(s))}{\sqrt{1-\nu}} \neq 0 \end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 2.2. Let $\gamma : I \longrightarrow S_1^3$ be a unit speed spacelike curve with $\langle t'(s), t'(s) \rangle \neq 1$. For $\mathbf{v}_0^\pm = LS_\gamma^\pm(s_0, u_0)$, we define a function $\mathcal{H} : S_1^3 \longrightarrow \mathbb{R}$ by $\mathcal{H}(\mathbf{u}) = \langle \mathbf{u}, \mathbf{v}_0^\pm \rangle - 1$. Then we have $\bar{h}_{v_0^\pm}(s) = \mathcal{H}(\gamma(s))$. Since $LC^2(\mathbf{v}_0^\pm) = \mathcal{H}^{-1}(0)$ and 0 is a regular value of $\mathcal{H}$, $\bar{h}_{v_0^\pm}$ has the A_k -singularity at s_0 if and only if γ and $LC^2(\mathbf{v}_0^\pm)$ have $k+1$ -point contact for s_0 . It follows from Proposition 3.1, Theorem 5.1 and Proposition 5.2 that the proof of Theorem 2.2 is completed. $\square$

6 Generic properties of spacelike curves

In this section we consider generic properties of spacelike curves in S_1^3 . The main tool is a kind of transversality theorems. Let $\text{Emb}_{sp}(I, S_1^3)$ be the space of spacelike embeddings $\gamma :$

$I \longrightarrow S_1^3$ with $\langle \mathbf{t}', \mathbf{t}' \rangle \neq 1$ equipped with Whitney C^∞ -topology. We also consider the function $\mathcal{H} : S_1^3 \times S_1^3 \longrightarrow \mathbb{R}$ defined by $\mathcal{H}(\mathbf{u}, \mathbf{v}) = \langle \mathbf{u}, \mathbf{v} \rangle - 1$. We claim that $\mathcal{H}_v$ is a submersion for any $\mathbf{v} \in S_1^3$, where $\mathcal{H}_v(\mathbf{u}) = \mathcal{H}(\mathbf{u}, \mathbf{v})$. For any $\gamma \in \text{Emb}_{sp}(I, S_1^3)$, we have $\overline{H} = \mathcal{H} \circ (\gamma \times id_{S_1^3})$. We also have the ℓ -jet extension

$$j_1^\ell \overline{H} : I \times S_1^3 \longrightarrow J^\ell(I, \mathbb{R})$$

defined by $j_1^\ell \overline{H}(s, \mathbf{v}) = j_1^\ell \overline{h}_v(s)$. We consider the trivialisation $J^\ell(I, \mathbb{R}) \equiv I \times \mathbb{R} \times J^\ell(1, 1)$. For any submanifold $Q \subset J^\ell(1, 1)$, we denote that $\widetilde{Q} = I \times \{0\} \times Q$. Then we have the following proposition as a corollary of Lemma 6 in Wassermann[9].

Proposition 6.1 *Let Q be a submanifold of $J^\ell(1, 1)$. Then the set*

$$T_Q = \{\gamma \in \text{Emb}_{sp}(I, S_1^3) \mid j_1^\ell \overline{H} \text{ is transversal to } \widetilde{Q}\}$$

is a residual subset of $\text{Emb}_{sp}(I, S_1^3)$. If Q is a closed subset, then T_Q is open.

Let $f : (\mathbb{R}, 0) \longrightarrow (\mathbb{R}, 0)$ be a function germ which has an A_k -singularity at 0. It is well-known that there exists a diffeomorphism germ $\phi : (\mathbb{R}, 0) \longrightarrow (\mathbb{R}, 0)$ such that $f \circ \phi(s) = \pm s^{k+1}$. This is the classification of A_k -singularities. For any $z = j^\ell f(0) \in J^\ell(1, 1)$, we have the orbit $L^\ell(z)$ given by the action of the Lie group of ℓ -jets of diffeomorphism germs. If f has an A_k -singularity, then the codimension of the orbit is k . There is another characterisation of versal unfoldings as follows(cf., [6]):

Proposition 6.2 *Let $F : (\mathbb{R} \times \mathbb{R}^r, 0) \longrightarrow (\mathbb{R}, 0)$ be an r -parameter unfolding of $f : (\mathbb{R}, 0) \longrightarrow (\mathbb{R}, 0)$ which has an A_k -singularity at 0. Then F is a versal unfolding if and only if $j_1^\ell F$ is transversal to the orbit $L^\ell(j^\ell f(0))$ for $\ell \geq k + 1$.*

Here, $j_1^\ell F : (\mathbb{R} \times \mathbb{R}^r, 0) \longrightarrow J^\ell(\mathbb{R}, \mathbb{R})$ is the ℓ -jet extension of F given by $j_1^\ell F(s, x) = j^\ell F_x(s)$.

The generic classification theorem is given as follows:

Theorem 6.3 *There exists an open and dense subset $\mathcal{O} \subset \text{Emb}_{sp}(I, S_1^3)$ such that for any $\gamma \in \mathcal{O}$, the lightlike surface $LS_\gamma^\pm$ of γ is locally diffeomorphic to the cuspidal edge or the swallowtail at any singular point.*

Proof. For $\ell \geq 4$, we consider the decomposition of the jet space $J^\ell(1, 1)$ into $L^\ell(1)$ orbits. We now define a semi-algebraic set by

$$\Sigma^\ell = \{z = j^\ell f(0) \in J^\ell(1, 1) \mid f \text{ has an } A_{\geq 4}\text{-singularity}\}.$$

Then the codimension of Σ^ℓ is 4. Therefore, the codimension of $\widetilde{\Sigma}_0 = I \times \{0\} \times \Sigma^\ell$ is 5. We have the orbit decomposition of $J^\ell(1, 1) - \Sigma^\ell$ into

$$J^\ell(1, 1) - \Sigma^\ell = L_0^\ell \cup L_1^\ell \cup L_2^\ell \cup L_3^\ell,$$

where L_k^ℓ is the orbit through an A_k -singularity. Thus, the codimension of $\widetilde{L}_k^\ell$ is $k + 1$. We consider the ℓ -jet extension $j_1^\ell \overline{H}$ of the extended spacelike height function $\overline{H}$. By Proposition 6.1, there exists an open and dense subset $\mathcal{O} \subset \text{Emb}(I, S_1^3)$ such that $j_1^\ell \overline{H}$ is transversal to $\widetilde{L}_k^\ell$ ($k = 0, 1, 2, 3$) and the orbit decomposition of $\widetilde{\Sigma}^\ell$. This means that $j_1^\ell \overline{H}(I \times S_1^3) \cap \widetilde{\Sigma}^\ell = \emptyset$ and $\overline{H}$ is a versal unfolding of h at any point (s_0, v_0) . By Theorem 5.1, the discriminant set of $\overline{H}$ (i.e., the lightlike surface of γ) is locally diffeomorphic to the cuspidal edge or the swallow tail if the point is singular. $\square$

References

- [1] M.A.Akivis and V.V.Goldberg, *Differential Geometry of Varieties with Degenerate Gauss Maps*, Canadian Mathematical Society, Springer, 2004.
- [2] J. W. Bruce and P. J. Giblin, *Curves and singularities (second edition)*. Cambridge University press (1992)
- [3] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, International Series of Monographs on Physics. 69 Oxford Univeristy press, 1983.
- [4] S.Izumiya, D.Pei and T.Sano, *The lightcone Gauss map and the lightcone developable of a spacelike curve in Minkowski 3-space*, Glasgow Math. J. **42** (2000), 75-89.
- [5] S. Izumiya, D-H. Pei and T. Sano, *Horospherical surfaces of curves in Hyperbolic space*, Publ. Math. (Debrecen) **64** (2004),1–13
- [6] J. Martinet, *Singularities of Smooth Functions and Maps*, London Math. Soc. Lecture Note Series, Cambridge Univ. Press,**58** (1982)
- [7] C. W. Misner, K. S. Thorpe and J. W. Wheeler, *Gravitation*, W. H. Freeman and Co., San Francisco, CA, 1973.
- [8] B. O'neil, *Semi-Riemannian Geometry*. Academic Press, New York (1983).
- [9] G. Wassermann, *Stability of Caustics*. Math. Ann., **210** (1975), 43–50

Takesi Fusho, Department of Mathematics, Hokkaido University, Sapporo 060-0810, Japan
 Shyuichi Izumiya, Department of Mathematics, Hokkaido University, Sapporo 060-0810, Japan
 e-mail:izumiya@math.sci.hokudai.ac.jp