



Title	Invariant Subspaces Of Toeplitz Operators And Uniform Algebras
Author(s)	Nakazi, Takahiko
Citation	Hokkaido University Preprint Series in Mathematics, 815, 1-9
Issue Date	2006
DOI	10.14943/83965
Doc URL	http://hdl.handle.net/2115/69623
Type	bulletin (article)
File Information	pre815.pdf



[Instructions for use](#)

Invariant Subspaces Of Toeplitz Operators And Uniform Algebras

By

Takahiko Nakazi

* This research was partially supported by No.17540176

2000 Mathematics Subject Classification : Primary 47 B 35, 47 A 15, Secondary
46 J 15

Key words and phrases : Toeplitz operator, invariant subspace, analytic symbol

Abstract Let T_ϕ be a Toeplitz operator on the one variable Hardy space H^2 . We show that if T_ϕ has a nontrivial invariant subspace in the set of invariant subspaces of T_z then ϕ belongs to H^∞ . In fact, we also study such a problem for the several variables Hardy space H^2 .

§1. Introduction

Let X be a compact Hausdorff space, let $C(X)$ be the algebra of complex-valued continuous functions on X , and let A be a uniform algebra on X . A probability measure m (on X) denotes a representing measure for some nonzero complex homomorphism. The abstract Hardy space $H^p = H^p(m)$, $1 \leq p \leq \infty$, determined by A is defined to be the closure of A in $L^p = L^p(m)$ when p is finite and to be the weak* closure of A in $L^\infty = L^\infty(m)$ when $p = \infty$.

Let P be the orthogonal projection from L^2 onto H^2 . For ϕ in L^∞ , put

$$T_\phi f = P(\phi f) \quad (f \in H^2)$$

and then T_ϕ is called a Toeplitz operator. In this paper, we are interested in invariant subspaces of Toeplitz operators. Put $\mathcal{A} = \{T_\phi ; \phi \in H^\infty\}$ and $\mathcal{A}^* = \{T_\phi^* ; \phi \in H^\infty\}$. $\text{Lat } T_\phi$ denotes the set of all invariant subspaces of T_ϕ , $\text{Lat } \mathcal{A} = \cap \{\text{Lat } T_\phi ; \phi \in H^\infty\}$ and $\text{Lat } \mathcal{A}^* = \cap \{\text{Lat } T_\phi^* ; \phi \in H^\infty\}$. We don't know whether arbitrary T_ϕ has a nontrivial invariant subspace. When ϕ is in H^∞ and H^∞ has a nonconstant unimodular function q , T_ϕ has a nontrivial invariant subspace $M = qH^2$. Hence $\text{Lat } T_\phi \neq \{\langle 0 \rangle, H^2\}$.

Let K be the orthogonal complement of \bar{H}^2 in L^2 . Then $L^2 = H^2 \oplus \bar{K}$. $I(H^\infty)$ denotes the set of all unimodular functions in H^∞ . A function in $I(H^\infty)$ is called an inner function. For a subset Y in L^∞ , Y^\perp denotes $\{g \in L^1 ; \int g \bar{f} dm = 0 \quad (f \in Y)\}$.

In this paper we study the following four natural questions :

Question 1. *If $\text{Lat } T_\phi \supseteq \text{Lat } \mathcal{A}$ then does T_ϕ belong to \mathcal{A} ?*

Question 2. *Suppose that H^∞ is a weak* closed maximal algebra in L^∞ . If $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$ then is $\text{Lat } T_\phi = \{\langle 0 \rangle, H^2\}$?*

Question 3. *Is $\text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$?*

Question 4. *Can we describe $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$ or equivalently $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A}^*$?*

In this paper, we will answer these four questions positively when A is the disc algebra. In fact, for Question 1 we can do it for more general uniform algebras. However for Question 2 we could not answer even for simple uniform algebras. Question 3 can be answered for almost all uniform algebras.

In this paper $H^p(D^n)$ denotes the Hardy space on the polydisc D^n and $H^p(\Omega)$ denotes the Hardy space on a finitely connected domain Ω . $L_a^p(D)$ denotes the Bergman space on D and put $N^2 = L^2(D) \ominus \{L_a^2(D) \oplus \bar{z}\bar{L}_a^2(D)\}$. H_0^p denotes the set of $\{f \in H^p ; \int f dm = 0\}$. $H^p(\Gamma)$ denotes the usual Hardy space on the dual group $\hat{\Gamma}$ where Γ is an ordered subgroup of the reals.

§2. Lat $\mathcal{A} \subseteq \text{Lat } T_\phi$

In this section we study Question 1. Theorem 1 shows that Question 1 can be answered positively for very general uniform algebras.

Lemma 1. *Let M be a closed subspace of H^2 . $M \in \text{Lat } T_\phi$ if and only if $\phi M \subset M \oplus \bar{K}$.*

Proof. By definition of a Toeplitz operator, this is clear.

Lemma 2. *If ϕ is a function in L^∞ and $\text{Lat } \mathcal{A} \subseteq \text{Lat } T_\phi$ then $\phi = \phi_0 + \bar{k}_0$ where $\phi_0 \in H^2$ and $\bar{k}_0 \in \cap\{\bar{q}\bar{K} ; q \in I(H^\infty)\}$.*

Proof. Since $L^2 = H^2 \oplus \bar{K}$, there exist $h \in H^2$ and $k \in K$ such that $\phi = h + \bar{k}$. If $q \in I(H^\infty)$ then $qH^2 \in \text{Lat } \mathcal{A}$ and so by Lemma 1 $\phi q = qh + q\bar{k} \in qH^2 + \bar{K}$. Since $T_\phi q \in qH^2$ and $qh \in qH^2$, $P(q\bar{k}) \in qH^2$. Hence $q\bar{k} = q\ell + \bar{t}$ where $\ell \in H^2$ and $\bar{t} \in \bar{K}$. Therefore $\bar{k} = \ell + \bar{q}\bar{t}$ and $\ell = \bar{k} - \bar{q}\bar{t} \in H^2 \cap \bar{K} = \langle 0 \rangle$. Hence $\ell = 0$ and $\bar{k} = \bar{q}\bar{t}$. This implies that k belongs to $\bar{q}\bar{K}$ for any $q \in I(H^\infty)$.

Theorem 1. *Suppose that $\cap\{\bar{q}\bar{K} ; q \in I(H^\infty)\} = \langle 0 \rangle$. If ϕ is a function in L^∞ and $\text{Lat } \mathcal{A} \subseteq \text{Lat } T_\phi$ then ϕ belongs to H^∞ .*

Proof. Lemmas 1 and 2 imply the theorem trivially.

Corollary 1. *Suppose that $H^2 = H^2(\mathbf{T}^N)$. If ϕ is a function in L^∞ and $\text{Lat } \mathcal{A} \subseteq \text{Lat } T_\phi$ then ϕ belongs to H^∞ .*

Proof. K is an invariant subspace under multiplications by the coordinates functions z_1, \dots, z_n . $\cap\{z_1^{\ell_1} \dots z_n^{\ell_n} K ; (\ell_1, \dots, \ell_n) \geq (0, \dots, 0)\}$ is a reducing subspace and so $\cap z_1^{\ell_1} \dots z_n^{\ell_n} K = \chi_E L^2$ for some characteristic function χ_E . Since $\chi_E L^2$ is orthogonal to \bar{H}^2 , $\chi_E = 0$ and so $\langle 0 \rangle = \cap \bar{z}_1^{\ell_1} \dots \bar{z}_n^{\ell_n} \bar{K} = \cap\{\bar{q}\bar{K} ; q \in I(H^\infty)\}$.

Corollary 2. *Suppose that $H^2 = H^2(\Omega)$. If ϕ is a function in L^∞ and $\text{Lat } \mathcal{A} \subseteq \text{Lat } T_\phi$ then ϕ belongs to H^∞ .*

Proof. Let Z be the Ahlfors function for Ω then $|Z| = 1$ on $\partial\Omega = X$ (see [3]).

$\bigcap_{n=0}^{\infty} \bar{Z}^n \bar{K}$ is invariant under the multiplications by Z and \bar{H}^∞ . Since H^∞ is a weak* maximal subalgebra of L^∞ , $\bigcap_{n=0}^{\infty} \bar{Z}^n \bar{K} = \chi_E L^2$. Since $\chi_E L^2$ is orthogonal to H^2 , $\chi_E = 0$ and so $\cap\{\bar{q}\bar{K} ; q \in I(H^\infty)\} = \{0\}$.

Corollary 3. *Let A be a Dirichlet algebra (see [4]). If ϕ is a function in L^∞ and $\text{Lat } \mathcal{A} \subseteq \text{Lat } T_\phi$ then ϕ belongs to H^∞ .*

Proof. Since H^∞ is a uniform algebra which has the annulus property ([2],[6]) on a totally disconnected space, by [2, Theorem 1] the set of quotients of inner functions is norm dense in the set of unimodular functions in L^∞ . In this situation, $\bar{K} = \bar{H}_0^2$ and $Y = \cap\{\bar{q}\bar{K} ; q \in I(H^\infty)\} \subset \bar{H}^2$. $\bar{q}Y = Y$ for any q in $I(H^\infty)$ and so $\bar{q}_1\bar{q}_2Y \subseteq Y$ for any q_1, q_2 in $I(H^\infty)$. Hence $\phi Y \subseteq Y$ for any unimodular function ϕ in L^∞ . Hence $Y = \chi_E L^2$ for the characteristic function χ_E for some set E . Since $Y \subset \bar{H}^2$, Y must be $\{0\}$.

Proposition 1. *Suppose that $H^2 = L_a^2(D)$, ϕ is a function in L^∞ and $\text{Lat } \mathcal{A} \subseteq \text{Lat } T_\phi$. Then the following are valid.*

(1) ϕ belongs to $L_a^2(D) + N^2$.

(2) If $\phi = f + \ell$ where $f \in H^\infty$ and $\ell \in N^2$ then $\text{Lat } T_\ell \supseteq \text{Lat } \mathcal{A}$.

Proof. (1) Since $zL_a^2 \in \text{Lat } T_\phi$ by hypothesis, $\mathcal{C} \in \text{Lat } T_\phi^* = \text{Lat } T_{\bar{\phi}}$ and so $\bar{\phi} = \bar{c} + \bar{k}$ where $c \in \mathcal{C}$ and $k \in zL_a^2(D) + N^2$. Hence $\phi \in L_a^2(D) + N^2$. (2) If $\phi = f + \ell$ and $M \in \text{Lat } \mathcal{A}$ then $\phi M \subset M + \bar{K}$. Hence $(f + \ell)g = fg + \ell g \in M + \bar{K}$ for any $g \in M$. Since $fg \in M$, $\ell g \in M + \bar{K}$ for any $g \in M$ and so $\ell M \subset M + \bar{K}$. Thus $M \in \text{Lat } T_\ell$.

A bounded operator B is called reflexive if whenever C is a bounded operator and $\text{Lat } B \subseteq \text{Lat } C$ then C belongs to the closed algebra (in weak operator topology) generated by B . When B is subnormal, it is known that B is reflexive [7]. Hence if f is a nonzero function in H^∞ and $\text{Lat } T_\phi \supseteq \text{Lat } T_f$ then T_ϕ belongs to the closed algebra generated by T_f . Hence T_ϕ belongs to \mathcal{A} . Usually $\text{Lat } \mathcal{A} \subsetneq \text{Lat } T_f$ and so this does not answer Question 1. However if there exists a function f in H^∞ such that $\text{Lat } T_f = \text{Lat } \mathcal{A}$ then the above result about subnormal operators answers Question 1. Hence when $H^2 = H^2(T)$, if $\text{Lat } T_\phi \supseteq \text{Lat } \mathcal{A}$ then T_ϕ belongs to \mathcal{A} because $\text{Lat } T_z = \text{Lat } \mathcal{A}$. Therefore Corollary 1 is not new for $N = 1$. Similarly Question 1 can be answered for $H^2 = L_a^2(D)$. Hence Proposition 1 is a very weak result.

§3. $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$

In this section we study Question 2. Theorem 2 shows that Question 2 can be answered positively for the disc algebra. In fact, it gives a few results for more general uniform algebras about Question 2.

Lemma 3. *Let Q be a function in $I(H^\infty)$. Then $\bar{K} = \sum_{n=0}^{\infty} \oplus (\bar{K} \ominus \bar{Q}\bar{K})\bar{Q}^n \oplus \bigcap_{n=0}^{\infty} \bar{Q}^n \bar{K}$.*

Proof. Since $|Q| = 1$ a.e. and $\bar{Q}\bar{K} \subset \bar{K}$, \bar{Q} is an isometry on \bar{K} . Hence this is well known and called a Wold decomposition.

Theorem 2. *Suppose that $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$. If $M \in \text{Lat } T_\phi$ and $\cap \{\bar{Q}^n \bar{K} ; Q \in \mathcal{I}\} = \{0\}$ for some subset \mathcal{I} in $I(H^\infty)$ then there exists a nonconstant Q in \mathcal{I} such that $M \cap (H^2 \ominus QH^2) \neq \langle 0 \rangle$ or $\phi M \subseteq M$.*

Proof. If $M \in \text{Lat } T_\phi$ then by Lemma 1 there exist $f \in M$, $g \in M$ and $k \in K$ such that $\phi f = g + \bar{k}$. If $\phi M \not\subseteq M$ then we may assume that $k \neq 0$. For any fixed $Q \in \mathcal{I}$, by Lemma 3 $\bar{K} = \left\{ \sum_{n=0}^{\infty} \oplus (\bar{K} \ominus \bar{Q}\bar{K})\bar{Q}^n \right\} \oplus \bigcap_{n=0}^{\infty} \bar{Q}^n \bar{K}$ and so

$$\bar{k} = \sum_{n=0}^{\infty} k_n \bar{Q}^n + k_\infty$$

where $k_n \in \bar{K} \ominus \bar{Q}\bar{K}$ ($n = 0, 1, 2, \dots$) and $k_\infty \in \bigcap_{n=0}^{\infty} \bar{Q}^n \bar{K}$. Then $Q\bar{k} = Qk_0 + \sum_{n=1}^{\infty} k_n \bar{Q}^{n-1} + Qk_\infty$ and by Lemma 1 $Q\bar{k}$ belongs to $M + \bar{K}$ because $\phi f = g + \bar{k}$ and $QM \subset M$.

Suppose that there does not exist a nonconstant function Q in \mathcal{I} such that $M \cap (H^2 \ominus QH^2) \neq \langle 0 \rangle$. Then we will get a contradiction. By what was proved above, Qk_0 belongs to $M \cap (H^2 \ominus QH^2) = \{0\}$. Hence $k_0 \equiv 0$. Next we consider $Q^2 \bar{k}$ and then $k_1 \equiv 0$ follows. Proceeding similarly we can show that $\bar{k} = k_\infty$. By hypothesis, this implies that $\bar{k} \equiv 0$ because Q is arbitrary in \mathcal{I} . This contradiction implies that there exists $Q \in \mathcal{I}$ such that $M \cap (H^2 \ominus QH^2) \neq \langle 0 \rangle$.

Corollary 4. *Suppose that $H^2 = H^2(\mathbf{T}^N)$, ϕ is a function in L^∞ and $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$. If $M \in \text{Lat } T_\phi$ and $M \neq \langle 0 \rangle$ then M contains a nonzero function which is $(N - 1)$ -variable. Hence if $N = 1$ then $M = H^2$.*

Proof. It is known that if $\phi M \subseteq M$ then $\phi \in H^\infty$. Hence we may assume that $\phi M \not\subseteq M$. Put $\mathcal{I} = \{z_1, \dots, z_N\}$ then \mathcal{I} satisfies the condition of Theorem 2. By Theorem 2, there exists z_j such that $1 \leq j \leq N$ and $(H^2 \ominus z_j H^2) \cap M \neq \{0\}$. Since $H^2 \ominus z_j H^2 = H^2(z'_j, T^{N-1})$ where $z = (z_j, z'_j)$, M contains a nonzero $(N - 1)$ -variable function.

Corollary 5. *Suppose that $H^2 = H^2(\Omega)$, $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$ and Z is the Alfors function for Ω (see [3]). If $M \in \text{Lat } T_\phi$ and $M \neq \langle 0 \rangle$ then $M \cap (H^2 \ominus ZH^2) \neq \langle 0 \rangle$.*

Proof. Put $\mathcal{I} = \{Z\}$ then \mathcal{I} satisfies the condition of Theorem 2. It is known that if $\phi M \subseteq M$ then $\phi \in H^\infty$. Hence we may assume that $\phi M \not\subseteq M$.

Proposition 2. *If T_ϕ is subnormal and $\text{Lat } T_\phi \subseteq \text{Lat } \mathcal{A}$ then T_ϕ commutes with \mathcal{A} and so $T_\phi f = P(\phi_0 f)$ ($f \in H^\infty$) for some ϕ_0 in H^2 . If \mathcal{A} is a uniform algebra which approximates in modulus on X then ϕ belongs to $H^2 \cap L^\infty$.*

Proof. If T_ϕ is subnormal and $\text{Lat } T_\phi \subseteq \text{Lat } \mathcal{A}$ then it is known [7] that \mathcal{A} is contained in the closed algebra generated by T_ϕ . Hence T_ϕ commutes with \mathcal{A} . Let $\phi_0 =$

$T_\phi 1$ then $T_\phi f = T_\phi T_f 1 = T_f T_\phi 1 = P(\phi_0 f)$ for $f \in H^\infty$. Since $\|\phi_0 f\|_2 \leq \|T_\phi\| \|f\|_2$ ($f \in H^\infty$),

$$\left| \int_X \phi_0 f \bar{g} dm \right| \leq \|\phi\|_\infty \|f\|_2 \|g\|_2 \quad (f, g \in H^\infty).$$

Hence

$$\left| \int_X \phi_0 |f|^2 dm \right| \leq \|\phi\|_\infty \|f^2\|_1.$$

Since A approximates in modulus on X , ϕ_0 belongs to $H^2 \cap L^\infty$. It is easy to see that $\phi = \phi_0$.

Corollary 6. *Suppose that $H^2 = H^2(\mathbf{T}^N)$ or $H^2 = H^2(\Omega)$. If T_ϕ is subnormal then $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$ or ϕ belongs to H^∞ .*

Proof. A uniform algebra A approximates in modulus on X , that is, for every positive continuous function g on X and $\varepsilon > 0$, there is an f in A with $|g - |f|| < \varepsilon$ if the set of unimodular elements of A separates points of X (see [6, Lemma 4.12]). Since the coordinate functions z_1, \dots, z_n separate T^N , the polydisc algebra approximates in modulus on T^N . If T_ϕ is subnormal on $H^2(T^N)$ and $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$ then by Proposition 2 ϕ belongs to $H^2(T^N) \cap L^\infty = H^\infty(T^N)$. If $A = H^\infty(\Omega)$ then by [3, Lemma 4.8] $I(H^\infty(\Omega))$ separates $X =$ the maximal ideal space of $L^\infty(\partial D)$. Hence Corollary 6 for $H^2 = H^2(\Omega)$ follows from Proposition 2.

Proposition 3. *If $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$, then $\text{Lat } T_\phi^* \cap \text{Lat } T_\phi \subset \text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A}$.*

Proof. If $M \in \text{Lat } T_\phi^*$ then $M^\perp \in \text{Lat } T_\phi$ and so $M^\perp \in \text{Lat } \mathcal{A}$ because $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$. Hence $M \in \text{Lat } \mathcal{A}^*$ and so $\text{Lat } T_\phi^* \subsetneq \text{Lat } \mathcal{A}^*$.

By Proposition 3, when $\text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$, if $\text{Lat } T_\phi \subsetneq \text{Lat } \mathcal{A}$ then T_ϕ does not have a nontrivial reducing subspace. Hence if T_ϕ is normal then $\text{Lat } T_\phi \not\subsetneq \text{Lat } \mathcal{A}$. Therefore it is important to know that $\text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$, that is, \mathcal{A} is irreducible.

§4. $\text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A}$

In this section we study Question 3. Theorem 3 shows that Question 3 can be answered positively for usual uniform algebras. Recall $\mathcal{A}^* = \{T_\phi^* ; \phi \in H^\infty\}$.

Theorem 3. *If $M \in \text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A}$ then $M \subset \chi_E L^2 \subset M + \bar{K}$ where $E = \cup\{\text{supp } f ; f \in M\}$. Hence if $E = X$ then $M = H^2$.*

Proof. If $\phi \in L^\infty$ then by the Stone-Weierstrass theorem for any $\varepsilon > 0$ there exist f_1, \dots, f_n and g_1, \dots, g_n in H^∞ such that $\|\phi - \sum_{j=1}^n f_j \bar{g}_j\|_\infty < \varepsilon$. Since $T_{f_j \bar{g}_j} M \subset M$ for

$j = 1, \dots, n$, $T_\phi M \subset M$. By Lemma 1 $\phi M \subset M \oplus \bar{K}$. Thus $\chi_E L^2 \subset M \oplus \bar{K}$. If $E = X$ then $L^2 = M \oplus \bar{K}$ and so $M = H^2$.

Corollary 7. *Suppose that there does not exist a nonzero function in H^2 such that $m(\{x \in X ; f(x) = 0\}) \neq 0$. If $M \in \text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A}$ then $M = \langle 0 \rangle$ or H^2 .*

§5. Lat $T_\phi \cap \text{Lat } \mathcal{A}$

In this section we study Question 4. We don't know whether $\text{Lat } T_\phi \neq \{\langle 0 \rangle, H^2\}$. However we show that $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$ if $\phi \notin H^\infty$ and $H^2 = H^2(\mathbf{T})$. For any M in $\text{Lat } T_\phi$, put

$$K_M = \{k \in K ; \bar{k} = \phi f - g \text{ for some } f \text{ and } g \in M\}$$

, then $K_M \subseteq K$ and $\phi M \subset M + \bar{K}_M$ (see Lemma 1).

Theorem 4. *If $M \in \text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$ then $K_M \times (H^2 \ominus M) \subseteq (H^\infty)^\perp$ and $T_k^*(H^\infty) \subseteq M$ for any k in K_M .*

Proof. By the remark above, if $M \in \text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$ then $\phi M \subset M + \bar{K}_M$. If $k \in K_M$ then by its definition there exist f and g such that $\phi f = g + \bar{k}$. For any $\ell \in H^\infty$, $\phi f \ell = g \ell + \bar{k} \ell \in M + \bar{K}_M$ and so $P(\bar{k} \ell) \in M$. Since

$$\bar{k} \ell = P(\bar{k} \ell) + (I - P)(\bar{k} \ell) \in M + \bar{K}_M,$$

if $s \in H^2 \ominus M$ then $\langle \bar{k} \ell, s \rangle = \langle P(\bar{k} \ell), s \rangle = 0$. Hence $k s$ belongs to $(H^\infty)^\perp$ and so $K_M \times (H^2 \ominus M) \subseteq (H^\infty)^\perp$. The above proof implies that $T_k^*(H^\infty) \subseteq M$.

Corollary 8. *Suppose that $H^2 = H^2(\Omega)$, $\mathcal{C} \setminus \Omega$ has n components and $\phi \notin H^\infty$. If $M \in \text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$ then $\dim(H^2 \ominus M) \leq n$.*

Proof. By Theorem 4

$$K_M \times (H^2 \ominus M) \subseteq (H^\infty)^\perp \cap (\bar{H}^\infty)^\perp = (H^\infty + \bar{H}^\infty)^\perp \cap L^1$$

and $\dim(H^\infty + \bar{H}^\infty)^\perp \cap L^1 = n$ because $\mathcal{C} \setminus \Omega$ has n components. If $K_M = \langle 0 \rangle$ then $\phi M \subset M$. It is known [4] that L^∞ is generated by ϕ and H^∞ in the weak* topology. Hence $M \in \text{Lat } \mathcal{A} \cap \text{Lat } \mathcal{A}^* = \{\langle 0 \rangle, H^2\}$ by Corollary 7 and so $M = H^2$. It is clear that if $K_M \neq \langle 0 \rangle$ then $\dim(H^2 \ominus M) \leq n$.

Corollary 9. *If $H^2 = H^2(\mathbf{T})$ and $\phi \notin H^\infty$ then $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$.*

Proof. When Ω is the open unit disc, $H^2(\Omega) = H^2(\mathbf{T})$ and so by Corollary 8 $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$.

Corollary 10. *Let A be a Dirichlet algebra. If $\phi \notin H^\infty$ then $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$.*

Proof. It is known that $(\bar{H}^\infty)^\perp \cap (H^\infty)^\perp = \langle 0 \rangle$. The corollary is a result of Theorem 4.

In general, it seems to be difficult to describe $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$. When $H^2 = H^2(\Omega)$ and $\bar{\phi} \in H^\infty$, $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$ by Corollary 8. In fact, if $M \in \text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$ then $\bar{\phi}(H^2 \ominus M) \subseteq H^2 \ominus M$. Since $\dim(H^2 \ominus M) < \infty$ by Corollary 8, M must be equal to H^2 . When $H^2 = H^2(T^2)$ and $\phi = \bar{z}$, $\text{Lat } T_\phi \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, qH^2(w, T); q = q(w) \text{ is a one variable inner function}\}$ where z and w are the independent variables on T^2 . In fact, if $M \in \text{Lat } T_\phi \cap \text{Lat } \mathcal{A}$ then $T_z^* M_1$ is orthogonal to M where $M_1 = M \ominus zM$. Since $T_z^* M_1 \subset M$, $T_z^* M_1 = \langle 0 \rangle$ and so $M_1 \subset H^2(w, T)$. Corollary 10 shows that $\text{Lat } \mathcal{A}^* \cap \text{Lat } \mathcal{A} = \{\langle 0 \rangle, H^2\}$ if A is a Dirichlet algebra.

References

1. R. G. Douglas, Banach Algebra Techniques in Operator Theory, Academic Press, 1972
2. R. G. Douglas and W. Rudin, Approximation by inner functions, Pacific J.M. 31(1969), 313-320
3. S. Fisher, Function Theory On Planar Domains, John Wiley and Son, New York, 1983
4. T. Gamelin, Uniform Algebras, Prentice-Hall, Englewood Cliffs, N.J.. 1969
5. T. W. Gamelin, J.B.Garnett, L.A.Rubel, and A.L.Shields, On badly approximable functions, J.Approx.Theory 17(1976), 280-296
6. I. Glicksberg, Measures orthogonal to algebras and sets of antisymmetry, Trans. Amer. Math.Soc. 105(1962), 415-435
7. R. F. Olin and J.E.Thomson, Algebras of subnormal operators, J.Funct.Anal. 37(1980), 271-301

Department of Mathematics
Faculty of Science
Hokkaido University
Sapporo 060-0810, Japan
nakazi@math.sci.hokudai.ac.jp