



| | |
|------------------|---|
| Title | Remarks on the relativistic Hartree equations |
| Author(s) | Cho, Yonggeun; Ozawa, Tohru; Sasaki, Hironobu; Shim, Yong-sun |
| Citation | Hokkaido University Preprint Series in Mathematics, 827, 1-19 |
| Issue Date | 2007 |
| DOI | 10.14943/83977 |
| Doc URL | http://hdl.handle.net/2115/69636 |
| Type | bulletin (article) |
| File Information | pre827.pdf |



[Instructions for use](#)

REMARKS ON THE RELATIVISTIC HARTREE EQUATIONS

YONGGEUN CHO, TOHRU OZAWA, HIRONOBU SASAKI, AND YONG-SUN SHIM

ABSTRACT. We study the global well-posedness (GWP) and small data scattering of radial solutions of the relativistic Hartree type equations with nonlocal nonlinearity $F(u) = \lambda(|\cdot|^{-\gamma} * |u|^2)u$, $\lambda \in \mathbb{R} \setminus \{0\}$, $0 < \gamma < n$, $n \geq 3$. We establish a weighted L^2 Strichartz estimate applicable to non-radial functions and some fractional integral estimates for radial functions.

1. INTRODUCTION

In this paper, we consider the Cauchy problems concerning the relativistic Hartree equations:

$$(1) \quad \begin{cases} i\partial_t u = \sqrt{1 - \Delta}u + F(u) & \text{in } \mathbb{R}^n \times \mathbb{R}, \ n \geq 3, \\ u(0) = \varphi, \end{cases}$$

$$(2) \quad \begin{cases} \partial_t^2 u + (1 - \Delta)u = F(u) & \text{in } \mathbb{R}^n \times \mathbb{R}, \ n \geq 3, \\ u(0) = \varphi_1, \quad \partial_t u(0) = \varphi_2. \end{cases}$$

The nonlinear part $F(u)$ is of Hartree type such that $F(u) = V_\gamma(u)u$, where

$$V_\gamma(u)(x) = \lambda(|\cdot|^{-\gamma} * |u|^2)(x) = \lambda \int_{\mathbb{R}^n} \frac{|u(y)|^2}{|x - y|^\gamma} dy.$$

Here λ is a non-zero real number and γ is a positive number less than the space dimension n .

The first equation (1) is called the semi-relativistic equation which describes the Boson stars [6, 7, 13] and the second one (2) is the well-known Klein-Gordon equation whose physical model is originated from the helium atom [10, 14, 17]. For the simplicity of presentation, the mass, speed of light and Planck constant of both equations have been normalized.

The equations (1) and (2) can be rewritten in the form of the integral equations

$$(3) \quad u(t) = U(t)\varphi - i \int_0^t U(t - t')F(u)(t')dt',$$

$$(4) \quad u(t) = (\cos t\omega)\varphi_1 + \omega^{-1}(\sin t\omega)\varphi_2 - \int_0^t \omega^{-1}(\sin(t - t')\omega)F(u) dt',$$

1991 *Mathematics Subject Classification.* Primary: 35Q40, 35Q55; Secondary: 47J35.

Key words and phrases. relativistic Hartree type equations, global well-posedness, scattering, radial solutions.

where $\omega = \sqrt{1 - \Delta}$ and the associated unitary group $U(t)$ is realized by the Fourier transform as

$$U(t)\varphi = (e^{-it\omega}\varphi)(x) \equiv \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{ix \cdot \xi} e^{-it\sqrt{1+|\xi|^2}} \widehat{\varphi}(\xi) d\xi,$$

where \widehat{g} denotes the Fourier transform of g defined by $\widehat{g}(\xi) = \int_{\mathbb{R}^n} e^{-ix \cdot \xi} g(x) dx$. The operators $\cos t\omega$ and $\sin t\omega$ are defined by replacing $e^{-it\sqrt{1+|\xi|^2}}$ with

$$\cos(t\sqrt{1+|\xi|^2}) \quad \text{and} \quad \sin(t\sqrt{1+|\xi|^2}),$$

respectively.

If the solution u of (1) or (3) has a decay at infinity and smoothness, it satisfies two conservation laws:

$$(5) \quad \begin{aligned} \|u(t)\|_{L^2} &= \|\varphi\|_{L^2}, \\ E_1(u) &\equiv K_1(u) + V(u) = E_1(\varphi), \\ K(u) &= \frac{1}{2} \langle \sqrt{1 - \Delta} u, u \rangle, V(u) = \frac{1}{4} \langle F(u), u \rangle, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ is the complex inner product in L^2 . Also the solution of (2) or (4) satisfies the conservation law:

$$(6) \quad \begin{aligned} E_2(u, \partial_t u) &\equiv K_2(u, \partial_t u) + V(u) = E_2(\varphi_1, \varphi_2), \\ K_2(u, \partial_t u) &= \frac{1}{2} (\langle \partial_t u, \partial_t u \rangle + \langle \sqrt{1 - \Delta} u, \sqrt{1 - \Delta} u \rangle). \end{aligned}$$

The main concern of this paper is to establish the global well-posedness and scattering of radial solutions of the equations (1) and (2).

The study of the global well-posedness (GWP) and scattering for the semi-relativistic equation (1) has not been long before. In [15], GWP was considered with a three dimensional Coulomb type potential which corresponds to $\gamma = 1$. In [4], the first and second authors of the present paper showed GWP for $0 < \gamma \leq 1$ if $n \geq 2$ and $0 < \gamma < 1$ if $n = 1$, for $0 < \gamma < \frac{2n}{n+1}$ if $n \geq 2$, and small data scattering for $\gamma > 2$ if $n \geq 3$. In this paper we tried to fill the gap $1 < \gamma \leq 2$ for GWP under the assumption of radial symmetry. For further study like blowup of solutions, solitary waves, mean field limit problem for semi-relativistic equation, see the references [13, 6, 7, 8, 9].

The first result is on the GWP for radial solutions of (3).

Theorem 1. *Let $1 < \gamma < \frac{3}{2}$ for $n = 3$ and $1 < \gamma < 2$ for $n \geq 4$. Let $\varphi \in H^{\frac{1}{2}}$ be radially symmetric and assume that $\|\varphi\|_{L^2}$ is sufficiently small if $\lambda < 0$. Then there exists a unique radial solution $u \in C_b H^{\frac{1}{2}}$ such that $|x|^{-1}u \in L_{loc}^2 L^2$ of (3) satisfying the energy and L^2 conservations (5).*

Here $C_b = C \cap L^\infty$, $H_r^s = (1 - \Delta)^{-s/2} L^r$ and $\dot{H}_r^s = (-\Delta)^{-s/2} L^r$ are the usual and homogeneous Sobolev spaces, respectively. We mean H_2^s by H^s and \dot{H}_2^s by \dot{H}^s . Hereafter, the space $L_T^q(B)$ denotes $L^q(-T, T; B)$ for $T > 0$ and $\|\cdot\|_{L_T^q B}$ its norm for some Banach space B . If $T = \infty$, we use $L^q(B)$ for $L^q(\mathbb{R}; B)$ with norm $\|\cdot\|_{L^q B}$, $1 \leq q \leq \infty$. We also denote $v \in L_T^q(B)$ for all $T < \infty$ by $v \in L_{loc}^q(B)$.

The next result is on the small data scattering of radial solutions of (3) for $n \geq 4$.

Theorem 2. *Let $\gamma = 2$ and $n \geq 4$. Let $\varphi \in H^1$ be radially symmetric. If $\|\varphi\|_{H^1}$ is sufficiently small, then there exists a unique radial solution $u \in C_b H^1$ such that $|x|^{-1}u \in L^2 L^2$ to (3). Moreover, there exist radial functions φ^+ and φ^- such that*

$$\|u(t) - U(t)\varphi^\pm\|_{H^1} \rightarrow 0 \quad \text{as } t \rightarrow \pm\infty.$$

In [4], the authors used the $L^q L^r$ type Strichartz estimates of the Klein-Gordon equation to prove GWP and scattering for the equation (1). Contrary to the case of Klein-Gordon equation, semi-relativistic equation preserves a regularity in a contraction argument based on the Strichartz estimate, from which the gap $(\frac{2n}{n+1} \leq \gamma \leq 2)$ arises naturally in the range of γ for GWP. To tide over this difficulty, we assume the radial symmetry for data and solutions, which enables us to estimate fractional integrals associated with the nonlinearity $V_\gamma(u)u$. Then we establish an L^2 Strichartz estimate for $n \geq 2$ with weight $|x|^{-a}$ which is useful to treat radial functions but also applicable to non-radial functions (a gain of angular regularity is achieved in the non-radial case). The one dimensional analog is attainable. See Remark 4 below.

For GWP we use a fractional integral estimate on the unit sphere such that

$$\int_{S^{n-1}} |re_1 - \rho\sigma|^{-\theta} d\sigma \leq M(r, \rho) < \infty,$$

where $e_1 = (1, 0, \dots, 0)$. The result of Theorem 1 corresponds to the case $\theta = \gamma + \frac{1}{2}$. If $n = 3$, then the finiteness of integrals enforces γ to be less than $\frac{3}{2}$ as in Theorem 1, since the integral is finite only when $n - 2 - \theta > -1$. For details see Lemma 2 and Lemma 3. In Theorem 2, we treated the case $\theta = 2$ for which the integral is not finite if $n = 3$. However, the three dimensional GWP can be slightly improved up to $\frac{5}{3}$ by using another Strichartz estimate on a hybrid Sobolev space (for this see [5]). It will be worthy of trying to fill the gap $\frac{5}{3} \leq \gamma \leq 2$ for $n = 3$.

The Klein-Gordon equation (2) was initially studied by [26] (see also [18]). In [21], the GWP is considered for $\lambda < 0$ and $0 < \gamma \leq 4$. It was proved in [26, 20, 23] that the scattering operator for (2) is well-defined on some domain if $n \geq 2$, $4/3 < \gamma \leq 4n/(n+1)$ and $\gamma < n$. Furthermore, [19] showed that if $n \geq 3$, $2 \leq \gamma \leq 4$ and $\gamma < n$, then the scattering operator can be defined on some neighborhood near zero in the energy space.

In this paper the small data scattering of radial solutions is successfully treated below energy space, provided $\frac{3}{2} < \gamma < 2$. To state precisely, let us define a weighted spaces $W_{s,\varepsilon}$ and a data space $D_{\alpha,\beta}$ by

$$W_{s,\varepsilon} = \{\psi \in L^2 : \|\psi\|_{W_{s,\varepsilon}}^2 \equiv \|\cdot\|^{-s-\varepsilon}\psi\|_{L^2(|x|\leq 1)}^2 + \|\cdot\|^{-s+\varepsilon}\psi\|_{L^2(|x|>1)}^2 < \infty\}$$

$$\text{and } D_{\alpha,\beta} = H^{\alpha-\frac{1}{2}} \cap L^{\frac{2n}{n+2-2\beta}},$$

respectively, where $\varepsilon > 0$ is sufficiently small.

Theorem 3. *Let $\frac{3}{2} < \gamma < 2$ for $n = 3$ and $\frac{3}{2} < \gamma \leq 2$ for $n \geq 4$. Then there is a real number s and ε such that*

$$(7) \quad \frac{1}{2} < s < \frac{\gamma}{2}, \quad 0 < \varepsilon < \min\left(\frac{\gamma}{2} - s, s - \frac{1}{2}\right), \quad 1 + s - \varepsilon < \gamma < 1 + s + \varepsilon.$$

For fixed such s and ε , let $(\varphi_1, \varphi_2) \in D_{s+\varepsilon, s+\varepsilon} \times D_{s+\varepsilon-1, s+\varepsilon}$ be radially symmetric data. Then if $\|\varphi_1\|_{D_{s+\varepsilon, s+\varepsilon}} + \|\varphi_2\|_{D_{s+\varepsilon-1, s+\varepsilon}}$ is sufficiently small, then there exists a unique radial solution $u \in C_b H^{s-\frac{1}{2}+\varepsilon} \cap L^2 W_{s, \varepsilon}$ to (4). Moreover, there exist radial functions $\varphi_1^\pm \in H^{s-\frac{1}{2}+\varepsilon}$ and $\varphi_2^\pm \in H^{s-\frac{3}{2}+\varepsilon}$ such that

$$\|u(t) - u^\pm(t)\|_{H^{s-\frac{1}{2}+\varepsilon}} \rightarrow 0 \quad \text{as } t \rightarrow \pm\infty,$$

where u^\pm is the solutions to the Cauchy problem

$$(8) \quad \begin{cases} \partial_t^2 u^\pm + (1 - \Delta)u^\pm = 0, \\ u^\pm(0) = \varphi_1^\pm, \partial_t u^\pm(0) = \varphi_2^\pm. \end{cases}$$

In the definition of initial data space $D_{\alpha, \beta}$ the space $L^{\frac{2n}{n+2-2\beta}}$ can be slightly weakened by the homogeneous Sobolev space $\dot{H}^{-(1-\beta)}$. In fact, $L^{\frac{2n}{n+2-2\beta}} \hookrightarrow \dot{H}^{-(1-\beta)}$ for $0 < \beta < 1$. See the proof of Theorem 3 below. Let $\tilde{D}_{\alpha, \beta}$ be the weakened space $H^{\alpha-\frac{1}{2}} \cap \dot{H}^{-(1-\beta)}$. Then one can easily show that the solution $u \in C_b(\mathbb{R}; \dot{H}^{-(1-(s-\varepsilon))})$ and then the existence of scattering operator of (2) on a small neighborhood of the origin in $\tilde{D}_{s+\varepsilon, s-\varepsilon} \times \tilde{D}_{s+\varepsilon-1, s-\varepsilon}$. For details see Remark 6 below.

The lower bound $\frac{3}{2}$ of γ is caused by the condition (7) which follows from the relation between the weight $|x|^{-a}$ and the L^2 estimate of Bessel function such that

$$\int_0^\infty r^{1-2a} |J_{\frac{n-2}{2}}(r)|^2 dr < \infty.$$

For the finiteness, the assumption $\frac{1}{2} < a < \frac{n}{2}$ is inevitable because $J_{\frac{n-2}{2}}(r) = O(r^{\frac{n-2}{2}})$ as $r \rightarrow 0$ and $J_{\frac{n-2}{2}}(r) = O(r^{-\frac{1}{2}})$ as $r \rightarrow \infty$. For more explicit formula, see the identity (13) below. Hence for the present it seems hard to improve the range of γ for the small data scattering. From the perspective of negative result for the scattering¹, it will be very interesting to show the scattering up to the value of γ greater than 1.

This paper is organized as follows. In Section 2 we introduce a weighted Strichartz estimate for $n \geq 2$. In Section 3 some fractional integral estimates are considered under radial symmetry. All the proofs of theorems are shown in Section 4.

If not specified, throughout this paper, the notation $A \lesssim B$ and $A \gtrsim B$ denote $A \leq CB$ and $A \geq C^{-1}B$, respectively. Different positive constants possibly depending on n, λ, γ and a might be denoted by the same letter C . $A \sim B$ means that both $A \lesssim B$ and $A \gtrsim B$ hold.

2. WEIGHTED L^2 STRICHARTZ ESTIMATE

In this section, we show the following weighted L^2 Strichartz estimate.

¹The non-existence of the asymptotically free solutions occurs when $\gamma \leq 1$. For instance see the last section of [4].

Proposition 1. *Let $\frac{1}{2} < a < \frac{n}{2}$ and $n \geq 2$. Then for any $\varphi \in H^s$ and $F \in L_T^1 H^s$, $s \geq 0$, we have*

$$(9) \quad \begin{aligned} & \|U(\cdot)\varphi\|_{L_T^2(\tilde{H}_a^s H_\sigma^{a-\frac{1}{2}})} \lesssim \|\varphi\|_{H^s}, \\ & \left\| \int_0^\bullet U(\cdot - t')F(t') dt' \right\|_{L_T^2(\tilde{H}_a^s H_\sigma^{a-\frac{1}{2}})} \lesssim \|F\|_{L_T^1 H^s}. \end{aligned}$$

The constants in the estimates can be chosen independently of T .

Here we denote the weighted Sobolev space \tilde{H}_a^s by

$$\tilde{H}_a^s = \{v : \|v\|_{\tilde{H}_a^s} \equiv \| |\cdot|^{-a} L_a(\Delta)(1 - \Delta)^{\frac{s}{2}} v \|_{L^2} < \infty\},$$

where a is a positive real number and $L_a(\Delta) = (-\Delta)^{\frac{1}{2}(1-a)}(1 - \Delta)^{-\frac{1}{4}}$. The Sobolev space $H_\sigma^\alpha = (1 - \Delta_\sigma)^{-\frac{\alpha}{2}} L^2(S^{n-1})$ is defined on the unit sphere S^{n-1} , where Δ_σ is the Laplace-Beltrami operator on the unit sphere S^{n-1} (see [11, 16] for instance). The mixed norm $\tilde{H}_a^s H_\sigma^\alpha$ is defined as follows.

$$\|v\|_{\tilde{H}_a^s H_\sigma^\alpha}^2 = \int_{\mathbb{R}^n} |x|^{-2a} |L_a(\Delta)(1 - \Delta)^{\frac{s}{2}}(1 - \Delta_\sigma)^{\frac{\alpha}{2}} v|^2 dx.$$

Remark 1. If φ and F are radially symmetric, then the angular regularity H_σ^α is not necessary.

Remark 2. If we use Theorem 3.4 of [4] for small data GWP, then from the Strichartz estimates above, we readily observe that if $2 < \gamma < n, n \geq 3, s > \frac{\gamma}{2} - \frac{n-2}{2}$, and $\|\varphi\|_{H^s}$ is sufficiently small, then for $\frac{1}{2} < a < \frac{n}{2}$ the solution u of (1) is in $L^2(\tilde{H}_a^s H_\sigma^{a-\frac{1}{2}})$. In fact, in view of the proof of Theorem 3.4 of [4] we have $\|F(u)\|_{L^1 H^s} \lesssim \|\varphi\|_{H^s}$ and hence

$$\|u\|_{L^2(\tilde{H}_a^s H_\sigma^{a-\frac{1}{2}})} \lesssim \|\varphi\|_{H^s} + \|F(u)\|_{L^1 H^s} \lesssim \|\varphi\|_{H^s}.$$

Similarly, we have by using Lemma 2.4 of [23] that if $\frac{4}{3} < \gamma < 2, n \geq 2$ and $\|\varphi_1\|_{H^s} + \|\varphi_2\|_{H^{s-1}}$ is sufficiently small for $s \geq 1$, then the solution u of (2) is in $L^2(\tilde{H}_a^s H_\sigma^{a-\frac{1}{2}})$ for any $\frac{1}{2} < a < \frac{n}{2}$.

Proof of Proposition 1. Without loss of generality we may assume $s = 0$. Let us first define an operator $W_\nu(t)$ by

$$(10) \quad (W_\nu(t)f)(r) = r^{-a} \int_0^\infty e^{-it\sqrt{1+\rho^2}}(r\rho)^{\frac{1}{2}} J_\nu(r\rho) \frac{\rho^{1-a}}{(1+\rho^2)^{\frac{1}{4}}} f(\rho) d\rho,$$

where ν is a real number greater than equal to $\frac{n-2}{2}$ and J_ν is the Bessel function of order ν . We claim that for any $\nu \geq \frac{n-2}{2}$ and $\frac{1}{2} < a < \frac{n}{2}$

$$(11) \quad \|W_\nu(\cdot)f\|_{L^2(\mathbb{R}_+ \times (-T, T))} \leq \|W_\nu(\cdot)f\|_{L^2(\mathbb{R}_+^2)} \lesssim (1 + \nu)^{-(a-\frac{1}{2})} \|f\|_{L^2}.$$

In fact, using the change of variables $\sqrt{1+\rho^2} \mapsto \rho$, we have

$$\begin{aligned} (W_\nu(t)f)(r) &= r^{\frac{1}{2}-a} \int_{-\infty}^{\infty} e^{-it\rho} \chi_{(1,\infty)}(\rho) J_\nu(r\sqrt{\rho^2-1}) \sqrt{\rho} (\rho^2-1)^{\frac{1}{4}-\frac{a}{2}} f(\sqrt{\rho^2-1}) d\rho, \end{aligned}$$

where $\chi_{(1,\infty)}$ is the characteristic function on the interval $(1, \infty)$. Now from the Plancherel theorem with respect to the time variable and the change of variables $\sqrt{\rho^2-1} \mapsto \rho$, it follows that

$$\begin{aligned} (12) \quad \|W_\nu(\cdot)f\|_{L^2(\mathbb{R}_+^2)}^2 &= 2\pi \int_0^\infty \rho^{2-2a} |f(\rho)|^2 \int_0^\infty r^{1-2a} |J_\nu(r\rho)|^2 dr d\rho \\ &= 2\pi \left(\int_0^\infty r^{1-2a} |J_\nu(r)|^2 dr \right) \|f\|_{L^2}^2. \end{aligned}$$

For the estimate of inner integral, we use the known formula on Bessel function (see p. 402 of [27]) such that for any $\nu \geq \frac{n-2}{2}$ and $\frac{1}{2} < a < \frac{n}{2}$

$$(13) \quad \int_0^\infty r^{1-2a} |J_\nu(r)|^2 dr = \frac{\Gamma(2a-1)\Gamma(\nu+1-a)}{2^{2a-1}\Gamma(a)^2\Gamma(\nu+a)},$$

which has no singularity at $\nu \geq \frac{n-2}{2}$. We note that the numerator on the RHS is finite as far as $\frac{1}{2} < a < \frac{n}{2} \leq \nu+1$. By Stirling's formula such that $\Gamma(s) \sim s^{s-\frac{1}{2}} e^{-(s-1)}$ for large s (for instance, see [2]), we get (11).

From now on, we prove the proposition. We expand φ with radial functions and spherical harmonic functions as follows:

$$\varphi(r\sigma) = \sum_{k \geq 0} \sum_{1 \leq l \leq d(k)} g_{k,l}(r) Y_{k,l}(\sigma), \quad (r, \sigma) \in (0, \infty) \times S^{n-1},$$

where $g_{k,l}$ are radial functions such that

$$\int_0^\infty |g_{k,l}(r)|^2 r^{n-1} dr < \infty,$$

$Y_{k,l}$ are orthonormal spherical harmonics of order k , and $d(k)$ is the dimension of the space of spherical harmonics of order k . See [3, 11, 16].

By the orthonormality, we have

$$\|\varphi\|_{L^2}^2 \sim \sum_{k \geq 0} \sum_{1 \leq l \leq d(k)} \int_0^\infty |g_{k,l}(r)|^2 r^{n-1} dr.$$

Using the Fourier transform of spherical harmonic functions (see for instance [25]), we have

$$\widehat{g_{k,l} Y_{k,l}}(\rho\sigma) = G_{k,l}(\rho) Y_{k,l}(\sigma),$$

where

$$\begin{aligned} G_{k,l}(\rho) &= c_n \int_0^\infty g_{k,l}(r) r^{n-1} (r\rho)^{-\frac{n-2}{2}} J_{\nu(k)}(r\rho) dr, \\ \text{and } \nu(k) &= \frac{2k+n-2}{2}. \end{aligned}$$

The constant c_n is independent of k . By the Plancherel theorem, one can easily observe that

$$\int_0^\infty |g_{k,l}(r)|^2 r^{n-1} dr = \int_0^\infty |G_{k,l}(\rho)|^2 \rho^{n-1} d\rho.$$

Now let us define functions $f_{k,l}$ by $f_{k,l}(\rho) = G_{k,l}(\rho)\rho^{\frac{n-1}{2}}$. Then from the Fourier transform of spherical harmonic functions, we have

$$r^{-a} L_a(\Delta) U(t) \varphi(r\sigma) = r^{-\frac{n-1}{2}} \sum_{k,l} (W_{\nu(k)}(t) f_{k,l})(r) Y_{k,l}(\sigma).$$

By the fact $-\Delta_\sigma Y_{k,l} = k(k+n-2)Y_{k,l}$, the orthonormality of spherical harmonics and the estimate (11), we get

$$\begin{aligned} \|U(\cdot)\varphi\|_{L^2 \tilde{H}_a^0 H_\sigma^{a-\frac{1}{2}}}^2 &\lesssim \sum_{k,l} (1 + \nu(k))^{2a-1} \|W_{\nu(k)}(\cdot) f_{k,l}\|_{L^2(\mathbb{R}_+^2)}^2 \\ &\lesssim \sum_{k,l} \|f_{k,l}\|_{L^2}^2 = \sum_{k,l} \int_0^\infty |G_{k,l}(\rho)|^2 \rho^{n-1} d\rho \\ &= \sum_{k,l} \int_0^\infty |g_{k,l}(r)|^2 r^{n-1} dr \\ &\sim \|\varphi\|_{L^2}^2. \end{aligned}$$

For the proof of the second inequality of (9) we introduce a lemma for low-diagonal operator estimate (see [1, 24]).

Lemma 1. *Let \mathcal{A} and \mathcal{B} be Banach spaces. Let K be an operator such that $\|KG\|_{L_T^q(\mathcal{A})} \leq C\|G\|_{L_T^p(\mathcal{B})}$ for $1 \leq p \leq q \leq \infty$ with kernel k defined by $KG(t) = \int_0^T k(t-t')G(t') dt'$, where C does not depend on T . If $p < q$, then the low-diagonal operator \tilde{K} defined by $\tilde{K}G = \int_0^t k(t-t')G(t') dt'$ satisfies that $\|\tilde{K}G\|_{L_T^q(\mathcal{A})} \leq \tilde{C}\|G\|_{L_T^p(\mathcal{B})}$, where \tilde{C} is C times a constant depending only on p, q .*

In view of Lemma 1 with kernel $k(t) = U(t)$, $\mathcal{A} = \tilde{H}_a^0 H_\sigma^{a-\frac{1}{2}}$ and $\mathcal{B} = L^2$, it suffices to show that

$$(14) \quad \left\| \int_0^T U(\cdot - t') F(t') dt' \right\|_{L_T^2(\tilde{H}_a^0 H_\sigma^{a-\frac{1}{2}})} \lesssim \|F\|_{L_T^1 L^2}.$$

In fact, by the first Strichartz estimate (9), we have

$$\begin{aligned} \left\| \int_0^T U(\cdot - t') F(t') dt' \right\|_{L_T^2(\tilde{H}_a^0 H_\sigma^{a-\frac{1}{2}})} &= \left\| U(\cdot) \int_0^T U(-t') F(t') dt' \right\|_{L_T^2(\tilde{H}_a^0 H_\sigma^{a-\frac{1}{2}})} \\ &\lesssim \|F\|_{L_T^1 L^2}. \end{aligned}$$

This yields the second inequality of (9). \square

Remark 3. From the proof of Proposition 1, one can see the identity

$$\|e^{-it\omega}\varphi\|_{L^2\tilde{H}_a^s}^2 = \frac{2\pi\Gamma(2a-1)\Gamma(\frac{n}{2}-a)}{2^{2a-1}\Gamma(a)^2\Gamma(\frac{n-2}{2}+a)}\|\varphi\|_{H^s}^2$$

for any radial function $\varphi \in H^s$ for some $s \geq 0$. Thus the weighted Strichartz estimate is sharp as far as $\frac{1}{2} < a < \frac{n}{2}$.

Remark 4. If $n = 1$, then a modified weighted Strichartz estimate is possible. To state precisely, we take any L^2 function w as a weight and define a weighted Sobolev space \tilde{H}_w^s as follows

$$\tilde{H}_w^s = \{v : \|v\|_{\tilde{H}_w^s} \equiv \|wL_{\frac{1}{2}}(\Delta)(1-\Delta)^{\frac{s}{2}}v\|_{L^2} < \infty\}.$$

Then for any $T > 0$ we have

$$\begin{aligned} \|U(\cdot)\varphi\|_{L_T^2\tilde{H}_w^s} &\lesssim \|\varphi\|_{H^s} \\ \left\| \int_0^\bullet U(\cdot-t')F(t') dt' \right\|_{L_T^2\tilde{H}_w^s} &\lesssim \|F\|_{L_T^1H^s}. \end{aligned}$$

To prove these estimates we have only to show that $\|W(\cdot)f\|_{L^2L^2} \lesssim \|f\|_{L^2}$, where

$$W(t)f(x) = w(x) \int_{-\infty}^{\infty} e^{i(x\xi-t\sqrt{1+\xi^2})} \frac{|\xi|^{\frac{1}{2}}}{(1+\xi^2)^{\frac{1}{4}}} f(\xi) d\xi.$$

By the change of variables $\xi \mapsto \sqrt{1+\xi^2}$ and by applying Plancherel theorem w.r.t. the time variable, and then using the change of variables $\xi \mapsto \sqrt{\xi^2-1}$ again, one can readily have that

$$\|W(\cdot)f\|_{L_T^2L^2} \leq \|W(\cdot)f\|_{L^2L^2} \leq \sqrt{2}\|w\|_{L^2}\|f\|_{L^2}.$$

The inhomogeneous Strichartz estimate can be treated by the same way as in the proof of Proposition 1.

3. FRACTIONAL INTEGRAL ESTIMATES FOR RADIAL FUNCTIONS

We prove some fractional integral estimates for radial functions.

Lemma 2. *Let $n \geq 3$ and $0 < \gamma < n-1$.*

(i) *If f and g are radial functions with $f, |x|^{-\delta}f, |x|^{-(\gamma-\delta)}g \in L^2$ for some $0 < \delta \leq \gamma$, then*

$$(15) \quad \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(y)||g(y)|}{|x-y|^\gamma} dy \lesssim \| |x|^{-\delta}f \|_{L^2} \| |x|^{-(\gamma-\delta)}g \|_{L^2}.$$

(ii) *If f, g are radially symmetric and $f, |x|^{-(\gamma-\delta)}g \in L^2$ for some $0 < \delta \leq \gamma$, then*

$$(16) \quad \sup_{x \in \mathbb{R}^n} |x|^\delta \int_{\mathbb{R}^n} \frac{|f(y)||g(y)|}{|x-y|^\gamma} dy \lesssim \|f\|_{L^2} \| |x|^{-(\gamma-\delta)}g \|_{L^2}.$$

Remark 5. In [8] Fröhlich and Lenzmann showed that if $\varphi \in C_0^\infty(\mathbb{R}^3)$ is radially symmetric and satisfies $E_1(\varphi) < 0$, then the radial solution u of (1) with $\gamma = 1$ blows up within a finite time. To lead to the blowup they used a variance type estimate where the estimates $|V_1(u)(x)| \leq \|\varphi\|_{L^2}^2/|x|$ and $|\nabla V_1(u)(x)| \leq \|\varphi\|_{L^2}^2/|x|^2$ are crucial. The lemma above leads us to the same estimates for $n \geq 4$ and hence to the finite time blowup.

Proof of Lemma 2. We revisit the proof of Lemma 3 of [5]. Fixing x , we divide the integration into three parts as follows

$$\int_{\mathbb{R}^n} \frac{|f(y)||g(y)|}{|x-y|^\gamma} dy = \int_{|y|>2|x|} + \int_{\frac{|x|}{2} \leq |y| \leq 2|x|} + \int_{|y|<\frac{|x|}{2}} \equiv I + II + III.$$

For I , since $|x-y| \geq \frac{|y|}{2}$ for $|y| > 2|x|$, we have

$$\begin{aligned} I &\lesssim \int_{|y|>2|x|} \frac{|f(y)|}{|y|^\delta} \frac{|g(y)|}{|y|^{\gamma-\delta}} dy \\ &\lesssim \| |\cdot|^{-\delta} f \|_{L^2} \| |x|^{-(\gamma-\delta)} g \|_{L^2} \quad \text{or} \quad |x|^{-\delta} \|f\|_{L^2} \| |x|^{-(\gamma-\delta)} g \|_{L^2}. \end{aligned}$$

Since f and g are radially symmetric, we may assume that $x = |x|e_1 = re_1 = r(1, 0, \dots, 0, 0)$. Using the spherical coordinates $(\rho, \theta_1, \theta_2, \dots, \theta_{n-1}) \in (0, \infty) \times [0, \pi] \times [0, \pi] \times \dots \times [0, 2\pi]$ for y variable, the integrals II and III are converted into

$$(17) \quad II + III = \left(\int_{\frac{r}{2}}^{2r} + \int_0^{\frac{r}{2}} \right) \rho^{n-1} |f(\rho)||g(\rho)| \Omega(r, \rho) d\rho,$$

where

$$\begin{aligned} \Omega(r, \rho) &= \int_0^{2\pi} \int_0^\pi \dots \int_0^\pi (r^2 + \rho^2 - 2r\rho \cos \theta_1)^{-\frac{\gamma}{2}} \\ &\quad \times \sin^{n-2} \theta_1 \sin^{n-3} \theta_2 \dots \sin \theta_{n-2} d\theta_1 \dots d\theta_{n-2} d\theta_{n-1}. \end{aligned}$$

If $\frac{r}{2} \leq \rho \leq 2r$, then by the fact $n-2-\gamma > -1$ and

$$\sqrt{r^2 + \rho^2 - 2r\rho \cos \theta_1} \geq \rho \sin \theta_1$$

we have

$$(18) \quad \Omega(r, \rho) \lesssim \rho^{-\gamma} \int_0^\pi \sin^{n-2-\gamma} \theta_1 d\theta_1 \lesssim \rho^{-\gamma}.$$

If $\rho < \frac{r}{2}$, then

$$(19) \quad \Omega(r, \rho) \lesssim r^{-\gamma},$$

since $r^2 + \rho^2 - 2r\rho \cos \theta_1 \geq r^2$. Therefore by the Hölder inequality we have

$$II + III \lesssim \| |\cdot|^{-\delta} f \|_{L^2} \| |\cdot|^{-(\gamma-\delta)} g \|_{L^2} \quad \text{or} \quad |x|^{-\delta} \|f\|_{L^2} \| |\cdot|^{-(\gamma-\delta)} g \|_{L^2}.$$

This completes the proof. \square

Lemma 3. *Let $1 < \gamma < 2$ if $n \geq 4$ and $1 < \gamma < \frac{3}{2}$ if $n = 3$. Let ε satisfy $0 < \varepsilon < \min(\gamma - 1, 2 - \gamma)$. Then for any radial functions $f, g \in L^2$ with $|x|^{-1}f, |x|^{-1}g \in L^2$*

$$(20) \quad \left\| | \cdot |^{-\gamma-\frac{1}{2}} * (fg) \right\|_{L^{2n}} \lesssim (\|f\|_{L^2}^{2-\gamma-\varepsilon} \| | \cdot |^{-1}f \|_{L^2}^{\gamma-1+\varepsilon} + \|f\|_{L^2}^{2-\gamma+\varepsilon} \| | \cdot |^{-1}f \|_{L^2}) \| | \cdot |^{-1}g \|_{L^2}.$$

Proof of Lemma 3. As in the proof of Lemma 2 we split the fractional integral and estimate pointwise as

$$\begin{aligned} \int \frac{|f(y)g(y)|}{|x-y|^{\gamma+\frac{1}{2}}} dy &= \int_{|x|>2|y|} + \int_{\frac{1}{2}|y|\leq|x|\leq 2|y|} + \int_{|x|<\frac{1}{2}|y|} \\ &= I + II + III. \end{aligned}$$

In case that $|x| < 1$, from similar estimates to (17), (18) and (19), and from the Hölder inequality and interpolation it follows that for small ε with $\gamma + \varepsilon < 2$

$$\begin{aligned} I &\lesssim |x|^{-\frac{1}{2}+\varepsilon} \int \frac{|f(y)g(y)|}{|y|^{\gamma+\varepsilon}} dy \lesssim |x|^{-\frac{1}{2}+\varepsilon} \|f\|_{L^2}^{2-\gamma-\varepsilon} \| | \cdot |^{-1}f \|_{L^2}^{\gamma-1+\varepsilon} \| | \cdot |^{-1}g \|_{L^2}, \\ II + III &\lesssim |x|^{-\frac{1}{2}+\varepsilon} \|f\|_{L^2}^{2-\gamma-\varepsilon} \| | \cdot |^{-1}f \|_{L^2}^{\gamma-1+\varepsilon} \| | \cdot |^{-1}g \|_{L^2} \int_0^\pi \sin^{n-\frac{5}{2}-\gamma} \theta d\theta. \end{aligned}$$

Since $n - \frac{5}{2} - \gamma > -1$ if $1 < \gamma < 2$ for $n \geq 4$ and $1 < \gamma < \frac{3}{2}$ for $n = 3$, the last integral is finite. Hence

$$\| | \cdot |^{-\gamma-\frac{1}{2}} * (fg) \|_{L^{2n}(|x|<1)} \lesssim \|f\|_{L^2}^{2-\gamma-\varepsilon} \| | \cdot |^{-1}f \|_{L^2}^{\gamma-1+\varepsilon} \| | \cdot |^{-1}g \|_{L^2}.$$

If $|x| \geq 1$, then choosing ε such that $\gamma - \varepsilon > 1$, by the same argument as above we have

$$I + II + III \lesssim |x|^{-\frac{1}{2}-\varepsilon} \|f\|_{L^2}^{2-\gamma+\varepsilon} \| | \cdot |^{-1}f \|_{L^2}^{\gamma-1-\varepsilon} \| | \cdot |^{-1}g \|_{L^2}.$$

Hence

$$\| | \cdot |^{-\gamma-\frac{1}{2}} * (fg) \|_{L^{2n}(|x|\geq 1)} \lesssim \|f\|_{L^2}^{2-\gamma+\varepsilon} \| | \cdot |^{-1}f \|_{L^2}^{\gamma-1-\varepsilon} \| | \cdot |^{-1}g \|_{L^2}.$$

The proof has been completed. \square

Lemma 4. *Let $n \geq 3$ and $1 < \gamma < n - 1$. Let $f, g \in W_{s,\varepsilon}$ be radial functions for some s, ε satisfying the condition (7). Then it follows that*

$$(21) \quad \left\| | \cdot |^{-\gamma} * (fg) \right\|_{L^{\frac{n}{1-(s-\varepsilon)}} \cap L^{\frac{n}{1-(s+\varepsilon)}}} \lesssim \|f\|_{W_{s,\varepsilon}} \|g\|_{W_{s,\varepsilon}}.$$

Proof of Lemma 4. By the same spirit as in the proof of Lemma 3, we split the fractional integral into three parts I, II, III and estimate them using radial symmetry. We also divide each part into two regions of x ; inside the unit ball and its outside

If $|x| < 1$, then since $\varepsilon < \frac{\gamma}{2} - s$, we have

$$I + II + III \lesssim |x|^{-(\gamma-2(s+\varepsilon))} \| | \cdot |^{-(s+\varepsilon)} f \|_{L^2} \| | \cdot |^{-(s+\varepsilon)} g \|_{L^2}.$$

Since $\| |\cdot|^{-(s+\varepsilon)} f \|_{L^2} \leq \|f\|_{W_{s,\varepsilon}}$ and $(\gamma - 2(s + \varepsilon)) \frac{n}{1-(s \pm \varepsilon)} < n$, we have

$$\| |\cdot|^{-\gamma} * (fg) \|_{(L^{\frac{n}{1-(s-\varepsilon)}} \cap L^{\frac{n}{1-(s+\varepsilon)}})_{(|x|<1)}} \lesssim \|f\|_{W_{s,\varepsilon}} \|g\|_{W_{s,\varepsilon}}.$$

If $|x| \geq 1$, then

$$I + II + III \lesssim |x|^{-(\gamma-2(s-\varepsilon))} \| |\cdot|^{-(s-\varepsilon)} f \|_{L^2} \| |\cdot|^{-(s-\varepsilon)} g \|_{L^2}.$$

Since $(\gamma - 2(s - \varepsilon)) \frac{n}{1-(s \pm \varepsilon)} > n$, we have

$$\| |\cdot|^{-\gamma} * (fg) \|_{(L^{\frac{n}{1-(s-\varepsilon)}} \cap L^{\frac{n}{1-(s+\varepsilon)}})_{(|x|\geq 1)}} \lesssim \|f\|_{W_{s,\varepsilon}} \|g\|_{W_{s,\varepsilon}}.$$

This completes the proof of the lemma. \square

4. PROOFS OF THE THEOREMS

4.1. Proof of Theorem 1. We only consider the positive time because the proof for negative time can be treated in the same way.

Let us first define a complete metric space $X_{T,\rho}$ with metric $d(u, v) = \|u - v\|_{X_T}$, where $X_T = C([0, T]; H^{\frac{1}{2}}) \cap L_T^2 \tilde{H}_1^{\frac{1}{2}}$ by

$$X_{T,\rho} \equiv \{v \in X_T : v \text{ is radially symmetric and } \|v\|_{X_T} \leq \rho\}.$$

Here let us observe that the space $\tilde{H}_1^{\frac{1}{2}}$ is exactly the same as $\{v : \| |\cdot|^{-1} v \|_{L^2} < \infty\}$.

Now we define a mapping $N : u \mapsto N(u)$ on $X_{T,\rho}$ by

$$(22) \quad N(u)(t) = U(t)\varphi - i \int_0^t U(t-t')F(u)(t') dt'.$$

For any $u \in X_{T,\rho}$, $N(u)$ is radially symmetric. By the Strichartz estimate (9) with $a = 1, b = 0$ and $s = \frac{1}{2}$, we have

$$\|N(u)\|_{X_{T,\rho}} \lesssim \|\varphi\|_{H^{\frac{1}{2}}} + \|F\|_{L^1 H^{\frac{1}{2}}}.$$

For the second term, we use the generalized Leibniz rule (see Lemma A1 \sim Lemma A4 in Appendix of [12]) such that for any $s \geq 0$

$$(23) \quad \|D^s(uv)\|_{L^r} \lesssim \|D^s u\|_{L^{r_1}} \|v\|_{L^{q_2}} + \|u\|_{L^{q_1}} \|D^s v\|_{L^{r_2}},$$

where $D^s = (-\Delta)^{s/2}$

$$\text{and } \frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2} = \frac{1}{q_1} + \frac{1}{q_2}, \quad r_i \in (1, \infty), \quad q_i \in (1, \infty], \quad i = 1, 2.$$

From (23), we have

$$(24) \quad \begin{aligned} & \|N(u)\|_{X_{T,\rho}} \\ & \lesssim \|\varphi\|_{H^{\frac{1}{2}}} + \|V_\gamma(u)\|_{L_T^1 L^\infty} \|u\|_{L_T^\infty H^{\frac{1}{2}}} + \|(-\Delta)^{\frac{1}{4}} V_\gamma(u)\|_{L_T^1 L^{2n}} \|u\|_{L_T^\infty L^{\frac{2n}{n-1}}}. \end{aligned}$$

To estimate the last two terms, we use Lemma 2. Using (15) with $f = g = u$ and interpolation we have

$$\|V_\gamma(u)\|_{L^\infty} \lesssim \|u\|_{L^2}^{2-\gamma} \|u\|_{\tilde{H}_1^{\frac{1}{2}}}^\gamma.$$

and using (21) with $f = g = u$ for some small ε as in Lemma 3

$$\begin{aligned} \|(-\Delta)^{\frac{1}{4}} V_\gamma(u)\|_{L^{2n}} &\lesssim \|V_{\gamma+\frac{1}{2}}(u)\|_{L^{2n}} \\ &\lesssim \|u\|_{L^2}^{2-\gamma-\varepsilon} \|u\|_{\tilde{H}_T^{\frac{1}{2}}}^{\gamma+\varepsilon} + \|u\|_{L^2}^{2-\gamma+\varepsilon} \|u\|_{\tilde{H}_T^{\frac{1}{2}}}^{\gamma-\varepsilon}. \end{aligned}$$

Hence the nonlinear estimate (24) has the following: for some positive number ε

$$\begin{aligned} \|N(u)\|_{X_{T,\rho}} &\lesssim \|\varphi\|_{H^{\frac{1}{2}}} + (T^{\frac{2}{2-\gamma}} + T^{\frac{2}{2-\gamma-\varepsilon}} + T^{\frac{2}{2-\gamma+\varepsilon}}) \|u\|_{L_T^\infty H^{\frac{1}{2}}}^2 \|u\|_{L_T^2 \tilde{H}_1^{\frac{1}{2}}} \\ &\lesssim \|\varphi\|_{H^{\frac{1}{2}}} + (T^{\frac{2}{2-\gamma}} + T^{\frac{2}{2-\gamma-\varepsilon}} + T^{\frac{2}{2-\gamma+\varepsilon}}) \rho^3. \end{aligned}$$

From the choice of T and ρ satisfying that

$$\begin{aligned} C\|\varphi\|_{H^1} &\leq \frac{\rho}{2}, \\ C(T^{\frac{2}{2-\gamma}} + T^{\frac{2}{2-\gamma-\varepsilon}} + T^{\frac{2}{2-\gamma+\varepsilon}}) \rho^3 &\leq \frac{\rho}{2} \end{aligned}$$

for some constant C , it follows that N maps $X_{T,\rho}$ to itself.

For any $u, v \in X_{T,\rho}$, we have

$$\begin{aligned} d(N(u), N(v)) &\lesssim \|F(u) - F(v)\|_{L_T^1 H^{\frac{1}{2}}} \\ &\lesssim \|(V_\gamma(u) - V_\gamma(v))u\|_{L_T^1 H^{\frac{1}{2}}} + \|V_\gamma(v)(u - v)\|_{L_T^1 H^{\frac{1}{2}}}. \end{aligned}$$

Using Lemma 2, Lemma 3 and the Leibniz rule (23) again, we have that

$$\begin{aligned} &\|(V_\gamma(u) - V_\gamma(v))u\|_{L_T^1 H^{\frac{1}{2}}} \\ &\lesssim \|(V_\gamma(u) - V_\gamma(v))u\|_{L_T^1 L^2} + \|(-\Delta)^{\frac{1}{4}}((V_\gamma(u) - V_\gamma(v))u)\|_{L_T^1 L^2} \\ &\lesssim \left(\|V_\gamma(u) - V_\gamma(v)\|_{L_T^1 L^\infty} + \|(-\Delta)^{\frac{1}{4}}(V_\gamma(u) - V_\gamma(v))\|_{L_T^1 L^{2n}} \right) \|u\|_{L_T^\infty H^{\frac{1}{2}}} \\ &\lesssim \int_0^T (\|u\|_{L^2} + \|v\|_{L^2})^{2-\gamma} \|u - v\|_{\tilde{H}_1^{\frac{1}{2}}}^\gamma dt \|u\|_{L_T^\infty H^{\frac{1}{2}}} \\ &\quad + \| |\cdot|^{-(\gamma+\frac{1}{2})} * (|u|^2 - |v|^2) \|_{L_T^1 L^{2n}} \|u\|_{L_T^\infty H^{\frac{1}{2}}} \\ &\lesssim (T^{\frac{2}{2-\gamma}} + T^{\frac{2}{2-(\gamma+\varepsilon)}} + T^{\frac{2}{2-(\gamma-\varepsilon)}}) \rho^2 \|u - v\|_{L_T^2 \tilde{H}_1^{\frac{1}{2}}} \end{aligned}$$

and

$$\begin{aligned} &\|V_\gamma(v)(u - v)\|_{L_T^1 H^{\frac{1}{2}}} \\ &\lesssim \|V_\gamma(v)\|_{L_T^1 L^\infty} \|u - v\|_{L_T^\infty H^{\frac{1}{2}}} + \|(-\Delta)^{\frac{1}{4}} V_\gamma(v)\|_{L_T^1 L^{2n}} \|u\|_{L_T^\infty H^{\frac{1}{2}}} \\ &\lesssim (T^{\frac{2}{2-\gamma}} + T^{\frac{2}{2-(\gamma+\varepsilon)}} + T^{\frac{2}{2-(\gamma-\varepsilon)}}) \rho^2 \|u - v\|_{L_T^\infty H^{\frac{1}{2}}}. \end{aligned}$$

Thus by the choice of T, ρ as above

$$(25) \quad d(N(u), N(v)) \leq C(T^{\frac{2}{2-\gamma}} + T^{\frac{2}{2-\gamma-\varepsilon}} + T^{\frac{2}{2-\gamma+\varepsilon}}) \rho^2 d(u, v) \leq \frac{1}{2} d(u, v).$$

Therefore N becomes a contraction mapping on $X_{T,\rho}$. This proves the local existence. The energy and L^2 conservations follow from the Strichartz estimate (9) and the method of [22].

Now we consider the global well-posedness. To do so, we need the following energy inequality that for $\lambda > 0$,

$$\frac{1}{2} \|u(t)\|_{\dot{H}^{\frac{1}{2}}}^2 \leq E(u) = E(\varphi).$$

If $\lambda < 0$, then for any φ with $\|\varphi\|_{L^2} \leq 1$

$$\begin{aligned} \|u(t)\|_{\dot{H}^{\frac{1}{2}}}^2 &\leq 2|E(u)| + 2|V(u)| \\ &\leq 2|E(\varphi)| + C\|u\|_{L^{\frac{2n}{n-\gamma+1}}}^2 \|u\|_{\dot{H}^{\frac{1}{2}}}^2 \\ &\leq C(1 + \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^2)^\gamma + C\|\varphi\|_{L^2}^{4-2\gamma} \|u\|_{\dot{H}^{\frac{1}{2}}}^{2\gamma}. \end{aligned}$$

Here for the third inequality we used the fact that

$$2|E(\varphi)| \leq 1 + \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^2 + C\|\varphi\|_{L^2}^{4-2\gamma} \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^{2\gamma} \leq C(1 + \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^2)^\gamma.$$

Hence especially in the case of negative λ , with assumption of the smallness of $\|\varphi\|_{L^2}$ for $\gamma < 2$ such that

$$\|\varphi\|_{L^2} \leq \min \left(1, (8^\gamma C^\gamma (1 + \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^2)^{\gamma(\gamma-1)})^{-\frac{1}{4-2\gamma}} \right),$$

we have

$$(26) \quad \|u(t)\|_{\dot{H}^{\frac{1}{2}}}^2 \leq 2C(1 + \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^2)^\gamma.$$

Now let us denote $E(\varphi)$ for $\lambda < 0$ and $(1 + \|\varphi\|_{\dot{H}^{\frac{1}{2}}}^2)^\gamma$ for $\lambda < 0$ by $\mathcal{E}(\varphi)$. Then from the Strichartz estimate (9), (26) and Lemma 2, 3, we have for some small time $0 < \delta < 1$ and small $\varepsilon > 0$

$$\begin{aligned} \|u\|_{L_\delta^2 \tilde{H}_{1,0}^{\frac{1}{2}}} &\lesssim (1 + \mathcal{E}(\varphi))^{\frac{1}{2}} + \delta^{\frac{2}{2-\gamma-\varepsilon}} (1 + \|u\|_{L_\delta^\infty \dot{H}^{\frac{1}{2}}})^{3-\gamma+\varepsilon} \|u\|_{L_\delta^2 \tilde{H}_1^{\frac{1}{2}}} \\ &\lesssim (1 + \mathcal{E}(\varphi))^{\frac{1}{2}} + \delta^{\frac{2}{2-\gamma-\varepsilon}} (1 + \mathcal{E}(\varphi))^{\frac{3-\gamma+\varepsilon}{2}} \|u\|_{L_\delta^2 \tilde{H}_1^{\frac{1}{2}}}. \end{aligned}$$

Thus for some δ so small that $\frac{1}{4} \leq \delta^{\frac{2}{2-\gamma-\varepsilon}} (1 + \mathcal{E}(\varphi))^{\frac{3-\gamma+\varepsilon}{2}} \leq \frac{1}{2}$, we have

$$\|u\|_{L^2(T_{j-1}, T_j; \tilde{H}_1^{\frac{1}{2}})} \leq C(1 + \mathcal{E}(\varphi))^{\frac{1}{2}},$$

where $T_j - T_{j-1} = \delta$ for $j \leq k-1$, $T_k = T$ and $T_k - T_{k-1} \sim \delta$. This implies that

$$\begin{aligned} (27) \quad \|u\|_{L_T^2 \tilde{H}_1^{\frac{1}{2}}}^2 &\leq \sum_{1 \leq j \leq k} \|u\|_{L^2(T_{j-1}, T_j; \tilde{H}_1^{\frac{1}{2}})}^2 \\ &\lesssim k\delta(1 + \mathcal{E}(\varphi))^{1 + \frac{(3-\gamma+\varepsilon)(2-\gamma-\varepsilon)}{4}} \\ &\lesssim T(1 + \mathcal{E}(\varphi))^{1 + \frac{(3-\gamma+\varepsilon)(2-\gamma-\varepsilon)}{4}}. \end{aligned}$$

From (26) and (27) we conclude that $u \in C_b(\mathbb{R}_+; \dot{H}^{\frac{1}{2}}) \cap L_{loc}^2 \tilde{H}_1^{\frac{1}{2}}$. This completes the proof of Theorem 1.

4.2. Proof of Theorem 2. Let Y_ρ be a complete metric space with metric $d(u, v) = \|u - v\|_Y$, where $Y = C_b(\mathbb{R}; H^1) \cap L^2 \tilde{H}_1^{\frac{1}{2}}$ by

$$Y_\rho \equiv \{v \in Y : v \text{ is radially symmetric and } \|v\|_Y \leq \rho\}.$$

Then we claim that the map N defined as (22) is a contraction on Y , provided ρ is sufficiently small.

From the Strichartz estimate (9) and the fractional integral estimates (15) and (16), we have for any $u \in Y_\rho$

$$\begin{aligned} \|u\|_Y &\lesssim \|\varphi\|_{H^1} + \|V_2(u)\|_{L^1 L^\infty} \|u\|_{L^\infty H^1} + \|(\nabla V_2(u))u\|_{L^1 L^2} \\ &\lesssim \|\varphi\|_{H^1} + \|u\|_{L^2 \tilde{H}_1^{\frac{1}{2}}}^2 \|u\|_{L^\infty H^1} \\ &\lesssim \|\varphi\|_{H^1} + \rho^3. \end{aligned}$$

Hence choosing ρ so small that $C\|\varphi\|_{H^1} \leq \frac{\rho}{2}$ and $C\rho^3 \leq \frac{\rho}{2}$, the mapping N maps Y to itself. We also have

$$\begin{aligned} d(N(u), N(v)) &\lesssim \|V_2(u)u - V_2(v)v\|_{L^1 H^1} \\ &\lesssim \|(V_2(u) - V_2(v))u\|_{L^1 H^1} + \|V_2(v)(u - v)\|_{L^1 H^1}. \end{aligned}$$

Using Lemma 2, we have

$$\begin{aligned} &\|(V_2(u) - V_2(v))u\|_{L^1 H^1} \\ &\lesssim \|(V_2(u) - V_2(v))\|_{L^1 L^\infty} \|u - v\|_{L^\infty H^1} + \|\nabla((V_2(u) - V_2(v))u)\|_{L^1 L^2} \\ &\lesssim (\|u\|_{L^2 \tilde{H}_1^{\frac{1}{2}}} + \|v\|_{L^2 \tilde{H}_1^{\frac{1}{2}}}) \|u - v\|_{L^2 \tilde{H}_1^{\frac{1}{2}}} \|u\|_{L^\infty H^1} \\ &\quad + (\|u\|_{L^\infty H^1} + \|v\|_{L^\infty H^1}) \|u - v\|_{L^2 \tilde{H}_1^{\frac{1}{2}}} \|u\|_{L^2 \tilde{H}_1^{\frac{1}{2}}} \end{aligned}$$

and

$$\begin{aligned} &\|V_2(v)(u - v)\|_{L^1 H^1} \\ &\lesssim \|V_2(v)\|_{L^1 L^\infty} \|u - v\|_{L^1 H^1} + \|\nabla(V_2(v)(u - v))\|_{L^1 L^2} \\ &\lesssim \|v\|_{L^2 \tilde{H}_1^{\frac{1}{2}}}^2 \|u - v\|_{L^\infty H^1} + \|v\|_{L^\infty H^1} \|v\|_{L^2 \tilde{H}_1^{\frac{1}{2}}} \|u - v\|_{L^2 \tilde{H}_1^{\frac{1}{2}}}. \end{aligned}$$

Hence from the condition of u and v

$$d(N(u), N(v)) \lesssim \rho^2 d(u, v).$$

Thus by the choice of ρ such that $C\rho^2 \leq \frac{1}{2}$, N becomes a contraction.

As for the scattering, let us define functions φ_\pm by

$$\varphi_\pm = \varphi - i \int_0^{\pm\infty} U(-s)F(u)(s) ds.$$

Then clearly $\varphi_\pm \in H^1$ and one can show that

$$\begin{aligned} \|u(t) - U(t)\varphi_\pm\|_{H^1} &\lesssim \|u\|_{L^2(I_t^\pm; \tilde{H}_1^{\frac{1}{2}})}^2 \|u\|_{L^\infty H^1} \\ &\rightarrow 0 \quad \text{as } t \rightarrow \pm\infty, \end{aligned}$$

where $I_t^+ = (t, \infty)$ and $I_t^- = (-\infty, t)$. This proves Theorem 2.

4.3. Proof of Theorem 3. Let Z_ρ be a complete metric space with metric $d(u, v) = \|u - v\|_Z$, where $Z = C_b(\mathbb{R}; H^s) \cap L^2 W_{s,\varepsilon}$ by

$$Z_\rho \equiv \{v \in Y : v \text{ is radially symmetric and } \|v\|_Z \leq \rho\}.$$

Here from the definition we easily observe that the space $W_{s,\varepsilon}$ contains the space

$$L_{s\pm\varepsilon}(\Delta)\tilde{H}_{s\pm\varepsilon}^0 = \{v : (L_{s\pm\varepsilon}(\Delta))^{-1}v \in \tilde{H}_{s\pm\varepsilon}^0\}.$$

Now we define a nonlinear map \tilde{N} by

$$\tilde{N}(u) = (\cos tw)\varphi_1 + \omega^{-1}(\sin tw)\varphi_2 - \int_0^t \omega^{-1}(\sin(t-t')\omega)F(u) dt'.$$

Then we claim that the map \tilde{N} is a contraction on Z , provided that ρ is sufficiently small.

We first observe from the Strichartz estimate (9)² and fractional integration that

$$\begin{aligned} \|(L_{s\pm\varepsilon}(\Delta))^{-1}(\cos(\cdot)\omega)\varphi_1\|_{L^2\tilde{H}_{s\pm\varepsilon}^0} &\lesssim \|(L_{s\pm\varepsilon}(\Delta))^{-1}\varphi_1\|_{L^2} \\ &\lesssim \|(-\Delta)^{-\frac{1-(s\pm\varepsilon)}{2}}\varphi_1\|_{L^2} + \|(-\Delta)^{\frac{s-\frac{1}{2}+\varepsilon}{2}}\varphi_1\|_{L^2} \\ &\lesssim \|\varphi_1\|_{L^{\frac{2n}{n+2-2(s-\varepsilon)}} \cap H^{s+\varepsilon-\frac{1}{2}}} \end{aligned}$$

and

$$\begin{aligned} \|(L_{s\pm\varepsilon}(\Delta))^{-1}\omega^{-1}(\sin(\cdot)\omega)\varphi_2\|_{L^2\tilde{H}_{s\pm\varepsilon}^0} &\lesssim \|(L_{s\pm\varepsilon}(\Delta))^{-1}\omega^{-1}\varphi_2\|_{L^2} \\ &\lesssim \|\varphi_2\|_{L^{\frac{2n}{n+2-2(s-\varepsilon)}} \cap H^{s+\varepsilon-\frac{3}{2}}} \end{aligned}$$

Since

$$\|\psi\|_{W_{s,\varepsilon}} \lesssim \|\psi\|_{L_{s+\varepsilon}(\Delta)\tilde{H}_{s+\varepsilon}^0} + \|\psi\|_{L_{s-\varepsilon}(\Delta)\tilde{H}_{s-\varepsilon}^0},$$

we have

$$\|(\cos tw)\varphi_1 + \omega^{-1}(\sin tw)\varphi_2\|_{L^\infty H^s \cap L^2 W_{s,\varepsilon}} \lesssim \|\varphi_1\|_{D_{s+\varepsilon,s+\varepsilon}} + \|\varphi_2\|_{D_{s+\varepsilon-1,s+\varepsilon}}.$$

Now we estimate the nonlinear part. From the Strichartz estimate (9) and the boundedness of $(-\Delta)^{\frac{1-(s\pm\varepsilon)}{2}}(1-\Delta)^{-\frac{3-(s+\varepsilon)}{2}}$ in L^2 , it follows that

$$\begin{aligned} &\left\| \int_0^t \omega^{-1}(\sin(t-t')\omega)F(u) dt' \right\|_{L^\infty H^{s-\frac{1}{2}+\varepsilon} \cap L^2 W_{s,\varepsilon}} \\ (28) \quad &\lesssim \|(-\Delta)^{-\frac{1-(s+\varepsilon)}{2}}F(u)\|_{L^1 L^2} + \|(-\Delta)^{-\frac{1-(s-\varepsilon)}{2}}F(u)\|_{L^1 L^2} \\ &\lesssim \|F(u)\|_{L^1 L^{\frac{2n}{n+2-(s+\varepsilon)}}} + \|F(u)\|_{L^1 L^{\frac{2n}{n+2-(s-\varepsilon)}}} \\ &\lesssim (\|V_\gamma(u)\|_{L^1 L^{\frac{n}{1-(s+\varepsilon)}}} + \|V_\gamma(u)\|_{L^1 L^{\frac{n}{1-(s-\varepsilon)}}}) \|u\|_{L^\infty L^2}. \end{aligned}$$

Using Lemma 4 with $f = g = u$, the last term on the RHS of (28) is bounded by a constant multiple of

$$(29) \quad \|u\|_{L^2 W_{s,\varepsilon}}^2 \|u\|_{L^\infty L^2}.$$

Therefore, for any $u \in Z_\rho$ we have

$$\|\tilde{N}(u)\|_Z \leq C(\|\varphi_1\|_{D_{s+\varepsilon,s+\varepsilon}} + \|\varphi_2\|_{D_{s+\varepsilon-1,s+\varepsilon}}) + C\rho^3$$

²To apply (9) we need the condition $s - \varepsilon > \frac{1}{2}$.

and this implies that \tilde{N} maps from Z_ρ to itself, provided ρ and the norm of the initial data are sufficiently small.

On the other hand, by the Strichartz estimate (9) and fractional integral estimate we have that

$$\begin{aligned}
& d(\tilde{N}(u), \tilde{N}(v)) \\
& \lesssim \|(-\Delta)^{\frac{1-(s+\varepsilon)}{2}}(V_\gamma(u)u - V_\gamma(v)v)\|_{L^1L^2} + \|(-\Delta)^{\frac{1-(s-\varepsilon)}{2}}(V_\gamma(u)u - V_\gamma(v)v)\|_{L^1L^2} \\
& \lesssim \|(-\Delta)^{\frac{1-(s+\varepsilon)}{2}}((V_\gamma(u) - V_\gamma(v))u)\|_{L^1L^2} + \|(-\Delta)^{\frac{1-(s-\varepsilon)}{2}}((V_\gamma(u) - V_\gamma(v))u)\|_{L^1L^2} \\
& \quad + \|(-\Delta)^{\frac{1-(s+\varepsilon)}{2}}V_\gamma(v)(u-v)\|_{L^1L^2} + \|(-\Delta)^{\frac{1-(s-\varepsilon)}{2}}V_\gamma(v)(u-v)\|_{L^1L^2} \\
& \lesssim \|(V_\gamma(u) - V_\gamma(v))u\|_{L^1L^{\frac{2n}{n+2-\frac{2n}{s+\varepsilon}}}} + \|(V_\gamma(u) - V_\gamma(v))u\|_{L^1L^{\frac{2n}{n+2-\frac{2n}{s-\varepsilon}}}} \\
& \quad + \|V_\gamma(v)(u-v)\|_{L^1L^{\frac{2n}{n+2-\frac{2n}{s+\varepsilon}}}} + \|V_\gamma(v)(u-v)\|_{L^1L^{\frac{2n}{n+2-\frac{2n}{s-\varepsilon}}}}.
\end{aligned}$$

Applying Lemma 4 with $f = u - v, g = \bar{u}$ or $f = v, g = \overline{u-v}$ or $f = g = v$ to the last four terms in the above estimate, we have

$$\begin{aligned}
& d(\tilde{N}(u), \tilde{N}(v)) \\
& \lesssim (\|u\|_{L^2W_{s,\varepsilon}} + \|v\|_{L^2W_{s,\varepsilon}})\|u-v\|_{L^2W_{s,\varepsilon}}\|u\|_{L^\infty L^2} + \|v\|_{L^2W_{s,\varepsilon}}^2\|u-v\|_{L^\infty L^2} \\
& \lesssim \rho^2 d(u, v).
\end{aligned}$$

Hence the smallness of ρ and of the norms of initial data makes \tilde{N} a contraction.

Now we consider the scattering. Let us define four functions $\varphi_i^\pm, i = 1, 2$ by

$$\begin{aligned}
\widehat{\varphi}_1^\pm(\xi) &= \widehat{\varphi}_1(\xi) + \int_0^{\pm\infty} (\sqrt{1+|\xi|^2})^{-1} \sin(t'\sqrt{1+|\xi|^2}) \widehat{F}(u)(\xi, t') dt' \\
\widehat{\varphi}_2^\pm(\xi) &= \widehat{\varphi}_2(\xi) - \int_0^{\pm\infty} \cos(t'\sqrt{1+|\xi|^2}) \widehat{F}(u)(\xi, t') dt'.
\end{aligned}$$

Then it follows from the regularity of the solution u that $\varphi_1^\pm \in H^{s-\frac{1}{2}+\varepsilon}$ and $\varphi_2^\pm \in H^{s-\frac{3}{2}+\varepsilon}$. Furthermore, since $u \in L^2W_{s,\varepsilon}$, for the linear solution u^\pm of (8) we conclude from the estimate (28) and (29) that

$$\begin{aligned}
& \|u(t) - u^\pm(t)\|_{H^s} \\
& \lesssim \|(-\Delta)^{-\frac{1-(s+\varepsilon)}{2}}F(u)\|_{L^1(I_t^\pm; L^2)} + \|(-\Delta)^{\frac{1-(s-\varepsilon)}{2}}F(u)\|_{L^1(I_t^\pm; L^2)} \\
& \lesssim (\|V_\gamma(u)\|_{L^1(I_t^\pm; L^{\frac{n}{1-(s+\varepsilon)}})} + \|V_\gamma(u)\|_{L^1(I_t^\pm; L^{\frac{n}{1-(s-\varepsilon)}})})\|u\|_{L^\infty L^2} \\
& \lesssim \|u\|_{L^2(I_t^\pm; W_{s,\varepsilon})}^2\|u\|_{L^\infty L^2} \rightarrow 0 \quad \text{as } t \rightarrow \pm\infty.
\end{aligned}$$

This completes the proof of the theorem.

Remark 6. If the initial data $(\varphi_1, \varphi_2) \in \tilde{D}_{s+\varepsilon, s-\varepsilon} \times \tilde{D}_{s+\varepsilon-1, s-\varepsilon}$ and their norm is sufficiently small, then the solution u is in $C_b(\mathbb{R}; \dot{H}^{-(1-(s-\varepsilon))})$. In fact, the existence and uniqueness in $C_b(\mathbb{R}; H^{s-\frac{1}{2}+\varepsilon}) \cap L^2W_{s,\varepsilon}$ follows immediately from the previous proof. Hence we have only to show $\|u\|_{L^\infty \dot{H}^{-(1-(s-\varepsilon))}} < \infty$. From (28) and (29) we

have

$$\begin{aligned}
& \|u\|_{L^\infty \dot{H}^{-(1-(s-\varepsilon))}} \\
& \leq \|\varphi_1\|_{\dot{H}^{-(1-(s-\varepsilon))}} + \|\varphi_2\|_{\dot{H}^{-(1-(s-\varepsilon))}} + \int_0^t \|(-\Delta)^{-\frac{1-(s-\varepsilon)}{2}} F(u)\|_{L^2} dt' \\
& \lesssim \|\varphi_1\|_{\dot{H}^{-(1-(s-\varepsilon))}} + \|\varphi_2\|_{\dot{H}^{-(1-(s-\varepsilon))}} + \|V_\gamma(u)\|_{L^1 L^{\frac{n}{1-(s-\varepsilon)}}} \|u\|_{L^\infty L^2} \\
& \lesssim \|\varphi_1\|_{\dot{H}^{-(1-(s-\varepsilon))}} + \|\varphi_2\|_{\dot{H}^{-(1-(s-\varepsilon))}} + \|u\|_{L^2 W_{s,\varepsilon}}^2 \|u\|_{L^\infty L^2} < \infty.
\end{aligned}$$

Let u^- be the radial solution of the linear equation (8) with radially symmetric initial data $(\varphi_1^-, \varphi_2^-) \in \tilde{D}_{s+\varepsilon, s-\varepsilon} \times \tilde{D}_{s+\varepsilon-1, s-\varepsilon}$. Then from the proof as above we can find a unique solution $u \in C_b(\mathbb{R}; H^{s+\varepsilon-\frac{1}{2}} \cap \dot{H}^{-(1-(s-\varepsilon))}) \cap L^2 W_{s,\varepsilon}$ satisfying

$$u(t) = u^-(t) + \int_t^{-\infty} \omega^{-1} \sin((t-t')\omega) F(u) dt',$$

provided that $\|\varphi_1^-\|_{\tilde{D}_{s+\varepsilon, s-\varepsilon}} + \|\varphi_2^-\|_{\tilde{D}_{s+\varepsilon-1, s-\varepsilon}}$ is sufficiently small. Here the solution u satisfies (1) with initial data $\varphi \in \tilde{D}_{s+\varepsilon, s-\varepsilon}$ such that

$$\varphi = u(0) = \varphi_1^- - \int_0^{-\infty} \omega^{-1} \sin(t'\omega) F(u) dt'.$$

Now in turn there are radial functions φ_1^+ and φ_2^+ as in Theorem 3. Actually, they are uniquely determined under a smallness condition of initial data. Hence we conclude that there exists a scattering operator S maps $(\varphi_1^-, \varphi_2^-)$ in a small neighborhood of $\tilde{D}_{s+\varepsilon, s-\varepsilon} \times \tilde{D}_{s+\varepsilon-1, s-\varepsilon}$ to $(\varphi_1^+, \varphi_2^+)$ in a small neighborhood of $\tilde{D}_{s+\varepsilon, s-\varepsilon} \times \tilde{D}_{s+\varepsilon-1, s-\varepsilon}$ and that S is injective.

REFERENCES

- [1] C. Ahn and Y. Cho, *Lorentz space extension of Strichartz estimate*, Proc. Amer. Math. Soc. **133** (2005), 3497-3503.
- [2] G. B. Arfken and H. J. Weber, *Mathematical Methods for Physicists*, Academic Press, 1995.
- [3] N. Burq, F. Planchon, J.G. Stalker and A.S. Tahvildar-Zadeh, *Strichartz estimate for the wave and Schrödinger equations with the inverse square potential*, J. Func. Anal., **203** (2003), 519-549.
- [4] Y. Cho and T. Ozawa, *On the semi-relativistic Hartree type equation*, SIAM J. Math. Anal., **38** (2006), 1060-1074.
- [5] Y. Cho and T. Ozawa, *On radial solutions of semi-relativistic Hartree equations*, Hokkaido University Preprint Series in Math #792.
- [6] A. Elgart and B. Schlein, *Mean field dynamics of boson stars*, to appear in Comm. Pure Appl. Math., arXiv:math-ph/0504051.
- [7] J. Fröhlich and E. Lenzmann, *Mean-field limit of quantum bose gases and nonlinear Hartree equation*, Sémin. Equ. Dériv., Partielles XIX (2003-2004), 1-26, arXiv:math-ph/0409019.
- [8] J. Fröhlich and E. Lenzmann, *Blow-up for nonlinear wave equations describing Boson stars*, to appear in Comm. Pure Appl. Math., arXiv:math-ph/0511003.

- [9] J. Fröhlich, B. L. G. Jonsson and E. Lenzmann, *Boson stars as solitary waves*, to appear in Comm. Pure Appl. Math., arXiv:math-ph/0512040.
- [10] E. Gross, *Quantum theory of interacting bosons*, Ann. Physics, **9** (1960), 292-324.
- [11] J. Kato, M. Nakamura and T. Ozawa, *A generalization of the weighted Strichartz estimates for wave equations and an application to self-similar solutions*, Comm. Pure Appl. Math., **60** (2007), 164-186.
- [12] T. Kato, *On nonlinear Schrödinger equations II. H^s -solutions and unconditional well-posedness*, J. Anal. Math., **67** (1995), 281-306.
- [13] E. H. Lieb and H.-T. Yau, *The Chandrasekhar theory of stellar collapse as the limit of quantum mechanics*, Commun. Math. Phys., **112** (1987), 147-174.
- [14] P.-L. Lions, *The Choquard equation and related questions*, Nonlinear Anal., TMA, **4** (1980), 1073-1079.
- [15] E. Lenzmann, *Well-posedness for semi-relativistic Hartree equations of critical type*, to appear in Mathematical Physics, Analysis and Geometry; arXiv:math.AP/0505456.
- [16] S. Machihara, M. Nakamura, K. Nakanashi and T. Ozawa, *Endpoint Strichartz estimates and global solutions for the nonlinear Dirac equation*, J. Func. Anal., **219** (2005), 1-20.
- [17] G. Menzala, *On regular solutions of a nonlinear equation of Choquard's type*, Proc. Royal Soc. Edinburgh Sect. A, **86** (1980), 291-301.
- [18] G. Menzala and W. Strauss, *On a wave equation with a cubic convolution*, J. Differential Equations, **43** (1982), 93-105.
- [19] K. Mochizuki, *On small data scattering with cubic convolution nonlinearity*, J. Math. Soc. Japan, **41** (1989), 143-160.
- [20] K. Mochizuki and T. Motai, *On small data scattering for some nonlinear wave equations*, in "Patterns and Waves -qualitative analysis of nonlinear differential equations -," 543-560, Stud. Math. Appl., 18, North-Holland, Amsterdam, 1986.
- [21] T. Motai, *On the Cauchy problem for the nonlinear Klein-Gordon equation with a cubic convolution*, Tsukuba J. Math., **12** (1988), 353-369.
- [22] T. Ozawa, *Remarks on proofs of conservation laws for nonlinear Schrödinger equations*, Cal. Var. PDE., **25** (2006), 403-408.
- [23] H. Sasaki, *Small data scattering for the Klein-Gordon equation with cubic convolution nonlinearity*, Discrete Contin. Dynam. Systems, **15** (2006), 973-981.
- [24] H. Smith and C. D. Sogge, *Global Strichartz estimates for nontrapping perturbations of the Laplacian*, Comm. Partial Differential Equations, **25** (2000), 2171-2183.
- [25] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, 1971.
- [26] W. Strauss, *Nonlinear scattering theory at low energy: sequel*, J. Funct. Anal., **43** (1981), 281-293.
- [27] G. Watson, *A Treatise on the Theory of Bessel Functions*, Reprint of the second (1944) edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1995.

DEPARTMENT OF MATHEMATICS, POSTECH, POHANG 790-784, REPUBLIC OF KOREA
E-mail address: `changocho@gmail.com`

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN
E-mail address: `ozawa@math.sci.hokudai.ac.jp`

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN
E-mail address: `hisasaki@math.sci.hokudai.ac.jp`

DEPARTMENT OF MATHEMATICS, POSTECH, POHANG 790-784, REPUBLIC OF KOREA
E-mail address: `shim@postech.ac.kr`