



Title	Sobolev inequalities with symmetry
Author(s)	Cho, Yonggeun; Ozawa, Tohru
Citation	Hokkaido University Preprint Series in Mathematics, 859, 1-8
Issue Date	2007
DOI	10.14943/84009
Doc URL	http://hdl.handle.net/2115/69668
Type	bulletin (article)
File Information	pre859.pdf



[Instructions for use](#)

SOBOLEV INEQUALITIES WITH SYMMETRY

YONGGEUN CHO AND TOHRU OZAWA

ABSTRACT. In this paper we derive some Sobolev inequalities for radially symmetric functions in \dot{H}^s with $\frac{1}{2} < s < \frac{n}{2}$. We show the end point case $s = \frac{1}{2}$ on the homogeneous Besov space $\dot{B}_{2,1}^{\frac{1}{2}}$. These results are extensions of the well-known Strauss' inequality [11]. Also non-radial extensions are given, which show that compact embeddings are possible in some L^p spaces if a suitable angular regularity is imposed.

1. INTRODUCTION

In this paper we derive Sobolev inequalities with symmetry. We first consider several Sobolev inequalities for radially symmetric functions in $\dot{H}^s(\mathbb{R}^n)$ with $\frac{1}{2} < s < \frac{n}{2}$. There is a sharp result by Sickel and Skrzypczak [8], although the argument below is much simpler and direct and a constant in the inequality in Proposition 1 below is given explicitly in terms of s .

Definition 1.

$$\dot{H}_{\text{rad}}^s = \{u \in \dot{H}^s(\mathbb{R}^n) : u \text{ is radially symmetric}\}, \quad s \geq 0.$$

$$\dot{B}_{p,q,\text{rad}}^s = \{u \in \dot{B}_{p,q}^s(\mathbb{R}^n) : u \text{ is radially symmetric}\}, \quad s \geq 0, \quad 1 \leq p, q \leq \infty.$$

The inhomogeneous spaces of radially symmetric functions are defined by the same way with spaces H^s and $B_{p,q}^s$.

Proposition 1. *Let $n \geq 2$ and let s satisfy $1/2 < s < n/2$.*

Then

$$\sup_{x \in \mathbb{R}^n \setminus \{0\}} |x|^{n/2-s} |u(x)| \leq C(n, s) \|(-\Delta)^{s/2} u\|_{L^2} \quad (1)$$

for all $u \in \dot{H}_{\text{rad}}^s$, where

$$C(n, s) = \left(\frac{\Gamma(2s-1)\Gamma(\frac{n}{2}-s)\Gamma(\frac{n}{2})}{2^{2s}\pi^{n/2}\Gamma(s)^2\Gamma(\frac{n}{2}-1+s)} \right)^{1/2}$$

and Γ is the gamma function.

Remark 1. For $s = 1$ with $n \geq 3$, the inequality (1) reduces to Ni's inequality [6, 7].

Remark 2. The restriction $1/2 < s < n/2$ is necessary for $C(n, s)$ to be finite.

2000 *Mathematics Subject Classification.* Primary 46E35; Secondary 42C15.

Keywords and phrases. Sobolev inequality, function space with radial symmetry, angular regularity.

Remark 3. The inequality (1) fails for $s = n/2$. Indeed, $u(x) = \mathcal{F}^{-1} \left(\frac{1}{(1+|\xi|)^n (1+\log(1+|\xi|))} \right)$ satisfies $u \in H_{\text{rad}}^{n/2}$, and $u \notin L^\infty$ where \mathcal{F} is the Fourier transform [12] and \mathcal{F}^{-1} is its inverse.

Remark 4. The inequality (1) fails if $0 \leq s < 1/2$ and $n \geq 3$. Indeed, $u = \mathcal{F}^{-1}(J_{\frac{n}{2}-1}(|\xi|)|\xi|^{-n/2})$ satisfies $u \in \dot{H}_{\text{rad}}^s$ and $u(x) = \infty$ for all $x \in S^{n-1}$, where we note that $u \in \dot{H}_{\text{rad}}^s$ if and only if $1 - n/2 < s < 1/2$, since

$$\|(-\Delta)^{s/2}u\|_{L^2}^2 = c_n \int_0^\infty |J_{\frac{n}{2}-1}(\rho)|^2 \rho^{2s-1} d\rho$$

and that by the asymptotic behavior of Bessel function (10)

$$u(x) = \int_0^\infty |J_{\frac{n}{2}-1}(\rho)|^2 d\rho = \infty, \quad x \in S^{n-1}.$$

See also the proof of Proposition 1 below.

In the endpoint case $s = 1/2$, we have the following proposition.

Proposition 2. *Let $n \geq 2$. Then there exists a constant C such that*

$$\sup_{x \in \mathbb{R}^n \setminus \{0\}} |x|^{(n-1)/2} |u(x)| \leq C \|u\|_{\dot{B}_{2,1}^{1/2}} \quad (2)$$

for all $u \in \dot{B}_{2,1,\text{rad}}^{1/2}$.

Remark 5. The inhomogeneous version of (2) has been given in [8] whose proof is based on the atomic decomposition.

Proposition 3. *Let $n \geq 2$ and let s satisfy $1/2 \leq s < 1$. Then there exists C such that for all $u \in H_{\text{rad}}^1$*

$$\sup_{x \in \mathbb{R}^n \setminus \{0\}} |x|^{n/2-s} |u(x)| \leq C(n, s) \|u\|_{L^2}^{1-s} \|\nabla u\|_{L^2}^s. \quad (3)$$

Remark 6. For $s = 1/2$, the inequality (3) reduces to Strauss' inequality [11].

Remark 7. For $s = 0$, the inequality (3) holds for nonincreasing functions in $|x|$ [2]. For $s = 1$, the inequality (3) holds for $n \geq 3$ and fails for $n = 2$. See Proposition 1 and Remark 2.

Now we extend the results above on radial functions to the non-radial functions with additional angular regularity. For details, let us define function spaces $H_\omega^{s,m}$ and $B_{2,1,\omega}^{s,m}$, $s \geq 0, m \geq 0$ as follows.

Definition 2.

$$H_\omega^{s,m} = \left\{ u \in H^s : \|u\|_{H_\omega^{s,m}} \equiv \|(1 - \Delta_\omega)^{\frac{m}{2}} u\|_{H^s} < \infty \right\},$$

$$B_{2,1,\omega}^{s,m} = \left\{ u \in B_{2,1}^{\frac{1}{2}} : \|u\|_{B_{2,1,\omega}^{s,m}}^2 \equiv \|(1 - \Delta_\omega)^{\frac{m}{2}} u\|_{B_{2,1}^s} < \infty \right\},$$

where Δ_ω is the Laplace-Beltrami operator on S^{n-1} .

The homogeneous spaces $\dot{H}_\omega^{s,m}$ and $\dot{B}_{2,1,\omega}^{s,m}$ is similarly defined by the definition of \dot{H}^s and $\dot{B}_{2,1}^s$. Then we have the following.

Proposition 4. (1) *If $1/2 < s < n/2$ and $m > n - 1 - s$, then there exists a constant C such that for any $u \in H_\omega^{s,m}$*

$$\sup_{\mathbb{R}^n \setminus \{0\}} |x|^{n/2-s} |u(x)| \leq C \|u\|_{\dot{H}_\omega^{s,m}}. \quad (4)$$

(2) *If $m > n - \frac{3}{2}$, then there exists a constant C such that for any $u \in B_{2,1,\omega}^{\frac{1}{2},m}$*

$$\sup_{\mathbb{R}^n \setminus \{0\}} |x|^{(n-1)/2} |u(x)| \leq C \|u\|_{\dot{B}_{2,1,\omega}^{\frac{1}{2},m}}. \quad (5)$$

Remark 8. $H_\omega^{s,m}$ and $B_{2,1,\omega}^{\frac{1}{2},m}$ are closed subspaces of H^s and $B_{2,1}^s$, respectively and they contain H_{rad}^s and $B_{2,1,\text{rad}}^{\frac{1}{2}}$ naturally, respectively. We can identify the spaces H^s with $(1 - \Delta_\omega)^{m/2} H_\omega^{s,m}$ and also $B_{2,1}^{\frac{1}{2}}$ with $(1 - \Delta_\omega)^{m/2} B_{2,1,\omega}^{\frac{1}{2},m}$.

Remark 9. $H_\omega^{s,m}$ is a Hilbert space with the same inner product as L^2 space and its dual space is given by $H_\omega^{-s,-m}$.

From the decay at infinity we deduce compact embeddings of $H_\omega^{s,m}$ and $B_{2,1,\omega}^{\frac{1}{2},m}$ into some L^p spaces as follows. See [2, 3, 4, 8, 9] for the radial case.

Corollary 1. *The embedding $H_\omega^{s,m} \hookrightarrow L^p$ is compact for $1/2 < s < n/2$, $m > n - 1 - s$ and $2 < p < 2n/(n - 2s)$.*

Corollary 2. *The embedding $B_{2,1,\omega}^{\frac{1}{2},m} \hookrightarrow L^p$ is compact for $m > n - 3/2$ and $2 < p < 2n/(n - 1)$.*

2. PROOFS

2.1. Proof of Proposition 1. We use the following Fourier representation for radially symmetric functions as

$$u(x) = |x|^{1-\frac{n}{2}} \int_0^\infty J_{\frac{n}{2}-1}(|x|\rho) \widehat{u}(\rho) \rho^{\frac{n}{2}} d\rho, \quad (6)$$

where J_ν is the Bessel function of order ν , \widehat{u} is the Fourier transform normalized as

$$\widehat{u}(\xi) = (2\pi)^{-n/2} \int e^{-ix \cdot \xi} u(x) dx,$$

and we have identified radially symmetric functions on \mathbb{R}^n with the corresponding functions on $(0, \infty)$.

By the Cauchy-Schwarz inequality and the Plancherel formula, we have

$$\begin{aligned}
& |x|^{\frac{n}{2}-s}|u(x)| \\
& \leq |x|^{1-s} \left(\int_0^\infty |J_{\frac{n}{2}-1}(|x|\rho)|^2 \rho^{1-2s} d\rho \right)^{1/2} \left(\int_0^\infty |\widehat{u}(\rho)|^2 \rho^{2s+n-1} d\rho \right)^{1/2} \\
& = \left(\int_0^\infty |J_{\frac{n}{2}-1}(\rho)|^2 \rho^{1-2s} d\rho \right)^{1/2} \left(\frac{\Gamma(\frac{n}{2})}{2\pi^{n/2}} \int |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi \right)^{1/2} \\
& = C(n, s) \|(-\Delta)^{s/2} u\|_{L^2},
\end{aligned} \tag{7}$$

as required.

2.2. Proof of Proposition 2. We use the following estimates on Bessel functions:

$$\sup_{r \geq 0} |J_{\frac{n}{2}-1}(r)| \leq 1. \tag{8}$$

$$\sup_{r \geq 0} r^{1/2} |J_{\frac{n}{2}-1}(r)| \leq C. \tag{9}$$

The first inequality (8) follows from the integral representation (see [13])

$$\begin{aligned}
J_{\frac{n}{2}-1}(r)^2 &= \frac{2}{\pi} \int_0^{\pi/2} J_{n-2}(2r \cos \theta) d\theta, \\
J_m(t) &= \frac{1}{2\pi} \int_0^{2\pi} \cos(m\theta - t \sin \theta) d\theta, \quad m \in \mathbb{Z}.
\end{aligned}$$

The second inequality (9) follows from the first and the well-known asymptotics (see [10])

$$J_\nu(r) \sim \sqrt{\frac{2}{\pi r}} \cos\left(r - \frac{(2\nu+1)\pi}{4}\right) \quad \text{as } r \rightarrow \infty. \tag{10}$$

We apply the Littlewood-Paley decomposition $\{\varphi_j\}_{j \in \mathbb{Z}}$ on $\mathbb{R}^n \setminus \{0\}$ to (4) to obtain

$$u(x) = |x|^{1-\frac{n}{2}} \sum_{j \in \mathbb{Z}} \int_0^\infty J_{\frac{n}{2}-1}(|x|\rho) \psi_j(\rho) \varphi_j(\rho) \widehat{u}(\rho) \rho^{\frac{n}{2}} d\rho, \tag{11}$$

where $\psi_j = \varphi_{j-1} + \varphi_j + \varphi_{j+1}$ and $\text{supp } \varphi_j \subset \{2^{j-1} \leq \rho \leq 2^{j+1}\}$.

As in (7), we have

$$\begin{aligned}
& |x|^{(n-1)/2} |u(x)| \\
& \leq |x|^{1/2} \sup_{j \in \mathbb{Z}} \left(\int_0^\infty |J_{\frac{n}{2}-1}(|x|\rho)|^2 \psi_j(\rho)^2 d\rho \right)^{1/2} \\
& \quad \cdot \sum_{j \in \mathbb{Z}} \left(\int_0^\infty |\varphi_j(\rho) \widehat{u}(\rho)|^2 \rho^n d\rho \right)^{1/2}.
\end{aligned}$$

By (9), we estimate

$$\begin{aligned}
& |x|^{1/2} \sup_{j \in \mathbb{Z}} \left(\int_0^\infty |J_{\frac{n}{2}-1}(|x|\rho)|^2 \psi_j(\rho)^2 d\rho \right)^{1/2} \\
& \leq C \sup_{j \in \mathbb{Z}} \left(\int_0^\infty \frac{1}{\rho} \psi_j(\rho)^2 d\rho \right)^{1/2} \\
& \leq C \sup_{j \in \mathbb{Z}} \left(\int_{2^{j-2}}^{2^{j+2}} \frac{1}{\rho} d\rho \right)^{1/2} \leq C.
\end{aligned} \tag{12}$$

This proves (2) since

$$\sum_{j \in \mathbb{Z}} \left(\int_0^\infty |\varphi_j(\rho) \widehat{u}(\rho)|^2 \rho^n d\rho \right)^{1/2}$$

is equivalent to the seminorm on $\dot{B}_{2,1,\text{rad}}^{1/2}$.

2.3. Proof of Proposition 3. If we use Cauchy-Schwartz inequality as in (7), we have for any $M > 0$

$$\begin{aligned}
& |x|^{n/2-s} |u(x)| \\
& \leq |x|^{1-s} \left(\int_0^{M|x|} |J_{\frac{n}{2}-1}(|x|\rho)|^2 \rho d\rho \right)^{1/2} \left(\int_0^{M|x|} |\widehat{u}(\rho)|^2 \rho^{n-1} d\rho \right)^{1/2} \\
& + |x|^{1-s} \left(\int_{M|x|}^\infty |J_{\frac{n}{2}-1}(|x|\rho)|^2 \rho^{-1} d\rho \right)^{1/2} \left(\int_{M|x|}^\infty |\widehat{u}(\rho)|^2 \rho^{n+1} d\rho \right)^{1/2} \\
& \leq |x|^{-s} \left(\int_0^M |J_{\frac{n}{2}-1}(r)|^2 r dr \right)^{1/2} \|u\|_{L^2} \\
& + |x|^{1-s} \left(\int_M^\infty |J_{\frac{n}{2}-1}(r)|^2 r^{-1} dr \right)^{1/2} \|\nabla u\|_{L^2}.
\end{aligned}$$

From (8) and (9) we deduce that $\sup_{r \geq 0} |r^{1-s} J_{\frac{n}{2}-1}(r)| \leq C$ for any $\frac{1}{2} \leq s < 1$. Hence we have for any $M > 0$

$$\begin{aligned}
& |x|^{n/2-s} |u(x)| \\
& \leq |x|^{-s} \left(\int_0^M |J_{\frac{n}{2}-1}(r)|^2 r dr \right)^{\frac{1}{2}} \|u\|_{L^2} \\
& + |x|^{1-s} \left(\int_M^\infty |J_{\frac{n}{2}-1}(r)|^2 r^{-1} dr \right)^{\frac{1}{2}} \|\nabla u\|_{L^2} \\
& \leq C |x|^{-s} M^s \|u\|_{L^2} + |x|^{1-s} M^{-(1-s)} \|\nabla u\|_{L^2}.
\end{aligned}$$

The minimization of the RHS of the last inequality with respect to M yields Proposition 3.

2.4. Proofs of Proposition 4 and Corollaries. The proof for (4) follows from the one of Proposition 1 and the spherical harmonic expansion of functions in $H_\omega^{s,m}$ [5, 10]. In fact, if we write $u(r\omega) = \sum_{k \geq 0} \sum_{1 \leq l \leq d(k)} f_{k,l}(r) Y_{k,l}(\omega)$, where $d(k)$ is the dimension of space of spherical harmonic functions of degree k and

$$d(k) \leq Ck^{n-2} \text{ for large } k. \quad (13)$$

Then we have

$$|x|^{\frac{n}{2}-s} u(|x|\omega) = c_n \sum_{k,l} |x|^{1-s} \int_0^\infty J_{\nu(k)}(|x|\rho) \rho^{\frac{n}{2}} g_{k,l}(\rho) d\rho Y_{k,l}(\omega), \quad (14)$$

where $\omega \in S^{n-1}$, $\nu(k) = \frac{n+2k-2}{2}$ and $\widehat{f_{k,l} Y_{k,l}}(\rho\omega) = g_{k,l}(\rho) Y_{k,l}(\omega)$. Here

$$g_{k,l}(\rho) = c_{n,k} \int_0^\infty f_{k,l}(r) r^{n-1} (r\rho)^{-\frac{n-2}{2}} J_{\nu(k)}(r\rho) dr.$$

The absolute value of $c_{n,k}$ is bounded by a constant depending only on n . See [10] for this.

Using the Cauchy-Schwarz inequality as in (7), we have

$$\begin{aligned} & |x|^{\frac{n}{2}-s} |u(|x|\omega)| \\ & \leq C \sum_{k,l} \|Y_{k,l}\|_{L^\infty(S^{n-1})} \left(\int_0^\infty |J_{\nu(k)}(|x|\rho)|^2 \rho^{1-2s} \right)^{\frac{1}{2}} \left(\int_0^\infty |g_{k,l}(\rho)|^2 \rho^{2s+n-1} d\rho \right)^{\frac{1}{2}} \\ & \leq C \sum_{k,l} k^{\frac{n-2}{2}} \left(\frac{\Gamma(\nu(k)+1-s)}{\Gamma(\nu(k)+s)} \right)^{\frac{1}{2}} \left(\int_0^\infty |g_{k,l}(\rho)|^2 \rho^{2s+n-1} d\rho \right)^{\frac{1}{2}} \|Y_{k,l}\|_{L^2(S^{n-1})}. \end{aligned}$$

Here we used the inequality that $\|Y_{k,l}\|_{L^\infty} \leq Ck^{\frac{n-2}{2}} \|Y_{k,l}\|_{L^2}$ (see for instance [10]). Using the Stirling's formula for gamma function that $\Gamma(t) \approx t^{t-\frac{1}{2}} e^{-(t-1)}$ for large t (for instance, see [1]) and the fact $-\Delta_\omega Y_{k,l} = k(k+n-2)Y_{k,l}$, we have from (13)

$$\begin{aligned} & |x|^{\frac{n}{2}-s} |u(|x|\omega)| \\ & \leq C \sum_k k^{\frac{n-2}{2}} d(k)^{\frac{1}{2}} \left(\frac{\Gamma(\nu(k)+1-s)}{\Gamma(\nu(k)+s)} \right)^{\frac{1}{2}} \\ & \quad \cdot \left(\sum_{1 \leq l \leq d(k)} \int_0^\infty |g_{k,l}(\rho)|^2 \rho^{2s+n-1} d\rho \|Y_{k,l}\|_{L^2(S^{n-1})}^2 \right)^{\frac{1}{2}} \\ & \leq C \left(\sum_k k^{2(n-\frac{3}{2}-s-m)} \right)^{\frac{1}{2}} \left(\sum_{k,l} k^{2m} \int_0^\infty |g_{k,l}(\rho)|^2 \rho^{2s+n-1} d\rho \|Y_{k,l}\|_{L^2(S^{n-1})}^2 \right)^{\frac{1}{2}} \\ & \leq C \left(\sum_{k,l} k^{2m} \int_0^\infty \int_{S^{n-1}} |\mathcal{F}(f_{k,l} Y_{k,l})(\rho\omega)|^2 \rho^{2s+n-1} d\rho d\omega \right)^{\frac{1}{2}} \\ & \leq C \left(\sum_{k,l} \int_0^\infty \int_{S^{n-1}} \left| \mathcal{F}((1-\Delta_\omega)^{\frac{m}{2}}(f_{k,l} Y_{k,l}))(\rho\omega) \right|^2 \rho^{2s+n-1} d\rho d\omega \right)^{\frac{1}{2}} \\ & \leq C \|u\|_{\dot{H}_\omega^{s,m}}, \end{aligned}$$

where \mathcal{F} is the Fourier transform. This proves part (1).

For the part (2), if we use the Littlewood-Paley decomposition $\{\varphi_j\}_{j \in \mathbb{Z}}$ as in the proof of Proposition 2, then we have

$$|x|^{\frac{n-1}{2}} u(|x|\omega) = c_n \sum_{j \in \mathbb{Z}} \sum_{k,l} |x|^{\frac{1}{2}} \int_0^\infty J_{\nu(k)}(|x|\rho) \rho^{\frac{n}{2}} \psi_j(\rho) \varphi_j(\rho) g_{k,l}(\rho) d\rho Y_{k,l}(\omega), \quad (15)$$

Since $m > n - 3/2$, by (12) we deduce that

$$\begin{aligned} & |x|^{\frac{n-1}{2}} |u(|x|\omega)| \\ & \leq C \sum_{j \in \mathbb{Z}} \sum_{k,l} \left(\int_0^\infty |\varphi_j(\rho) g_{k,l}(\rho)|^2 \rho^n d\rho \right)^{\frac{1}{2}} \|Y_{k,l}\|_{L^\infty(S^{n-1})} \\ & \leq C \sum_{j \in \mathbb{Z}} \left(\sum_k k^{2(n-2-m)} \right)^{\frac{1}{2}} \left(\sum_{k,l} k^{2m} \int_0^\infty |\varphi_j(\rho) g_{k,l}(\rho)|^2 \rho^n d\rho \|Y_{k,l}\|_{L^2(S^{n-1})}^2 \right)^{\frac{1}{2}} \\ & \leq C \sum_{j \in \mathbb{Z}} 2^{\frac{j}{2}} \|\varphi_j \mathcal{F}((1 - \Delta_\omega)^{\frac{m}{2}} u)\|_{L^2} = C \|u\|_{\dot{B}_{2,1,\omega}^{\frac{1}{2}}}. \end{aligned}$$

To show Corollary 1 we use the fact that $H_\omega^{s,m}$ is a Hilbert space. Hence any bounded sequence $\{u_j\}$ in $H_\omega^{s,m}$ satisfies $u_j(x) \rightarrow 0$ as $|x| \rightarrow \infty$ uniformly and has a subsequence converges to u in $H_\omega^{s,m}$ weakly. Let us denote the subsequence by u_{j_k} .

Now choose a smooth function φ supported in the ball of radius $R + 1$ and with value 1 in the ball of radius R . By the standard argument one can easily show that for each R the mapping $u \mapsto \varphi u$ is compact from H^t to $H^{t'}$ if $t' < t$. By the compactness above and Sobolev embedding we may assume that the sequence φu_{j_k} satisfies that for $2 \leq q < \frac{2n}{n-2s}$

$$\|\varphi u_{j_k} - \varphi u\|_{L^q} \rightarrow 0 \text{ as } k \rightarrow \infty. \quad (16)$$

Thus we have

$$\|u_{j_k} - u\|_{L^p} \leq \|\varphi(u_{j_k} - u)\|_{L^p} + \|(1 - \varphi)(u_{j_k} - u)\|_{L^p} \equiv I_k + II_k$$

with $I_k \rightarrow 0$ as $k \rightarrow \infty$ by (16) since $2 < p < \frac{2n}{n-2s}$. From the uniform convergence that $|u_{j_k}(x)| + |u(x)| \rightarrow 0$ as $|x| \rightarrow \infty$ it follows that

$$\limsup_{k \rightarrow \infty} II_k \leq \sup_k \|u_{j_k} - u\|_{L^\infty(|x| > R)}^{\frac{p-2}{p}} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

This proves the compactness of the embedding $H_\omega^{s,m} \hookrightarrow L^p$.

Since $B_{2,1,\omega}^{\frac{1}{2},m} \hookrightarrow H_\omega^{\frac{1}{2},m}$, one can adapt the same arguments (compactness of cut-off mapping and uniform convergence at infinity) as above except for weak-* convergence of u_{j_k} to u in $B_{2,1,\omega}^{\frac{1}{2},m}$ for the compactness of embedding $B_{2,1,\omega}^{\frac{1}{2},m}$ to L^p . This completes the proof.

REFERENCES

- [1] G. B. Arfken and H. J. Weber, *Mathematical Methods for Physicists*, Academic Press, 1995.
- [2] T. Cazenave, *Semilinear Schrödinger Equations*, Courant Lecture Notes in Mathematics. **10**, AMS, 2003.
- [3] J. Chabrowski, *On compact embeddings of radial Sobolev spaces and their applications*, Commun. Partial Differential Equations, **17**(1992), 941-952.
- [4] Y. Ebihara and T.P. Schonbek, *On the (non)compactness of the radial Sobolev spaces*, Hiroshima Math. J. **16**(1986), 665-669.
- [5] J. Kato, M. Nakamura and T. Ozawa, *A generalization of the weighted Strichartz estimates for wave equations and an application to self-similar solutions*, Commun. Pure. Appl. Math. **60**(2007), 164-186.
- [6] W.-M. Ni, *A nonlinear Dirichlet problem on the unit ball and its applications*, Indiana Univ. Math. J. **31**(1982), 801-807.
- [7] W. Rother, *Some existence results for the equation $\Delta U + K(x)U^p = 0$* , Commun. Partial Differential Equations. **15**(1990), 1461-1473.
- [8] W. Sickel and L. Skrzypczak, *Radial subspaces of Besov and Lizorkin-Triebel classes: extended Strauss lemma and compactness of embeddings*, The Journal of Fourier Analysis and Applications. **6**(2000), 639-662.
- [9] L. Skrzypczak, *Rotation invariant subspaces of Besov and Triebel-Lizorkin space: compactness of embeddings, smoothness and decay of functions*, Rev. Mat. Iberoamericana, **18**(2002), 267-299.
- [10] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, 1971.
- [11] W. A. Strauss, *Existence of solitary waves in higher dimensions*, Commun. Math. Phys. **55**(1977), 149-162.
- [12] M. E. Taylor, *Partial Differential Equations I, Basic Theory*, Applied Mathematical Sciences **115**, Springer, 1996.
- [13] G. Watson, *A Treatise on the Theory of Bessel Functions*, Reprint of the second (1944) edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1995.

DEPARTMENT OF MATHEMATICS, POSTECH, POHANG 790-784, REPUBLIC OF KOREA
E-mail address: changocho@postech.ac.kr

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, 060-0810, JAPAN
E-mail address: ozawa@math.sci.hokudai.ac.jp