



Title	On time analyticity of the Navier-Stokes equations in a rotating frame with spatially almost periodic data
Author(s)	Giga, Yoshikazu; Jo, Hideaki; Mahalov, Alex; Yoneda, Tsuyoshi
Citation	Hokkaido University Preprint Series in Mathematics, 860, 2-16
Issue Date	2007
DOI	10.14943/84010
Doc URL	http://hdl.handle.net/2115/69669
Type	bulletin (article)
File Information	pre860.pdf



[Instructions for use](#)

On time analyticity of the Navier-Stokes equations in a rotating frame with spatially almost periodic data

Yoshikazu Giga

*Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba,
Meguro-ku, Tokyo 153-8914, Japan*

Hideaki Jo

*Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba,
Meguro-ku, Tokyo 153-8914, Japan*

Alex Mahalov

*Department of Mathematics, Arizona State University, Tempe, AZ 85287-1804,
USA*

Tsuyoshi Yoneda

*Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba,
Meguro-ku, Tokyo 153-8914, Japan*

Abstract

We consider the Navier-Stokes equations with the Coriolis force when initial data may not decrease at spatial infinity so that almost periodic data is allowed. We prove that the local-in-time solution is analytic in time when initial data is in FM_0 , Fourier preimage of the space of all finite Radon measures with no point mass at the origin. When the initial data is almost periodic, this implies that the complex amplitude is analytic in time. In particular, a new mode cannot be created at any positive time.

Key words: Navier-Stokes equations, analyticity, frequency set, Coriolis force

Email addresses: labgiga@ms.u-tokyo.ac.jp (Yoshikazu Giga),
hideaki@ms.u-tokyo.ac.jp (Hideaki Jo), mahalov@asu.edu (Alex Mahalov),
yoneda@ms.u-tokyo.ac.jp (Tsuyoshi Yoneda).

1 Introduction and main results

We consider the Cauchy problem for the three-dimensional Navier-Stokes equations with the Coriolis force:

$$u_t + (u \cdot \nabla)u + \Omega e_3 \times u - \Delta u = -\nabla p, \quad \nabla \cdot u = 0, \quad u|_{t=0} = u_0, \quad (1)$$

where $u = u(x, t) = (u_1(x, t), u_2(x, t), u_3(x, t))$ is the unknown velocity vector field and $p = p(x, t)$ is the unknown scalar pressure at the point $x = (x_1, x_2, x_3) \in \mathbf{R}^3$ in space and time $t > 0$ while $u_0 = u_0(x)$ is the given initial velocity field. Here $\Omega \in \mathbf{R}$ is the Coriolis parameter, which is twice the angular velocity of the rotation around the vertical unit vector $e_3 = (0, 0, 1)$; the kinematic viscosity coefficient is normalized by one. By \times we denote the exterior product, hence, the Coriolis term is represented by $e_3 \times u = Ju$ with the corresponding skew-symmetric 3×3 matrix J .

We would like to solve (1) when initial data u_0 is almost periodic, i.e., u_0 is formally of the form

$$u_0(x) = \sum_{\lambda \in \Lambda} a_\lambda e^{i\lambda x}, \quad (2)$$

where Λ (called a frequency set) is a countable set in \mathbf{R}^3 and $a_\lambda \in \mathbf{C}^3 \setminus \{0\}$. We would like to discuss time evolution of frequency sets. A naive space for this purpose is BUC , the space of all bounded uniformly continuous functions since almost periodic functions in the sense of Bohr belong to BUC . The problem (1) is well-posed when $\Omega = 0$ as proved in [10]. However, when $\Omega \neq 0$ it is shown in [11] that the linearized problem around $u = 0$ is ill-posed in the space BUC . Sawada [20] and Hieber and Sawada [15] proved unique local existence for initial data in a homogeneous Besov space $B_{\infty,1}^0$, which is strictly smaller than BUC . The estimate for the existence time is improved by [11]. However, none of these results gives a uniform estimate for the existence time in the speed of rotation $\Omega \in \mathbf{R}$, the precondition to consider fast singular oscillating limits $\Omega \rightarrow \pm\infty$ and prove global existence for large fixed Ω .

To overcome this difficulty the space FM_0 (Fourier preimage of the space of all finite Radon measures with no point mass at the origin) is introduced in [12]. This space FM_0 is strictly smaller than $B_{\infty,1}^0$ as proved in [12]. However, the existence time can be taken uniformly in Ω as proved in [12].

The global existence of a regular solution of (1) for large $|\Omega|$ were obtained in L^2 -setting for the periodic domains such as cylinder with infinite length in the direction of the rotating axis (periodic or zero stress boundary conditions in the vertical direction) or spatial lattices [4], [5], [17]. In this regard, for the Euler equations, Nicolaenko, Bardos Golse and the third author [18] proved uniform local existence and long-time regularity for initial data in H^4 .

In FM_0 setting a global solvability is proved for small Reynolds number but all Ω under some conditions on frequency set of u_0 [13].

Our goal in this paper is to prove the analyticity of the solution u in time. Here is a typical result.

Theorem 1.1. Assume that $u_0 \in FM_0$ with $\operatorname{div} u_0 = 0$. Let u be the unique mild solution $u = u(t) \in C([0, T], FM_0)$ of (1) (see [12]). Then u is analytic in $(0, T)$ in the sense it can be extended to be a holomorphic function with values in FM_0 in a complex neighborhood of $(0, T)$.

For the case $\Omega = 0$ we may state in the framework of BUC .

Theorem 1.2. Assume that $\Omega = 0$ and $u_0 \in BUC$ with $\operatorname{div} u_0 = 0$. Let u be the unique mild solution $u = u(t) \in C([0, T], BUC)$ of (1). Then u is analytic in $(0, T)$ with values in BUC .

The analyticity in time for the Navier-Stokes equations has been discussed by many authors in various settings. The first result in this direction was established by Masuda[19]. He proved that if $u \in C([0, T]; H^2(D))$ is a weak solution of (1) with $\Omega = 0$, then u is analytic in $(0, T)$ with value in $H^2(D)$. In [19] D is assumed to be a bounded domain and the Dirichlet condition is imposed. Also, an analytic exterior force is allowed. (His result works for unbounded domains in L^2 -setting too.) This result is extended by the first author [9] for L^p -setting. Foias and Temam[8] and Iooss[16] also discuss the time analyticity when $\Omega = 0$ in a different setting. However, non of these results apply to the space containing nondecaying functions such as almost periodic functions.

We now consider an almost periodic initial data $u_0 \in FM_0$ of the form (2). As pointed out in [12] the solution $u(t) \in FM_0$ is almost periodic as proved in [14] for other spaces. It is interesting to study evolution of the frequency set.

Theorem 1.3. Let u be the mild solution of (1) in $[0, T]$ with initial data $u_0 \in FM_0$ which is almost periodic of the form (2). Then the frequency set $\Lambda(u(\cdot, t))$ of $u(\cdot, t)$ is contained in the additive free semigroup G generated by Λ . Thus u is of the form

$$u(x, t) = \sum_{\lambda \in G} a_\lambda(t) e^{i\lambda \cdot x}, \quad (3)$$

where the amplitude $a_\lambda(t) \in \mathbf{C}^3$ and satisfies

$$\sup_{0 < t < T} \sum_{\lambda \in G} |a_\lambda(t)| < \infty.$$

(The sum converges in absolute sense uniformly in $x \in \mathbf{R}^3$.)

Since the amplitude function $a_\lambda(t)$ can be calculated by

$$a_\lambda(t) = M[ue^{-i\lambda \cdot x}],$$

where $M[f] = \lim_{R \rightarrow \infty} |C_R|^{-1} \int_{C_R} f dx$ is the spatial averaging operator; C_R denote the cube $\{x = (x_1, x_2, x_3) \mid |x_i| \leq R, 1 \leq i \leq 3\}$; see [7]. By the time analyticity result we are able to prove the analyticity of the amplitude function.

Theorem 1.4. Assume the same hypotheses of either Theorem 1.1 or 1.2. Then the complex amplitude function $a_\lambda(t)$ is analytic in time $t > 0$.

This implies that there is no sudden creation of new mode at $t_0 > 0$ or there is no chance that some mode vanishes identically from some $t_0 > 0$. All modes appearing for $t > 0$ are created instantaneously. There is no waiting time for exciting amplitude dynamics corresponding to any frequency.

The work of the first author was partly supported by the Grant-in-Aid for Scientific Research, No. 18204011, No. 17654037, the Japan Society of the Promotion of Science (JSPS). He was also partly supported by the Grant-in-Aid for formation of COE 'Mathematics of Nonlinear Structures via Singularities' (Hokkaido University) sponsored by JSPS. The work of the third author was partly supported by AFOSR contract FA9550-05-1-0047. The work of the fourth author was partly supported by the Grant-in-Aid for JSPS Fellows.

2 Persistence of almost periodicity and frequency set

We begin by recalling the notion of almost periodicity in the sense of Bohr [1], [7]. Let f be in $BUC(\mathbf{R}^n)$. We say f is *almost periodic* if the set

$$\Sigma_f := \{f(\cdot + \xi) \mid \xi \in \mathbf{R}^n\} \subset L^\infty(\mathbf{R}^n)$$

is relatively compact in $L^\infty(\mathbf{R}^n)$. (Uniform continuity is redundant. In fact, if f must be uniformly continuous [1],[7].) If f is periodic in x , clearly, f is almost periodic since Σ_f is a torus.

We next recall the fact that if u_0 is almost periodic so is the solution $u(\cdot, t)$ of (1). To state more precisely we recall an integral equation which is formally equivalent to (1):

$$u(t) = e^{-A(\Omega)t} u_0 - \int_0^t \operatorname{div} e^{-A(\Omega)(t-s)} \mathbf{P}(u(s) \otimes u(s)) ds, \quad (4)$$

where $A(\Omega) = -\Delta + \Omega \mathbf{S}$, $\mathbf{S} = \mathbf{P} \mathbf{J} \mathbf{P}$ (Poincaré-Riesz operator) and $\mathbf{P} = (I + R_i R_j)_{i \leq i, j \leq 3}$ (Helmholtz projection); here, R_j denotes the Riesz operator $R_j = (-\Delta)^{-1/2} \partial / \partial x_j$. Instead of considering solutions of (1), we rather consider a solution of (4) in $C([0, T], FM_0)$ (or $C([0, T], BUC)$ for $\Omega = 0$) which is called a mild solution. The relation of (1) and (2) is discussed in [12] and [10]; we do not repeat it here.

Theorem 2.1.

- (i) Assume that $u_0 \in FM_0$ with $\operatorname{div} u_0 = 0$ is almost periodic. Let $u \in C([0, T], FM_0)$ be the unique mild solution of (4). Then $u(\cdot, t)$ is almost periodic in \mathbf{R}^3 for $t \in [0, T]$.
- (ii) Assume that $\Omega = 0$. Assume that $u_0 \in BUC(\mathbf{R}^3)$ with $\operatorname{div} u_0 = 0$ is almost periodic. Let $u \in C([0, T], BUC)$ be the unique mild solution of (4). Then $u(\cdot, t)$ is almost periodic in \mathbf{R}^3 for $t \in [0, T]$.

The second statement (ii) is found in [14] (Theorem 2.3). The first statement (i) is not explicitly written but the proof parallels that of (ii). We just give a sketch of the proof of (i) for the reader's convenience.

Sketch of the proof of (i). For the solution $u(t) = u(\cdot, t)$ let $S(t)$ denote the mapping $u_0 \mapsto u(t)$.

Since (1) is invariant under translation in the space variables, the solution u_η with initial data $u_{0\eta}(x) = u_0(x + \eta)$, $\eta \in \mathbf{R}^3$ fulfills $u_\eta(x, t) = u(x + \eta, t)$. Thus $S(t)$ maps Σ_{u_0} onto $\Sigma_{u(t)}$. We notice that

$$\sup_{0 < t < T} \|u_\eta\|_{FM}(t) = \sup_{0 < t < T} \|u\|_{FM}(t).$$

We now apply a uniform continuity dependence on initial data described below in Lemma 2.2 and observe that $S(t)$ is a well-defined continuous mapping from the closure $\bar{\Sigma}_{u_0}$ to $\bar{\Sigma}_{u(\cdot, t)}$ in FM_0 . Thus if $\bar{\Sigma}_{u_0}$ is compact in FM_0 , so is $\bar{\Sigma}_{u(\cdot, t)}$. One may replace FM_0 by BUC as proved below in Lemma 2.4. The proof is now complete. \square

Lemma 2.2.(Continuity with respect to initial data.) For a given N and T let $B(N)$ denote the set of all $u_0 \in FM_0$ such that

$$\sup_{0 < t < T} \|S(t)u_0\|_{FM} \leq N.$$

Then $S(t)$ is a Lipschitz continuous on $B(N)$ to FM_0 with a constant depending only on N and T .

The proof parallels that of [14] (Proposition 3.2). It is essentially contained in [12] (Section 3).

We conclude this section by proving Theorem 1.3. For this purpose we clarify the meaning of (2) when $u_0 \in FM$.

Lemma 2.3. Assume that f is almost periodic and belongs to $FM(\mathbf{R}^n)$. Then f is a Fourier sum of the form

$$f(x) = \sum_{\lambda \in \Lambda} a_\lambda e^{i\lambda x} \quad \text{with} \quad \sum_{\lambda \in \Lambda} |a_\lambda| < \infty, \quad (5)$$

where $a_\lambda \in \mathbf{C}^n$ and Λ is a countable subset of \mathbf{R}^n . Conversely, if f has the form (5), then $f \in FM(\mathbf{R}^n)$ and f is almost periodic.

Proof. Let μ be the Fourier image \hat{f} of f . We split $\mu \in M$ as

$$\mu = \mu_p + \mu_c$$

where μ_p is the point mass part and μ_c is the continuous part. The inverse Fourier image $F^{-1}\mu_p$ of μ_p is a Fourier sum as in (5). So $F^{-1}\mu_p$ is almost periodic [7]. We should prove that $f_c = F^{-1}\mu_c$ is zero if it is almost periodic. As in [7] it suffices to prove that all Fourier coefficients (complex amplitudes) of f_c must be zero which is obtained by calculating the average

$$M[f_c e^{-i\lambda \cdot x}] = \lim_{R \rightarrow \infty} (2R)^{-n} \int_{C_R} f_c e^{-i\lambda \cdot x} dx.$$

We observe that

$$\begin{aligned} (2\pi)^n \int_{C_R} f_c e^{-i\lambda \cdot x} dx &= \int_{\mathbf{R}^n} \int_{C_R} e^{ix \cdot \xi} e^{-i\lambda \cdot x} \mu_c(d\xi) dx = \int_{\mathbf{R}^n} \int_{C_R} e^{ix \cdot \eta} \mu_c^\lambda(d\eta) dx \\ &= R^n \int_{\mathbf{R}^n} \left(\int_{C_1} e^{iy \cdot R\eta} dy \right) \mu_c^\lambda(d\eta), \end{aligned}$$

where μ_c^λ is the translation of μ_c defined by $\mu_c^\lambda(B) = \mu_c(B + \{\lambda\})$ for Borel set B . Since

$$\lim_{R \rightarrow \infty} \int_{C_1} e^{iy \cdot R\eta} dy = 0 \quad \text{for all} \quad \eta \neq 0$$

and μ_c^λ has no point mass at the origin, by the Lebesgue dominant convergence theorem we conclude that $M[f_c e^{-i\lambda \cdot x}] = 0$ for all $\lambda \in \mathbf{R}^n$.

The converse is clear since $\sum |a_\lambda| = \|f\|_{FM}$ and the uniform limit of the Fourier sum is almost periodic. \square

Lemma 2.4. A function $f \in FM_0(\mathbf{R}^n)$ is almost periodic if and only if Σ_f is relatively compact in $FM_0(\mathbf{R}^n)$.

Proof. It suffices to prove ‘only if’ part since $FM_0 \subset L^\infty$ is continuous. We may assume that f is a scalar valued function. Suppose that $\Sigma_f(\subset FM_0)$ is relatively compact in $L^\infty(\mathbf{R}^n)$. Then any sequence $\{f_{n_k}\} \subset \Sigma_f$ has a convergent subsequence $\{f_\ell = f(\cdot + \eta_{k(\ell)})\} \rightarrow f_0$ in $L^\infty(\mathbf{R}^n)$ as $\ell \rightarrow \infty$. It suffices to prove that $f_\ell \rightarrow f_0$ in FM_0 . By (5) we have $f_0(x) = \sum_{\lambda \in \Lambda} a_\lambda e^{i\lambda x}$ and $f_\ell(x) = \sum_{\lambda \in \Lambda} a_\lambda e_{\lambda\ell} e^{i\lambda x}$, where $e_{\lambda\ell} := \exp(i\lambda \eta_{k(\ell)})$. Since $f_\ell \rightarrow f_0$ in $L^\infty(\mathbf{R}^n)$, we have

$$\lim_{\ell \rightarrow \infty} M[(f_\ell - f_0)e^{-i\mu \cdot x}] = 0 \quad \text{for all } \mu \in \Lambda.$$

In other words

$$a_\mu e_{\mu\ell} \rightarrow a_\mu \quad \text{as } \ell \rightarrow \infty$$

for all $\mu \in \Lambda$. Thus we conclude that

$$\|f_\ell - f_0\|_{FM} = \sum_{\lambda \in \Lambda} |a_\lambda - a_\lambda e_{\lambda\ell}| \rightarrow 0,$$

since $\sum_{\lambda \in \Lambda} |a_\lambda| = \sum_{\lambda \in \Lambda} |a_\lambda e_{\lambda\ell}| < \infty$. (The same statement holds even if we replace FM_0 by FM .) \square

Proof of Theorem 1.3. Assume that $f, g \in FM$ are almost periodic. Let $\Lambda(f)$ denote the frequency set, i.e., it is the set of all $\lambda \in \Lambda$ such that $a_\lambda \neq 0$. Then by Lemma 2.3 it is easy to see that

$$\Lambda(fg) \subset \Lambda(f) + \Lambda(g) = \{a + b | a \in \Lambda(f), b \in \Lambda(g)\}.$$

Since the solution of (4) is the limit of the successive approximation

$$u_{m+1}(t) = e^{-A(\Omega)t} u_0 - \int_0^t \operatorname{div} e^{-A(\Omega)(t-s)} \mathbf{P}(u_m(s) \otimes u_m(s)) ds \quad (m \geq 1),$$

$$u_1(t) = e^{-A(\Omega)t} u_0$$

and $e^{-A(\Omega)t}$, $\operatorname{div} e^{-A(\Omega)t} \mathbf{P}$ does not grow the frequency set, we observe that $u_m(t)$ is a Fourier sum of the form (5) and that

$$\begin{aligned} \Lambda(u_1(t)) &\subset \Lambda(u_0), \\ \Lambda(u_{m+1}(t)) &\subset \bigcup_{0 \leq \tau < t} \Lambda(u_m(\tau)) + \Lambda(u_m(\tau)), m = 1, 2, \dots \end{aligned}$$

This implies that $\Lambda(u_m(t)) \subset G$, where G is the free additive semigroup generated by Λ . Since the convergence of u_m in $C([0, T_0], FM_0)$ to the solution u for small T_0 has been established in [12] (Section 3), and since $\sum_{\lambda \in G} |a_\lambda(t)| = \|u(t)\|_{FM}$, we obtain (3) in $(0, T_0]$. We repeat this argument starting from $t = T_0$ with $\Lambda = G$ and obtain (1.3) in $[T_0, T_0 + T_1]$ with $T_1 > 0$.

We repeat this argument finitely many times to conclude (3) in $[0, T]$ since the amount of increment of time interval in each step can be taken uniformly since it depends only on the size of initial data in FM . \square

Remark 2.5.

- (i) Our terminology is slightly different from [13], where it gives an impression that frequency set is the support of \hat{f} , which is the closure of our frequency set of almost periodic function f .
- (ii) The assertion that the frequency set of u is contained in G is still valid in BUC setting when $\Omega = 0$ since we have $\Lambda(fg) \subset \Lambda(f) + \Lambda(g)$ for almost periodic $f, g \in BUC(\mathbf{R}^n)$; note that an almost periodic function can be approximated by trigonometric polynomials in uniform sense.

3 Time analyticity

We shall prove Theorem 1.1 and 1.2. Theorem 1.4 is an easy corollary of Theorem 1.1 and 1.2.

We recall the definition of holomorphic functions with values in a (complex) Banach space X . Let D be an open set in \mathbf{C} . We say that a function from D to X is (strongly) holomorphic (with values in X) in D if f is strongly differentiable at every point of D , i.e., for each $z \in D$ there exists an $L \in X$ such that

$$\lim_{h \rightarrow 0, h \in \mathbf{C}} \frac{1}{|h|} \|f(z+h) - f(z) - hL\|_X = 0.$$

For basic properties of Banach valued holomorphic functions the reader is referred to classical books by Yosida [22] and by Berger [3]. Let $H^\infty(D, X)$ be the space of all bounded holomorphic functions in D with values in X . It is a Banach space equipped with the supremum norm defined by

$$\|f\|_{H^\infty(D)} = \sup\{\|f(z)\|_X : z \in D\}.$$

We still use such notation when D is an open set in \mathbf{C}^2 ; in this case we say that $f : D \rightarrow X$ is holomorphic if $f = f(t, s)$ is holomorphic in each variable t and s . For $\sigma \in (0, \pi/2)$ and $T > 0$ we set

$$\begin{aligned} S(T, \sigma) &= \{t \in \mathbf{C} : |\arg t| < \sigma, \operatorname{Re} t < T\} \\ Z(T, \sigma) &= \{(t, s) \in S(T, \sigma) \times S(T, \sigma) : \operatorname{Re}(t - s) > 0\} \\ Z_\varepsilon(T, \sigma) &= \{(t, s) \in Z(T, \sigma) : \operatorname{Re}(t - s) > \varepsilon\} \text{ for } \varepsilon > 0, \end{aligned}$$

where $\operatorname{Re} z$ denotes the real part of a complex number z . The next Theorem is elementary but a key for our argument. It is implicitly used in the proof of [9] (Lemma 2).

Theorem 3.1. Assume that $f \in H^\infty(Z_\varepsilon(T, \sigma), X)$ for each $\varepsilon \in (0, T)$. Assume that there is an positive integrable functions φ on $(0, T)$ such that

$$\|f(t, s)\|_X \leq \varphi(\operatorname{Re}(t - s)) \text{ for } (t, s) \in Z(T, \sigma). \quad (6)$$

Then $F(t) := \int_0^t f(t, s)ds$ is in $H^\infty(S(T, \sigma), X)$ with

$$\|F\|_{H^\infty(S(T, \sigma))} \leq \int_0^T \varphi(\tau)d\tau. \quad (7)$$

Here, we interpret that the complex integration is made along the segment connecting two ends 0 and t .

Proof. We set

$$F_\delta(t) = \int_0^{t(1-\delta)} f(t, s) ds$$

for $\delta \in (0, 1)$. It suffices to prove that $F_\delta \in H^\infty(S(T, \sigma), X)$ since F is a uniform limit of F_δ as $\delta \rightarrow 0$ by the assumption (6); the estimate (7) easily follows from (6).

Since the boundedness of F_δ in $S(T, \sigma)$ is clear, it suffices to prove that F_δ is holomorphic in $S(T, \sigma)$. For $t \in S(T, \sigma)$ we take $h \in \mathbf{C}$ small so that $t + h \in S(T, \sigma)$. Applying Cauchy's integration theorem to change the path we observe that

$$\begin{aligned} F_\delta(t + h) - F_\delta(t) &= \int_0^{\alpha(t+h)} (f(t + h, s) - f(t, s))ds \\ &\quad + \int_0^{\alpha(t+h)} f(t, s)ds - \int_0^{\alpha t} f(t, s)ds \\ &= \int_0^{\alpha(t+h)} (f(t + h, s) - f(t, s))ds + \int_{\alpha t}^{\alpha(t+h)} f(t, s)ds \\ &=: J_1 + J_2 \end{aligned}$$

with $\alpha = 1 - \delta$. We recall a form of Taylor's expansion (See [2]):

$$g(z) = \sum_{k=0}^{n-1} \frac{g^{(k)}(a)}{k!} (z - a)^k + \left(\frac{1}{2\pi i} \int_C \frac{g(\zeta)}{(\zeta - a)^n (\zeta - z)} d\zeta \right) \cdot (z - a)^n. \quad (8)$$

Here g is an holomorphic function defined in a domain $D \subset \mathbf{C}$ and $a \in D$; $C(\subset D)$ is a circle (oriented counterclockwise) centered at a which encloses z .

We use (8) with $n = 2$, $a = t$, $z = t + h$, $g(z) = f(z, s)$. We fix the circle $C : |\zeta - t| = r$ so that $r < \text{dist}(t, \partial S(T, \sigma))$. We take h small so that $|h| < r$. Then

$$\frac{J_1}{h} - \int_0^{\alpha(t+h)} \frac{\partial f}{\partial t}(t, s) ds = h \int_0^{\alpha(t+h)} \left\{ \frac{1}{2\pi i} \int_C \frac{f(\zeta, s)}{(\zeta - (t+h))(\zeta - t)^2} d\zeta \right\} ds.$$

The norm of the right hand side in X is dominated by

$$\frac{h}{2\pi} \frac{M_\varepsilon}{(r - |h|)r^2} \cdot 2\pi r \text{ with } M_\varepsilon = \|f\|_{H^\infty(Z_\varepsilon(T, \sigma))}.$$

Similarly, we apply (8) with $n = 1$, $a = \alpha t$, $z = s$ and $g(z) = f(t, z)$ to observe that

$$\frac{J_2}{h} - \alpha f(t, \alpha t) = \int_{\alpha t}^{\alpha(t+h)} \left\{ \frac{1}{2\pi i} \int_C \frac{f(\zeta, s)}{(\zeta - \alpha t)(\zeta - s)} d\zeta \cdot (s - \alpha t) \right\} ds.$$

Here we fix the critical $C : |\zeta - \alpha t| = r$ so that $r < \text{dist}(\alpha t, \partial S(T, \sigma))$ and take h small so that $\alpha|h| < r$. The norm of the right hand side is dominated by

$$\frac{1}{2\pi} \frac{M_\varepsilon}{r(r - \alpha|h|)} 2\pi r \cdot \alpha|h|.$$

Thus we conclude that

$$\lim_{|h| \rightarrow 0} \frac{1}{|h|} \left\| F_\delta(t+h) - F_\delta(t) - h \left\{ \int_0^{\alpha t} \frac{\partial f}{\partial t}(t, s) ds + \alpha f(t, \alpha t) \right\} \right\|_X = 0,$$

that is to say that F_δ is (complex) differentiable at $t \in S(T, \sigma)$. \square

Lemma 3.2

If $u \in H^\infty(S(T, \sigma); FM)$, then

$$f(t, s) := \text{div} e^{-(t-s)A(\Omega)} \mathbf{P}u(s) \otimes u(s) \quad (9)$$

belongs to $H^\infty(Z_\varepsilon(T, \sigma); FM)$ for all $\varepsilon \in (0, T)$. Moreover,

$$\|f(t, s)\|_{FM} \leq \frac{C}{(\text{Re}(t-s))^{\frac{1}{2}}} \|u\|_{FM}^2 \quad (10)$$

holds with constant C independent of t, s and u .

Proof. First we show that the function $f(t, s)$ belongs to $H^\infty(Z_\varepsilon(T, \sigma); FM)$ if $u \in H^\infty(S(T, \sigma); FM)$ briefly. Let $\sigma^*(A)$ be the symbol of the operator A .

We shall recall the symbol of operators appearing in $\operatorname{div}e^{-(t-s)A(\Omega)}\mathbf{P}$. As in [11] the symbol of $e^{t\Omega\mathbf{S}}$ is given by

$$\sigma^*(e^{t\Omega\mathbf{S}}) = \exp(\Omega t \frac{\xi_3}{|\xi|} \mathbf{R}(\xi)), \quad (11)$$

where $\mathbf{R}(\xi)$ is the symbol of the vector Riesz operator, i.e.,

$$\mathbf{R}(\xi) = \frac{1}{|\xi|} \begin{pmatrix} 0 & -\xi_3 & \xi_2 \\ \xi_3 & 0 & -\xi_1 \\ -\xi_2 & \xi_1 & 0 \end{pmatrix}.$$

The symbol of $e^{-tA(\Omega)}$ and \mathbf{P} equal to

$$\sigma^*(e^{-A(\Omega)t}) = e^{-|\xi|^2 t} \sigma^*(e^{-t\Omega\mathbf{S}}) \quad \text{and} \quad \sigma^*(\mathbf{P}) = (\delta_{jk} - \frac{\xi_j \xi_k}{|\xi|^2}).$$

Thus the symbol of $\operatorname{div}e^{-(t-s)A(\Omega)}\mathbf{P}$ is holomorphic in (t, s) from $Z(T, \sigma) \rightarrow BUC(\mathbf{R}^3)$, since the symbol $\sigma^*(\operatorname{div}e^{-tA(\Omega)}\mathbf{P})$ can be extended to be continuous up to the origin. Moreover, $\sigma^*(\operatorname{div}e^{-(t-s)A(\Omega)}\mathbf{P}) \in H^\infty(Z_\varepsilon(T, \sigma), BUC(\mathbf{R}^3))$. Thus for $\mu \in M$, the associate measure $\mu[\sigma^*(\operatorname{div}e^{-(t-s)A(\Omega)}\mathbf{P})]$ is in $H^\infty(Z_\varepsilon(T, \sigma), M)$. We now conclude that $f \in H^\infty(Z_\varepsilon(T, \sigma), FM)$.

Next we show (10). Note that $\|\mathbf{P}u\|_{FM} \leq \|u\|_{FM}$ holds (see [12]). By [12] we have

$$\|e^{-t\Omega\mathbf{S}}\|_{FM \rightarrow FM} \leq \sup_{\xi \in \mathbf{R}^3} |\sigma^*(e^{-t\Omega\mathbf{S}})|,$$

where $|\cdot|$ denotes the operator norm of 3×3 matrix from the Euclidean space \mathbf{R}^3 into itself. Note that the symbol of $e^{-t\Omega\mathbf{S}}$ is expressed in (11) and it is unitary in real t . There exists a constant C^* depending only on Ω , σ and T satisfying

$$\|e^{-(t-s)\Omega\mathbf{S}}\|_{FM \rightarrow FM} \leq \sup_{\xi \in \mathbf{R}^3} |\sigma^*(e^{-\operatorname{Im}(t-s)\Omega\mathbf{S}})| \leq C^*$$

for all $(t, s) \in Z(\sigma, T)$, where $\operatorname{Im} z$ denotes the imaginary part of a complex number z .

Thus we have

$$\begin{aligned} \|f(t, s)\|_{FM} &\leq \|\operatorname{div}e^{(t-s)\Delta}\|_{FM \rightarrow FM} \|\mathbf{P}\|_{FM \rightarrow FM} \|e^{-(t-s)\Omega\mathbf{S}}(u_m \otimes u_m)\|_{FM} \\ &\leq C^* \|\operatorname{div}e^{(t-s)\Delta}\|_{FM \rightarrow FM} \|(u_m \otimes u_m)\|_{FM}. \end{aligned}$$

Since $F[\partial_{x_j} e^{t\Delta} g] = \hat{g} \cdot i\xi_j e^{-t|\xi|^2}$ for $g \in FM$, we have

$$\begin{aligned}\|\partial_{x_j} e^{t\Delta} g\|_{FM} &= \|i\xi_j e^{-t|\xi|^2}\|_{L^\infty} \|g\|_{FM} \\ &\leq \sup_{|\xi| \geq 0} \left| |\xi| e^{-\operatorname{Re} t |\xi|^2} \right| \|g\|_{FM} \leq \frac{1}{(2(\operatorname{Re} t)e)^{\frac{1}{2}}} \|g\|_{FM}\end{aligned}$$

(cf.[12]). Therefore we obtain

$$\|f(t, s)\|_{FM} \leq \frac{C^*}{(2(\operatorname{Re}(t-s))e)^{\frac{1}{2}}} \|u_m \otimes u_m\|_{FM} \leq \frac{C^*}{(2(\operatorname{Re}(t-s))e)^{\frac{1}{2}}} \|u_m\|_{FM}^2.$$

Thus, we have the desired estimate with $C = \frac{C^*}{\sqrt{2e}}$.

Proof of Theorem 1.1.

We use successive approximation method. Let $\{u_m\}_{m \in \mathbb{N}}$ be as follows ;

$$\begin{aligned}u_1(t) &= e^{-tA(\Omega)} u_0, \\ u_{m+1}(t) &= u_1(t) - \int_0^t e^{-(t-s)A(\Omega)} \mathbf{P} \operatorname{div}(u_m \otimes u_m)(s) ds.\end{aligned}$$

First, we prove $u_m \in H^\infty(S(T, \sigma), FM_0)$ by induction on m .

Since $e^{-tA(\Omega)}$ is an analytic semigroup, $u_1 \in H^\infty(S(T, \sigma), FM_0)$. Assume that $u_m \in H^\infty(S(T, \sigma), FM_0)$, then it follows immediately from Theorem 3.1 and Lemma 3.2 that $u_{m+1} \in H^\infty(S(T, \sigma), FM_0)$.

Next, we prove $\{u_m\}_{m \in \mathbb{N}}$ becomes a Cauchy sequence for sufficiently small T_0 . We now define

$$K_m(T_0) = \sup_{s \in S(T_0, \sigma)} \|u_m(s)\|_{FM}, \quad L_m(T_0) = \sup_{s \in S(T_0, \sigma)} \|u_m(s) - u_{m-1}(s)\|_{FM}.$$

Applying the above estimates, we have

$$K_{m+1}(T_0) \leq C^* \|u_0\|_{FM} + \left| \int_0^t \frac{C}{(\operatorname{Re}(t-s))^{\frac{1}{2}}} \|u_m\|_{FM}^2 ds \right|.$$

Since

$$\left| \int_0^t \frac{ds}{(\operatorname{Re}(t-s))^{\frac{1}{2}}} \right| = \frac{2|t|}{(\operatorname{Re} t)^{\frac{1}{2}}} < \frac{2T_0^{\frac{1}{2}}}{\cos \sigma},$$

we obtain $K_{m+1}(T_0) \leq C^* \|u_0\|_{FM} + C' K_m^2$, where $C' = \frac{2T_0^{\frac{1}{2}}}{\cos \sigma} C^*$. Now, we take T_0 so small that both $C^* \leq \frac{3}{2}$ and $C' \leq \frac{1}{6\|u_0\|_{FM}}$ holds. Under these assumptions, we have

$$K_{m+1}(T_0) \leq \frac{3}{2} \|u_0\|_{FM} + \frac{(K_m(T_0))^2}{6\|u_0\|_{FM}}$$

and one can easily check that $K_m(T_0) \leq 3\|u_0\|_{FM}$ holds for all $m \in \mathbb{N}$.

Next, we estimate $\|u_{m+1} - u_m\|_{FM}$. By the similar argument together with the uniform boundedness, we obtain

$$L_{m+1}(T_0) \leq \frac{36T_0^{\frac{1}{2}}C^*}{\cos \sigma} \|u_0\|_{FM} L_m(T_0).$$

It follows that if we take T_0 so small that $\frac{36T_0^{\frac{1}{2}}C^*}{\cos \sigma} \|u_0\|_{FM} < 1$, then $\{u_m\}_{m \in \mathbb{N}}$ becomes a Cauchy sequence in $H^\infty(S(T_0, \sigma), FM_0)$.

Since the solution in $[0, T_0]$ is unique, this implies that u is analytic with values in FM_0 . Since $u(s) \in FM_0$ for all $s \in (0, T)$, starting from $u(s)$ and conclude that u is analytic in some open interval $(s, s + T_0)$. Since u can be taken arbitrary, this implies that u is analytic in $(0, T)$. (Note that we need not to assume that u is continuous up to $t = 0$ and $t = T$. We may replace $[0, T]$ by $(0, T)$ to get analyticity in $(0, T)$.)

Proof of Theorem 1.2

It suffices to prove that the statement of Lemma 3.2 still holds even if we replace FM by BUC provided that we assume $\Omega = 0$. (The semigroup $e^{t\mathbf{S}}$ is not bounded in BUC .) The analyticity of $f(t, s)$ easily follows from the analyticity of the semigroup $e^{t\Delta}$ in BUC . To prove (10) we need an estimate for $\nabla e^{t\Delta}\mathbf{P}$ which is proved in the Appendix. The remaining proof is the same as that of Theorem 1.1.

Appendix.

We give a short direct proof of the key estimate for the derivative of the heat semigroup in the proof of Theorem 1.2.

Lemma A. There is a universal constant C such that

$$\|\nabla e^{t\Delta}\mathbf{P}f\|_\infty \leq C (\operatorname{Re} t)^{-1/2} \|f\|_\infty$$

for all $f \in L^\infty(\mathbf{R}^n)$ and $t \in \mathbf{C}$ with $\operatorname{Re} t > 0$.

Proof. It suffices to prove that

$$\|\partial_k e^{t\Delta} R_i R_j f\|_\infty \leq C (\operatorname{Re} t)^{-1/2} \|f\|_\infty, \quad (1 \leq i, j, k \leq n),$$

where $R_j = \partial_j(-\Delta)^{-1/2}$ is the Riesz operator and $\partial_k = \partial/\partial x_k$. Since

$$R_i R_j = \partial_i \partial_j (-\Delta)^{-1} = \partial_i \partial_j \int_0^\infty e^{s\Delta} ds,$$

we observe that

$$\partial_k e^{t\Delta} R_i R_j f = e^{t\Delta} \int_{\operatorname{Re} t}^\infty \partial_k \partial_i \partial_j e^{s\Delta} ds f + \partial_k \partial_i \partial_j e^{t\Delta} \int_0^{\operatorname{Re} t} e^{s\Delta} ds f.$$

Since

$$\|\partial_k \partial_i \partial_j e^{t\Delta} f\|_\infty \leq C_1 (\operatorname{Re} t)^{-3/2} \|f\|_\infty, \quad \|e^{t\Delta} f\|_\infty \leq \|f\|_\infty \text{ for } \operatorname{Re} t > 0$$

with some C_1 independent of t and $f \in L^\infty(\mathbf{R}^n)$, we have

$$\begin{aligned} \|\partial_k e^{t\Delta} R_i R_j f\| &\leq C_1 \int_{\operatorname{Re} t}^\infty s^{-3/2} ds \|f\|_\infty + C_1 (\operatorname{Re} t)^{-3/2} (\operatorname{Re} t) \|f\|_\infty \\ &\leq 3C_1 (\operatorname{Re} t)^{-1/2} \|f\|_\infty. \end{aligned}$$

□

Remark. Lemma A for $t > 0$ is proved in [10] by using a Hardy space estimate of the Gauss kernel found by Carpio [6]. A direct proof estimating integral kernels is found by Shibata and Shimizu [21]. The proof above is very short and elementary.

References

- [1] N. I. Akhiezer and I. M. Glazman, *Theory of linear operators in Hilbert space*. F. Unger Pub. Co., New York (1961).
- [2] L. Ahlfors, *Complex analysis. An introduction to the theory of analytic functions of one complex variable. Third edition*. International Series in Pure and Applied Mathematics. McGraw-Hill Book Co., New York (1978).
- [3] M. Berger, *Nonlinearity and functional analysis*. Academic Press, New York-London, (1977).
- [4] A. Babin, A. Mahalov and B. Nicolaenko, *Global regularity of the 3D Rotating Navier-Stokes Equations for resonant domains*. Indiana University Mathematics Journal, 48 (1999), 1133-1176.
- [5] A. Babin, A. Mahalov and B. Nicolaenko, *3D Navier-Stokes and Euler equations with initial data characterized by uniformly large vorticity*. Indiana University Mathematics Journal, 50 (2001), 1-35.

- [6] A. Carpio, *Large-time behaviour in incompressible Navier-Stokes equations*. SIAM J. Math. Anal., 27 (1996), 449-475.
- [7] C. Corduneanu, *Almost Periodic Functions*, Interscience Publishers, New York (1968).
- [8] C. Foias and R. Temam, *Some analytic and geometric properties of the solutions of the evolution Navier-Stokes equations*. J. Math. Pure Appl., 58 (1979), 339-366.
- [9] Y. Giga, *Time and spatial analyticity of solutions of the Navier-Stokes equations*, Comm. in Partial Differential Equations, 8 (1983), 929-948.
- [10] Y. Giga, K. Inui and S. Matsui, *On the Cauchy problem for the Navier-Stokes equations with nondecaying initial data*. Advances in fluid dynamics, 27-68, Quad. Mat., 4, Dept. Math., Seconda Univ. Napoli, Caserta, (1999).
- [11] Y. Giga, K. Inui, A. Mahalov and S. Matsui, *Navier-Stokes equations in a rotating frame in R^3 with initial data nondecaying at infinity*. Hokkaido Math. J., 35 (2006), 321-364.
- [12] Y. Giga, K. Inui, A. Mahalov and S. Matsui, *Uniform local solvability for the Navier-Stokes equations with the Coriolis force*. Methods Appl. Anal., 12 (2005), 381-393.
- [13] Y. Giga, K. Inui and A. Mahalov and J. Saal, *Global solvability of the Navier-Stokes equations in spaces based on sum-closed frequency sets*. Adv. Diff. Eq., 12 (2007), 721-736.
- [14] Y. Giga, A. Mahalov and B. Nicolaenko, *The Cauchy problem for the Navier-Stokes equations with spatially almost periodic initial data*. Hokkaido Univ. Preprint Series in Math., #683 (2004).
- [15] M. Hieber and O. Sawada, *The Navier-Stokes equations in \mathbb{R}^n with linearly growing initial data*. Arch. Rational. Mech. Anal., 175 (2005), 269-285.
- [16] G. Iooss, *Sur la deuxième bifurcation d'une solution stationnaire de systems du type Navier-Stokes equations*. Arch. Rational Mech. Anal., 64 (1977), 339-369.
- [17] A. Mahalov and B. Nicolaenko, *Global regularity of the 3D Navier-Stokes Equations with weakly aligned large initial vorticity*. Russian Math. Surveys, 58 (350), (2003), 287-318.
- [18] A. Mahalov, B. Nicolaenko, C. Bardos and F. Golse, *Regularity of Euler equations for a class of three-dimensional initial data*. Special Volume on 'Trends in Partial Differential Equations of Mathematical Physics', Nonlinear Analysis Series, Birkhauser-Verlag, 61 (2004), 161-185.
- [19] K. Masuda, *On the analyticity and the unique continuation theorem for solutions of the Navier-Stokes equation*. Proc. Japan Acad., 43 (1967), 827-832.
- [20] O. Sawada, *The Navier-Stokes flow with linearly growing initial velocity in the whole space*. Bol. Soc. Parana. Mat., (3) 22, (2004), 75-96.

- [21] Y. Shibata and S. Shimizu, *A decay property of the Fourier transform and its application to the Stokes problem*. J. Math. Fluid Mech., 3 (2001), 213–230.
- [22] K. Yosida, *Functional analysis. Sixth edition*. Springer-Verlag, Berlin-New York, (1980).