



Title	RANDOM PERTURBATIONS OF NON-SINGULAR TRANSFORMATIONS ON $[0; 1]$
Author(s)	IWATA, YUKIKO; OGIHARA, TOMOHIRO
Citation	Hokkaido University Preprint Series in Mathematics, 975, 1-18
Issue Date	2011-2-22
DOI	10.14943/84122
Doc URL	<a href="http://hdl.handle.net/2115/69782">http://hdl.handle.net/2115/69782</a>
Type	bulletin (article)
File Information	pre975.pdf



[Instructions for use](#)

# RANDOM PERTURBATIONS OF NON-SINGULAR TRANSFORMATIONS ON $[0, 1]$

YUKIKO IWATA<sup>1</sup>, TOMOHIRO OGIHARA<sup>2</sup>

ABSTRACT. We consider random perturbations of non-singular measurable transformations  $S$  on  $[0, 1]$ . By using the spectral decomposition theorem of Komornik and Lasota, we prove that the existence of the invariant densities for random perturbations of  $S$ . Moreover the densities for random perturbations with small noise strongly converges to the density for Perron-Frobenius operator corresponding to  $S$  with respect to  $L^1([0, 1])$ -norm.

AMS Classification: 34E10, 37A50, 37A30, 37H99, 60E05.

Key Words : random dynamical system, spectral decomposition theorem, random perturbations.

## 1. INTRODUCTION

It is known that every Markov process on a state space can be represented as a random dynamical system ([2]). There are many important Markov models in applications which are analysed as random dynamical systems. We focus on the following random dynamical system with additive noise : Let  $S : X \rightarrow X$  be a non-singular measurable transformation on a measurable space  $(X, \mathcal{B}, \lambda)$  and let  $(\Omega, \mathcal{F}, \mu)$  be a probability space. For a given random variable  $X_0$  and an i.i.d. sequence  $\{\xi_n\}_{n \geq 0}$  on  $\Omega$  with values in  $X$ , we define the following Markov process  $\{X_n\}_{n \geq 0}$  by

$$(1) \quad X_{n+1}(\omega) := S(X_n(\omega)) + \xi_n(\omega).$$

When  $X = \mathbb{R}$ , we call the above Markov process  $\{X_n(\omega)\}_{n \geq 0}$  *first-order nonlinear autoregressive model (NLAR(1))*. On the other hand, if we let  $Q(x, A)$  be a family of transition probabilities (from a point  $x \in X$  to a Borel set  $A \in \mathcal{B}$ ), then the Markov process on  $X$  defined by the transition probabilities  $Q(Sx, A)$  is called a *random perturbation* of the dynamical system  $(X, S)$ . In this paper, we consider NLAR(1) on  $[0, 1]$ , i.e. let  $X = [0, 1]$  for (1) and we identify  $X_n$  with  $X_n - [X_n]$  for all  $n \geq 0$ , where  $[x]$  is the largest integer less than or equal to  $x$ . Note that considering NLAR(1) on  $[0, 1]$  is coincident with considering a random perturbation of the dynamical system  $S$  on  $[0, 1]$  in our case.

A stability property of NLAR(1) can be derived from contraction assumptions by Lasota and Mackey ([15]) by using the spectral decomposition theorem of Komornik and Lasota (Theorem 2.5). This theorem is our main method in this paper. Vu Kuok Fong [5] and independently Sine [18] have

showed that the generalization of this spectral decomposition theorem of Komorník-Lasota is a simple corollary of the Jacobs-de Leeuw-Glicksberg theorem. We prove that for any non-singular transformation  $S : [0, 1] \rightarrow [0, 1]$ , there exists an invariant density of  $\{X_n\}_{n \geq 0}$  for NLAR(1) on  $[0, 1]$  by using the spectral decomposition theorem of Komorník-Lasota.

In this paper, we also study the limiting distribution of NLAR(1) on  $[0, 1]$  with small additive noise (or small perturbations of  $([0, 1], S)$ ) given by

$$(2) \quad X_{n+1}^\varepsilon(\omega) := S(X_n^\varepsilon(\omega)) + \varepsilon \xi_n(\omega) \pmod{1},$$

as  $\varepsilon \downarrow 0$ , where  $X_0^\varepsilon = X_0$ . Many authors observe the relation between deterministic dynamical systems and small perturbed random dynamical systems ([4],[6],[9],[11],[16]). For example, in [9], Katok and Kifer considered small random perturbations, where  $S$  is an endomorphism of the interval  $[0, 1]$  satisfying the conditions of Misiurewicz and small transition probabilities  $P^\varepsilon(x, A) = Q^\varepsilon(Sx, A)$  for sufficiently small  $\varepsilon > 0$ . They proved that the densities of  $X_n^\varepsilon$ -invariant measures  $\mu^\varepsilon$  converge weakly to a density of the invariant measure  $\mu_S$  corresponding to  $S$  as  $\varepsilon \rightarrow 0$  in  $L^1$  topology ([9]).

In [14], Lasota and Mackey showed that the density functions of  $\{X_n^\varepsilon\}_{n \geq 0}$  for NLAR(1) (on  $\mathbb{R}$ ) with small additive noise are given by

$$P_\varepsilon^n f(x) := \int_{\mathbb{R}} g(y) P_S f(x - \varepsilon y) dy,$$

where  $P_S$  is the Perron-Frobenius operator corresponding to  $S$ ,  $g$  is the density of  $\{\xi_n\}_{n \geq 0}$  and  $f$  is the density of  $X_0$ . They prove that

$$(3) \quad \lim_{\varepsilon \rightarrow 0} \|P_\varepsilon f - P_S f\|_{L^1(\mathbb{R})} = 0$$

for all  $f \in L^1(\mathbb{R})$  (see [14]). We obtain the same result for NLAR(1) on  $[0, 1]$ . Moreover since the existence of the densities of  $X_n^\varepsilon$ -invariant measures are guaranteed by the spectral decomposition theorem of Komorník-Lasota, under certain conditions, we prove that if there exists the limit  $f_*$  of the densities of  $X_n^\varepsilon$ -invariant measures in  $L^1$  as  $\varepsilon \downarrow 0$  then the limit function  $f_*$  is an invariant density corresponding to  $S$ . This implies that we gave the sufficient condition of the existence of an invariant density corresponding to  $S$ . On the other hand, in the sense of weak convergence of invariant probability measures for small random perturbations of a dynamical system  $S$ , the bounded variation case is first proved by Keller (see the condition S1 in [10]). Afterwards, Young and Baladi considered random perturbations of piecewise  $C^2$  expanding map  $S : [0, 1] \rightarrow [0, 1]$  for which there exists the unique invariant density  $f_*$ . Indeed, in [1], Young and Baladi proved that for any piecewise  $C^2$  expanding map which has no periodic turning points, there exists invariant densities of small random perturbations and they converges to the invariant density  $f_*$  corresponding to  $S$  with respect to  $L^1$ -norm as  $\varepsilon \rightarrow 0$  (see also [3]). In section 3, we can see that the spectral decomposition theorem of Komorník-Lasota and (3) hold for NLAR(1) on  $[0, 1]$  defined by

(1) with respect to intermittent maps  $S$  which have an infinite invariant density function.

## 2. MAIN THEOREMS

**2.1. Random perturbations of Dynamical systems.** Let  $(\Omega, \mathcal{F}, \mu)$  be a probability space, where  $\mathcal{F}$  denotes a Borel  $\sigma$ -field and  $\mu$  a probability measure. Let  $x_0, \xi_0, \xi_1, \dots$  be random variables on  $\Omega$  with values in  $[0, 1]$  and  $S : [0, 1] \rightarrow [0, 1]$  be a non-singular measurable transformation (i.e.  $\lambda(S^{-1}(A)) = 0$  for any Borel set  $A \subset [0, 1]$  with  $\lambda(A) = 0$ , where  $\lambda$  is the normalized Lebesgue measure on  $[0, 1]$ ).

Consider the following stochastic process defined by

$$(4) \quad x_{n+1}(\omega) = S(x_n(\omega)) + \xi_n(\omega) \pmod{1}$$

for each  $n \geq 0$ .

**Definition 2.1.** We say that a random dynamical system  $\{x_n\}_{n \geq 0}$  generated by (4) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$  is first-order nonlinear autoregressive model on  $[0, 1]$  (NLAR(1) on  $[0, 1]$ ) if the following conditions C1-C3 hold :

- C1:**  $x_0, \xi_0, \xi_1, \xi_2, \dots$  are independent random variables;
- C2:**  $x_0$  has the density function  $f_0 \in D$  (i.e.  $\mu(\{\omega : x_0(\omega) \in B\}) = \int_B f_0(x) dx$  for any Borel set  $B \subset [0, 1]$ ), where  $D := \{f \in L^1([0, 1]) : f \geq 0 \text{ and } \int_{[0,1]} f(x) dx = 1\}$ ;
- C3:** each  $\xi_n$  has the same density function  $g \in L^1(\mathbb{R})$  such that  $g \geq 0$ ,  
 $\text{supp}(g) := \overline{\{x \in [0, 1] : g(x) \neq 0\}} \subseteq [0, 1]$  and  $\int_{\mathbb{R}} g(x) dx = 1$ .

Under conditions C1-C3, there exists a Markov operator  $P : L^1([0, 1]) \rightarrow L^1([0, 1])$  such that

$$(5) \quad \mu_n(A) := \mu(\{\omega : x_n(\omega) \in A\}) = \int_A P^n f_0(x) dx$$

for all Borel set  $A$  on  $[0, 1]$  and  $n \geq 0$ .

**Proposition 2.2.** Let  $\{x_n\}_{n \geq 0}$  be a NLAR(1) on  $[0, 1]$  generated by (4) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$ . Then there exists a Markov operator  $P : L^1([0, 1]) \rightarrow L^1([0, 1])$  defined by

$$(6) \quad Pf(x) = \int_{[0,1]} f(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) dy,$$

which satisfies (5).

In our paper, the spectral decomposition theorem of Komorník and Lasota [13] plays a central role. We introduce the sufficient condition for this theorem :

**Definition 2.3.** Let  $(X, \mathcal{F}, \nu)$  be a finite measure space. A linear operator  $P : L^1(X, \nu) \rightarrow L^1(X, \nu)$  is constrictive if there exists  $\delta > 0$  and  $\kappa < 1$  such that for every  $f \in D$  there is an integer  $n_0(f)$  for which

$$(7) \quad \int_E P^n f(x) \nu(dx) \leq \kappa \quad \text{for all } n \geq n_0(f) \text{ and } E \text{ with } \nu(E) \leq \delta.$$

**Remark 2.4.** If the space  $(X, \mathcal{F}, \mu)$  is  $\sigma$ -finite, we can substitute the above condition by the following :

there exists  $\delta > 0$ ,  $\kappa < 1$  and a measurable set  $B$  with  $\nu(B) < \infty$  such that for every  $f \in D$  there is an integer  $n_0(f)$  for which

$$(8) \quad \int_{(X \setminus B) \cup E} P^n f(x) \nu(dx) \leq \kappa \quad \text{for all } n \geq n_0(f) \text{ and } E \text{ with } \nu(E) \leq \delta.$$

It is easy to see that this condition reduces to that of Definition 2.3 when  $X$  is finite and let  $X = B$ .

**Theorem 2.5.** (*spectral decomposition theorem* [13]) Let  $P : L^1(X, \mathcal{F}, \nu) \rightarrow L^1(X, \mathcal{F}, \nu)$  be a constrictive Markov operator. Then there is an integer  $r$ , non negative functions  $g_i \in D_0 := \{f \in L^1(X, \mathcal{F}, \nu) : \|f\|_{L^1} = 1, f \geq 0\}$  and  $k_i \in L^\infty(X, \mathcal{F}, \nu)$ ,  $i = 1, 2, \dots, r$  and a operator  $Q : L^1(X, \mathcal{F}, \nu) \rightarrow L^1(X, \mathcal{F}, \nu)$  such that for every  $f \in L^1(X, \mathcal{F}, \nu)$ ,  $Pf$  is represented by the form

$$(9) \quad Pf(x) = \sum_{i=1}^r \lambda_i(f) g_i(x) + Qf,$$

where

$$\lambda_i(f) = \int_X f(x) k_i(x) \nu(dx).$$

Moreover the functions  $g_i$  and the operator  $Q$  have the following properties:

- $g_i(x) g_j(x) = 0$  for all  $i \neq j$ .
- For each integer  $i$ , there exists an unique integer  $\sigma(i)$  such that  $Pg_i = g_{\sigma(i)}$ . Further  $\sigma(i) \neq \sigma(j)$  for  $i \neq j$ .
- $\lim_{n \rightarrow \infty} \|P^n Qf\| = 0$  for every  $f \in L^1(X, \mathcal{F}, \nu)$ .

**Remark 2.6.** The spectral decomposition theorem of Komorník and Lasota holds when the space  $(X, \mathcal{F}, \nu)$  is  $\sigma$ -finite space and Markov operator is constrictive.

**Remark 2.7.** If Theorem 2.5 holds for a Markov operator  $P$ , then there is an invariant density  $f^*$  defined by

$$f_* = \frac{1}{r} \sum_{i=1}^r g_i.$$

Indeed,

$$Pf_* = \frac{1}{r} \sum_{i=1}^r Pgi = \frac{1}{r} \sum_{i=1}^r gi = f_*.$$

Therefore  $Pf_* = f_*$ .

The following theorem is our main result.

**Theorem 2.8.** *The Markov operator  $P : L^1([0, 1]) \rightarrow L^1([0, 1])$  defined by (6) corresponding to a NLAR(1) on  $[0, 1]$  generated by (4) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$  is constrictive, that is, theorem 2.5 holds for  $P$ .*

Moreover when the density of noise  $g(x)$  is not zero for all  $x$ , we have the following result.

**Proposition 2.9.** *Let  $P : L^1([0, 1]) \rightarrow L^1([0, 1])$  be the Markov operator defined by (6) corresponding to a NLAR(1) on  $[0, 1]$  generated by (4) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$ . If  $g(x) > 0$  for all  $x \in [0, 1]$ , then there exists a unique  $f_* \in D$  such that  $Pf_* = f_*$  and*

$$\lim_{n \rightarrow \infty} \|P^n f - f_*\| = 0 \quad \text{for every } f \in D.$$

**Remark 2.10.** *A sequence  $\{P^n\}_{n \geq 1}$  satisfying (9) is called asymptotically periodic. Proposition 2.9 implies that  $r = 1$  for (9). In this case, the sequence  $\{P^n\}_{n \geq 1}$  is called asymptotically stable.*

**2.2. Small random perturbations of dynamical systems.** In this section, we observe limiting behaviour of density functions of a NLAR(1) on  $[0, 1]$  generated by (4) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$  parametrized by  $\varepsilon > 0$  as  $\varepsilon \rightarrow 0$ .

We consider the following first-order nonlinear autoregressive model  $\{x_n^\varepsilon\}_{n \geq 0}$  on  $[0, 1]$  with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$  parametrized by  $\varepsilon > 0$  :

$$(10) \quad x_{n+1}^\varepsilon(\omega) = S(x_n^\varepsilon(\omega)) + \varepsilon \xi_n(\omega) \quad \text{for } 0 < \varepsilon < 1,$$

where  $x_0^\varepsilon = x_0$ .

Since random variables  $\varepsilon \xi_n$  have the same density  $\frac{1}{\varepsilon} g(\frac{\cdot}{\varepsilon})$ , we have the Markov operator  $P_\varepsilon : L^1([0, 1]) \rightarrow L^1([0, 1])$  defined by

$$(11) \quad P_\varepsilon f(x) = \frac{1}{\varepsilon} \int_{[0, 1]} f(y) \left( \sum_{i=0}^1 g\left(\frac{x - S(y) + i}{\varepsilon}\right) \right) dy$$

which satisfies that  $f_{n+1}^\varepsilon = P_\varepsilon f_n^\varepsilon$ , where  $\{f_n^\varepsilon\}_{n \geq 0}$  is the sequence of the density function of  $x_n^\varepsilon$ . Since  $S$  is non-singular, there exists the Perron-Frobenius operator  $P_S : L^1([0, 1]) \rightarrow L^1([0, 1])$  with respect to  $S : [0, 1] \rightarrow$

$[0, 1]$ . Hence, if we let  $g_{x,i,\varepsilon}(y) := g\left(\frac{x+i-y}{\varepsilon}\right)$ , then we have that

$$\begin{aligned} P_\varepsilon f(x) &= \frac{1}{\varepsilon} \int_{[0,1]} f(y) \left( \sum_{i=0}^1 g_{x,i,\varepsilon}(S(y)) \right) dy \\ &= \frac{1}{\varepsilon} \int_{[0,1]} P_S f(y) \left( \sum_{i=0}^1 g_{x,i,\varepsilon}(y) \right) dy \\ &= \frac{1}{\varepsilon} \int_{[0,1]} P_S f(y) \left( \sum_{i=0}^1 g\left(\frac{x+i-y}{\varepsilon}\right) \right) dy \\ &= \sum_{i=0}^1 \int_{\left[\frac{x+i-1}{\varepsilon}, \frac{x+i}{\varepsilon}\right] \cap [0,1]} P_S f(x+i-\varepsilon y) g(y) dy \end{aligned}$$

by condition C3.

We should expect that in some sense  $\lim_{\varepsilon \rightarrow 0} P_\varepsilon f(x) = P_S f(x)$ .

Let  $\|f\|_\infty := \inf\{M : |f(x)| \leq M \text{ for } \lambda\text{-a.e. } x \in [0, 1]\}$ , where  $\lambda$  is the normalized Lebesgue measure on  $[0, 1]$ .

**Theorem 2.11.** *Let  $S : [0, 1] \rightarrow [0, 1]$  be a non-singular measurable transformation and  $P_\varepsilon$  be the Markov operator defined by (11) corresponding to a NLAR(1) on  $[0, 1]$  generated by (10) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$ . Suppose that  $\|P_S f\|_\infty < \infty$  for any continuous function  $f$  on  $[0, 1]$ . Then we have that*

$$(12) \quad \lim_{\varepsilon \rightarrow 0} \|P_\varepsilon f - P_S f\|_{L^1([0,1])} = 0$$

for all  $f \in L^1([0, 1])$ .

**Remark 2.12.** *There is a big class of dynamical systems  $S : [0, 1] \rightarrow [0, 1]$  satisfying  $\|P_S f\|_\infty < \infty$  for any continuous function  $f$  on  $[0, 1]$ . For example, piecewise monotonic maps (including unimodal maps) and piecewise convex maps satisfy the assumption of Theorem 2.11.*

It is obviously that  $\{P_\varepsilon^n\}_{n \geq 1}$  defined by (11) is asymptotically periodic for each  $\varepsilon > 0$ . Hence the function  $f_\varepsilon$  defined by

$$(13) \quad f_\varepsilon(x) = \frac{1}{r(\varepsilon)} \sum_{i=1}^{r(\varepsilon)} g_{i,\varepsilon}(x),$$

where  $r(\varepsilon)$  is a positive integer and  $g_{i,\varepsilon}(x)$  are density functions depending only on  $\varepsilon$ , satisfies that  $f_\varepsilon \in D$  and  $P_\varepsilon f_\varepsilon = f_\varepsilon$ . This implies that for each  $\varepsilon > 0$ , Markov operator  $P_\varepsilon$  has at least one invariant density.

**Corollary 2.13.** *Let  $S : [0, 1] \rightarrow [0, 1]$  be a non-singular measurable transformation,  $P_\varepsilon$  be the Markov operator defined by (11) corresponding to a NLAR(1) on  $[0, 1]$  generated by (10) with respect to  $(\Omega, [0, 1], S, x_0, \{\xi_n\}_{n \geq 0})$  and  $f_\varepsilon$  be an invariant density for  $P_\varepsilon$  defined by (13). Suppose that  $\|P_S f\|_\infty <$*

$\infty$  for any continuous function  $f$  on  $[0, 1]$ . If there exists an integrable function  $f_*$  on  $[0, 1]$  such that

$$\lim_{\varepsilon \rightarrow 0} \|f_\varepsilon - f_*\|_{L^1([0,1])} = 0,$$

then  $f_*$  is an invariant density for  $P_S$ , that is  $P_S f_* = f_*$ .

**Remark 2.14.** Corollary 2.13 holds for any continuous piecewise  $C^2$ , piecewise expanding map  $S : [0, 1] \rightarrow [0, 1]$  which has no periodic turning points. Indeed, by Theorem 1.1 in [3] (and see Theorem 3 in [1]), there exists a unique absolutely continuous invariant probability measure  $\mu_0 = f_* dx$  which satisfies that

$$\lim_{\varepsilon \rightarrow 0} \|f_\varepsilon - f_*\|_{L^1([0,1])} = 0.$$

### 3. EXAMPLES

It is obviously that Theorem 2.8 holds for all non-singular transformations. We give some examples of non-singular transformations which also satisfy the assumptions of Theorem 2.11.

**(1):  $m$ -adic transformation** [14].

Consider the transformation  $S : [0, 1] \rightarrow [0, 1]$  given by

$$Sx = mx \pmod{1},$$

where  $m \geq 1$  is an integer. Thus the Perron-Frobenius operator  $P_S : L^1([0, 1]) \rightarrow L^1([0, 1])$  corresponding to  $S$  is given by

$$P_S f(x) = \frac{1}{m} \sum_{i=0}^{m-1} f\left(\frac{i+x}{m}\right).$$

Since  $P_S \mathbf{1} = \mathbf{1}$ , the Borel measure on  $[0, 1]$  is invariant with respect to the  $m$ -adic transformation  $S$ . Moreover it is obviously that for any continuous function  $f$  on  $[0, 1]$ ,  $Pf(x)$  is equal to a continuous function, hence  $\|P_S f\|_\infty < \infty$ .

**(2): Maps with indifferent fixed points with infinite invariant measure** [19]

Let  $\alpha \in (0, \infty)$  be a real parameter and consider the one-parameter family of maps  $S_\alpha$  of the interval  $[0, 1]$  onto itself defined by


$$(14) \quad S_\alpha(x) := 2 \frac{e^{\alpha x} - 1}{e^\alpha - 1} \pmod{1}.$$

For every  $\alpha > 0$ ,  $S_\alpha$  is piecewise onto and  $C^\infty$ -class. When the parameter  $\alpha$  varies, the dynamics of the maps changes. Some properties of this family established in [17] are listed below :


- (1) For  $\alpha > 0$  with  $|S'_\alpha(0)| > 1$ ,  $S_\alpha$  is a piecewise expanding map (see Figure 1). Then there exists the unique absolutely continuous invariant probability measure with respect to the Lebesgue measure on  $[0, 1]$  by the Lasota-Yorke theorem.




- (2) For  $\alpha > 0$  with  $|S'_\alpha(0)| = 1$ ,  $S_\alpha$  admits an indifferent fixed point 0 (see Figure 2). For these maps, there is NO finite absolutely continuous invariant measure. However there exists a  $\sigma$ -finite infinite absolutely continuous invariant measure.
- (3) For  $\alpha > 0$  with  $|S'_\alpha(0)| < 1$ ,  $S_\alpha$  admits a stable fixed point 0 (see Figure 3). For these maps, almost all points converge to 0 by using the symbolic dynamics with 4-symbols(see [17] more details.). Therefore there is no absolutely continuous invariant measure with respect to the Lebesgue measure.



**Figure 1.**  $|S'_\alpha(0)| = 1.5$ ,  
 $\alpha \doteq 0.5502$ .



**Figure 2.**  $|S'_\alpha(0)| = 1$ ,  
 $\alpha \doteq 1.2564$ .



**Figure 3.**  $|S'_\alpha(0)| = 0.5$ ,  
 $\alpha \doteq 2.3366$ .

Next, we shall apply our results (Theorem 2.11) to this family. Because  $T_\alpha(0) = 0$ ,  $T_\alpha(1) = 2$ , where  $T_\alpha(x) := 2\frac{e^{\alpha x} - 1}{e^\alpha - 1}$  is monotonic continuous function for every  $\alpha > 0$ , there exists the unique point  $x_\alpha \in (0, 1)$  such that  $T_\alpha(x_\alpha) = 1$ . Let  $I_0 = [0, x_\alpha]$  and  $I_1 = [x_\alpha, 1]$ . Since  $C^\infty$ -extensions of the maps  $S_\alpha|_{I_0} : I_0 \rightarrow [0, 1]$  and  $S_\alpha|_{I_1} : I_1 \rightarrow [0, 1]$  are one-to-one and onto, there exist the local inverses  $u_{\alpha,j} = (S_\alpha|_{I_j})^{-1}$  for  $j = 0, 1$ , we get

$$(15) \quad u_{\alpha,j}(x) = \frac{1}{\alpha} \log\left(1 + \frac{e^\alpha - 1}{2}(x + j)\right).$$

Thus the Perron-Frobenius operator corresponding to  $S_\alpha$  is given by

$$(16) \quad P_{S_\alpha} f = f \circ u_{\alpha,0} \cdot u'_{\alpha,0} + f \circ u_{\alpha,1} \cdot u'_{\alpha,1}.$$

Therefore we have  $\|P_{S_\alpha} f(x)\|_\infty < \infty$  for any continuous function  $f$  on  $[0, 1]$ .

#### 4. PROOF

*Proof. Proof of Proposition 2.2*

We let the density of  $x_n$  be denoted by  $f_n \in D$  ( $n \geq 1$ ) and desire a relation connecting  $f_{n+1}$  and  $f_n$ .

We assume that  $f_n$  exists for some  $n \geq 0$ .

Let  $\bar{A} = A \setminus \{1\}$  for any Borel set  $A \subset [0, 1]$ . Note that since  $x_{n+1}(\Omega) \subset [0, 1)$  and  $S(x_n)$  and  $\xi_n$  are independent for all  $n \geq 0$ , we have that

$$\begin{aligned}
\text{(i): } & \mu(\{\omega \in \Omega : x_{n+1} \in A\}) = \mu(\{\omega \in \Omega : x_{n+1} \in \bar{A}\}), \\
\text{(ii): } & \bigcap_{i=0,1} \{\omega : S(x_n(\omega)) + \xi_n(\omega) \in \bar{A} + i\} \cap \{\omega : S(x_n(\omega)) + \xi_n(\omega) = 2\} = \phi, \\
\text{(iii): } & \\
& \mu(S(x_n(\omega)) + \xi_n(\omega) = 2) = \mu(S(x_n(\omega)) = 1 \text{ and } \xi_n(\omega) = 1) \\
& = \int_{S^{-1}(\{1\})} f_n(x) dx \int_{\{1\}} g(y) dy = 0.
\end{aligned}$$

From (i)-(iii), we have that for any Borel set  $A \subset [0, 1]$  and  $n \geq 0$ ,

$$\begin{aligned}
& \mu(\{\omega \in \Omega : x_{n+1} \in A\}) = \mu(\{\omega \in \Omega : x_{n+1} \in \bar{A}\}) \\
& = \mu(\{\omega \in \Omega : S(x_n(\omega)) + \xi_n(\omega) \pmod{1} \in \bar{A}\}) \\
& = \mu(\{\omega \in \Omega : S(x_n(\omega)) + \xi_n(\omega) \in \bar{A}\}) + \mu(\{\omega \in \Omega : S(x_n(\omega)) + \xi_n(\omega) \in \bar{A} + 1\}) \\
& \quad \left( + \mu(\{\omega \in \Omega : S(x_n(\omega)) + \xi_n(\omega) = 2\}) \quad \text{if } 0 \in A \right) \\
& = \int \int_{S(x)+y \in \bar{A}} f_n(x)g(y) dx dy + \int \int_{S(x)+y-1 \in \bar{A}} f_n(x)g(y) dx dy.
\end{aligned}$$

By a change of variables (see Lemma 5.2 in Appendix.), this can be written as

$$\begin{aligned}
\mu(\{\omega \in \Omega : x_{n+1} \in A\}) & = \int_{a \in \bar{A}} \left\{ \int_{B^0(a)} f_n(b)g(a - S(b)) db \right\} da \\
& \quad + \int_{a \in \bar{A}} \left\{ \int_{B^1(a)} f_n(b)g(a - S(b) + 1) db \right\} da,
\end{aligned}$$

where

$$B^0(a) := \{b \in [0, 1] : 0 \leq a - S(b) \leq 1\} = \{b \in [0, 1] : 0 \leq S(b) \leq a\}$$

and

$$B^1(a) := \{b \in [0, 1] : 0 \leq a - S(b) + 1 \leq 1\} = \{b \in [0, 1] : a \leq S(b) \leq 1\}$$

for each  $a \in [0, 1]$ . By condition C3, we have that

$$\begin{aligned}
g(x - S(y)) & = 0 & \text{for all } y \in \{b \in [0, 1] : x < S(b)\} = [0, 1] \setminus B^0(x) \\
g(x - S(y) + 1) & = 0 & \text{for all } y \in \{b \in [0, 1] : x > S(b)\} = [0, 1] \setminus B^1(x)
\end{aligned}$$

for each  $x \in [0, 1]$ . Hence we get that

$$\int_{[0,1] \setminus B^0(x)} f_n(y)g(x - S(y)) dy = 0 = \int_{[0,1] \setminus B^1(x)} f_n(y)g(x - S(y) + 1) dy$$

for each  $x \in [0, 1]$ . This implies that

$$\begin{aligned}
\int_{[0,1]} f_n(y)g(x - S(y)) dy & = \int_{B^0(x)} f_n(y)g(x - S(y)) dy \\
\int_{[0,1]} f_n(y)g(x - S(y) + 1) dy & = \int_{B^1(x)} f_n(y)g(x - S(y) + 1) dy.
\end{aligned}$$

Therefore we have that

$$\begin{aligned} \mu(\{\omega \in \Omega : x_{n+1} \in A\}) &= \int_{a \in \bar{A}} \int_{[0,1]} f_n(b)g(a - S(b))dbda \\ &\quad + \int_{a \in \bar{A}} \int_{[0,1]} f_n(b)g(a - S(b) + 1)dbda. \end{aligned}$$

Since  $\{1\}$  is a 1-point set and  $h(a) := \int_{b \in [0,1]} f_n(b)g(a - S(b) + i)db \in L^1([0, 1])$ , we have that for  $i = 0, 1$ ,

$$\int_{\{1\}} \left\{ \int_{[0,1]} f_n(b)g(a - S(b) + i)db \right\} da = \int_{\{1\}} h(a)da = 0.$$

Then we have that

$$\begin{aligned} \mu(\{\omega \in \Omega : x_{n+1} \in A\}) &= \sum_{i=0}^1 \int_{a \in \bar{A}} \int_{b \in [0,1]} f_n(b)g(a - S(b) + i)dbda \\ &= \sum_{i=0}^1 \int_{a \in A} \int_{b \in [0,1]} f_n(b)g(a - S(b) + i)dbda. \end{aligned}$$

Therefore using the fact that  $A$  was an arbitrary Borel set on  $[0, 1]$ , we get the density  $f_{n+1}$  of  $x_{n+1}$  defined by

$$f_{n+1}(x) = \sum_{i=0}^1 \int_{[0,1]} f_n(y)g(x - S(y) + i)dy \quad \text{a.e. } x \in [0, 1].$$

On the other hand, we get that

$$\int_{x \in [0,1]} \sum_{i=0}^1 g(x - S(y) + i)dx = \int_{[0,1]} g(x)dx = 1 \quad \text{for } \forall y \in [0, 1]$$

by condition C3. Then by Fubini's theorem, we have that

$$\begin{aligned} \int_{[0,1]} f_{n+1}(x)dx &= \sum_{i=0}^1 \int_{y \in [0,1]} \left\{ \int_{x \in [0,1]} f_n(y)g(x - S(y) + i)dx \right\} dy \\ &= \int_{y \in [0,1]} f_n(y)dy = 1. \end{aligned}$$

Moreover  $f_{n+1} \geq 0$  because of the positivity of  $g$  and  $f_n$ . Therefore if  $x_n$  has the density  $f_n \in D$ , then  $f_{n+1}$  also have to exist in  $D$ .

From this fact, we can define a linear operator  $P : L^1([0, 1]) \rightarrow L^1([0, 1])$  by

$$Pf(x) = \int_{y \in [0,1]} f(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) dy$$

which satisfies that

$$f_{n+1} = Pf_n \quad \text{a.e.}$$

for all  $n \geq 0$ . Next we shall show that  $P : L^1([0, 1]) \rightarrow L^1([0, 1])$  is a Markov operator, that is,  $P$  is a linear operator which satisfies that  $Pf \geq 0$  and  $\|Pf\|_{L^1([0,1])} = \|f\|_{L^1([0,1])}$  for any  $f \in L^1([0, 1])$  with  $f \geq 0$ . It is easy to see that  $P$  is a positive linear operator on  $L^1([0, 1])$  because  $g$  is positive. Moreover we have that for  $f \in L^1([0, 1])$  with  $f \geq 0$  by the Fubini's theorem,

$$\begin{aligned} \|Pf\|_{L^1([0,1])} &:= \int_{[0,1]} Pf(x)dx \\ &= \int_{x \in [0,1]} \int_{y \in [0,1]} f_n(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) dy dx \\ &= \int_{x \in [0,1]} \sum_{i=0}^1 g(x - S(y) + i) \left\{ \int_{[0,1]} f(y)dy \right\} dx \\ &= \int_{[0,1]} f(y)dy = \|f\|_{L^1([0,1])}. \end{aligned}$$

Therefore  $P$  is a Markov operator. □

*Proof. Proof of Theorem 2.8*

From the spectral decomposition theorem by Komorník and Lasota [14], it is enough to show that  $P$  is constrictive : there exists a  $\delta > 0$  and  $\kappa < 1$  such that for every  $f \in D$  there is an integer  $n_0(f)$  for which

$$\int_B P^n f(x)dx \leq \kappa \quad \text{for all } n \geq n_0(f) \text{ and } B \subset [0, 1] \text{ with } \lambda(B) \leq \delta,$$

where  $\lambda$  is the normalized Lebesgue measure on  $[0, 1]$ .

Since  $g$  is the integrable function on  $\mathbb{R}$  supported in  $[0, 1]$ , for any  $\varepsilon > 0$ , there exists  $0 < \delta(\varepsilon) \leq 1$  such that whenever  $\lambda(A) \leq \delta(\varepsilon)$ ,

$$\int_A g(x)dx \leq \varepsilon.$$

Take arbitrary  $0 < \varepsilon < 1$ , hence there exists  $\delta(\varepsilon) > 0$  which satisfies  $\int_A g(x)dx \leq \frac{\varepsilon}{2}$  for any Borel set  $A \subset [0, 1]$  with  $\lambda(A) \leq \delta(\varepsilon)$ . Thus we have that for each  $f \in D$  and  $n \geq 1$ ,

$$\begin{aligned} \int_A P^n f(x)dx &= \int_A \int_{[0,1]} P^{n-1} f(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) dy dx \\ &= \int_{[0,1]} \left\{ \sum_{i=0}^1 \int_{A-S(y)+i} g(x)dx \right\} P^{n-1} f(y)dy. \end{aligned}$$

Let  $\bar{\lambda}$  be the Lebesgue measure on  $\mathbb{R}$ . Since  $\bar{\lambda}(A - S(y) + i) = \bar{\lambda}(A) = \lambda(A) \leq \delta(\varepsilon)$  for each  $y \in [0, 1]$  and  $i = 0, 1$ , we obtain that

$$(17) \quad \int_A P^n f(x)dx \leq \varepsilon \int_{[0,1]} P^{n-1} f(y)dy = \varepsilon \quad \text{for all } n \geq 1,$$

which implies that  $P$  is constrictive. □

*Proof. Proof of Proposition 2.9*

From the theorem 5.6.1 in [14], it is enough to show that there exists a set  $A \subset [0, 1]$  of nonzero measure  $\lambda(A) > 0$  with the property that for every  $f \in D$ , there is an integer  $n_0(f)$  such that

$$(18) \quad P^n f(x) > 0 \quad \text{for a.e. } x \in A \quad \text{and} \quad \text{for all } n \geq n_0(f).$$

Let  $f \in D$  be arbitrary. From the assumption about  $g$ , there exists a positive number  $0 < \varepsilon < 1$  which satisfies that there exists  $\delta(\varepsilon) > 0$  such that for all  $\lambda(A) \leq \delta(\varepsilon)$ ,  $\int_A g(x) dx \leq \frac{\varepsilon}{2}$ . Take an arbitrarily  $0 < \delta < 1$  with  $1 - \delta < \delta(\varepsilon)$ . Since  $\lambda((\delta - S(y) + i, 1 - S(y) + i]) = 1 - \delta \leq \delta(\varepsilon)$  for each  $y \in [0, 1]$  and  $i = 0, 1$ , we have that

$$\begin{aligned} & \int_{\delta < x \leq 1} P^n f(x) dx \\ &= \int_{[0, 1]} \left\{ \sum_{i=0}^1 \int_{(\delta - S(y) + i, 1 - S(y) + i]} g(x) dx \right\} P^{n-1} f(y) dy \leq \varepsilon \end{aligned}$$

for all  $n \geq 1$ . From this inequality, we have that

$$(19) \quad \begin{aligned} \int_{0 \leq y \leq \delta} P^n f(y) dy &= \int_{[0, 1]} P^n f(y) dy - \int_{\delta < y \leq 1} P^n f(y) dy \\ &\geq 1 - \varepsilon > 0 \end{aligned}$$

for all  $n \geq 1$ .

On the other hand, we have that

$$(20) \quad \begin{aligned} P^{n+1} f(x) &= \int_{[0, 1]} P^n f(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) dy \\ &\geq \int_{0 \leq y \leq \delta} P^n f(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) dy. \end{aligned}$$

From the assumption about  $g$ , we have that

$$(21) \quad (g(x - S(y)) + g(x - S(y) + 1)) > 0 \quad \text{for all } x \in [0, 1] \text{ and } 0 \leq y \leq \delta.$$

From (19) and (21), we have that for a.e.  $x \in [0, 1]$ ,

$$P^n f(y) \left( \sum_{i=0}^1 g(x - S(y) + i) \right) \quad \text{for } n \geq 1$$

as a function of  $y$ , does not vanish in  $\{0 \leq y \leq \delta\}$ . As a consequence, inequality (20) implies (18) with respect to the set  $[0, 1]$ , thus completing the proof of the proposition. □

*Proof. Proof of Theorem 2.11*

Since the set of continuous functions on  $[0, 1]$  is dense in  $L^1([0, 1])$  and  $P_\varepsilon, P_S$  are Markov operators, it is enough to prove the theorem for continuous functions on  $[0, 1]$ . Indeed, for any  $f \in L^1([0, 1])$  and  $\eta > 0$ , there exists a continuous function  $f_\eta$  on  $[0, 1]$  such that  $\|f - f_\eta\|_{L^1([0,1])} \leq \eta$ . Thus if we have that  $\lim_{\varepsilon \rightarrow 0} \|P_\varepsilon f_\eta - P_S f_\eta\|_{L^1([0,1])} = 0$ , then we have that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \|P_\varepsilon f - P_S f\|_{L^1([0,1])} &= \lim_{\varepsilon \rightarrow 0} \|P_\varepsilon(f - f_\eta) - P_S(f - f_\eta) + P_\varepsilon f_\eta - P_S f_\eta\|_{L^1([0,1])} \\ &\leq 2\|f - f_\eta\|_{L^1([0,1])} + \lim_{\varepsilon \rightarrow 0} \|P_\varepsilon f_\eta - P_S f_\eta\|_{L^1([0,1])} \\ &\leq 2\eta. \end{aligned}$$

From the fact that  $\eta$  was an arbitrary positive number, we have that  $\lim_{\varepsilon \rightarrow 0} \|P_\varepsilon f - P_S f\|_{L^1([0,1])} = 0$ .

Fix an arbitrarily continuous function  $f$  on  $[0, 1]$ . We split the integral into two parts,

$$\begin{aligned} \|P_\varepsilon f - P_S f\|_{L^1([0,1])} &= \int_{[0,\varepsilon]} |P_\varepsilon f - P_S f| dx + \int_{(\varepsilon,1]} |P_\varepsilon f - P_S f| dx \\ &= C_1(\varepsilon) + C_2(\varepsilon) \quad \text{for } 0 < \varepsilon < 1. \end{aligned}$$

Firstly, we consider  $C_1(\varepsilon)$ . Let  $H_i(x, y) := P_S f(x+i-\varepsilon y)g(y)\mathbf{1}_{[\frac{x+i-1}{\varepsilon}, \frac{x+i}{\varepsilon}]}(y)$  for  $i = 0, 1$ . Note that the essential supremum of  $|P_S f|$  is finite (i.e.  $\|P_S f\|_\infty < \infty$ ) from the assumption about  $P_S f$ . Fix an arbitrarily point  $x_0 \in [0, 1]$ . Since

$$0 \leq x_0 + i - \varepsilon y \leq 1 \quad \text{for all } y \in \left[ \frac{x_0 + i - 1}{\varepsilon}, \frac{x_0 + i}{\varepsilon} \right],$$

we have that for each  $i = 0, 1$ ,

$$|P_S f(x_0 + i - \varepsilon y)| \leq \|P_S f\|_\infty \quad \text{for } \lambda\text{-a.e. } y \in \left[ \frac{x_0 + i - 1}{\varepsilon}, \frac{x_0 + i}{\varepsilon} \right].$$

Moreover we have that

$$[0, 1] \subset \bigcup_{i=\{0,1\}} \left[ \frac{x_0 + i - 1}{\varepsilon}, \frac{x_0 + i}{\varepsilon} \right] = \left[ \frac{x_0 - 1}{\varepsilon}, \frac{x_0}{\varepsilon} \right] \cup \left[ \frac{x_0}{\varepsilon}, \frac{x_0 + 1}{\varepsilon} \right]$$

for all  $0 < \varepsilon < 1$ . Then we have that,

$$\begin{aligned}
\left| \sum_{i=\{0,1\}} \int_{[0,1]} H_i(x_0, y) dy \right| &\leq \sum_{i=\{0,1\}} \int_{[0,1]} |P_S f(x_0 + i - \varepsilon y)| g(y) \mathbf{1}_{[\frac{x_0+i-1}{\varepsilon}, \frac{x_0+i}{\varepsilon}]}(y) dy \\
&\leq \sum_{i=\{0,1\}} \|P_S f\|_\infty \int_{[0,1]} g(y) \mathbf{1}_{[\frac{x_0+i-1}{\varepsilon}, \frac{x_0+i}{\varepsilon}]}(y) dy \\
&= \|P_S f\|_\infty \left\{ \int_{\bigcup_{i=0}^1 [\frac{x_0+i-1}{\varepsilon}, \frac{x_0+i}{\varepsilon}] \cap [0,1]} g(y) dy \right\} \\
(22) \qquad \qquad \qquad &= \|P_S f\|_\infty \left\{ \int_{[0,1]} g(y) dy \right\} = \|P_S f\|_\infty
\end{aligned}$$

by condition C3. Since  $x_0$  was an arbitrary point in  $[0, 1]$ , we have that

$$\begin{aligned}
\|P_\varepsilon f\|_{L^2([0,1])} &= \left( \int_{[0,1]} \left| \sum_{i=0}^1 \int_{[0,1]} H_i(x, y) dy \right|^2 dx \right)^{1/2} \\
&\leq \|P_S f\|_\infty < \infty.
\end{aligned}$$

This implies that the family  $\{P_\varepsilon f, 0 < \varepsilon < 1\}$  is uniformly integrable. Then we have that

$$(23) \qquad \lim_{\varepsilon \rightarrow 0} \sup_{0 < \eta < 1} \int_{[0, \varepsilon]} |P_\eta f| dx = 0$$

by Lemma 4.10 in [8]. Since

$$\int_{[0, \varepsilon]} |P_\varepsilon f| dx \leq \sup_{0 < \eta < 1} \int_{[0, \varepsilon]} |P_\eta f| dx \quad \text{for } 0 < \varepsilon < 1,$$

we have that

$$0 \leq \lim_{\varepsilon \rightarrow 0} \int_{[0, \varepsilon]} |P_\varepsilon f| dx \leq \overline{\lim}_{\varepsilon \rightarrow 0} \int_{[0, \varepsilon]} |P_\varepsilon f| dx \leq \overline{\lim}_{\varepsilon \rightarrow 0} \sup_{0 < \eta < 1} \int_{[0, \varepsilon]} |P_\eta f| dx = 0$$

by (23). Therefore we have that  $\lim_{\varepsilon \rightarrow 0} \int_{[0, \varepsilon]} |P_\varepsilon f| dx = 0$ . Moreover since the family  $\{P_S f\}$  consisting of only one function  $P_S f$  is obviously uniformly integrable, we also have that

$$\lim_{\varepsilon \rightarrow 0} \int_{[0, \varepsilon]} |P_S f| dx = 0.$$

Therefore we have that

$$(24) \qquad \lim_{\varepsilon \rightarrow 0} C_1(\varepsilon) \leq \lim_{\varepsilon \rightarrow 0} \int_{[0, \varepsilon]} |P_\varepsilon f| dx + \lim_{\varepsilon \rightarrow 0} \int_{[0, \varepsilon]} |P_S f| dx = 0.$$

Note that  $[0, 1] \subset [\frac{x-1}{\varepsilon}, \frac{x}{\varepsilon}]$  and  $[\frac{x}{\varepsilon}, \frac{x+1}{\varepsilon}] \subset (1, \infty)$  for each  $x \in (\varepsilon, 1]$ . Hence we have that

$$\begin{aligned} P_\varepsilon f(x) &= \sum_{i=0}^1 \int_{[\frac{x+i-1}{\varepsilon}, \frac{x+i}{\varepsilon}]} P_S f(x+i-\varepsilon y) g(y) dy \\ &= \int_{[0,1]} P_S f(x-\varepsilon y) g(y) dy. \end{aligned}$$

Thus we have that with respect to  $C_2(\varepsilon)$ ,

$$\begin{aligned} C_2(\varepsilon) &= \int_{(\varepsilon,1]} \left| \int_{[0,1]} P_S f(x-\varepsilon y) g(y) dy - P_S f(x) \right| dx \\ &= \int_{(\varepsilon,1]} \left| \int_{[0,1]} [P_S f(x-\varepsilon y) - P_S f(x)] g(y) dy \right| dx. \end{aligned}$$

Without loss of generality, we can assume that  $P_S f(x) = 0$  for all  $x \notin [0, 1]$  (for example set  $S(x) = x$ ,  $f(x) = 0$  for all  $x \notin [0, 1]$ ). Since  $P_S f$  is an integrable function and the set  $\{P_S f\}$  is compact in  $L^1(\mathbb{R})$ , we have that for an arbitrarily small  $\delta > 0$ , there exists  $\varepsilon_0$  such that for all  $\varepsilon \leq \varepsilon_0$ ,

$$\int_{[0,1]} |P_S f(x-\varepsilon y) - P_S f(x)| dx \leq \delta$$

for each  $y \in [0, 1]$ . Thus we have that

$$\begin{aligned} C_2(\varepsilon) &\leq \int_{[0,1]} \int_{[0,1]} |P_S f(x-\varepsilon y) - P_S f(x)| g(y) dy dx \\ &\leq \delta \int_{[0,1]} g(y) dy = \delta. \end{aligned}$$

Therefore  $\lim_{\varepsilon \rightarrow 0} C_2(\varepsilon) = 0$ . Then theorem is proved.  $\square$

*Proof. Proof of Corollary 2.13*

Since  $P_\varepsilon$  is the Markov operator, we have that

$$\|P_\varepsilon(f_* - f_\varepsilon)\|_{L^1([0,1])} \leq \|f_* - f_\varepsilon\|_{L^1([0,1])}.$$

Hence we have that

$$\begin{aligned} \|P_\varepsilon f_* - f_*\|_{L^1([0,1])} &= \|f_\varepsilon + P_\varepsilon(f_* - f_\varepsilon) - f_*\|_{L^1([0,1])} \\ &\leq \|f_\varepsilon - f_*\|_{L^1([0,1])} + \|P_\varepsilon(f_* - f_\varepsilon)\|_{L^1([0,1])} \\ &\leq 2\|f_\varepsilon - f_*\|_{L^1([0,1])} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Thus  $P_\varepsilon f_*$  converges to  $f_*$  in  $L^1([0, 1])$ -norm. On the other hand, from Theorem 2.11,  $P_\varepsilon f_*$  converges to  $P_S f_*$  in  $L^1([0, 1])$ -norm. Therefore  $P_S f_* = f_*$ .  $\square$



## 5. APPENDIX

In this section, we give a supplementary explanation of the change of variables theorem for the Lebesgue integral on  $\mathbb{R}$  which is applied in the proof of Proposition 2.2.

**Lemma 5.1.** ([7]) *If  $h(t) \geq 0$  is an integrable function on  $[\alpha, \beta]$  such that there exists a increasing function  $H(t)$  satisfying  $H(t) = \int_c^t h(t)dt$ , where  $c$  is a constant. Let  $a = H(\alpha), b = H(\beta)$ . Then we have that*

$$\int_a^b f(x)dx = \int_\alpha^\beta f(H(t))h(t)dt$$

for all integrable function  $f$  defined on  $[a, b]$ .

By using Lemm 5.1, we prove the following lemma.

**Lemma 5.2.** *Let  $X$  and  $Y$  are independent random variables on a probability space  $(\Omega, \mathcal{F}, \mu)$  with values in  $[0, 1]$  which satisfy the followings:*

- (1)  *$X$  has the density function  $f : [0, 1] \rightarrow \mathbb{R}$  with  $f \geq 0$  such that*

$$\int_{[0,1]} f(x)dx = 1,$$

- (2)  *$Y$  has the density function  $g : \mathbb{R} \rightarrow \mathbb{R}$  with  $g \geq 0$  such that*

$$\text{supp}(g) := \overline{\{x \in \mathbb{R} : g(x) \neq 0\}} \subset [0, 1] \quad \text{and} \quad \int_{[0,1]} g(x)dx = 1.$$

Then we have that for any Borel set  $A \subset [0, 1]$ ,

$$\mu(\{\omega \in \Omega : X(\omega) + Y(\omega) \in A\}) = \int_{x \in A} \int_{y \in B(x)} f(y)g(x - y)dydx,$$

where  $B(x) = \{y \in [0, 1] : 0 \leq x - y \leq 1\}$  for each  $x \in [0, 1]$ .

*Proof.* Since  $X$  and  $Y$  are independent,

$$\mu(\{\omega \in \Omega : X(\omega) + Y(\omega) \in A\}) = \int \int_{\{(x,y) \in [0,1] \times [0,1] : x+y \in A\}} f(x)g(y)dx dy.$$

Since  $f$  and  $g$  are positive integrable functions on  $[0, 1]$ , we have

$$\int \int_{\{(x,y) \in [0,1] \times [0,1] : x+y \in A\}} f(x)g(y)dx dy < \infty,$$

so, we can apply the Fubini's theorem to this integral. Indeed, we have that

$$\int \int_{\{(x,y) \in [0,1] \times [0,1] : x+y \in A\}} f(x)g(y)dx dy = \int_{x \in [0,1]} \int_{\{y \in [0,1] : x+y \in A\}} f(x)g(y)dy dx.$$

Let  $a := x + y$  and  $Z(a) := a - x$  for fixed  $x \in [0, 1]$ . Since  $Z(a)$  is absolutely continuous (i.e.  $Z(a) = \int_x^a \mathbf{1}(t)dt$ ), we have that by Lemma 5.1 and Fubini's theorem, we have that

$$\begin{aligned} & \int_{x \in [0, 1]} \int_{\{y \in [0, 1]: x+y \in A\}} f(x)g(y)dydx \\ &= \int_{x \in [0, 1]} \int_{\{a \in A: 0 \leq a-x \leq 1\}} f(x)g(a-x)dadx \quad (\text{change of variables}) \\ &= \int_{x \in [0, 1]} \int_{\{a \in [0, 1]: 0 \leq a-x \leq 1\}} f(x)g(a-x)\mathbf{1}_A(a)dadx \\ &= \int_{a \in [0, 1]} \int_{\{x \in [0, 1]: 0 \leq a-x \leq 1\}} f(x)g(a-x)\mathbf{1}_A(a)dxda \quad (\text{Fubini's theorem}) \\ &= \int_{a \in A} \int_{x \in B(a)} f(x)g(a-x)dxda. \end{aligned}$$

Therefore we have that

$$\mu(\{\omega \in \Omega : X(\omega) + Y(\omega) \in A\}) = \int_{a \in A} \int_{x \in B(a)} f(x)g(a-x)dxda.$$

□

#### ACKNOWLEDGEMENTS

It is a pleasure to thank Professor Michiko Yuri for her helping to improve the exposition and for many helpful and pleasant conversations.

#### REFERENCES

#### REFERENCES

- [1] Baladi V and Young L-S 1993 *On the spectra of randomly perturbed expanding maps*, Comm. Math. Phys. **156** 355–385
- [2] Bhattacharya R and Majumdar M 2007 *Random Dynamical Systems. Theory and Applications*(Cambridge: Cambridge University Press) p 463
- [3] Blank M and Keller G 1981 *Random perturbations of chaotic dynamical systems: stability of the spectrum*, Nonlinearity. **11** no. 5, 74–96
- [4] Cowieson W and Young L-S 2005 *SRB measures as zero-noise limits*, Ergodic Theory Dynam. Systems. **25** no. 4, 1115–1138
- [5] Vu Kuok Fong 1986 *Asymptotic almost periodicity and compactifying representations of semigroups*, Ukrain. Mat. Zh **38** 688–692
- [6] Freidlin M I and Wentzell A D 1993 *Random Perturbations of Dynamical Systems*(New York: Springer-Verlag) p 430
- [7] Jones F 1993 *Lebesgue Integration on Euclidean Space*(Boston: Jones and Bartlett Publishers) P 588
- [8] Kallenberg O 1997 *Foundations of modern probability, Probability and its Applications*(New York: Springer-Verlag) p 638
- [9] Katok A and Kifer Y 1986 *Random perturbations of transformations of an interval*, J. Analyse Math. **47** 193–237
- [10] Keller G 1982 *Stochastic stability in some chaotic dynamical systems*, Monatsh. Math. **94** 313–333

- [11] Kifer Y 1981 *The exit problem for small random perturbations of dynamical systems with a hyperbolic fixed point*, Israel J. Math. **40** no. 1, 74–96
- [12] Kifer Y 1998 *Limit theorems for random transformations and processes in random environments*, Trans. Amer. Math. Soc. **350** 1481–1518
- [13] Komornik J and Lasota A 1987 *Asymptotic decomposition of Markov operators*, Bull. Polish Acad. Sci. Math. **35** no. 5-6, 321–327
- [14] Lasota A and Mackey M 1994 *Chaos, fractals, and noise*(New York: Springer-Verlag) p 472
- [15] Lasota A and Mackey M 1987 *Noise and statical periodicity*, Physica D. **28** 143–154
- [16] Lin K 2005 *Convergence of invariant densities in the small-noise limit*, Nonlinearity. **18** no.2, 659–683
- [17] Ogiwara T *Statistical properties of a one-parameter family of interval maps exhibiting non-hyperbolic phenomena*, In preparation
- [18] Sine R 1991 *Constricted systems*, Rocky Mountain J. Math. **21** 1373–1383
- [19] Thaler M 2000 *The asymptotics of the Perron-Frobenius operator of a class of interval maps preserving infinite measures*, Studia Math. **143** no.2, 103–119

Authors' addresses:

<sup>1</sup>Iwata: Laboratory of Nonlinear Studies and Computation Research Institute for Electronic Science, Hokkaido University, Kita 12, Nishi 6, Sapporo, 060-0812, JP  
e-mail: iwata@math.sci.hokudai.ac.jp

<sup>2</sup>Ogiwara: Department of Mathematics, Hokkaido University Kita 10, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810, JP  
e-mail: ogi@math.sci.hokudai.ac.jp