



Title	KOECHER-MAASS SERIES OF A CERTAIN HALF-INTEGRAL WEIGHT MODULAR FORM RELATED TO THE DUKE-IMAMOGLU-IKEDA LIFT
Author(s)	Katsurada, Hidenori; Kawamura, Hisa-aki
Citation	Hokkaido University Preprint Series in Mathematics, 1038, 1-41
Issue Date	2013-6-28
DOI	10.14943/84182
Doc URL	http://hdl.handle.net/2115/69842
Type	bulletin (article)
File Information	pre1038.pdf



[Instructions for use](#)

KOECHER-MAASS SERIES OF A CERTAIN HALF-INTEGRAL WEIGHT MODULAR FORM RELATED TO THE DUKE-IMAMOĞLU-IKEDA LIFT

HIDENORI KATSURADA AND HISA-AKI KAWAMURA

ABSTRACT. Let k and n be positive even integers. For a cuspidal Hecke eigenform h in the Kohnen plus space of weight $k - n/2 + 1/2$ for $\Gamma_0(4)$, let f be the corresponding primitive form of weight $2k - n$ for $SL_2(\mathbf{Z})$ under the Shimura correspondence, and $I_n(h)$ the Duke-Imamoğlu-Ikeda lift of h to the space of cusp forms of weight k for $Sp_n(\mathbf{Z})$. Moreover, let $\phi_{I_n(h),1}$ be the first Fourier-Jacobi coefficient of $I_n(h)$ and $\sigma_{n-1}(\phi_{I_n(h),1})$ be the cusp form in the generalized Kohnen plus space of weight $k - 1/2$ corresponding to $\phi_{I_n(h),1}$ under the Ibukiyama isomorphism. We then give an explicit formula for the Koecher-Maass series $L(s, \sigma_{n-1}(\phi_{I_n(h),1}))$ of $\sigma_{n-1}(\phi_{I_n(h),1})$ expressed in terms of the usual L -functions of h and f .

1. INTRODUCTION

Let l be an integer or a half integer, and let F be a modular form of weight l for the congruence subgroup $\Gamma_0^{(m)}(N)$ of the symplectic group $Sp_m(\mathbf{Z})$. Then the Koecher-Maass series $L(s, F)$ of F is defined as

$$L(s, F) = \sum_A \frac{c_F(A)}{e(A)(\det A)^s},$$

where A runs over a complete set of representatives for the $SL_m(\mathbf{Z})$ -equivalence classes of positive definite half-integral matrices of degree m , $c_F(A)$ is the A -th Fourier coefficient of F , and $e(A)$ denotes the order of the special orthogonal group of A . We note that $L(s, F)$ can also be obtained by the Mellin transform of F , and thus, its analytic properties are relatively known. (As for this, see Maass [19] and Arakawa [1, 2, 3].) Now we are interested in an explicit form of the Koecher-Maass series for a specific choice of F . In particular, whenever F is a certain lift of an elliptic modular form h of either integral or half-integral weight, we may hope to express $L(s, F)$ in terms of certain Dirichlet series related to h . Indeed, this expectation is verified in the case where F is a lift of h such that the weight l is an integer (cf. [8, 9, 10]). In this paper, we discuss a similar problem for a lift of elliptic modular forms to half-integral weight Siegel modular forms.

2010 *Mathematics Subject Classification*. Primary 11F67; Secondary 11F46.

Key words and phrases. Koecher-Maass series, Duke-Imamoğlu-Ikeda lift.

Let us explain our main result briefly. Let k and n be positive even integers. For a cuspidal Hecke eigenform h in the Kohnen plus space of weight $k - n/2 + 1/2$ for $\Gamma_0(4)$, let f be the primitive form of weight $2k - n$ for $SL_2(\mathbf{Z})$ corresponding to h under the Shimura correspondence, and let $I_n(h)$ be the Duke-Imamog̃lu-Ikeda lift of h (or of f) to the space of cusp forms of weight k for $Sp_n(\mathbf{Z})$. We note that $I_2(h)$ is nothing but the Saito-Kurokawa lift of h . Let $\phi_{I_n(h),1}$ be the first coefficient of the Fourier-Jacobi expansion of $I_n(h)$ and $\sigma_{n-1}(\phi_{I_n(h),1})$ the cusp form in the generalized Kohnen plus space of weight $k - 1/2$ for $\Gamma_0^{(n-1)}(4)$ corresponding to $\phi_{I_n(h),1}$ under the Ibukiyama isomorphism σ_{n-1} . (For the precise definitions of the Duke-Imamog̃lu-Ikeda lift, the generalized Kohnen plus space and the Ibukiyama isomorphism, see Section 2.) Then our main result expresses $L(s, \sigma_{n-1}(\phi_{I_n(h),1}))$ in terms of $L(s, h)$ and $L(s, f)$ (cf. Theorem 2.1). To prove Theorem 2.1, for a fundamental discriminant d_0 and a prime number p we define certain formal power series $P_{n-1,p}^{(1)}(d_0, \varepsilon^l, X, t) \in \mathbf{C}[X, X^{-1}][[t]]$ associated with some local Siegel series appearing in the p -factor of the Fourier coefficient of $\sigma_{n-1}(\phi_{I_n(h),1})$. Here ε is the Hasse invariant defined on the set of nondegenerate symmetric matrices with entries in \mathbf{Q}_p . We then rewrite $L(s, \sigma_{n-1}(\phi_{I_n(h),1}))$ in terms of the Euler products $\prod_p P_{n-1,p}^{(1)}(d_0, \varepsilon^l, \beta_p, p^{-s+k/2+n/4-1/4})$ with $l = 0, 1$, where β_p is the Satake p -parameter of f (cf. Theorem 3.2). By using a method similar to those in [9, 10], in Section 4, we get an explicit formula for the formal power series $P_{n-1,p}^{(1)}(d_0, \varepsilon^l, X, t)$ (cf. Theorem 4.4.1), which yields the desired formula for $L(s, \sigma_{n-1}(\phi_{I_n(h),1}))$ immediately. The result is very simple, and the proof proceeds similarly to the one of [10], where we gave an explicit formula for the Koecher-Maass series of the Siegel Eisenstein series of integral weight. However, it is more elaborate than the preceding one. For instance, we should be careful in dealing with the argument for $p = 2$, which divides the level of $\sigma_{n-1}(\phi_{I_n(h),1})$. We also note that the method in this paper is useful for giving an explicit formula for the Rankin-Selberg series of $\sigma_{n-1}(\phi_{I_n(h),1})$, and as a consequence, we can prove a conjecture of Ikeda [13] concerning the period of the Duke-Imamog̃lu-Ikeda lift, which we will discuss in [15].

Notation. Let R be a commutative ring. We denote by R^\times and R^* the semigroup of non-zero elements of R and the unit group of R , respectively. We also put $S^\square = \{a^2 \mid a \in S\}$ for a subset S of R . We denote by $M_{ml}(R)$ the set of $m \times l$ matrices with entries in R . In particular put $M_m(R) = M_{mm}(R)$. Put $GL_m(R) = \{A \in M_m(R) \mid \det A \in R^*\}$, and $SL_m(R) =$

$\{A \in GL_m(R) \mid \det A = 1\}$, where $\det A$ denotes the determinant of a square matrix A . For an $m \times l$ matrix X and an $m \times m$ matrix A , we write $A[X] = {}^tXAX$, where tX denotes the transpose of X . Let $S_m(R)$ denote the set of symmetric matrices of degree m with entries in R . Furthermore, if R is an integral domain of characteristic different from 2, let $\mathcal{L}_m(R)$ denote the set of half-integral matrices of degree m over R , that is, $\mathcal{L}_m(R)$ is the subset of symmetric matrices of degree m with entries in the field of fractions of R , whose (i, j) -th entry belongs to R or $\frac{1}{2}R$ according as $i = j$ or not. In particular, we put $\mathcal{L}_m = \mathcal{L}_m(\mathbf{Z})$ and $\mathcal{L}_{m,p} = \mathcal{L}_m(\mathbf{Z}_p)$ for a prime number p . For a subset S of $M_m(R)$ we denote by S^\times the subset of S consisting of all nondegenerate matrices. If S is a subset of $S_m(\mathbf{R})$ with \mathbf{R} , the field of real numbers, we denote by $S_{>0}$ (resp. $S_{\geq 0}$) the subset of S consisting of positive definite (resp. semi-positive definite) matrices. The group $GL_m(R)$ acts on the set $S_m(R)$ as $GL_m(R) \times S_m(R) \ni (g, A) \mapsto {}^tgAg \in S_m(R)$. Let G be a subgroup of $GL_m(R)$. For a G -stable subset \mathcal{B} of $S_m(R)$, we denote by \mathcal{B}/G the set of equivalence classes of \mathcal{B} under the action of G . We sometimes identify \mathcal{B}/G with a complete set of representatives of \mathcal{B}/G . We abbreviate $\mathcal{B}/GL_m(R)$ as \mathcal{B}/\sim if there is no fear of confusion. For a given ring R' , two symmetric matrices A and A' with entries in R are said to be *equivalent over R'* to each other and write $A \sim_{R'} A'$ if there is an element X of $GL_m(R')$ such that $A' = A[X]$. We also write $A \sim A'$ if there is no fear of confusion. For square matrices X and Y we write $X \perp Y = \begin{pmatrix} X & O \\ O & Y \end{pmatrix}$.

For an integer D with $D \equiv 0$ or $1 \pmod{4}$, let \mathfrak{d}_D be the discriminant of $\mathbf{Q}(\sqrt{D})$, and put $\mathfrak{f}_D = \sqrt{\frac{D}{\mathfrak{d}_D}}$. We call D a *fundamental discriminant* if it is either 1 or the discriminant of some quadratic field extension of \mathbf{Q} . For a fundamental discriminant D , let $\left(\frac{D}{*}\right)$ be the character corresponding to $\mathbf{Q}(\sqrt{D})/\mathbf{Q}$. Here we make the convention that $\left(\frac{D}{*}\right) = 1$ if $D = 1$.

We put $\mathbf{e}(x) = \exp(2\pi\sqrt{-1}x)$ for $x \in \mathbf{C}$. For a prime number p we denote by $\nu_p(*)$ the additive valuation of \mathbf{Q}_p normalized so that $\nu_p(p) = 1$, and by $\mathbf{e}_p(*)$ the continuous additive character of \mathbf{Q}_p such that $\mathbf{e}_p(x) = \mathbf{e}(x)$ for $x \in \mathbf{Q}$.

2. MAIN RESULT

Put $J_m = \begin{pmatrix} O_m & -1_m \\ 1_m & O_m \end{pmatrix}$, where 1_m and O_m denotes the unit matrix and the zero matrix of degree m , respectively. Furthermore, put $\Gamma^{(m)} = Sp_m(\mathbf{Z}) = \{M \in GL_{2m}(\mathbf{Z}) \mid J_m[M] = J_m\}$. Let l be an integer or a half integer. For

a congruence subgroup Γ of $\Gamma^{(m)}$, we denote by $\mathfrak{M}_l(\Gamma)$ the space of holomorphic modular forms of weight l for Γ . We denote by $\mathfrak{S}_l(\Gamma)$ the subspace of $\mathfrak{M}_l(\Gamma)$ consisting of all cusp forms. For a positive integer N , put $\Gamma_0^{(m)}(N) = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma^{(m)} \mid C \equiv O_m \pmod{N} \right\}$. Let $F(Z)$ be an element of $\mathfrak{M}_l(\Gamma_0^{(m)}(N))$. Then $F(Z)$ has the following Fourier expansion:

$$F(Z) = \sum_{A \in (\mathcal{L}_m)_{\geq 0}} c_F(A) \mathbf{e}(\mathrm{tr}(AZ)),$$

where $\mathrm{tr}(X)$ denotes the trace of a matrix X . We then define the Koecher-Maass series $L(s, F)$ of F as

$$L(s, F) = \sum_{A \in (\mathcal{L}_m)_{>0}/SL_m(\mathbf{Z})} \frac{c_F(A)}{e(A)(\det A)^s},$$

where $e(A) = \#\{X \in SL_m(\mathbf{Z}) \mid A[X] = A\}$. We note that $L(s, F)$ is nothing but Hecke's L -function of F if $m = 1$ and l is an integer.

Now put

$$\mathcal{L}'_m = \{A \in \mathcal{L}_m \mid A \equiv -{}^t r r \pmod{4\mathcal{L}_m} \text{ for some } r \in \mathbf{Z}^m\}.$$

For $A \in \mathcal{L}'_m$, the integral vector $r \in \mathbf{Z}^m$ in the above definition is uniquely determined modulo $2\mathbf{Z}^m$ by A , and is denoted by r_A . Moreover it is easily shown that the matrix

$$\begin{pmatrix} 1 & r_A/2 \\ {}^t r_A/2 & ({}^t r_A r_A + A)/4 \end{pmatrix},$$

which will be denoted by $A^{(1)}$ in the sequel, belongs to \mathcal{L}_{m+1} , and that its $SL_{m+1}(\mathbf{Z})$ -equivalence class is uniquely determined by A . Suppose that l is a positive even integer. We define the *generalized Kohnen plus space* of weight $l - 1/2$ for $\Gamma_0^{(m)}(4)$ as

$$\mathfrak{M}_{l-1/2}^+(\Gamma_0^{(m)}(4)) = \{F \in \mathfrak{M}_{l-1/2}(\Gamma_0^{(m)}(4)) \mid c_F(A) = 0 \text{ unless } A \in \mathcal{L}'_m\},$$

and put $\mathfrak{S}_{l-1/2}^+(\Gamma_0^{(m)}(4)) = \mathfrak{M}_{l-1/2}^+(\Gamma_0^{(m)}(4)) \cap \mathfrak{S}_{l-1/2}(\Gamma_0^{(m)}(4))$. Then there exists an isomorphism from the space of Jacobi forms of index 1 to the generalized Kohnen plus space due to Ibukiyama. To explain this, let $\Gamma_J^{(m)} = \Gamma^{(m)} \times H_m(\mathbf{Z})$ be the Jacobi group, where $\Gamma^{(m)}$ is identified with its image inside $\Gamma^{(m+1)}$ via the natural embedding

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \mapsto \left(\begin{array}{c|c} 1 & 0 \\ \hline A & B \\ 0 & 1 \\ \hline C & D \end{array} \right)$$

and

$$H_m(\mathbf{Z}) = \left\{ \left(\begin{array}{cc|cc} 1 & 0 & \kappa & \mu \\ 0 & 1_m & {}^t\mu & O_m \\ \hline & & 1 & 0 \\ & & 0 & 1_m \end{array} \right) \left(\begin{array}{cc|cc} 1 & \lambda & & \\ 0 & 1_m & & \\ \hline & & 1 & 0 \\ & & -{}^t\lambda & 1_m \end{array} \right) \middle| \begin{array}{l} (\lambda, \mu) \in \mathbf{Z}^m \oplus \mathbf{Z}^m, \\ \kappa \in \mathbf{Z} \end{array} \right\}.$$

Let $J_{l,1}(\Gamma_J^{(m)})$ denote the space of Jacobi forms of weight l and index 1 for $\Gamma_J^{(m)}$, and $J_{l,1}^{\text{cusp}}(\Gamma_J^{(m)})$ the subspace of $J_{l,1}(\Gamma_J^{(m)})$ consisting of all cusp forms. Let $\phi(Z, z) \in J_{l,1}(\Gamma_J^{(m)})$. Then we have the following Fourier expansion:

$$\phi(Z, z) = \sum_{\substack{T \in \mathcal{L}_m, r \in \mathbf{Z}^m, \\ 4T - {}^t r r \geq 0}} c_\phi(T, r) \mathbf{e}(\text{tr}(TZ) + r^t z).$$

We then define $\sigma_m(\phi)$ as

$$\sigma_m(\phi) = \sum_{A \in (\mathcal{L}'_m)_{\geq 0}} c_\phi((A + {}^t r_A r_A)/4, r_A) \mathbf{e}(\text{tr}(AZ)),$$

where $r = r_A$ denotes an element of \mathbf{Z}^m such that $A + {}^t r_A r_A \in 4\mathcal{L}_m$. This r_A is uniquely determined modulo $2\mathbf{Z}^m$, and $c_\phi((A + {}^t r_A r_A)/4, r_A)$ does not depend on the choice of the representative of $r_A \pmod{2\mathbf{Z}^m}$. Then Ibukiyama [7] showed that the mapping σ_m gives a \mathbf{C} -linear isomorphism $J_{l,1}(\Gamma_J^{(m)}) \simeq \mathfrak{M}_{l-1/2}^+(\Gamma_0^{(m)}(4))$, and in particular, $\sigma_m(J_{l,1}^{\text{cusp}}(\Gamma_J^{(m)})) = \mathfrak{S}_{l-1/2}^+(\Gamma_0^{(m)}(4))$. We call σ_m the *Ibukiyama isomorphism*.

Let p be a prime number. For a non-zero element $a \in \mathbf{Q}_p$ we put $\chi_p(a) = 1, -1$, or 0 according as $\mathbf{Q}_p(a^{1/2}) = \mathbf{Q}_p$, $\mathbf{Q}_p(a^{1/2})$ is an unramified quadratic extension of \mathbf{Q}_p , or $\mathbf{Q}_p(a^{1/2})$ is a ramified quadratic extension of \mathbf{Q}_p . We note that $\chi_p(D) = \left(\frac{D}{p}\right)$ if D is a fundamental discriminant. For the rest of this section, let n be a positive even integer. For an element T of $\mathcal{L}_{n,p}^\times$, put $\xi_p(T) = \chi_p((-1)^{n/2} \det T)$. Let T be an element of \mathcal{L}_n^\times . Then $(-1)^{n/2} \det(2T) \equiv 0$ or $1 \pmod{4}$, and we define \mathfrak{d}_T and \mathfrak{f}_T as $\mathfrak{d}_T = \mathfrak{d}_{(-1)^{n/2} \det(2T)}$ and $\mathfrak{f}_T = \mathfrak{f}_{(-1)^{n/2} \det(2T)}$, respectively. For an element T of $\mathcal{L}_{n,p}^\times$, there exists an element \tilde{T} of \mathcal{L}_n^\times such that $\tilde{T} \sim_{\mathbf{Z}_p} T$. We then put $\mathfrak{e}_p(T) = \nu_p(\mathfrak{f}_{\tilde{T}})$, and $[\mathfrak{d}_T] = \mathfrak{d}_{\tilde{T}} \pmod{\mathbf{Z}_p^{*\square}}$. They do not depend on the choice of \tilde{T} . We note that $(-1)^{n/2} \det(2T)$ can be expressed as $(-1)^{n/2} \det(2T) = dp^{2\mathfrak{e}_p(T)} \pmod{\mathbf{Z}_p^{*\square}}$ for any $d \in [\mathfrak{d}_T]$.

For each $T \in \mathcal{L}_{n,p}^\times$ we define the *local Siegel series* $b_p(T, s)$ by

$$b_p(T, s) = \sum_{R \in S_n(\mathbf{Q}_p)/S_n(\mathbf{Z}_p)} \mathbf{e}_p(\text{tr}(TR)) p^{-\nu_p(\mu_p(R))s},$$

where $\mu_p(R) = [R\mathbf{Z}_p^n + \mathbf{Z}_p^n : \mathbf{Z}_p^n]$. We remark that there exists a unique polynomial $F_p(T, X)$ in X such that

$$b_p(T, s) = F_p(T, p^{-s}) \frac{(1 - p^{-s}) \prod_{i=1}^{n/2} (1 - p^{2i-2s})}{1 - \xi_p(T) p^{n/2-s}}$$

(cf. Kitaoka [16]). We then define a polynomial $\tilde{F}_p(T, X)$ in X and X^{-1} as

$$\tilde{F}_p(B, X) = X^{-\epsilon_p(T)} F_p(T, p^{-(n+1)/2} X).$$

We remark that $\tilde{F}_p(B, X^{-1}) = \tilde{F}_p(B, X)$ (cf. [14]). Now, for a positive even integer k , let

$$h(z) = \sum_{\substack{m \in \mathbf{Z}_{>0}, \\ (-1)^{n/2} m \equiv 0, 1 \pmod{4}}} c_h(m) \mathbf{e}(mz)$$

be a Hecke eigenform in the Kohnen plus space $\mathfrak{S}_{k-n/2+1/2}^+(\Gamma_0(4))$ and

$$f(z) = \sum_{m=1}^{\infty} c_f(m) \mathbf{e}(mz)$$

the primitive form in $\mathfrak{S}_{2k-n}(\Gamma^{(1)})$ corresponding to h under the Shimura correspondence (cf. Kohnen [18]). Let $\beta_p \in \mathbf{C}$ such that $\beta_p + \beta_p^{-1} = p^{-k+n/2+1/2} c_f(p)$, which we call the *Satake p -parameter* of f . We define a Fourier series $I_n(h)(Z)$ on \mathbf{H}_n by

$$I_n(h)(Z) = \sum_{T \in (\mathcal{L}_n)_{>0}} c_{I_n(h)}(T) \mathbf{e}(\text{tr}(TZ)),$$

where

$$c_{I_n(h)}(T) = c_h(|\mathfrak{d}_T|) \mathfrak{f}_T^{k-n/2-1/2} \prod_p \tilde{F}_p(T, \beta_p).$$

Then Ikeda [12] showed that $I_n(h)(Z)$ is a Hecke eigenform in $\mathfrak{S}_k(\Gamma^{(n)})$ whose standard L -function coincides with $\zeta(s) \prod_{i=1}^n L(s+k-i, f)$, where $\zeta(s)$ is Riemann's zeta function. The existence of such a Hecke eigenform was conjectured by Duke and Imamoglu in their unpublished paper. We call $I_n(h)$ the *Duke-Imamoglu-Ikeda lift* of h (or of f) as in Section 1. Let $\phi_{I_n(h),1}$ be the first coefficient of the Fourier-Jacobi expansion of $I_n(h)$, that is,

$$I_n(h) \left(\begin{pmatrix} w & z \\ t & Z \end{pmatrix} \right) = \sum_{N=1}^{\infty} \phi_{I_n(h),N}(Z, z) \mathbf{e}(Nw),$$

where $Z \in \mathbf{H}_{n-1}$, $z \in \mathbf{C}^{n-1}$ and $w \in \mathbf{H}_1$. We easily see that $\phi_{I_n(h),1}$ belongs to $J_{k-1/2,1}^{\text{cusp}}(\Gamma_J^{(n-1)})$ and

$$\phi_{I_n(h),1}(Z, z) = \sum_{\substack{T \in \mathcal{L}_{n-1}, r \in \mathbf{Z}^{n-1}, \\ 4T^{-t} r r > 0}} c_{I_n(h)} \left(\begin{pmatrix} 1 & r/2 \\ t & T \end{pmatrix} \right) \mathbf{e}(\text{tr}(TZ) + r^t z).$$

Moreover we have

$$\sigma_{n-1}(\phi_{I_n(h),1})(Z) = \sum_{T \in (\mathcal{L}'_{n-1})_{>0}} c_{I_n(h)}(T^{(1)}) \mathbf{e}(\mathrm{tr}(TZ)).$$

Put $\Gamma_{\mathbf{C}}(s) = 2(2\pi)^{-s}\Gamma(s)$, and $\tilde{\xi}(s) = \Gamma_{\mathbf{C}}(s)\zeta(s)$. Then our main result in this paper is stated as follows:

Theorem 2.1. *Let h and f be as above. Then we have*

$$\begin{aligned} L(s, \sigma_{n-1}(\phi_{I_n(h),1})) &= 2^{-\delta_{2,n}-s(n-2)-(n-2)/2} \prod_{i=1}^{(n-2)/2} \tilde{\xi}(2i) \\ &\times \left\{ L(s - n/2 + 1, h) \prod_{i=1}^{(n-2)/2} L(2s - n + 2i + 1, f) \right. \\ &\quad \left. + (-1)^{n(n-2)/8} L(s, h) \prod_{i=1}^{(n-2)/2} L(2s - n + 2i, f) \right\}, \end{aligned}$$

where $\delta_{2,n}$ denotes Kronecker's delta.

In the case of $n = 2$, the modular form $\sigma_{n-1}(\phi_{I_n(h),1})$ is h itself, and then the above formula is trivial. We note that, unlike the cases of [8, 9, 10], there does not appear any convolution product of modular forms in the above theorem. However, the proof may not be so simple because the nature of Fourier coefficients of the modular form $\sigma_{n-1}(\phi_{I_n(h),1})$ is much more complicated rather than those in the papers cited above.

3. REDUCTION TO LOCAL COMPUTATIONS

By definition, it turns out that the Fourier coefficient of $\sigma_{n-1}(\phi_{I_n(h),1})$ can be expressed as a product of local Siegel series, and therefore we can reduce the problem to local computations. To explain this, we recall some terminologies and notation. For given $a, b \in \mathbf{Q}_p^\times$ let $(a, b)_p$ denote the Hilbert symbol over \mathbf{Q}_p . Following Kitaoka [17], we define the *Hasse invariant* $\varepsilon(A)$ of $A \in S_m(\mathbf{Q}_p)^\times$ by

$$\varepsilon(A) = \prod_{1 \leq i < j \leq m} (a_i, a_j)_p$$

if A is equivalent to $a_1 \perp \cdots \perp a_m$ over \mathbf{Q}_p with some $a_1, a_2, \dots, a_m \in \mathbf{Q}_p^\times$. We note that this definition does not depend on the choice of a_1, a_2, \dots, a_m .

Now put

$$\mathcal{L}'_{m,p} = \{A \in \mathcal{L}_{m,p} \mid A \equiv -{}^t r r \pmod{4\mathcal{L}_{m,p}} \text{ for some } r \in \mathbf{Z}_p^m\}.$$

Furthermore we put $S_m(\mathbf{Z}_p)_e = 2\mathcal{L}_{m,p}$ and $S_m(\mathbf{Z}_p)_o = S_m(\mathbf{Z}_p) \setminus S_m(\mathbf{Z}_p)_e$. We note that $\mathcal{L}'_{m,p} = \mathcal{L}_{m,p} = S_m(\mathbf{Z}_p)$ if $p \neq 2$. Let $T \in \mathcal{L}'_{m-1,p}$. Then there

exists an element $r \in \mathbf{Z}_p^{m-1}$ such that $\begin{pmatrix} 1 & r/2 \\ t_{r/2} & (T+t_{rr})/4 \end{pmatrix} \in \mathcal{L}_{m,p}$. As is easily shown, r is uniquely determined by T , modulo $2\mathbf{Z}_p^{m-1}$, and is denoted by r_T . Moreover as will be shown in the next lemma, $\begin{pmatrix} 1 & r_T/2 \\ t_{r_T/2} & (T+t_{r_T r_T})/4 \end{pmatrix}$ is uniquely determined by T , up to $GL_m(\mathbf{Z}_p)$ -equivalence, and is denoted by $T^{(1)}$.

Lemma 3.1. *Let m be a positive integer.*

(1) *Let A and B be elements of $\mathcal{L}'_{m-1,p}$. Then $\begin{pmatrix} 1 & r_A/2 \\ t_{r_A/2} & (A+t_{r_A r_A})/4 \end{pmatrix} \sim \begin{pmatrix} 1 & r_B/2 \\ t_{r_B/2} & (B+t_{r_B r_B})/4 \end{pmatrix}$ if $A \sim B$.*

(2) *Let $A \in \mathcal{L}'_{m-1,p}$.*

(2.1) *Let $p \neq 2$. Then $A^{(1)} \sim \begin{pmatrix} 1 & 0 \\ 0 & A \end{pmatrix}$.*

(2.2) *Let $p = 2$. If $r_A \equiv 0 \pmod{2}$, then $A \sim 4B$ with $B \in \mathcal{L}_{m-1,2}$, and $A^{(1)} \sim \begin{pmatrix} 1 & 0 \\ 0 & B \end{pmatrix}$. In particular, $\nu_2((\det B)) \geq m$ or $m+1$ according as m is even or odd. If $r_A \not\equiv 0 \pmod{2}$, then $A \sim a \perp 4B$ with $a \equiv -1 \pmod{4}$ and $B \in \mathcal{L}_{m-2,2}$ and we have $A^{(1)} \sim \begin{pmatrix} 1 & 1/2 & 0 \\ 1/2 & (a+1)/4 & 0 \\ 0 & 0 & B \end{pmatrix}$. In particular, $\nu_2((\det B)) \geq m$ or $m-1$ according as m is even or odd.*

Proof. The assertion can easily be proved. \square

Now let m be a positive even integer. For $T \in (\mathcal{L}'_{m-1})^\times$, put $\mathfrak{d}_T^{(1)} = \mathfrak{d}_{T^{(1)}}$, and $\mathfrak{f}_T^{(1)} = \mathfrak{f}_{T^{(1)}}$, and for $T \in (\mathcal{L}'_{m-1,p})^\times$, we define $[\mathfrak{d}_T^{(1)}]$ and $\mathfrak{e}_T^{(1)}$ as $[\mathfrak{d}_{T^{(1)}}]$ and $\mathfrak{e}_{T^{(1)}}$, respectively. These do not depend on the choice of r_T . We note that $(-1)^{m/2} \det T = 2^{m-2} \mathfrak{f}_T^{(1)2} \mathfrak{d}_T^{(1)}$ for $T \in (\mathcal{L}'_{m-1})^\times$. We define a polynomial $F_p^{(1)}(T, X)$ in X , and a polynomial $\tilde{F}_p^{(1)}(T, X)$ in X and X^{-1} by

$$F_p^{(1)}(T, X) = F_p(T^{(1)}, X),$$

and

$$\tilde{F}_p^{(1)}(T, X) = X^{-\mathfrak{e}_p^{(1)}(T)} F_p^{(1)}(T, p^{-(m+1)/2} X).$$

Let B be an element of $(\mathcal{L}'_{m-1,p})^\times$. Let $p \neq 2$. Then

$$\tilde{F}_p^{(1)}(B, X) = \tilde{F}_p(1 \perp B, X).$$

Let $p = 2$. Then

$$\begin{aligned} & \tilde{F}_2^{(1)}(B, X) \\ &= \begin{cases} \tilde{F}_2\left(\begin{pmatrix} 1 & 1/2 \\ 1/2 & (a+1)/4 \end{pmatrix} \perp B', X\right) & \text{if } B = a \perp 4B' \text{ with} \\ & a \equiv -1 \pmod{4}, B' \in \mathcal{L}_{m-2,2}, \\ \tilde{F}_2(1 \perp B', X) & \text{if } B = 4B' \text{ with } B' \in \mathcal{L}_{m-1,2}. \end{cases} \end{aligned}$$

Now let m and l be positive integers such that $m \geq l$. Then for nondegenerate symmetric matrices A and B of degree m and l respectively with entries in \mathbf{Z}_p we define the *local density* $\alpha_p(A, B)$ and the *primitive local density* $\beta_p(A, B)$ representing B by A as

$$\alpha_p(A, B) = 2^{-\delta_{m,l}} \lim_{a \rightarrow \infty} p^{a(-ml+l(l+1)/2)} \# \mathcal{A}_a(A, B)$$

and

$$\beta_p(A, B) = 2^{-\delta_{m,l}} \lim_{a \rightarrow \infty} p^{a(-ml+l(l+1)/2)} \# \mathcal{B}_a(A, B),$$

where

$$\mathcal{A}_a(A, B) = \{X \in M_{ml}(\mathbf{Z}_p)/p^a M_{ml}(\mathbf{Z}_p) \mid A[X] - B \in p^a S_l(\mathbf{Z}_p)_e\}$$

and

$$\mathcal{B}_a(A, B) = \{X \in \mathcal{A}_a(A, B) \mid \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p}(X \bmod p) = l\}.$$

In particular we write $\alpha_p(A) = \alpha_p(A, A)$. Put

$$\mathcal{F}_p = \{d_0 \in \mathbf{Z}_p \mid \nu_p(d_0) \leq 1\}$$

if p is an odd prime, and

$$\mathcal{F}_2 = \{d_0 \in \mathbf{Z}_2 \mid d_0 \equiv 1 \pmod{4}, \text{ or } d_0/4 \equiv -1 \pmod{4}, \text{ or } \nu_2(d_0) = 3\}.$$

Let m be a positive integer. For $d \in \mathbf{Z}_p^\times$ put

$$S_m(\mathbf{Z}_p, d)$$

$$= \{T \in S_m(\mathbf{Z}_p) \mid (-1)^{[(m+1)/2]} \det T = p^{2i} d \pmod{\mathbf{Z}_p^{*\square}} \text{ with some } i \in \mathbf{Z}\},$$

and $S_m(\mathbf{Z}_p, d)_x = S_m(\mathbf{Z}_p, d) \cap S_m(\mathbf{Z}_p)_x$ for $x = e$ or o . Put $\mathcal{L}_{m,p}^{(0)} = S_m(\mathbf{Z}_p)_e^\times$ and $\mathcal{L}_{m,p}^{(1)} = (\mathcal{L}'_{m,p})^\times$. We also define $\mathcal{L}_{m,p}^{(j)}(d) = S_m(\mathbf{Z}_p, d) \cap \mathcal{L}_{m,p}^{(j)}$ for $j = 0, 1$. Let $\iota_{m,p}$ be the constant function on $\mathcal{L}_{m,p}^\times$ taking the value 1, and $\varepsilon_{m,p}$ the function on $\mathcal{L}_{m,p}^\times$ assigning the Hasse invariant of A for $A \in \mathcal{L}_{m,p}^\times$. We sometimes drop the suffix and write $\iota_{m,p}$ as ι_p or ι and the others if there is no fear of confusion. From now on, we sometimes write $\omega = \varepsilon^l$ with $l = 0$ or 1 according as $\omega = \iota$ or ε . Let n be a positive even integer. For $d_0 \in \mathcal{F}_p$ and $\omega = \varepsilon^l$ with $l = 0$ or 1 , we define a formal power series $P_{n-1,p}^{(1)}(d_0, \omega, X, t)$ in t by

$$P_{n-1,p}^{(1)}(d_0, \omega, X, t) = \kappa(d_0, n-1, l)^{-1} t^{\delta_{2,p}(2-n)} \sum_{B \in \mathcal{L}_{n-1,p}^{(1)}(d_0)} \frac{\tilde{F}_p^{(1)}(B, X)}{\alpha_p(B)} \omega(B) t^{\nu(\det B)},$$

where

$$\begin{aligned} \kappa(d_0, r-1, l) &= \kappa(d_0, r-1, l)_p \\ &= \{(-1)^{lr(r-2)/8} 2^{-(r-2)(r-1)/2}\}^{\delta_{2,p}} \cdot ((-1)^{r/2}, (-1)^{r/2} d_0)_p^l p^{-(r/2-1)l\nu(d_0)} \end{aligned}$$

for a positive even integer r . This type of formal power series appears in an explicit formula for the Koecher-Maass series associated with the Siegel Eisenstein series and the Duke-Imamog̃lu-Ikeda lift (cf. [9], [10]). Therefore we say that this formal power series is of *Koecher-Maass type*. An explicit formula for $P_{n-1,p}^{(1)}(d_0, \omega, X, t)$ will be given in the next section. Let \mathcal{F} denote the set of fundamental discriminants, and for $l = \pm 1$, put

$$\mathcal{F}^{(l)} = \{d_0 \in \mathcal{F} \mid ld_0 > 0\}.$$

Now let h be a Hecke eigenform in $\mathfrak{S}_{k-n/2+1/2}^+(T_0(4))$, and $f, I_n(h), \phi_{I_n(h),1}$ and $\sigma_{n-1}(\phi_{I_n(h),1})$ be as in Section 2.

Theorem 3.2. *Let the notation and the assumption be as above. Then for $\operatorname{Re}(s) \gg 0$, we have*

$$\begin{aligned} L(s, \sigma_{n-1}(\phi_{I_n(h),1})) &= \kappa_{n-1} 2^{-(n-2)s - (n-2)/2 - \delta_{2,n}} \\ &\times \left\{ \sum_{d_0 \in \mathcal{F}^{((-1)^{n/2}}} c_h(|d_0|) |d_0|^{n/4 - k/2 + 1/4} \prod_p P_{n-1,p}^{(1)}(d_0, \iota_p, \beta_p, p^{-s+k/2+n/4-1/4}) \right. \\ &\quad + (-1)^{n(n-2)/8} \sum_{d_0 \in \mathcal{F}^{((-1)^{n/2}}} c_h(|d_0|) |d_0|^{-n/4 - k/2 + 5/4} \\ &\quad \left. \times \prod_p P_{n-1,p}^{(1)}(d_0, \varepsilon_p, \beta_p, p^{-s+k/2+n/4-1/4}) \right\}, \end{aligned}$$

where $\kappa_{n-1} = \prod_{i=1}^{(n-2)/2} \Gamma_{\mathbf{C}}(2i)$.

Proof. Let $T \in (\mathcal{L}'_{n-1})_{>0}$. Then it follows from Lemma 4.1 that the T -th Fourier coefficient $c_{\sigma_{n-1}(\phi_{I_n(h),1})}(T)$ of $\sigma_{n-1}(\phi_{I_n(h),1})$ is uniquely determined by the genus to which T belongs, and by definition, it can be expressed as

$$c_{\sigma_{n-1}(\phi_{I_n(h),1})}(T) = c_{I_n(h)}(T^{(1)}) = c_h(|\mathfrak{d}_T^{(1)}|) (\mathfrak{f}_T^{(1)})^{k-n/2-1/2} \prod_p \widetilde{F}^{(1)}(T, \beta_p).$$

We note that

$$(\mathfrak{f}_T^{(1)})^{k-n/2-1/2} = |\mathfrak{d}_T^{(1)}|^{-(k/2-n/4-1/4)} (\det T)^{(k/2-n/4-1/4)} 2^{-(n-2)(k/2-n/4-1/4)}$$

for $T \in (\mathcal{L}_{n-1})_{>0}$. We also note that

$$\sum_{T' \in \mathcal{G}(T)} \frac{1}{e(T')} = \kappa_{n-1} 2^{3-n-\delta_{2,n}} \det T^{n/2} \prod_p \alpha_p(A)^{-1}$$

for $T \in S_{n-1}(\mathbf{Z})_{>0}$, where $\mathcal{G}(T)$ denotes the set of $SL_{n-1}(\mathbf{Z})$ -equivalence classes belonging to the genus of T (cf. Theorem 6.8.1 in [17]). Hence we

have

$$\sum_{T' \in \mathcal{G}(T)} \frac{c_{\sigma_{n-1}(\phi_{I_n(h),1})}(T)}{e(T')} = \kappa_{n-1} 2^{3-n-\delta_{2,n}-(n-2)(k/2-n/4-1/4)}$$

$$\times \det T^{k/2+n/4-1/4} |\mathfrak{d}_T^{(1)}|^{-k/2+n/4+1/4} \prod_p \frac{\widetilde{F}_p^{(1)}(T, \beta_p)}{\alpha_p(T)}.$$

Thus, by using the same method as in Proposition 2.2 of [11], similarly to Theorem 3.3 (1) of [8], and Theorem 3.2 of [9], we obtain

$$L(s, \sigma_{n-1}(\phi_{I_n(h),1}))$$

$$= \kappa_{n-1} 2^{-(k/2-n/4-1/4)(n-2)+2-n-\delta_{2,n}} \sum_{d_0 \in \mathcal{F}((-1)^{n/2})} c_h(|d_0|) |d_0|^{n/4-k/2+1/4}$$

$$\times \left\{ 2^{(-s+k/2+n/4-1/4)(n-2)} \prod_p \kappa_p(d_0, n-1, 0) P_{n-1,p}^{(1)}(d_0, \iota_p, \beta_p, p^{-s+k/2+n/4-1/4}) \right.$$

$$\left. + 2^{(-s+k/2+n/4-1/4)(n-2)} \prod_p \kappa_p(d_0, n-1, 1) P_{n-1,p}^{(1)}(d_0, \varepsilon_p, \beta_p, p^{-s+k/2+n/4-1/4}) \right\}.$$

We note that $\prod_p ((-1)^{n/2}, (-1)^{n/2} d_0)_p = 1$. Hence $\prod_p \kappa_p(d_0, n-1, 0) = 2^{-(n-2)(n-1)/2}$, and $\prod_p \kappa_p(d_0, n-1, 1) = (-1)^{n(n-2)/8} |d_0|^{-n/2+1} 2^{-(n-2)(n-1)/2}$. This proves the assertion. \square

4. FORMAL POWER SERIES ASSOCIATED WITH LOCAL SIEGEL SERIES

Throughout this section we fix a positive even integer n . We also simply write ν_p as ν and the others if the prime number p is clear from the context. In this section, we give an explicit formula for $P_{n-1}^{(1)}(d_0, \omega, X, t)$ with $\omega = \varepsilon^l$ ($l = 0, 1$) to prove Theorem 3.2 (cf. Theorem 4.4.1). The idea of the proof is to express the power series in question as a sum of certain subseries (cf. Proposition 4.4.3). Henceforth, for a $GL_m(\mathbf{Z}_p)$ -stable subset \mathcal{B} of $S_m(\mathbf{Q}_p)$, we simply write $\sum_{T \in \mathcal{B}}$ instead of $\sum_{T \in \mathcal{B}/\sim}$ if there is no fear of confusion.

4.1. Local densities.

Put $\mathcal{D}_{m,i} = GL_m(\mathbf{Z}_p) \begin{pmatrix} 1_{m-i} & 0 \\ 0 & p1_i \end{pmatrix} GL_m(\mathbf{Z}_p)$. For two elements S and T of $S_m(\mathbf{Z}_p)^\times$ and an nonnegative integer $i \leq m$, we define $\alpha_p(S, T, i)$ as

$$\alpha_p(S, T, i) = 2^{-1} \lim_{e \rightarrow \infty} p^{-e(m-1)m/2} \mathcal{A}_e(S, T, i),$$

where

$$\mathcal{A}_e(S, T, i) = \{\overline{X} \in \mathcal{A}_e(S, T) \mid X \in \mathcal{D}_{m,i}\}.$$

Lemma 4.1.1. *Let S and T be elements of $S_m(\mathbf{Z}_p)^\times$.*

(1) *Let $\Omega(S, T) = \{W \in M_m(\mathbf{Z}_p)^\times \mid S[W] \sim T\}$, and $\Omega(S, T, i) = \Omega(S, T) \cap \mathcal{D}_{m,i}$. Then*

$$\frac{\alpha_p(S, T)}{\alpha_p(T)} = \#(\Omega(S, T)/GL_m(\mathbf{Z}_p))p^{-m(\nu(\det T) - \nu(\det S))/2},$$

and

$$\frac{\alpha_p(S, T, i)}{\alpha_p(T)} = \#(\Omega(S, T, i)/GL_m(\mathbf{Z}_p))p^{-m(\nu(\det T) - \nu(\det S))/2}.$$

(2) *Let $\tilde{\Omega}(S, T) = \{W \in M_m(\mathbf{Z}_p)^\times \mid S \sim T[W^{-1}]\}$, and $\tilde{\Omega}(S, T, i) = \tilde{\Omega}(S, T) \cap \mathcal{D}_{m,i}$. Then*

$$\frac{\alpha_p(S, T)}{\alpha_p(S)} = \#(GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}(S, T))p^{(\nu(\det T) - \nu(\det S))/2},$$

and

$$\frac{\alpha_p(S, T, i)}{\alpha_p(S)} = \#(GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}(S, T, i))p^{(\nu(\det T) - \nu(\det S))/2}.$$

Proof. The assertion (1) follows from Lemma 2.2 of [4]. Now by Proposition 2.2 of [14] we have

$$\alpha_p(S, T) = \sum_{W \in GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}(S, T)} \beta_p(S, T[W^{-1}])p^{\nu(\det W)}.$$

Then $\beta_p(S, T[W^{-1}]) = \alpha_p(S)$ or 0 according as $S \sim T[W^{-1}]$ or not. Thus the assertion (2) holds. \square

A nondegenerate $m \times m$ matrix $D = (d_{ij})$ with entries in \mathbf{Z}_p is said to be *reduced* if D satisfies the following two conditions:

- (a) For $i = j$, $d_{ii} = p^{e_i}$ with a nonnegative integer e_i ;
- (b) For $i \neq j$, d_{ij} is a nonnegative integer satisfying $d_{ij} \leq p^{e_j} - 1$ if $i < j$ and $d_{ij} = 0$ if $i > j$.

It is well known that we can take the set of all reduced matrices as a complete set of representatives of $GL_m(\mathbf{Z}_p) \backslash M_m(\mathbf{Z}_p)^\times$. Let $j = 0$ or 1 according as m is even or odd. For $B \in \mathcal{L}_{m,p}^{(j)}$ put

$$\tilde{\Omega}^{(j)}(B) = \{W \in M_m(\mathbf{Z}_p)^\times \mid B[W^{-1}] \in \mathcal{L}_{m,p}^{(j)}\}.$$

Furthermore put $\tilde{\Omega}^{(j)}(B, i) = \tilde{\Omega}^{(j)}(B) \cap \mathcal{D}_{m,i}$. Let $n_0 \leq m$, and $\psi_{n_0, m}$ be the mapping from $GL_{n_0}(\mathbf{Q}_p)$ into $GL_m(\mathbf{Q}_p)$ defined by $\psi_{n_0, m}(D) = 1_{m-n_0} \perp D$.

Lemma 4.1.2. (1) *Suppose that $p \neq 2$. Let $\Theta \in GL_{n_0}(\mathbf{Z}_p) \cap S_{n_0}(\mathbf{Z}_p)$, and $B_1 \in S_{m-n_0}(\mathbf{Z}_p)^\times$.*

(1.1) Let n_0 be even. Then $\psi_{m-n_0,m}$ induces a bijection

$$GL_{m-n_0}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(B_1) \simeq GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(\Theta \perp B_1),$$

where $j = 0$ or 1 according as m is even or odd. In particular, we have

$$GL_{m-n_0}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(pB_1) \simeq GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(\Theta \perp pB_1).$$

(1.2) Let n_0 be odd. Then $\psi_{m-n_0,m}$ induces a bijection

$$GL_{m-n_0}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j')}(B_1) \simeq GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(\Theta \perp B_1),$$

where $j = 0$ or 1 according as m is even or odd, and $j' = 1$ or 0 according as m is even or odd. In particular, we have

$$GL_{m-n_0}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j')}(pB_1) \simeq GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(\Theta \perp pB_1),$$

(2) Suppose that $p = 2$. Let m be a positive integer, n_0 an even integer not greater than m , and $\Theta \in GL_{n_0}(\mathbf{Z}_2) \cap S_{n_0}(\mathbf{Z}_2)_e$.

(2.1) Let $B_1 \in S_{m-n_0}(\mathbf{Z}_2)^\times$. Then $\psi_{m-n_0,m}$ induces a bijection

$$GL_{m-n_0}(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(j)}(2^{j+1}B_1) \simeq GL_m(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(j)}(2^j \Theta \perp 2^{j+1}B_1),$$

where $j = 0$ or 1 according as m is even or odd.

(2.2) Suppose that m is even. Let $a \in \mathbf{Z}_2$ such that $a \equiv -1 \pmod{4}$, and $B_1 \in S_{m-n_0-2}(\mathbf{Z}_2)^\times$. Then $\psi_{m-n_0-1,m}$ induces a bijection

$$\begin{aligned} & GL_{m-n_0-1}(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(1)}(a \perp 4B_1) \\ & \simeq GL_m(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(0)}(\Theta \perp \begin{pmatrix} 2 & 1 \\ 1 & 2+a \end{pmatrix} \perp 2B_1). \end{aligned}$$

(2.3) Suppose that m is even, and let $B_1 \in S_{m-n_0-1}(\mathbf{Z}_2)^\times$. Then there exists a bijection $\tilde{\psi}_{m-n_0-1,m}$

$$GL_{m-n_0-1}(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(1)}(4B_1) \simeq GL_m(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(0)}(\Theta \perp 2 \perp 2B_1).$$

(As will be seen later, $\tilde{\psi}_{m-n_0-1,m}$ is not induced from $\psi_{m-n_0-1,m}$.)

(3) The assertions (1), (2) remain valid if one replaces $\tilde{\Omega}^{(j)}(B)$ by $\tilde{\Omega}^{(j)}(B, i)$.

Proof. (1) Clearly $\psi_{m-n_0,m}$ induces an injection from $GL_{m-n_0}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(B_1)$ to $GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(\Theta \perp B_1)$, which is denoted also by the same symbol $\psi_{m-n_0,m}$. To prove the surjectivity of $\psi_{m-n_0,m}$, take a representative D of an element of $GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(j)}(\Theta \perp B_1)$. Without loss of generality we may suppose that D is a reduced matrix. Since $(\Theta \perp B_1)[D^{-1}] \in S_m(\mathbf{Z}_p)$, we have $D = \begin{pmatrix} 1_{n_0} & 0 \\ 0 & D_1 \end{pmatrix}$ with $D_1 \in \tilde{\Omega}^{(j)}(B_1)$. This proves the assertion (1.1). The assertion (1.2) can be proved in the same way as above.

(2) First we prove (2.1). As in (1), the mapping $\psi_{m-n_0,m}$ induces an injection from $GL_{m-n_0}(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(j)}(2^{j+1}B_1)$ to $GL_m(\mathbf{Z}_2) \backslash \tilde{\Omega}^{(j)}(2^j\Theta \perp 2^{j+1}B_1)$, which is denoted also by the same symbol $\psi_{m-n_0,m}$. Then the surjectivity of $\psi_{m-n_0,m}$ in case $j = 0$ can be proved in the same manner as (1). To prove the surjectivity of $\psi_{m-n_0,m}$ in case $j = 1$, take a reduced matrix $D = \begin{pmatrix} D_1 & D_{12} \\ 0 & D_2 \end{pmatrix}$ with $D_1 \in M_{n_0}(\mathbf{Z}_2)^\times, D_2 \in M_{m-n_0}(\mathbf{Z}_2)^\times, D_{12} \in M_{n_0, m-n_0}(\mathbf{Z}_2)$. If $(2\Theta \perp 4B_1)[D^{-1}] \in \mathcal{L}_{m,2}^{(1)}$, then there exists an element $(r_1, r_2) \in \mathbf{Z}_2^{n_0} \times \mathbf{Z}_2^{m-n_0}$ such that

$$2\Theta[D_1^{-1}] \equiv -{}^t r_1 r_1 \pmod{4\mathcal{L}_{n_0,2}},$$

$$-2\Theta[D_1^{-1}]D_{12}D_2^{-1} \equiv -{}^t r_2 r_1 \pmod{2M_{n_0, m-n_0}(\mathbf{Z}_2)},$$

and

$$2\Theta[D_1^{-1}D_{12}D_2^{-1}] + 4B_1[D_2^{-1}] \equiv -{}^t r_2 r_2 \pmod{4\mathcal{L}_{m-n_0,2}}.$$

We have $\nu(\det(2\Theta[D_1^{-1}])) \geq n_0$, and $\nu(2\Theta) = n_0$. Hence we have $D_1 = 1_{n_0}$, and $r_1 \equiv 0 \pmod{2}$. Hence $4B_1[D_2^{-1}] \in \mathcal{L}_{m-n_0}^{(1)}$, and $D_{12}D_2^{-1} \in M_{n_0, m-n_0}(\mathbf{Z}_2)$. Hence $D = U \begin{pmatrix} 1_{n_0} & 0 \\ 0 & D_2 \end{pmatrix}$ with $U \in GL_m(\mathbf{Z}_p)$. Thus the surjectivity of $\psi_{m-n_0,m}$ can be proved in the same way as above. The assertion (2.2) can be proved in the same way as above.

To prove (2.3), we may suppose that $n_0 = 0$ in view of (2.1). Let $D \in \tilde{\Omega}^{(1)}(4B_1)$. Then

$$4B_1[D^{-1}] = -{}^t r_0 r_0 + 4B'$$

with $r_0 \in \mathbf{Z}_2^{m-1}$ and $B' \in \mathcal{L}_{m-1,2}$. Then we can take $r \in \mathbf{Z}_2^{m-1}$ such that

$$4{}^t D^{-1} {}^t r r D^{-1} \equiv {}^t r_0 r_0 \pmod{4\mathcal{L}_{m-1,2}}.$$

Furthermore, $2rD^{-1}$ is uniquely determined modulo $2\mathbf{Z}_2^{m-1}$ by r_0 . Put $\tilde{D} = \begin{pmatrix} 1 & r \\ 0 & D \end{pmatrix}$. Then \tilde{D} belongs to $\tilde{\Omega}^{(0)}(2 \perp 2B_1)$, and the mapping $D \mapsto \tilde{D}$ induces a bijection in question. \square

Corollary. *Suppose that m is even. Let $B \in \mathcal{L}_{m-1,p}^{(1)}$, and*

$$B^{(1)} = \begin{pmatrix} 1 & r_B/2 \\ {}^t r_B/2 & (B + {}^t r_B r_B)/4 \end{pmatrix}$$

with $r_B \in \mathbf{Z}_p^{m-1}$ as defined in Section 3. Then there exists a bijection

$$\psi : GL_{m-1}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(1)}(B) \simeq GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(0)}(2^{\delta_{2,p}} B^{(1)})$$

such that $\nu(\det(\psi(W))) = \nu(\det(W))$ for any $W \in GL_{m-1}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(1)}(B)$.

Moreover, this induces a bijection ψ_i from $GL_{m-1}(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(1)}(B, i)$ to $GL_m(\mathbf{Z}_p) \backslash \tilde{\Omega}^{(0)}(2^{\delta_{2,p}} B^{(1)}, i)$ for $i = 0, \dots, m-1$.

Proof. Let $p \neq 2$. Then we may suppose $r_B = 0$, and the assertion follows from (1.2). Let $p = 2$. If $r_B \equiv 0 \pmod{2}$ we may suppose that $r_B = 0$, and the assertion follows from (2.3). If $r_B \not\equiv 0 \pmod{2}$, we may suppose that $B = a \perp 4B_1$ with $B_1 \in \mathcal{L}_{m-2,2}^\times$ and $r_B = (1, 0, \dots, 0)$. Thus the assertion follows from (2.2). \square

Lemma 4.1.3. *Suppose that $p \neq 2$.*

(1) *Let $B \in S_m(\mathbf{Z}_p)^\times$. Then*

$$\alpha_p(p^r dB) = p^{rm(m+1)/2} \alpha_p(B)$$

for any nonnegative integer r and $d \in \mathbf{Z}_p^$.*

(2) *Let $U_1 \in GL_{n_0}(\mathbf{Z}_p) \cap S_{n_0}(\mathbf{Z}_p)$ and $B_1 \in S_{m-n_0}(\mathbf{Z}_p)^\times$. Then*

$$\begin{aligned} \alpha_p(pB_1 \perp U_1) &= 2^{r(n_0)} \alpha_p(pB_1) \\ &\times \begin{cases} \prod_{i=1}^{n_0/2} (1 - p^{-2i})(1 + \chi((-1)^{n_0/2} \det U_1) p^{-n_0/2})^{-1} & \text{if } n_0 \text{ even,} \\ \prod_{i=1}^{(n_0-1)/2} (1 - p^{-2i}) & \text{if } n_0 \text{ odd,} \end{cases} \end{aligned}$$

for $n_0 \geq 1$, where $r(n_0) = 0$ or 1 according as $n_0 = m$ or not.

Proof. The assertion (1) follows from Theorem 5.6.4 (a) of [17]. The assertion (2) follows from the formula on Page 110, line 4 from the bottom of Kitaoka [17]. \square

Lemma 4.1.4. (1) *Let $B \in S_m(\mathbf{Z}_2)^\times$. Then*

$$\alpha_2(2^r dB) = 2^{rm(m+1)/2} \alpha_2(B)$$

for any nonnegative integer r and $d \in \mathbf{Z}_2^$.*

(2) *Let n_0 be even, and $U_1 \in GL_{n_0}(\mathbf{Z}_2) \cap S_{n_0}(\mathbf{Z}_2)_e$. Then for $B_1 \in S_{m-n_0}(\mathbf{Z}_2)^\times$ we have*

$$\begin{aligned} \alpha_2(U_1 \perp 2B_1) &= 2^{r(n_0)} \alpha_2(2B_1) \\ &\times \begin{cases} \prod_{i=1}^{n_0/2} (1 - 2^{-2i})(1 + \chi((-1)^{n_0/2} \det U_1) 2^{-n_0/2})^{-1} & \text{if } B_1 \in S_{m-n_0}(\mathbf{Z}_2)_e, \\ \prod_{i=1}^{(n_0-1)/2} (1 - 2^{-2i}) & \text{if } B_1 \in S_{m-n_0}(\mathbf{Z}_2)_o, \end{cases} \end{aligned}$$

and for $u_0 \in \mathbf{Z}_2^$ and $B_2 \in S_{m-n_0-1}(\mathbf{Z}_2)^\times$ we have*

$$\alpha_2(u_0 \perp 2U_1 \perp 4B_2) = \alpha_2(2B_2) 2^{m(m-1)/2+1} \prod_{i=1}^{n_0/2} (1 - 2^{-2i}).$$

(3) *Let $u_0 \in \mathbf{Z}_2^*$ and $B_1 \in S_{m-1}(\mathbf{Z}_2)^\times$. Then we have*

$$\alpha_2(u_0 \perp 5B_1) = \alpha_2(u_0 \perp B_1).$$

Proof. The assertion (1) follows from Theorem 5.6.4 (a) of [17]. The assertion (2) follows from (4) on Page 111 of [17]. For a nondegenerate half-integral matrix A , let W_A be the quadratic space over \mathbf{Z}_p associated with A , and $n_{A,j}$, $q_{A,j}$ and $E_{A,j}$ be the quantities n_j , q_j and E_j , respectively, on Page 109 of [17] defined for W_A . Then the transformation $u_0 \perp B_1 \mapsto u_0 \perp 5B_1$ does not change these quantities. This proves the assertion (3). \square

Now let R be a commutative ring. Then the group $GL_m(R) \times R^*$ acts on $S_m(R)$. We write $B_1 \approx_R B_2$ if $B_2 \sim_R \xi B_1$ with some $\xi \in R^*$. Let m be a positive integer. Then for $B \in S_m(\mathbf{Z}_p)$ let $\tilde{\mathcal{S}}_{m,p}(B)$ denote the set of elements of $S_m(\mathbf{Z}_p)$ such that $B' \approx_{\mathbf{Z}_p} B$, and let $\mathcal{S}_{m-1,p}(B)$ denote the set of elements of $S_{m-1}(\mathbf{Z}_p)$ such that $1 \perp B' \approx_{\mathbf{Z}_p} B$.

Lemma 4.1.5. *Let m be a positive even integer. Let $B \in S_m(\mathbf{Z}_2)_o^\times$. Then*

$$\sum_{B' \in \mathcal{S}_{m-1,2}(B)/\sim} \frac{1}{\alpha_2(B')} = \frac{\#(\tilde{\mathcal{S}}_{m,2}(B)/\sim)}{2\alpha_2(B)}.$$

Proof. For a positive integer l let $l = l_1 + \cdots + l_r$ be the partition of l by positive integers, and $\{s_i\}_{i=1}^r$ the set of nonnegative integers such that $0 \leq s_1 < \cdots < s_r$. Then for a positive integer e let $S_l^0(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{l_i\}, \{s_i\})$ be the subset of $S_l(\mathbf{Z}_2/2^e\mathbf{Z}_2)$ consisting of symmetric matrices of the form $2^{s_1}U_1 \perp 2^{s_2}U_2 \perp \cdots \perp 2^{s_r}U_r$ with $U_i \in S_{l_i}(\mathbf{Z}_2/2^e\mathbf{Z}_2)$ unimodular. Let $B \in S_m(\mathbf{Z}_2)_o$ and $\det B = (-1)^{m/2}d$. Then B is equivalent, over \mathbf{Z}_2 , to a matrix of the following form:

$$2^{t_1}W_1 \perp 2^{t_2}W_2 \perp \cdots \perp 2^{t_r}W_r,$$

where $0 = t_1 < t_1 < \cdots < t_r$ and W_1, \dots, W_{r-1} , and W_r are unimodular matrices of degree n_1, \dots, n_{r-1} , and n_r , respectively, and in particular, W_1 is odd unimodular. Then by Lemma 3.2 of [11], similarly to (3.5) of [11], for a sufficiently large integer e , we have

$$\begin{aligned} \frac{\#(\tilde{\mathcal{S}}_{m,2}(B)/\sim)}{\alpha_2(B)} &= \sum_{\tilde{B} \in \tilde{\mathcal{S}}_{m,2}(B)/\sim} \frac{1}{\alpha_2(\tilde{B})} \\ &= 2^{m-1} 2^{-\nu(d) + \sum_{i=1}^r n_i(n_i-1)e/2 - (r-1)(e-1) - \sum_{1 \leq j < i \leq r} n_i n_j t_j} \\ &\quad \times \prod_{i=1}^r \#(SL_{n_i}(\mathbf{Z}_2/2^e\mathbf{Z}_2))^{-1} \# \tilde{S}_m^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n_i\}, \{t_i\}, B), \end{aligned}$$

where $\tilde{S}_m^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n_i\}, \{t_i\}, B)$ is the subset of $S_m^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n_i\}, \{t_i\})$ consisting of matrices A such that $A \approx_{\mathbf{Z}_2/2^e\mathbf{Z}_2} B$. We note that our local density $\alpha_2(\tilde{B})$ is 2^{-m} times that in [11] for $\tilde{B} \in S_m(\mathbf{Z}_2)$. If $n_1 \geq 2$, put

$r' = r, n'_1 = n_1 - 1, n'_2 = n_2, \dots, n'_r = n_r$, and $t'_i = t_i$ for $i = 1, \dots, r'$, and if $n_1 = 1$, put $r' = r - 1, n'_i = n_{i+1}$ and $t'_i = t_{i+1}$ for $i = 1, \dots, r'$. Let $S_{m-1}^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n'_i\}, \{t'_i\}, B)$ be the subset of $S_{m-1}^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n'_i\}, \{t'_i\})$ consisting of matrices $B' \in S_{m-1}(\mathbf{Z}_2/2^e\mathbf{Z}_2)$ such that $1 \perp B' \approx_{\mathbf{Z}_2/2^e\mathbf{Z}_2} B$. Then, similarly, we obtain

$$\sum_{B' \in S_{m-1,2}(B)/\sim} \frac{1}{\alpha_2(B')} = 2^{m-2} 2^{-\nu(d) + \sum_{i=1}^{r'} n'_i(n'_i-1)e/2 - (r'-1)(e-1) - \sum_{1 \leq j < i \leq r'} n'_i n'_j t'_j} \\ \times \prod_{i=1}^{r'} \#(SL_{n'_i}(\mathbf{Z}_2/2^e\mathbf{Z}_2))^{-1} \#S_{m-1}^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n'_i\}, \{t'_i\}, B).$$

Take an element A of $\tilde{S}_m^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n_i\}, \{t_i\}, B)$. Then

$$A = 2^{t_1} U_1 \perp 2^{t_2} U_2 \perp \dots \perp 2^{t_r} U_r$$

with $U_i \in S_{n_i}(\mathbf{Z}_2/2^e\mathbf{Z}_2)$ unimodular. Put $U_1 = (u_{\lambda\mu})_{n_1 \times n_1}$. Then by the assumption there exists an integer $1 \leq \lambda \leq n_1$ such that $u_{\lambda\lambda} \in \mathbf{Z}_2^*$. Let λ_0 be the least integer such that $u_{\lambda_0\lambda_0} \in \mathbf{Z}_2^*$, and V_1 be the matrix obtained from U_1 by interchanging the first and λ_0 -th rows and the first and λ_0 -th columns. Write V_1 as $V_1 = \begin{pmatrix} v_1 & \mathbf{v}_1 \\ {}^t\mathbf{v}_1 & V' \end{pmatrix}$ with $v_1 \in \mathbf{Z}_2^*, \mathbf{v}_1 \in M_{1, n_1-1}(\mathbf{Z}_2)$, and $V' \in S_{n_1-1}(\mathbf{Z}_2)$. Here we understand that $V' - {}^t\mathbf{v}_1\mathbf{v}_1$ is the empty matrix if $n_1 = 1$. Then

$$\tilde{V}_1 \sim \begin{pmatrix} v_1 & 0 \\ 0 & V' - {}^t\mathbf{v}_1 v_1^{-1} \mathbf{v}_1 \end{pmatrix}.$$

Then the map $A \mapsto v_1^{-1}(2^{t_1}(V' - {}^t\mathbf{v}_1 v_1^{-1} \mathbf{v}_1) \perp 2^{t_2} U_2 \perp \dots \perp 2^{t_r} U_r)$ induces a map Υ from $\tilde{S}_m^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n_i\}, \{t_i\}, B)$ to $S_{m-1}^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n'_i\}, \{t'_i\}, B)$. By a simple calculation, we obtain

$$\#\Upsilon^{-1}(B') = 2^{(e-1)n_1} (2^{n_1} - 1)$$

for any $B' \in S_{m-1}^{(0)}(\mathbf{Z}_2/2^e\mathbf{Z}_2, \{n'_i\}, \{t'_i\}, B)$. We also note that

$$\#SL_{n_1}(\mathbf{Z}_2/2^e\mathbf{Z}_2) = 2^{(e-1)(2n_1-1)} 2^{n_1-1} (2^{n_1} - 1) \#(SL_{n_1-1}(\mathbf{Z}_2/2^e\mathbf{Z}_2)) \text{ or } 1$$

according as $n_1 \geq 2$ or $n_1 = 1$, and

$$\sum_{i=1}^r n_i(n_i - 1)e/2 - (r-1)(e-1) - \sum_{1 \leq j < i \leq r} n_i n_j t_j \\ = e_{n_1} + \sum_{i=1}^{r'} n'_i(n'_i - 1)e/2 - (r'-1)(e-1) + \sum_{1 \leq j < i \leq r'} n'_i n'_j t'_j,$$

where $e_{n_1} = (n_1 - 1)e$ or $e_{n_1} = 1 - e$ according as $n_1 \geq 2$ or $n_1 = 1$. Hence

$$\begin{aligned} & 2^{m-1} 2^{-\nu(d) + \sum_{i=1}^r n_i(n_i-1)e/2 - (r-1)(e-1) - \sum_{1 \leq j < i \leq r} n_i n_j t_j} \\ & \quad \times \prod_{i=1}^r \#(SL_{n_i}(\mathbf{Z}_2/2^e \mathbf{Z}_2))^{-1} \# \widetilde{S}_m^{(0)}(\mathbf{Z}_2/2^e \mathbf{Z}_2, \{n_i\}, \{t_i\}, B) \\ & = 2 \cdot 2^{m-2} 2^{-\nu(d) + \sum_{i=1}^{r'} n'_i(n'_i-1)e/2 - (r'-1)(e-1) - \sum_{1 \leq j \leq i \leq r'} n'_i n'_j t'_j} \\ & \quad \times \prod_{i=1}^{r'} \#(SL_{n'_i}(\mathbf{Z}_2/2^e \mathbf{Z}_2))^{-1} \# S_{m-1}^{(0)}(\mathbf{Z}_2/2^e \mathbf{Z}_2, \{n'_i\}, \{t'_i\}, B). \end{aligned}$$

This proves the assertion. \square

4.2. Siegel series.

For a half-integral matrix B of degree m over \mathbf{Z}_p , let $(\overline{W}, \overline{q})$ denote the quadratic space over $\mathbf{Z}_p/p\mathbf{Z}_p$ defined by the quadratic form $\overline{q}(\mathbf{x}) = B[\mathbf{x}] \pmod{p}$, and define the radical $R(\overline{W})$ of \overline{W} by

$$R(\overline{W}) = \{\mathbf{x} \in \overline{W} \mid \overline{B}(\mathbf{x}, \mathbf{y}) = 0 \text{ for any } \mathbf{y} \in \overline{W}\},$$

where \overline{B} denotes the associated symmetric bilinear form of \overline{q} . We then put $l_p(B) = \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} R(\overline{W})^\perp$, where $R(\overline{W})^\perp$ is the orthogonal complement of $R(\overline{W})$ in \overline{W} . Furthermore, in case $l_p(B)$ is even, put $\overline{\xi}_p(B) = 1$ or -1 according as $R(\overline{W})^\perp$ is hyperbolic or not. In case $l_p(B)$ is odd, we put $\overline{\xi}_p(B) = 0$. Here we make the convention that $\xi_p(B) = 1$ if $l_p(B) = 0$. We note that $\overline{\xi}_p(B)$ is different from the $\xi_p(B)$ in general, but they coincide if $B \in \mathcal{L}_{m,p} \cap \frac{1}{2}GL_m(\mathbf{Z}_p)$.

Let m be a positive even integer. For $B \in \mathcal{L}_{m-1,p}^{(1)}$ put

$$B^{(1)} = \begin{pmatrix} 1 & r/2 \\ {}^t r/2 & (B + {}^t r r)/4 \end{pmatrix},$$

where r is an element of \mathbf{Z}_p^{m-1} such that $B + {}^t r r \in 4\mathcal{L}_{m-1,p}$. Then we put $\xi^{(1)}(B) = \xi(B^{(1)})$ and $\overline{\xi}^{(1)}(B) = \overline{\xi}(B^{(1)})$. These do not depend on the choice of r , and we have $\xi^{(1)}(B) = \chi((-1)^{m/2} \det B)$.

Let $p \neq 2$. Let $j = 0$ or 1 . Then an element B of $\mathcal{L}_{m-j,p}^{(j)}$ is equivalent, over \mathbf{Z}_p , to $\Theta \perp pB_1$ with $\Theta \in GL_{m-n_1-j}(\mathbf{Z}_p) \cap S_{m-n_1-j}(\mathbf{Z}_p)$ and $B_1 \in S_{n_1}(\mathbf{Z}_p)^\times$. Then $\overline{\xi}^{(j)}(B) = 0$ if n_1 is odd, and $\overline{\xi}^{(1)}(B) = \chi((-1)^{(m-n_1)/2} \det \Theta)$ if n_1 is even.

Let $p = 2$. Then an element $B \in \mathcal{L}_{m-1,2}^{(1)}$ is equivalent, over \mathbf{Z}_2 , to a matrix of the form $2\Theta \perp B_1$, where $\Theta \in GL_{m-n_1-2}(\mathbf{Z}_2) \cap S_{m-n_1-2}(\mathbf{Z}_2)_e$ and B_1 is one of the following three types:

- (I) $B_1 = a \perp 4B_2$ with $a \equiv -1 \pmod{4}$, and $B_2 \in S_{n_1}(\mathbf{Z}_2)_e^\times$;

(II) $B_1 \in 4S_{n_1+1}(\mathbf{Z}_2)^\times$;

(III) $B_1 = a \perp 4B_2$ with $a \equiv -1 \pmod{4}$, and $B_2 \in S_{n_1}(\mathbf{Z}_2)_o$.

Then $\bar{\xi}^{(1)}(B) = 0$ if B_1 is of type (II) or (III). If B_1 is of type (I), then $(-1)^{(m-n_1)/2} a \det \Theta \pmod{(\mathbf{Z}_2^*)^\square}$ is uniquely determined by B , and we have $\bar{\xi}^{(1)}(B) = \chi((-1)^{(m-n_1)/2} a \det \Theta)$. Moreover, an element $B \in \mathcal{L}_{m,2}^{(0)}$ is equivalent, over \mathbf{Z}_2 , to a matrix of the form $\Theta \perp 2B_1$, where $\Theta \in GL_{m-n_1-2}(\mathbf{Z}_2) \cap S_{m-n_1-2}(\mathbf{Z}_2)_e$ and $B_1 \in S_{n_1}(\mathbf{Z}_2)^\times$. Suppose that $p \neq 2$, and let $\mathcal{U} = \mathcal{U}_p$ be a complete set of representatives of $\mathbf{Z}_p^*/(\mathbf{Z}_p^*)^\square$. Then, for each positive integer l and $d \in \mathcal{U}_p$, there exists a unique, up to \mathbf{Z}_p -equivalence, element of $S_l(\mathbf{Z}_p) \cap GL_l(\mathbf{Z}_p)$ whose determinant is $(-1)^{[(l+1)/2]} d$, which will be denoted by $\Theta_{l,d}$. Suppose that $p = 2$, and put $\mathcal{U} = \mathcal{U}_2 = \{1, 5\}$. Then for each positive even integer l and $d \in \mathcal{U}_2$ there exists a unique, up to \mathbf{Z}_2 -equivalence, element of $S_l(\mathbf{Z}_2)_e \cap GL_l(\mathbf{Z}_2)$ whose determinant is $(-1)^{l/2} d$, which will be also denoted by $\Theta_{l,d}$. In particular, if p is any prime number and l is even, we put $\Theta_l = \Theta_{l,1}$. We make the convention that $\Theta_{l,d}$ is the empty matrix if $l = 0$. For an element $d \in \mathcal{U}$ we use the same symbol d to denote the coset $d \pmod{(\mathbf{Z}_p^*)^\square}$.

For $T \in \mathcal{L}_{n-1,p}^{(1)}$, let $\tilde{F}_p^{(1)}(T, X)$ be the polynomial in X and X^{-1} defined in Section 3. We also define a polynomial $G_p^{(1)}(T, X)$ in X by

$$\begin{aligned} G_p^{(1)}(T, X) &= \sum_{i=0}^{n-1} (-1)^i p^{i(i-1)/2} (X^2 p^n)^i \sum_{D \in GL_{n-1}(\mathbf{Z}_p) \setminus \mathcal{D}_{n-1,i}} F_p^{(1)}(T[D^{-1}], X). \end{aligned}$$

Lemma 4.2.1. *Let n be the fixed positive even integer. Let $B \in \mathcal{L}_{n-1,p}^{(1)}$ and put $\xi_0 = \chi((-1)^{n/2} \det B)$.*

(1) *Let $p \neq 2$, and suppose that $B = \Theta_{n-n_1-1,d} \perp pB_1$ with $d \in \mathcal{U}$ and $B_1 \in S_{n_1}(\mathbf{Z}_p)^\times$. Then*

$$G_p^{(1)}(B, Y) = \begin{cases} \frac{1 - \xi_0 p^{n/2} Y}{1 - p^{n_1/2+n/2} \bar{\xi}^{(1)}(B) Y} \prod_{i=1}^{n_1/2} (1 - p^{2i+n} Y^2) & \text{if } n_1 \text{ is even,} \\ (1 - \xi_0 p^{n/2} Y) \prod_{i=1}^{(n_1-1)/2} (1 - p^{2i+n} Y^2) & \text{if } n_1 \text{ is odd.} \end{cases}$$

(2) Let $p = 2$. Suppose that n_1 is even and that $B = 2\Theta \perp B_1$ with $\Theta \in S_{n-n_1-2}(\mathbf{Z}_2)_e \cap GL_{n-n_1-2}(\mathbf{Z}_2)$ and $B_1 \in S_{n_1+1}(\mathbf{Z}_2)^\times$. Then

$$G_2^{(1)}(B, Y) = \begin{cases} \frac{1 - \xi_0 2^{n/2} Y}{1 - 2^{n_1/2+n/2} \bar{\xi}^{(1)}(B) Y} \prod_{i=1}^{n_1/2} (1 - 2^{2i+n} Y^2) & \text{if } B_1 \text{ is of type (I),} \\ (1 - \xi_0 2^{n/2} Y) \prod_{i=1}^{n_1/2} (1 - 2^{2i+n} Y^2) & \text{if } B_1 \text{ is of type (II) or (III).} \end{cases}$$

Proof. By Corollary to Lemma 4.1.2 and by definition we have $G_p^{(1)}(B, Y) = G_p(B^{(1)}, Y)$. Thus the assertion follows from Lemma 9 of [16]. \square

Remark. Throughout the above lemma, $\bar{\xi}^{(1)}(B) = \xi_0(B)$ if $n_1 = 0$. Hence we have $G_p^{(1)}(B, Y) = 1$ in this case.

Lemma 4.2.2. Let $B \in \mathcal{L}_{n-1,p}^{(1)}$. Then

$$\begin{aligned} \tilde{F}_p^{(1)}(B, X) &= \sum_{B' \in \mathcal{L}_{n-1,p}^{(1)}/GL_{n-1}(\mathbf{Z}_p)} X^{-\epsilon^{(1)}(B')} \frac{\alpha_p(B', B)}{\alpha_p(B')} \\ &\quad \times G_p^{(1)}(B', p^{(-n-1)/2} X) (p^{-1} X)^{(\nu(\det B) - \nu(\det B'))/2}. \end{aligned}$$

Proof. We have

$$\begin{aligned} \tilde{F}_p^{(1)}(B, X) &= \sum_{W \in GL_{n-1}(\mathbf{Z}_p) \setminus \tilde{\Omega}^{(1)}(B)} X^{-\epsilon^{(1)}(B)} G_p^{(1)}(B[W^{-1}], p^{(-n-1)/2} X) X^{2\nu(\det W)} \\ &= \sum_{B' \in \mathcal{L}_{n-1,p}^{(1)}/GL_{n-1}(\mathbf{Z}_p)} \sum_{W \in GL_{n-1}(\mathbf{Z}_p) \setminus \tilde{\Omega}(B', B)} X^{-\epsilon^{(1)}(B')} G_p^{(1)}(B', p^{(-n-1)/2} X) X^{2\nu(\det W)} \\ &= \sum_{B' \in \mathcal{L}_{n-1,p}^{(1)}/GL_{n-1}(\mathbf{Z}_p)} X^{-\epsilon^{(1)}(B')} \#(GL_{n-1}(\mathbf{Z}_p) \setminus \tilde{\Omega}(B', B)) p^{(\nu(\det B) - \nu(\det B'))/2} \\ &\quad \times G_p^{(1)}(B', p^{(-n-1)/2} X) (p^{-1} X)^{(\nu(\det B) - \nu(\det B'))/2}. \end{aligned}$$

Thus the assertion follows from (2) of Lemma 4.1.1. \square

4.3. Certain reduction formulas.

To give an explicit formula for the power series $P_{n-1}^{(1)}(d_0, \omega, X, t)$, we give certain reduction formulas, by which we can express $P_{n-1}^{(1)}(d_0, \omega, X, t)$ in terms of the power series defined in [11]. First we review the notion of canonical forms of the quadratic forms over \mathbf{Z}_2 in the sense of Watson [20].

Let $B \in \mathcal{L}_{m,2}^\times$. Then B is equivalent, over \mathbf{Z}_2 , to a matrix of the following form:

$$\perp_{i=0}^r 2^i (V_i \perp U_i),$$

where $V_i = \perp_{j=1}^{k_i} c_{ij}$ with $0 \leq k_i \leq 2$, $c_{ij} \in \mathbf{Z}_2^*$ and $U_i = \frac{1}{2} \Theta_{m_i, d}$ with $0 \leq m_i, d \in \mathcal{U}$. The degrees k_i and m_i of the matrices are uniquely determined by B . Furthermore we can take the matrix $\perp_{i=0}^r 2^i (V_i \perp U_i)$ uniquely so that it satisfies the following conditions:

- (c.1) $c_{i1} = \pm 1$ or ± 3 if $k_i = 1$ and $(c_{i1}, c_{i2}) = (1, \pm 1), (1, \pm 3), (-1, -1)$, or $(-1, 3)$ if $k_i = 2$;
- (c.2) $k_{i+2} = k_i = 0$ if $U_{i+2} = \frac{1}{2} \Theta_{m_{i+2}, 5}$ with $m_{i+2} > 0$;
- (c.3) $-\det V_i \equiv 1 \pmod{4}$ if $k_i = 2$ and $U_{i+1} = \frac{1}{2} \Theta_{m_{i+1}, 5}$ with $m_{i+1} > 0$;
- (c.4) $(-1)^{k_i-1} \det V_i \equiv 1 \pmod{4}$ if $k_i, k_{i+1} > 0$;
- (c.5) $V_i \neq \begin{pmatrix} -1 & 0 \\ 0 & c_{i2} \end{pmatrix}$ if $k_{i-1} > 0$;
- (c.6) $V_i = \phi, (\pm 1), \begin{pmatrix} 1 & 0 \\ 0 & \pm 1 \end{pmatrix}$, or $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ if $k_{i+2} > 0$.

The matrix satisfying the conditions (c.1), \dots , (c.6) is called the *canonical form* of B , and denote it by $C(B)$. Now for $V = \perp_{j=1}^k c_j$ with $1 \leq k \leq 2$, put $\tilde{V} = 5c_1$ or $\tilde{V} = 5c_1 \perp c_2$ according as $V = c_1$ or $V = c_1 \perp c_2$.

Lemma 4.3.1. *For $B \in S_m(\mathbf{Z}_2)_o^\times$, let $C(B) = V_0 \perp \perp_{i=1}^r (U_i \perp V_i)$ be the canonical form of B stated as above. Let $l = l_B$ be the smallest integer such that $k_{2l+2} = 0$. Then we have*

$$C(\tilde{V}_0 \perp \perp_{i=1}^r (U_i \perp V_i)) = V_0 \perp \perp_{i=1}^{2l-1} (U_i \perp V_i) \perp U_{2l} \perp C(\tilde{V}_{2l}) \perp \perp_{i=2l+1}^r (U_i \perp V_i).$$

Proof. We note that $5a_1 \perp 4a_2 \sim a_1 \perp 4 \cdot 5a_2$ for $a_1, a_2 \in \mathbf{Z}_2^*$. Hence we have

$$\tilde{V}_0 \perp \perp_{i=1}^l V_{2i} \sim V_0 \perp \perp_{i=1}^{l-1} V_{2i} \perp \tilde{V}_{2l}.$$

This proves the assertion. \square

Corollary. *For $B, B' \in S_{2m+1}(d_0)_o$, let $C(B) = V_0 \perp \perp_{i=1}^r (U_i \perp V_i)$ and $C(B') = V'_0 \perp \perp_{i=1}^{r'} (U'_i \perp V'_i)$ with $V_0 = \perp_{j=1}^{k_0} c_{0j}$ and $V'_0 = \perp_{j=1}^{k'_0} c'_{0j}$. Put*

$$B_1 = \perp_{j=2}^{k_0} c_{0j} \perp_{i=1}^r (U_i \perp V_i) \text{ and } B'_1 = \perp_{j=2}^{k'_0} c'_{0j} \perp_{i=1}^{r'} (U'_i \perp V'_i).$$

Then $B \sim B'$ if and only if $c_{01} \perp 5B_1 \sim c'_{01} \perp 5B'_1$.

Proof. We note that $c_{01} \perp 5B_1 \sim c'_{01} \perp 5B'_1$ if and only if $5c_{01} \perp B_1 \sim 5c'_{01} \perp B'_1$. Hence the assertion follows from the lemma. \square

The following lemma follows from Theorem 3.4.2 of [17].

Lemma 4.3.2. *Let m and r be integers such that $0 \leq r \leq m$, and $d_0 \in \mathbf{Z}_p^\times$.*

(1) Let $p \neq 2$, and $T \in S_r(\mathbf{Z}_p, d_0)$. Then for any $d \in \mathcal{U}$ we have

$$\varepsilon(\Theta_{m-r,d} \perp T) = ((-1)^{\lfloor (m-r+1)/2 \rfloor} d, d_0)_p \varepsilon(T).$$

Furthermore we have

$$\varepsilon(pT) = \begin{cases} (p, d_0)_p \varepsilon(T) & \text{if } r \text{ even,} \\ (p, (-1)^{(r+1)/2})_p \varepsilon(T) & \text{if } r \text{ odd,} \end{cases}$$

and $\varepsilon(aT) = (a, d_0)_p^{r+1} \varepsilon(T)$ for any $a \in \mathbf{Z}_p^*$.

(2) Let $p = 2$, and $T \in S_r(\mathbf{Z}_2, d_0)$. Suppose that $m-r$ is even, and let $d \in \mathcal{U}$.

Then for $\Theta = 2\Theta_{m-r,d}$ or $2\Theta_{m-r-2} \perp (-d)$, we have

$$\varepsilon(\Theta \perp T) = (-1)^{(m-r)(m-r+2)/8} ((-1)^{(m-r)/2} d, (-1)^{\lfloor (r+1)/2 \rfloor} d_0)_2 \varepsilon(T)$$

and

$$\varepsilon(\Theta_{m-r,d} \perp T) = (-1)^{(m-r)(m-r+2)/8} (2, d)_2 ((-1)^{(m-r)/2} d, (-1)^{\lfloor (r+1)/2 \rfloor} d_0)_2 \varepsilon(T).$$

Furthermore we have $\varepsilon(2T) = (2, d_0)_2^{r+1} \varepsilon(T)$,

$$\varepsilon(a \perp T) = (a, (-1)^{\lfloor (r+1)/2 \rfloor + 1} d_0)_2 \varepsilon(T)$$

for any $a \in \mathbf{Z}_2^*$, and

$$\varepsilon(aT) = \begin{cases} (a, d_0)_2 \varepsilon(T) & \text{if } r \text{ even,} \\ (a, (-1)^{(r+1)/2})_2 \varepsilon(T) & \text{if } r \text{ odd} \end{cases}$$

for any $a \in \mathbf{Z}_2^*$.

Henceforth, we sometimes abbreviate $S_r(\mathbf{Z}_p)$ and $S_r(\mathbf{Z}_p, d)$ as $S_{r,p}$ and $S_{r,p}(d)$, respectively. Furthermore we abbreviate $S_r(\mathbf{Z}_2)_x$ and $S_r(\mathbf{Z}_2, d)_x$ as $S_{r,2;x}$ and $S_{r,2}(d)_x$, respectively, for $x = e, o$. Let R be a commutative ring. A function H defined on a subset \mathcal{S} of $S_m(\mathbf{Q}_p)$ with values in R is said to be $GL_m(\mathbf{Z}_p)$ -invariant if $H(A[U]) = H(A)$ for any $U \in GL_m(\mathbf{Z}_p)$ and $A \in \mathcal{S}$. Let $p \neq 2$. Let $\{H_{2r+j,\xi}^{(j)} \mid j \in \{0,1\}, 1-j \leq r \leq n/2-j, \xi = \pm 1\}$ be a set of $GL_{2r+j}(\mathbf{Z}_p)$ -invariant functions on $S_{2r+j}(\mathbf{Z}_p)^\times$ with values in R satisfying the following conditions for any positive even integer $m \leq n$:

- (H-p-0) $H_{m,\xi}^{(0)}(\Theta_{m,d}) = 1$ and $H_{m-1,\xi}^{(1)}(\Theta_{m-1,d}) = 1$ for $d \in \mathcal{U}$;
- (H-p-1) $H_{m,\xi}^{(0)}(\Theta_{m-2r,d} \perp pB) = H_{2r,\xi\chi(d)}^{(0)}(pB)$ for any $r \leq m/2-1, \xi = \pm 1, d \in \mathcal{U}$ and $B \in S_{2r}(\mathbf{Z}_p)^\times$;
- (H-p-2) $H_{m-1,\xi}^{(1)}(\Theta_{m-2r-2,d} \perp pB) = H_{2r+1,\xi}^{(1)}(pdB)$ for any $r \leq m/2-2, \xi = \pm 1, d \in \mathcal{U}$ and $B \in S_{2r+1}(\mathbf{Z}_p)^\times$;
- (H-p-3) $H_{m,\xi}^{(0)}(\Theta_{m-2r-1,d} \perp pB) = H_{2r+1,\xi}^{(1)}(-pdB)$ for any $r \leq m/2-2, \xi = \pm 1$, and $B \in S_{2r+1}(\mathbf{Z}_p)^\times$;
- (H-p-4) $H_{m-1,\xi}^{(1)}(\Theta_{m-2r-1,d} \perp pB) = H_{2r,\xi\chi(d)}^{(0)}(pB)$ for any $r \leq m/2-2, \xi = \pm 1, d \in \mathcal{U}$ and $B \in S_{2r}(\mathbf{Z}_p)^\times$;

(H-p-5) $H_{2r,\xi}^{(0)}(dB) = H_{2r,\xi}^{(0)}(B)$ for any $r \leq m/2, \xi = \pm 1, d \in \mathbf{Z}_p^*$ and $B \in S_{2r}(\mathbf{Z}_p)^\times$.

Let $d_0 \in \mathcal{F}_p$, and m be a positive even integer such that $m \leq n$. Then for each $0 \leq r \leq m/2 - 1$ we put

$$Q^{(1)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \epsilon^l, t) = \kappa(d_0, m-1, l)^{-1} \\ \times \sum_{d \in \mathcal{U}} \sum_{B \in p^{-1}S_{2r+1,p}(d_0d) \cap S_{2r+1,p}} \frac{H_{m-1,\xi}^{(1)}(\Theta_{m-2r-2,d} \perp pB) \epsilon(\Theta_{m-2r-2,d} \perp pB)^l}{\alpha_p(\Theta_{m-2r-2,d} \perp pB)} t^{\nu(\det(pB))}.$$

Let $d \in \mathcal{U}$. Then we put

$$Q^{(1)}(d_0, d, H_{m-1,\xi}^{(1)}, 2r, \epsilon^l, t) = \kappa(d_0, m-1, l)^{-1} \\ \times \sum_{B \in S_{2r,p}(d_0d)} \frac{H_{m-1,\xi}^{(1)}(\Theta_{m-2r-1,d} \perp pB) \epsilon(\Theta_{m-2r-1,d} \perp pB)^l}{\alpha_p(\Theta_{m-2r-1,d} \perp pB)} t^{\nu(\det(pB))}$$

for each $1 \leq r \leq m/2 - 1$, and

$$Q^{(0)}(d_0, d, H_{m,\xi}^{(0)}, 2r, \epsilon^l, t) \\ = \sum_{B \in S_{2r,p}(d_0d)} \frac{H_{m,\xi}^{(0)}(\Theta_{m-2r,d} \perp pB) \epsilon(\Theta_{m-2r,d} \perp pB)^l}{\alpha_p(\Theta_{m-2r,d} \perp pB)} t^{\nu(\det(pB))}$$

for each $1 \leq r \leq m/2$. Here we make the convention that

$$Q^{(0)}(d_0, 1, H_{m,\xi}^{(0)}, m, \epsilon^l, t) = \sum_{B \in S_{m,p}(d_0)} \frac{H_{m,\xi}^{(0)}(pB) \epsilon(pB)^l}{\alpha_p(pB)} t^{\nu(\det(pB))}.$$

We also define

$$Q^{(1)}(d_0, d, H_{m-1,\xi}^{(1)}, 0, \epsilon^l, t) = Q^{(0)}(d_0, d, H_{m,\xi}^{(0)}, 0, \epsilon^l, t) = \delta(d, d_0),$$

where $\delta(d, d_0) = 1$ or 0 according as $d = d_0$ or not. Furthermore put

$$Q^{(0)}(d_0, H_{m,\xi}^{(0)}, 2r+1, \epsilon^l, t) \\ = \sum_{d \in \mathcal{U}} \sum_{B \in p^{-1}S_{2r+1,p}(d_0d) \cap S_{2r+1,p}} \frac{H_{m,\xi}^{(0)}(-\Theta_{m-2r-1,d} \perp pB) \epsilon(-\Theta_{m-2r-1,d} \perp pB)^l}{\alpha_p(-\Theta_{m-2r-1,d} \perp pB)} t^{\nu(\det(pB))}$$

for each $0 \leq r \leq m/2 - 1$.

Let $\{H_{2r+j,\xi}^{(j)} \mid j \in \{0, 1\}, 1-j \leq r \leq n/2-j, \xi = \pm 1\}$ be a set of $GL_{2r+j}(\mathbf{Z}_2)$ -invariant functions on $S_{2r+j}(\mathbf{Z}_2)^\times$ with values in R satisfying the following conditions for any positive even integer $m \leq n$:

(H-2-0) $H_{m,\xi}^{(0)}(\Theta_{m,d}) = H_{m-1,\xi}^{(1)}(-d \perp 2\Theta_{m-2}) = 1$ for $d \in \mathcal{U}$;

(H-2-1) $H_{m,\xi}^{(0)}(\Theta_{m-2r,d} \perp 2B) = H_{2r,\xi\chi(d)}^{(0)}(2B)$ for any $r \leq m/2 - 1, \xi = \pm 1, d \in \mathcal{U}$ and $B \in S_{2r}(\mathbf{Z}_2)^\times$;

- (H-2-2) $H_{m-1,\xi}^{(1)}(2\Theta_{m-2r-2,d}\perp 4B) = H_{2r+1,\xi}^{(1)}(4dB)$ for any $r \leq m/2 - 2$, $\xi = \pm 1$, $d \in \mathcal{U}$ and $B \in S_{2r+1}(\mathbf{Z}_2)^\times$;
- (H-2-3) $H_{m,\xi}^{(0)}(2\perp\Theta_{m-2r-2}\perp 2B) = H_{2r+1,\xi}^{(1)}(4B)$ for any $r \leq m/2 - 2$, $\xi = \pm 1$, and $B \in S_{2r+1}(\mathbf{Z}_2)^\times$;
- (H-2-4) $H_{m-1,\xi}^{(1)}(-a\perp 2\Theta_{m-2r-2}\perp 4B) = H_{2r,\xi\chi(a)}^{(0)}(2B)$ for any $r \leq m/2 - 2$, $\xi = \pm 1$, $a \in \mathcal{U}$ and $B \in S_{2r}(\mathbf{Z}_2)^\times$;
- (H-2-5) $H_{2r,\xi}^{(0)}(dB) = H_{2r,\xi}^{(0)}(B)$ for any $r \leq m/2$, $\xi = \pm 1$, $d \in \mathbf{Z}_2^*$ and $B \in S_{2r}(\mathbf{Z}_2)^\times$;
- (H-2-6) $H_{2r+1,\xi}^{(1)}(4(u_0\perp B)) = H_{2r+1,\xi}^{(1)}(4(u_0\perp 5B))$ for any $r \leq m/2 - 1$, $\xi = \pm 1$ and $u_0 \in \mathbf{Z}_2^*$, $B \in S_{2r}(\mathbf{Z}_2)^\times$.

Let $d_0 \in \mathcal{F}_2$, and m be a positive even integer such that $m \leq n$. Then for each $0 \leq r \leq m/2 - 1$, we put

$$Q^{(11)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) = \kappa(d_0, m-1, l)^{-1} t^{2-m} \\ \times \sum_{d \in \mathcal{U}} \sum_{B \in S_{2r+1,2}(d_0 d)_e} \frac{H_{m-1,\xi}^{(1)}(2\Theta_{m-2r-2,d}\perp 4B) \varepsilon^l (2\Theta_{m-2r-2,d}\perp 4B)}{\alpha_2(2\Theta_{m-2r-2,d}\perp 4B)} \\ \times t^{m-2r-2+\nu(\det(4B))},$$

$$Q^{(12)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) = \kappa(d_0, m-1, l)^{-1} t^{2-m} \\ \times \sum_{B \in S_{2r+1,2}(d_0)_o} \frac{H_{m-1,\xi}^{(1)}(2\Theta_{m-2r-2}\perp 4B) \varepsilon^l (2\Theta_{m-2r-2}\perp 4B)}{\alpha_2(2\Theta_{m-2r-2}\perp 4B)} \\ \times t^{m-2r-2+\nu(\det(4B))},$$

and

$$Q^{(13)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) = \kappa(d_0, m-1, l)^{-1} t^{2-m} \\ \times \sum_{B \in S_{2r+2,2}(d_0)_o} H_{m-1,\xi}^{(1)}(-1\perp 2\Theta_{m-2r-4}\perp 4B) \\ \times \frac{\varepsilon^l(-1\perp 2\Theta_{m-2r-4}\perp 4B)}{\alpha_2(-1\perp 2\Theta_{m-2r-4}\perp 4B)} t^{m-2r-4+\nu(\det(4B))}.$$

Moreover put

$$Q^{(1)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) = Q^{(11)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) \\ + Q^{(12)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) + Q^{(13)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t).$$

We note that

$$Q^{(1)}(d_0, H_{m-1,\xi}^{(1)}, m-1, \varepsilon^l, t) = \kappa(d_0, m-1, l)^{-1} t^{2-m} \\ \times \sum_{B \in S_{m-1,2}(d_0)} H_{m-1,\xi}^{(1)}(4B) \frac{\varepsilon(4B)^l}{\alpha_2(4B)} t^{\nu(\det(4B))}.$$

Let $d \in \mathcal{U}$. Then we put

$$\begin{aligned} Q^{(1)}(d_0, d, H_{m-1, \xi}^{(1)}, 2r, \epsilon^l, t) &= \kappa(d_0, m-1, l)^{-1} t^{2-m} \\ &\times \sum_{B \in S_{2r, 2}(d_0 d)_e} H_{m-1, \xi}^{(1)}(-d \perp 2\Theta_{m-2r-2} \perp 4B) \\ &\times \frac{\epsilon(-d \perp 2\Theta_{m-2r-2} \perp 4B)^l}{\alpha_2(-d \perp 2\Theta_{m-2r-2} \perp 4B)} t^{m-2r-2+\nu(\det(4B))}, \end{aligned}$$

for each $1 \leq r \leq m/2 - 1$, and

$$\begin{aligned} Q^{(0)}(d_0, d, H_{m, \xi}^{(0)}, 2r, \epsilon^l, t) &= \kappa(d_0, m, l)^{-1} \\ &\times \sum_{B \in S_{2r, 2}(d_0 d)_e} \frac{H_{m, \xi}^{(0)}(\Theta_{m-2r, d} \perp 2B) \epsilon(\Theta_{m-2r, d} \perp 2B)^l}{\alpha_2(\Theta_{m-2r, d} \perp 2B)} t^{\nu(\det(2B))} \end{aligned}$$

for each $1 \leq r \leq m/2$, where $\kappa(d_0, m, l) = \{(-1)^{m(m+2)/8}((-1)^{m/2} 2, d_0)_2\}^l$. Here we make the convention that

$$Q^{(0)}(d_0, 1, H_{m, \xi}^{(0)}, m, \epsilon^l) = \kappa(d_0, m, l)^{-1} \sum_{B \in S_{m, 2}(d_0)_e} \frac{H_{m, \xi}^{(0)}(2B) \epsilon(2B)^l}{\alpha_2(2B)} t^{\nu(\det(2B))}.$$

We also define

$$Q^{(1)}(d_0, d, H_{m-1, \xi}^{(1)}, 0, \epsilon^l, t) = Q^{(0)}(d_0, d, H_{m, \xi}^{(0)}, 0, \epsilon^l, t) = \delta(d, d_0).$$

Furthermore put

$$\begin{aligned} Q^{(0)}(d_0, H_{m, \xi}^{(0)}, 2r+1, \epsilon^l, t) &= \kappa(d_0, m, l)^{-1} \\ &\times \sum_{B \in S_{2r+2, 2}(d_0)_o} \frac{H_{m, \xi}^{(0)}(\Theta_{m-2r-2} \perp 2B) \epsilon(\Theta_{m-2r-2} \perp 2B)^l}{\alpha_2(\Theta_{m-2r-2} \perp 2B)} t^{\nu(\det(2B))} \end{aligned}$$

for $0 \leq r \leq m/2 - 1$. Henceforth, for $d_0 \in \mathcal{F}_p$ and nonnegative integers m, r such that $r \leq m$, put $\mathcal{U}(m, r, d_0) = \{1\}, \mathcal{U} \cap \{d_0\}$, or \mathcal{U} according as $r = 0$, $r = m \geq 1$, or $1 \leq r \leq m - 1$.

Proposition 4.3.3. *Let the notation be as above.*

(1) *For $0 \leq r \leq (m-2)/2$, we have*

$$Q^{(0)}(d_0, H_{m, \xi}^{(0)}, 2r+1, \epsilon^l, t) = \frac{Q^{(1)}(d_0, H_{2r+1, \xi}^{(1)}, 2r+1, \epsilon^l, t)}{\phi_{(m-2r-2)/2}(p^{-2})}$$

if $l\nu(d_0) = 0$, and

$$Q^{(0)}(d_0, H_{m, \xi}^{(0)}, 2r+1, \epsilon, t) = 0$$

if $\nu(d_0) > 0$.

(2) For $1 \leq r \leq m/2$ and $d \in \mathcal{U}(m, m - 2r, d_0)$, we have

$$\begin{aligned} & Q^{(0)}(d_0, d, H_{m,\xi}^{(0)}, 2r, \varepsilon^l, t) \\ &= \frac{(1 + p^{-(m-2r)/2} \chi(d)) Q^{(0)}(d_0 d, 1, H_{2r,\xi\chi(d)}^{(0)}, 2r, \varepsilon^l, t)}{2\phi_{(m-2r)/2}(p^{-2})} \end{aligned}$$

if $l\nu(d_0) = 0$, and

$$Q^{(0)}(d_0, d, H_{m,\xi}^{(0)}, 2r, \varepsilon, t) = 0$$

if $\nu(d_0) > 0$.

Proof. First suppose that $p \neq 2$. We note that

$$(-\Theta_{m-2r-1,d}) \perp pB \sim d(-\Theta_{m-2r-1}) \perp pB \approx (-\Theta_{m-2r-1}) \perp dpB$$

for $d \in \mathcal{U}$ and $B \in p^{-1}S_{2r+1,p}(d_0d)$, and the mapping

$$p^{-1}S_{2r+1,p}(d_0d) \cap S_{2r+1,p} \ni B \mapsto dB \in p^{-1}S_{2r+1,p}(d_0) \cap S_{2r+1,p}$$

is a bijection. By Lemma 4.3.2 we have $\varepsilon((-\Theta_{m-2r-1,d}) \perp pB) = (d, d_0)_p \varepsilon(pB)$, and $\varepsilon(dpB) = \varepsilon(pB)$ for $B \in p^{-1}S_{2r+1,p}(d_0d)$. Thus the assertion (1) follows from (H-p-3), (H-p-5) and Lemma 4.1.3. By (H-p-2) and Lemmas 4.1.3 and 4.3.2, we have

$$\begin{aligned} Q^{(0)}(d_0, d, H_{m,\xi}^{(0)}, 2r, \varepsilon^l, t) &= \frac{(1 + p^{-(m-2r)/2} \chi(d)) ((-1)^{(m-2r)/2} d, d_0)_p^l}{2\phi_{(m-2r)/2}(p^{-2})} \\ &\quad \times Q^{(0)}(d_0 d, 1, H_{2r,\xi\chi(d)}^{(0)}, 2r, \varepsilon^l, t). \end{aligned}$$

Thus the assertion (2) immediately follows in case $l\nu(d_0) = 0$. Now suppose that $l = 1$ and $\nu(d_0) = 1$. Take an element $a \in \mathbf{Z}_p^*$ such that $(a, p)_p = -1$. Then the mapping $S_{2r}(\mathbf{Z}_p) \ni B \mapsto aB \in S_{2r}(\mathbf{Z}_p)$ induces a bijection from $S_{2r,p}(dd_0)$ to itself, and $\varepsilon(apB) = -\varepsilon(pB)$ and $\alpha_p(apB) = \alpha_p(pB)$ for $B \in S_{2r,p}(dd_0)$. Furthermore by (H-p-5) we have

$$\begin{aligned} Q^{(0)}(d_0 d, 1, H_{2r,\xi\chi(d)}^{(0)}, 2r, \varepsilon^l, t) &= \sum_{B \in S_{2r}(dd_0)} \frac{H_{2r,\xi\chi(d)}^{(0)}(apB) \varepsilon(apB)}{\alpha_p(apB)} \\ &= -Q^{(0)}(d_0 d, 1, H_{2r,\xi\chi(d)}^{(0)}, 2r, \varepsilon^l, t). \end{aligned}$$

Hence $Q^{(0)}(d_0 d, 1, H_{2r,\xi\chi(d)}^{(0)}, 2r, \varepsilon^l, t) = 0$. This proves the assertion.

Next suppose that $p = 2$. First suppose that $l = 0$, or $l = 1$ and $d_0 \equiv 1 \pmod{4}$. Fix a complete set \mathcal{B} of representatives of $(S_{2r+2,2}(d_0)_o) / \approx$. For $B \in \mathcal{B}$, let $\mathcal{S}_{2r+1,2}(B)$ and $\tilde{\mathcal{S}}_{2r+2,2}(B)$ be those defined in Subsection 4.1.

Then, by (H-2-1) and (H-2-5) we have

$$\begin{aligned} & Q^{(0)}(d_0, H_{m,\xi}^{(0)}, 2r+1, t, t) \\ &= \sum_{B \in \mathcal{B}} \frac{H_{2r+2,\xi}^{(0)}(2B)}{\phi_{(m-2r-2)/2}(2^{-2})2^{(r+1)(2r+3)}\alpha_2(B)} \#(\tilde{\mathcal{S}}_{2r+2,2}(B)/\sim) t^{\nu(\det(2B))}. \end{aligned}$$

We have $S_{2r+1,2}(d_0) = \cup_{B \in \mathcal{B}} \mathcal{S}_{2r+1,2}(B)$, and for any $B' \in \mathcal{S}_{2r+1,2}(B)$, we have $1 \perp B' \approx B$. Hence $\nu(\det(2B)) = \nu(\det(4B')) - 2r$ and $H_{2r+2,\xi}^{(0)}(2B) = H_{2r+2,\xi}^{(0)}(2 \perp 2B') = H_{2r+1,\xi}^{(1)}(4B')$. Hence by Lemma 4.1.5 we have

$$\begin{aligned} & Q^{(0)}(d_0, H_{m,\xi}^{(0)}, 2r+1, \varepsilon^l, t) \\ &= 2 \sum_{B' \in S_{2r+1,2}(d_0)} \frac{H_{2r+1,\xi}^{(1)}(4B')}{2^{(r+1)(2r+3)}\phi_{(m-2r-2)/2}(2^{-2})\alpha_2(B')} t^{\nu(\det(4B'))-2r} \\ &= 2^{(2r+1)r} t^{-2r} \sum_{B' \in 2^{-1}S_{2r+1,2}(d_0) \cap S_{2r+1,2}} \frac{H_{2r+1,\xi}^{(1)}(4B')}{\phi_{(m-2r-2)/2}(2^{-2})\alpha_2(4B')} t^{\nu(\det(4B'))}. \end{aligned}$$

This proves the assertion for $l = 0$. Now let $d_0 \equiv 1 \pmod{4}$, and put $\xi_0 = (2, d_0)_2$. Then by Lemma 4.3.2 we have

$$\varepsilon(\Theta_{m-2r-2} \perp 2B) = (-1)^{m(m+2)/8+r(r+1)/2+(r+1)^2} \xi_0 \varepsilon(B).$$

Furthermore for any $a \in \mathbf{Z}_2^*$ we have $\varepsilon(aB)^l = \varepsilon(B)^l$, and $\alpha_2(aB) = \alpha_2(B)$.

Thus, by using the same argument as above we obtain

$$\begin{aligned} & Q^{(0)}(d_0, H_{m,\xi}^{(0)}, 2r+1, \varepsilon, t) = (-1)^{m(m+2)/8} \xi_0 \\ & \times \sum_{B \in \mathcal{B}} \frac{H_{2r+2,\xi}^{(0)}(2B)(-1)^{m(m+2)/8+r(r+1)/2+(r+1)^2} \xi_0 \varepsilon(B)}{\phi_{(m-2r-2)/2}(2^{-2})2^{(r+1)(2r+3)}\alpha_2(B)} \#(\tilde{\mathcal{S}}_{2r+2,2}(B)/\sim) t^{\nu(\det(2B))}. \end{aligned}$$

We note that $\varepsilon(1 \perp B') = \varepsilon(4B')$ for $B' \in S_{2r+1,2}$. Hence, again by Lemma 4.1.5, we have

$$\begin{aligned} & Q^{(0)}(d_0, H_{m,\xi}^{(0)}, 2r+1, \varepsilon^l, t) = (-1)^{r(r+1)/2} ((-1)^{r+1}, (-1)^{r+1})_2 2^{(2r+1)r} t^{-2r} \\ & \times \sum_{B' \in S_{2r+1,2}(d_0)} \frac{H_{2r+1,\xi}^{(1)}(4B') \varepsilon(B)}{\phi_{(m-2r-2)/2}(2^{-2})\alpha_2(4B')} t^{\nu(\det(4B'))}. \end{aligned}$$

This proves the assertion for $l = 1$ and $d_0 \equiv 1 \pmod{4}$.

Next suppose that $l = 1$ and $4^{-1}d_0 \equiv -1 \pmod{4}$, or $l = 1$ and $8^{-1}d_0 \in \mathbf{Z}_2^*$. Then there exists an element $a \in \mathbf{Z}_2^*$ such that $(a, d_0)_2 = -1$. Then the map $2B \mapsto 2aB$ induces a bijection of $2S_{2r+2,2}(d_0)_o$ to itself. Furthermore $H_{2r+2,\xi}^{(0)}(2aB) = H_{2r+2,\xi}^{(0)}(2B)$, $\varepsilon(2aB) = -\varepsilon(2B)$, and $\alpha_2(2aB) = \alpha_2(2B)$. Thus the assertion can be proved by using the same argument as in the proof of (2) for $p \neq 2$. The assertion (2) for $p = 2$ can be proved by using (H-2-1), Lemmas 4.1.4 and 4.3.2 similarly to (2) for $p \neq 2$. \square

Proposition 4.3.4. *Let the notation be as above.*

(1) For $0 \leq r \leq (m-2)/2$ we have

$$Q^{(1)}(d_0, H_{m-1, \xi}^{(1)}, 2r+1, \varepsilon^l, t) = \frac{Q^{(1)}(d_0, H_{2r+1, \xi}^{(1)}, 2r+1, \varepsilon^l, t)}{\phi_{(m-2r-2)/2}(p^{-2})}.$$

(2) For $1 \leq r \leq (m-2)/2$ and $d \in \mathcal{U}(m-1, m-2r-1, d_0)$ we have

$$Q^{(1)}(d_0, d, H_{m-1, \xi}^{(0)}, 2r, \varepsilon^l, t) = \frac{Q^{(0)}(d_0 d, 1, H_{2r, \xi \chi(d)}^{(0)}, 2r, \varepsilon^l, t)}{2\phi_{(m-2r-2)/2}(p^{-2})}$$

if $l\nu(d_0) = 0$, and

$$Q^{(1)}(d_0, d, H_{m-1, \xi}^{(0)}, 2r, \varepsilon^l, t) = 0$$

otherwise.

Proof. We may suppose that $r < (m-2)/2$. First suppose that $p \neq 2$. As in the proof of Proposition 4.3.3 (1), we have a bijection $p^{-1}S_{2r+1, p}(d_0 d) \cap S_{2r+1, p} \ni B \mapsto dB \in p^{-1}S_{2r+1, p}(d_0) \cap S_{2r+1, p}$. We also note that $\varepsilon(dB) = \varepsilon(B)$, and $\alpha_p(dB) = \alpha_p(B)$. Hence, by (H-p-2), Lemmas 4.1.3 and 4.3.2, similarly to Proposition 4.3.3 (2), we have

$$\begin{aligned} Q^{(1)}(d_0, H_{m, \xi}^{(1)}, 2r+1, \varepsilon^l, t) &= p^{(m/2-1)\nu(d_0)}((-1)^{m/2}d_0, (-1)^{lm/2})_p \\ &\times \sum_{B \in p^{-1}S_{2r+1, p}(d_0) \cap S_{2r+1, p}} \frac{H_{2r+1, \xi}^{(1)}(pB)\varepsilon(pB)^l}{2\phi_{(m-2r-2)/2}(p^{-2})\alpha_p(pB)} t^{\nu(\det(pB))} \\ &\times \sum_{d \in \mathcal{U}} (1 + p^{-(m-2r-2)/2}\chi(d))((-1)^{(m-2r-2)/2}d, (-1)^{r+1}d_0 d)_p^l. \end{aligned}$$

Thus the assertion clearly holds if $l\nu(d_0) = 0$. Suppose that $l = 1$ and $\nu(d_0) = 1$. Then

$$\begin{aligned} &((-1)^{(m-2r-2)/2}d, (-1)^{r+1}d_0 d)_p \\ &= \chi(d)((-1)^{r+1}, (-1)^{r+1}d_0 d)_p((-1)^{m/2}, (-1)^{m/2}d_0)_p, \end{aligned}$$

and therefore

$$\begin{aligned} &\sum_{d \in \mathcal{U}} (1 + p^{-(m-2r-2)/2}\chi(d))((-1)^{(m-2r-2)/2}d, (-1)^{r+1}d_0)_p \\ &= 2p^{-(m-2r-2)/2}((-1)^{r+1}, (-1)^{r+1}d_0 d)_p((-1)^{m/2}, (-1)^{m/2}d_0)_p. \end{aligned}$$

This proves the assertion (1).

By (H-p-4) and by Lemmas 4.1.3 and 4.3.2, we have

$$\begin{aligned} &Q^{(1)}(d_0, d, H_{m-1, \xi}^{(1)}, 2r, \varepsilon^l, t) \\ &= \frac{Q^{(0)}(d_0 d, 1, H_{2r, \xi \chi(d)}^{(0)}, 2r, \varepsilon^l, t)}{2\phi_{(m-2r-2)/2}(p^{-2})}((-1)^{(m-2r)/2}d, d_0)_p^l. \end{aligned}$$

Thus the assertion (2) immediately follows if $l\nu(d_0) = 0$. The assertion for $l = 1$ and $\nu(d_0) = 1$ follows from Proposition 4.3.3 (2).

Next suppose that $p = 2$. We have

$$\begin{aligned} \varepsilon(2\Theta_{m-2r-2,d}\perp 4B) &= (-1)^{m(m-2)/8}(-1)^{r(r+1)/2}((-1)^{m/2}, (-1)^{m/2}d_0)_2 \\ &\quad \times ((-1)^{r+1}, (-1)^{r+1}d_0d)_2(d_0, d)_2 \varepsilon(4B) \end{aligned}$$

for $d \in \mathcal{U}$ and $B \in S_{2r+1,2}(dd_0)$. Thus, similarly to (1) for $p \neq 2$, we obtain

$$\begin{aligned} Q^{(11)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) &= (-1)^{r(r+1)l/2}t^{-2r}((-1)^{r+1}, (-1)^{r+1}d_0)_2^l \\ &\quad \times 2^{(m/2-1)l\nu(d_0)} \sum_{B \in S_{2r+1,2}(d_0)_e} \frac{2^{r(2r+1)}H_{2r+1,\xi}^{(1)}(4B)\varepsilon(4B)^l}{2 \cdot 2^{m-2r-2}\phi_{(m-2r-2)/2}(2^{-2})\alpha_2(4B)} t^{\nu(\det(4B))} \\ &\quad \times \sum_{d \in \mathcal{U}} (1 + 2^{-(m-2r-2)/2}\chi(d))(d, d_0)_2^l \\ &= \sum_{d \in \mathcal{U}} (1 + 2^{-(m-2r-2)/2}\chi(d))(d, d_0)_2^l \frac{Q^{(11)}(d_0, H_{2r+1,\xi}^{(1)}, 2r+1, \varepsilon^l, t)}{2^{1+(m-2r-2)(1-l\nu(d_0)/2)}\phi_{(m-2r-2)/2}(2^{-2})}. \end{aligned}$$

In the same manner as above, we obtain

$$\begin{aligned} Q^{(12)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) &= (-1)^{r(r+1)l/2}t^{-2r}((-1)^{r+1}, (-1)^{r+1}d_0)_2^l \\ &\quad \times 2^{(m/2-1)l\nu(d_0)} \sum_{B \in S_{2r+1,2}(d_0)_o} \frac{2^{r(2r+1)}H_{2r+1,\xi}^{(1)}(4B)\varepsilon(4B)^l}{2^{m-2r-2}\phi_{(m-2r-2)/2}(2^{-2})\alpha_2(4B)} t^{\nu(\det(4B))} \\ &= \frac{Q^{(11)}(d_0, H_{2r+1,\xi}^{(1)}, 2r+1, \varepsilon^l, t)}{2^{(m-2r-2)(1-l\nu(d_0)/2)}\phi_{(m-2r-2)/2}(2^{-2})}. \end{aligned}$$

Furthermore we have

$$\begin{aligned} \varepsilon(-1\perp 2\Theta_{m-2r-4}\perp 4B) &= (-1)^{m(m-2)/8}(-1)^{r(r+1)/2}((-1)^{m/2}, (-1)^{m/2}d_0)_2 \\ &\quad \times ((-1)^{r+1}, (-1)^{r+1}d_0)_2(2, d_0)_2\varepsilon(2B) \end{aligned}$$

for $d \in \mathcal{U}$ and $B \in S_{2r+2,2}(dd_0)_o$. Hence

$$\begin{aligned} Q^{(13)}(d_0, H_{m-1,\xi}^{(1)}, 2r+1, \varepsilon^l, t) &= (-1)^{r(r+1)l/2}t^{-2r-2}((-1)^{r+1}, (-1)^{r+1}d_0)_2^l \\ &\quad \times (2, d_0)_2^l 2^{(m/2-1)l\nu(d_0)} \sum_{B \in S_{2r+2,2}(d_0)_o} \frac{H_{2r+2,\xi}^{(0)}(2B)\varepsilon(4B)^l}{\phi_{(m-2r-4)/2}(2^{-2})\alpha_2(2B)} t^{\nu(\det(4B))} \\ &= (((-1)^{r+1}2, d_0)_2(-1)^{(r+1)(r+2)/2})^l 2^{(m/2-1)l\nu(d_0)} \\ &\quad \times \sum_{B \in S_{2r+2,2}(d_0)_o} \frac{H_{2r+2,\xi}^{(0)}(2B)\varepsilon(2B)^l}{\phi_{(m-2r-4)/2}(2^{-2})\alpha_2(2B)} t^{\nu(\det(2B))} \\ &= \frac{Q^{(0)}(d_0, H_{2r+2,\xi}^{(0)}, 2r+1, \varepsilon^l, t)}{\phi_{(m-2r-4)/2}(2^{-2})} 2^{(m/2-1)l\nu(d_0)}. \end{aligned}$$

First suppose that $l = 0$ or $\nu(d_0)$ is even. Then $(d, d_0)_2^l = 1$. Hence

$$\begin{aligned} & Q^{(11)}(d_0, H_{m-1, \xi}^{(1)}, 2r+1, \varepsilon^l, t) + Q^{(12)}(d_0, H_{m-1, \xi}^{(1)}, 2r+1, \varepsilon^l, t) \\ &= \frac{Q^{(1)}(d_0, H_{2r+1, \xi}^{(1)}, 2r+1, \varepsilon^l, t)}{2^{(m-2r-2)(1-\nu(d_0)l/2)} \phi_{(m-2r-2)/2}(2^{-2})}. \end{aligned}$$

Furthermore by Proposition 4.3.3 (2), we have

$$Q^{(13)}(d_0, H_{m-1, \xi}^{(1)}, 2r+1, \varepsilon^l, t) = \frac{Q^{(1)}(d_0, H_{2r+1, \xi}^{(1)}, 2r+1, \varepsilon^l, t)}{\phi_{(m-2r-4)/2}(2^{-2})}$$

if $l\nu(d_0) = 0$, and

$$Q^{(13)}(d_0, H_{m-1, \xi}^{(1)}, 2r+1, \varepsilon, t) = 0$$

if $4^{-1}d_0 \equiv -1 \pmod{4}$. Thus summing up these two quantities, we prove the assertion. Next suppose that $l = 1$ and $\nu(d_0) = 3$. Then, we have

$$Q^{(13)}(d_0, H_{m-1, \xi}^{(1)}, 2r+1, \varepsilon, t) = 0.$$

We prove

$$Q^{(12)}(d_0, H_{2r+1, \xi}^{(1)}, 2r+1, \varepsilon, t) = 0.$$

If $r = 0$, then clearly $S_{2r+1, 2}(d_0)_o$ is empty. Suppose that $r \geq 1$. Then for $B \in 4S_{2r+1, 2; o}$ take a canonical form $4c_{01} \perp 4B_1$ with $c_{01} \in \mathbf{Z}_2^*$, $B_1 \in S_{2r, 2}$, and put $B' = 4c_{01} \perp 4 \cdot 5B_1$. Then, by Corollary to Lemma 4.3.1, the mapping $B \mapsto B'$ induces a bijection from $4S_{2r+1, 2}(d_0)_o / \sim$ to itself, and $\varepsilon(B') = -\varepsilon(B)$. Then, by (H-2-6), and Lemma 4.1.4 (3), we can prove the above equality in the same way as in the proof of (1) for $p \neq 2$. We also note that $\sum_{d \in \mathcal{U}} (1 + 2^{-(m-2r-2)/2} \chi(d))(d, d_0)_2 = 2^{1-(m-2r-2)/2}$. This proves the assertion.

The assertion (2) for $p = 2$ can be proved in the same manner as in (2) for $p \neq 2$. \square

4.4. Proof of the main result.

In this section, we prove our main result. First we give an explicit formula for the power series of Koecher-Maass type.

Theorem 4.4.1. *Let $d_0 \in \mathcal{F}_p$, and put $\xi_0 = \chi(d_0)$. Then we have the following:*

$$\begin{aligned} (1) \quad P_{n-1}^{(1)}(d_0, t, X, t) &= \frac{(p^{-1}t)^{\nu(d_0)}(1 - \xi_0 t^2 p^{-5/2})}{\phi_{(n-2)/2}(p^{-2})(1 - t^2 p^{-2} X)(1 - t^2 p^{-2} X^{-1})} \\ &\quad \times \frac{1}{\prod_{i=1}^{(n-2)/2} (1 - t^2 p^{-2i-1} X)(1 - t^2 p^{-2i-1} X^{-1})}. \end{aligned}$$

$$(2) P_{n-1}^{(1)}(d_0, \varepsilon, X, t) = \frac{(p^{-1}t)^{\nu(d_0)}(1 - \xi_0 t^2 p^{-1/2-n})}{\phi_{(n-2)/2}(p^{-2})} \\ \times \frac{1}{\prod_{i=1}^{(n-2)/2} (1 - t^2 p^{-2i-1} X)(1 - t^2 p^{-2i-1} X^{-1})}.$$

To prove the above theorem, we define another formal power series. Namely, for $l = 0, 1$ we define $K_{n-1}^{(1)}(d_0, \varepsilon^l, X, t)$ as

$$K_{n-1}^{(1)}(d_0, \varepsilon^l, X, t) = \kappa(d_0, n-1, l)^{-1} t^{\delta_{2,p}(2-n)} \\ \times \sum_{B' \in \mathcal{L}_{n-1,p}^{(1)}(d_0)} \frac{G_p^{(1)}(B', p^{-(n+1)/2} X) \varepsilon(B')^l}{\alpha_p(B')} X^{-\epsilon^{(1)}(B')} t^{\nu(\det B')}.$$

Proposition 4.4.2. *Let d_0 be as above. Then, we have*

$$P_{n-1}^{(1)}(d_0, \omega, X, t) = \prod_{i=1}^{n-1} (1 - t^2 X p^{i-n-1})^{-1} K_{n-1}^{(1)}(d_0, \omega, X, t).$$

Proof. We note that B' belongs to $\mathcal{L}_{n-1,p}^{(1)}(d_0)$ if B belongs to $\mathcal{L}_{n-1,p}^{(1)}(d_0)$ and $\alpha_p(B', B) \neq 0$. Hence by Lemma 4.2.2 for $\omega = \varepsilon^l$ with $l = 0, 1$ we have

$$P_{n-1}^{(1)}(d_0, \omega, X, t) = \kappa(d_0, n-1, l)^{-1} t^{\delta_{2,p}(2-n)} \\ \times \sum_{B \in \mathcal{L}_{n-1,p}^{(1)}(d_0)} \frac{1}{\alpha_p(B)} \sum_{B'} \frac{G_p^{(1)}(B', p^{-(n+1)/2} X) X^{-\epsilon^{(1)}(B')} \alpha_p(B', B) \omega(B')}{\alpha_p(B')} \\ \times (p^{-1} X)^{(\nu(\det B) - \nu(\det B'))/2} t^{\nu(\det B)} \\ = \kappa(d_0, n-1, l)^{-1} t^{\delta_{2,p}(2-n)} \sum_{B' \in \mathcal{L}_{n-1,p}^{(1)}(d_0)} \frac{G_p^{(1)}(B', p^{-(n+1)/2} X) \omega(B')}{\alpha_p(B')} X^{-\epsilon^{(1)}(B')} \\ \times \sum_{B \in \mathcal{L}_{n-1,p}^{(1)}(d_0)} \frac{\alpha_p(B', B)}{\alpha_p(B)} (p^{-1} X)^{(\nu(\det B) - \nu(\det B'))/2} t^{\nu(\det B)}.$$

Hence by Theorem 5 of [4], and by (1) of Lemma 4.1.1, we have

$$\sum_B \frac{\alpha_p(B', B)}{\alpha_p(B)} (p^{-1} X)^{(\nu(\det B) - \nu(\det B'))/2} t^{\nu(\det B)} \\ = \sum_{W \in M_{n-1}(\mathbf{Z}_p)^\times / GL_{n-1}(\mathbf{Z}_p)} (t^2 X p^{-1} p^{-n+1})^{\nu(\det W)} t^{\nu(\det B')} \\ = \prod_{i=1}^{n-1} (1 - t^2 X p^{i-n-1})^{-1} t^{\nu(\det B')}.$$

Thus the assertion holds. \square

For a variable X we introduce the symbol $X^{1/2}$ so that $(X^{1/2})^2 = X$, and for an integer a write $X^{a/2} = (X^{1/2})^a$. Under this convention, we can write

$X^{-\epsilon^{(1)}(T)}t^{\nu(\det T)}$ as $X^{\delta_{2,p}(n-2)/2}X^{\nu(d_0)/2}(X^{-1/2}t)^{\nu(\det T)}$ if $T \in \mathcal{L}'_{r-1,p}(d_0)$, and hence we can write $K_{n-1}^{(1)}(d_0, \epsilon^l, X, t)$ as

$$K_{n-1}^{(1)}(d_0, \epsilon^l, X, t) = \kappa(d_0, n-1, l)^{-1}(tX^{-1/2})^{\delta_{2,p}(2-n)}X^{\nu(d_0)/2} \\ \times \sum_{B' \in \mathcal{L}'_{n-1,p}(d_0)} \frac{G_p^{(1)}(B', p^{-(n+1)/2}X)\epsilon(B')^l}{\alpha_p(B')} (tX^{-1/2})^{\nu(\det B')}.$$

In order to prove Theorem 4.4.1, we introduce some power series. Let m be an integer and $l = 0$ or 1 . Then for $d_0 \in \mathbf{Z}_p^\times$ put

$$\zeta_m(d_0, \epsilon^l, u) = \sum_{T \in S_{m,p}(d_0)/\sim} \frac{\epsilon(T)^l}{\alpha_p(T)} u^{\nu(\det T)},$$

and for $d_0 \in \mathbf{Z}_2^\times$ put

$$\zeta_m^*(d_0, \epsilon^l, u) = \sum_{T \in S_{m,2}(d_0)_e/\sim} \frac{\epsilon(T)^l}{\alpha_2(T)} u^{\nu(\det T)}.$$

We make the convention that $\zeta_0(d_0, \epsilon^l, u) = \zeta_m^*(d_0, \epsilon^l, u) = 1$ or 0 according as $d_0 \in \mathbf{Z}_p^*$ or not. Now for $d \in \mathbf{Z}_p^\times$, let $Z_m(u, \epsilon^l, d)$ and $Z_m^*(u, \epsilon^l, d)$ be the formal power series in Theorems 5.1, 5.2, and 5.3 of [11], which are given by

$$Z_m(u, \epsilon^l, d) = 2^{-\delta_{2,p}m} \sum_{i=0}^{\infty} \sum_{T \in \mathbf{S}_m(\mathbf{Z}_p, p^i d)/\sim} \frac{\epsilon(T)^l}{\alpha_p(T)} (\eta_m^l p^{(m+1)/2} u)^i$$

and

$$Z_m^*(u, \epsilon^l, d) = 2^{-m} \sum_{i=0}^{\infty} \sum_{T \in \mathbf{S}_m(\mathbf{Z}_2, 2^i d)_e/\sim} \frac{\epsilon(T)^l}{\alpha_2(T)} (\eta_m^l 2^{(m+1)/2} u)^i,$$

where $\mathbf{S}_m(\mathbf{Z}_p, a) = \{T \in S_m(\mathbf{Z}_p) \mid \det T = a \pmod{\mathbf{Z}_p^{\times\Box}}\}$, $\mathbf{S}_m(\mathbf{Z}_p, a)_e = \mathbf{S}_m(\mathbf{Z}_p, a) \cap S_m(\mathbf{Z}_p)_e$, and $\eta_m = ((-1)^{(m+1)/2}, p)_p$ or 1 according as m is odd or even. Here we recall that the local density for $T \in S_m(\mathbf{Z}_p)$ in our paper is $2^{-\delta_{2,p}m}$ times that in [11]. Put

$$Z_{m,e}(u, \epsilon^l, d) = \frac{1}{2}(Z_m(u, \epsilon^l, d) + Z_m(-u, \epsilon^l, d)), \\ Z_{m,o}(u, \epsilon^l, d) = \frac{1}{2}(Z_m(u, \epsilon^l, d) - Z_m(-u, \epsilon^l, d)).$$

We also define $Z_{m,e}^*(u, \epsilon^l, d)$ and $Z_{m,o}^*(u, \epsilon^l, d)$ in the same way. Furthermore put $x(i) = e$ or o according as i is even or odd. Let $d_0 \in \mathcal{F}_p$. Let $p \neq 2$. Then

$$\zeta_m(d_0, \epsilon^l, u) = Z_{m,x(\nu(d_0))}(p^{-(m+1)/2}((-1)^{(m+1)/2}, p)_p u, \epsilon^l, p^{-\nu(d_0)}(-1)^{(m+1)/2}d_0)$$

or

$$\zeta_m(d_0, \epsilon^l, u) = Z_{m,x(\nu(d_0))}(p^{-(m+1)/2}u, \epsilon^l, p^{-\nu(d_0)}(-1)^{\lfloor (m+1)/2 \rfloor}d_0)$$

according as m is odd and $l = 1$, or not. Let $p = 2$, and m is odd. Then

$$\zeta_m(d_0, \varepsilon^l, u) = 2^m Z_{m, x(\nu(d_0))}(2^{-(m+1)/2}u, \varepsilon^l, 2^{-\nu(d_0)}(-1)^{(m+1)/2}d_0).$$

Let $p = 2$ and m is even. Then

$$\zeta_m^*(d_0, \varepsilon^l, u) = 2^m Z_{m, x(\nu(d_0))}^*(2^{-(m+1)/2}u, \varepsilon^l, (-1)^{m/2}2^{-\nu(d_0)}d_0).$$

Proposition 4.4.3. *Let $d_0 \in \mathcal{F}_p$. For a positive even integer r and $d \in \mathcal{U}$ put*

$$c(r, d_0, d, X) = (1 - \chi(d_0)p^{-1/2}X) \prod_{i=1}^{r/2-1} (1 - p^{2i-1}X^2)(1 + \chi(d)p^{r/2-1/2}X),$$

and put $c(0, d_0, d, X) = 1$. Furthermore, for a positive odd integer r put

$$c(r, d_0, X) = (1 - \chi(d_0)p^{-1/2}X) \prod_{i=1}^{(r-1)/2} (1 - p^{2i-1}X^2).$$

(1) Suppose that $p \neq 2$.

(1.1) Let $l = 0$ or $\nu(d_0) = 0$. Then

$$\begin{aligned} & K_{n-1}^{(1)}(d_0, \varepsilon^l, X, t) \\ &= X^{\nu(d_0)/2} \left\{ \sum_{r=0}^{(n-2)/2} \sum_{d \in \mathcal{U}(n-1, n-2r-1, d_0)} \frac{p^{-r(2r+1)}(tX^{-1/2})^{2r} c(2r, d_0, d, X)}{2^{1-\delta_{0,r}} \phi_{(n-2r-2)/2}(p^{-2})} \right. \\ & \quad \times (p, d_0 d)_p^l \zeta_{2r}(d_0 d, \varepsilon^l, tX^{-1/2}) \\ & \quad \left. + \sum_{r=0}^{(n-2)/2} \frac{p^{-(r+1)(2r+1)}(tX^{-1/2})^{2r+1} c(2r+1, d_0, X)}{\phi_{(n-2r-2)/2}(p^{-2})} \zeta_{2r+1}(p^* d_0, \varepsilon^l, tX^{-1/2}) \right\}, \end{aligned}$$

where $p^* d_0 = pd_0$ or $p^{-1}d_0$ according as $\nu(d_0) = 0$ or $\nu(d_0) = 1$.

(1.2) Let $\nu(d_0) = 1$. Then

$$\begin{aligned} & K_{n-1}^{(1)}(d_0, \varepsilon, X, t) \\ &= X^{\nu(d_0)/2} \sum_{r=0}^{(n-2)/2} \frac{p^{-(r+1)(2r+1)-r}(tX^{-1/2})^{2r+1} c(2r+1, d_0, X)}{\phi_{(n-2r-2)/2}(p^{-2})} \\ & \quad \times \zeta_{2r+1}(p^{-1}d_0, \varepsilon, tX^{-1/2}). \end{aligned}$$

(2) Suppose that $p = 2$.

(2.1) Let $l = 0$ or $d_0 \equiv 1 \pmod{4}$. Then

$$\begin{aligned}
& K_{n-1}^{(1)}(d_0, \varepsilon^l, X, t) \\
&= X^{\nu(d_0)/2} \left\{ \sum_{r=0}^{(n-2)/2} \sum_{d \in \mathcal{U}(n-1, n-2r-1, d_0)} (tX^{-1})^{2r} 2^{-r(2r+1)} \frac{c(2r, d_0, d, X)}{2^{1-\delta_{0,r}} \phi_{(n-2r-2)/2}(2^{-2})} \right. \\
&\quad \times ((-1)^{(r+1)r/2} (2, d_0 d)_2)^l \zeta_{2r}^*(d_0 d, \varepsilon, tX^{-1/2}) \\
&\quad + \sum_{r=0}^{(n-2)/2} (tX^{-1/2})^{2r+1} 2^{-(r+1)(2r+1)} \frac{c(2r+1, d_0, X)}{\phi_{(n-2r-2)/2}(2^{-2})} \\
&\quad \left. \times ((-1)^{(r+1)r/2} ((-1)^{r+1}, (-1)^{r+1} d_0)_2)^l \zeta_{2r+1}(d_0, \varepsilon^l, tX^{-1/2}) \right\}.
\end{aligned}$$

(2.2) Suppose that $4^{-1}d_0 \equiv -1 \pmod{4}$ or $8^{-1}d_0 \in \mathbf{Z}_2^*$. Then

$$\begin{aligned}
& K_{n-1}^{(1)}(d_0, \varepsilon, X, t) \\
&= X^{\nu(d_0)/2} \sum_{r=0}^{(n-2)/2} (tX^{-1/2})^{2r+1} 2^{-(r+1)(2r+1)-r\nu(d_0)} \frac{c(2r+1, d_0, X)}{\phi_{(n-2r-2)/2}(2^{-2})} \\
&\quad \times (-1)^{(r+1)r/2} ((-1)^{r+1}, (-1)^{r+1} d_0)_2 \zeta_{2r+1}(d_0, \varepsilon, tX^{-1/2}).
\end{aligned}$$

Proof. Put $H_{2r+j, \xi}^{(j)}(B) = 1$ for $j \in \{0, 1\}$, $1-j \leq r \leq m/2 - j$, $\xi = \pm 1$, and $B \in S_{2r+j, p}$. Then clearly the set $\{H_{2r+j, \xi}^{(j)} \mid j \in \{0, 1\}, 1-j \leq r \leq n/2 - j, \xi = \pm 1\}$ satisfy the conditions (H-p-0), \dots , (H-p-5) in Subsection 4.3 for any positive even integer $m \leq n$. Hence by Lemma 4.2.1 and Proposition 4.3.4, and by using the same argument as in Lemma 3.1 (1) of [10], we have

$$\begin{aligned}
& K_{n-1}^{(1)}(d_0, \varepsilon^l, X, t) \\
&= \gamma_{l, d_0} X^{\nu(d_0)/2} \sum_{r=0}^{(n-2)/2} \sum_{d \in \mathcal{U}(n-1, n-2r-1, d_0)} \frac{c(2r, d_0, d, X)}{2^{1-\delta_{0,r}} \phi_{(n-2r-2)/2}(p^{-2})} \\
&\quad \times \sum_{B \in S_{2r, p}(d_0 d)} \frac{\varepsilon(pB)^l}{\alpha_p(pB)} (tX^{-1/2})^{\nu(\det(pB))} \\
&+ X^{\nu(d_0)/2} \sum_{r=0}^{(n-2)/2} \frac{c(2r+1, d_0, d, X)}{\phi_{(n-2r-2)/2}(p^{-2})} \\
&\quad \times \sum_{B \in p^{-1}S_{2r+1, p}(d_0) \cap S_{2r+1, p}} \frac{((-1)^{(r+1)/2}, (-1)^{(r+1)/2} d_0)_p^l p^{-l\nu(d_0)} \varepsilon(pB)^l}{\alpha_p(pB)} \\
&\quad \times (tX^{-1/2})^{\nu(\det(pB))},
\end{aligned}$$

where $\gamma_{l, d_0} = 1$ or 0 according as $\nu(d_0)l = 0$ or 1 . Thus the assertion (1.1) follows from Lemmas 4.1.3 and 4.3.2 by remarking that $p^{-1}S_{2r+1, p}(d_0) \cap$

$S_{2r+1,p} = S_{2r+1}(p^*d_0)$. Similarly the assertion (1.2) can be proved by remarking that $\varepsilon(pB) = ((-1)^{r+1}, p)\varepsilon(B)$ for $B \in p^{-1}S_{2r+1,p}(d_0) \cap S_{2r+1,p}$. The assertion for $p = 2$ can also be proved in the same manner as above. \square

Remark. As seen above, to prove Proposition 4.4.3, we have only to prove Propositions 4.3.2 and 4.3.3 for the simplest case where $\{H_{2r+j,\xi}^{(j)}\}$ are constant functions. However a similar statement for more general $\{H_{2r+j,\xi}^{(j)}\}$ will be necessary for giving an explicit formula for the Rankin-Selberg series of $\sigma_{n-1}(\phi_{I_n(h),1})$ (cf. [15]). Indeed, the proofs are essentially the same as those for the simplest case. This is why we formulate and prove those propositions in more general settings.

Proof of Theorem 4.4.1 in case $p \neq 2$. (1) First let $d_0 \in \mathbf{Z}_p^*$. Then by Proposition 4.4.3 (1.1), we have

$$\begin{aligned} K_{n-1}^{(1)}(d_0, \iota, X, t) &= \frac{1}{\phi_{(n-2)/2}(p^{-2})} \\ &+ \sum_{r=1}^{(n-2)/2} \sum_{d \in \mathcal{U}} \frac{p^{-r(2r+1)}(t^2 X^{-1})^r \prod_{i=1}^{r-1} (1 - p^{2i-1} X^2)}{2\phi_{(n-2r-2)/2}(p^{-2})} \\ &\quad \times (1 - p^{-1/2}\xi_0 X)(1 + \eta_d p^{r-1/2} X) \zeta_{2r}(d_0 d, \iota, tX^{-1/2}) \\ &+ \sum_{r=0}^{(n-2)/2} \frac{p^{-(2r+1)(r+1)}(t^2 X^{-1})^{r+1/2} \prod_{i=1}^r (1 - p^{2i-1} X^2)}{\phi_{(n-2r-2)/2}(p^{-2})} \\ &\quad \times (1 - p^{-1/2}\xi_0 X) \zeta_{2r+1}(pd_0, \iota, tX^{-1/2}). \end{aligned}$$

Here we put $\eta_d = \chi(d)$ for $d \in \mathcal{U}$. By Theorem 5.1 of [11], we have

$$\zeta_{2r+1}(pd_0, \iota, tX^{-1/2}) = \frac{p^{-1}tX^{-1/2}}{\phi_r(p^{-2})(1 - p^{-2}t^2 X^{-1}) \prod_{i=1}^r (1 - p^{2i-3-2r}t^2 X^{-1})},$$

and

$$\zeta_{2r}(d_0 d, \iota, tX^{-1/2}) = \frac{(1 + \xi_0 \eta_d p^{-r})(1 - \xi_0 \eta_d p^{-r-2}t^2 X^{-1})}{\phi_r(p^{-2})(1 - p^{-2}t^2 X^{-1}) \prod_{i=1}^r (1 - p^{2i-3-2r}t^2 X^{-1})}.$$

Hence the assertion for $n = 2$ can be proved by a direct calculation. Suppose that $n \geq 4$. Then $K_{n-1}^{(1)}(d_0, \iota, X, t)$ can be expressed as

$$K_{n-1}^{(1)}(d_0, \iota, X, t) = \frac{S(d_0, \iota, X, t)}{\phi_{(n-2)/2}(p^{-2})(1 - p^{-2}t^2 X^{-1}) \prod_{i=1}^{(n-2)/2} (1 - p^{2i-n-1}t^2 X^{-1})},$$

where $S(d_0, \iota, X, t)$ is a polynomial in t of degree n . We have

$$\begin{aligned} &2^{-1}(1 - p^{-1/2}\xi_0 X) \sum_{\eta=\pm 1} (1 + \eta p^{(n-2)/2-1/2} X)(1 + \xi_0 \eta p^{-(n-2)/2})(1 - \xi_0 \eta p^{-(n-2)/2-2}t^2 X^{-1}) \\ &= (1 - \xi_0 p^{-1/2} X)(1 + \xi_0 p^{-1/2} X - \xi_0 p^{-5/2} t^2 - p^{-n} t^2 X^{-1}). \end{aligned}$$

Hence

$$\begin{aligned}
& 2^{-1} \sum_{d \in \mathcal{U}} p^{(n-1)(-n+2)/2} (t^2 X^{-1})^{(n-2)/2} \prod_{i=1}^{(n-2)/2-1} (1 - p^{2i-1} X^2) \\
& \quad \times (1 - p^{-1/2} \xi_0 X) (1 + \eta_d p^{(n-2)/2-1/2} X) \zeta_{n-2}(d_0 d, \iota, t X^{-1/2}) \\
& + p^{-(n-1)n/2} (t^2 X^{-1})^{(n-2)/2+1/2} \prod_{i=1}^{(n-2)/2} (1 - p^{2i-1} X^2) (p^{-2})^{-1} \\
& \quad \times (1 - p^{-1/2} \xi_0 X) \zeta_{n-1}(p d_0, \iota, t X^{-1/2}) \\
& = \frac{(p^{-(n-1)} X^{-1} t^2)^{(n-2)/2} (1 - \xi_0 p^{-5/2} t^2) \prod_{i=0}^{(n-2)/2-1} (1 - p^{2i-1} X^2)}{\phi_{(n-2)/2}(p^{-2}) (1 - p^{-2} t^2 X^{-1}) \prod_{i=1}^{(n-2)/2} (1 - p^{2i-n-1} t^2 X^{-1})},
\end{aligned}$$

and therefore $S(d_0, \iota, X, t)$ can be expressed as

$$\begin{aligned}
& \text{(A)} \quad S(d_0, \iota, X, t) \\
& = (p^{-(n-1)} X^{-1} t^2)^{(n-2)/2} \prod_{i=0}^{(n-2)/2-1} (1 - p^{2i-1} X^2) (1 - p^{-5/2} \xi_0 t^2) \\
& \quad + (1 - p^{-n+1} t^2 X^{-1}) U(X, t),
\end{aligned}$$

where $U(X, t)$ is a polynomial in X, X^{-1} and t . Now by Proposition 4.4.2, we have

$$\begin{aligned}
& P_{n-1}^{(1)}(d_0, \iota, X, t) \\
& = \frac{S(d_0, \iota, X, t)}{\phi_{(n-2)/2}(p^{-2}) (1 - p^{-2} t^2 X^{-1}) \prod_{i=1}^{(n-2)/2} (1 - p^{2i-n-1} t^2 X^{-1}) \prod_{i=1}^{n-1} (1 - p^{i-n-1} X t^2)}.
\end{aligned}$$

Hence the power series $P_{n-1}^{(1)}(d_0, \iota, X, t)$ is a rational function in X and t . Since we have $\tilde{F}_p^{(1)}(T, X^{-1}) = \tilde{F}_p^{(1)}(T, X)$ for any $T \in \mathcal{L}_{n-1, p}^{(1)}$, we have $P_{n-1}^{(1)}(d_0, \iota, X^{-1}, t) = P_{n-1}^{(1)}(d_0, \iota, X, t)$. This implies that the reduced denominator of the rational function $P_{n-1}^{(1)}(d_0, \iota, X, t)$ in t is at most

$$(1 - p^{-2} t^2 X^{-1}) (1 - p^{-2} t^2 X) \prod_{i=1}^{(n-2)/2} \{(1 - p^{2i-n-1} t^2 X^{-1}) (1 - p^{2i-n-1} t^2 X)\}.$$

Hence we have

$$\text{(B)} \quad S(d_0, \iota, X, t) = \prod_{i=1}^{(n-2)/2} (1 - p^{2i-n-2} t^2 X) (a_0(X) + a_1(X) t^2)$$

with some polynomials $a_0(X), a_1(X)$ in $X + X^{-1}$. We easily see $a_0(X) = 1$. By substituting $p^{(n-1)/2} X^{1/2}$ for t in (A) and (B), and comparing them we see $a_1(X) = -p^{-5/2} \xi_0$. This proves the assertion.

Next let $d_0 \in p\mathbf{Z}_p^*$. Then by Proposition 4.4.3 (1.1), we have

$$\begin{aligned} K_{n-1}^{(1)}(d_0, \iota, X, t) &= X^{1/2} \left\{ 2^{-1} \sum_{r=1}^{(n-2)/2} \sum_{d \in \mathcal{U}} \frac{p^{-r(2r+1)} (t^2 X^{-1})^r \prod_{i=1}^{r-1} (1 - p^{2i-1} X^2)}{\phi_{(n-2r-2)/2}(p^{-2})} \right. \\ &\quad \times (1 + \eta_d p^{r-1/2} X) \zeta_{2r}(d_0 d, \iota, tX^{-1/2}) \\ &\quad \left. + \sum_{r=0}^{(n-2)/2} \frac{p^{-(2r+1)(r+1)} (t^2 X^{-1})^{r+1/2} \prod_{i=1}^r (1 - p^{2i-1} X^2)}{\phi_{(n-2r-2)/2}(p^{-2})} \right. \\ &\quad \left. \times \zeta_{2r+1}(p^{-1} d_0, \iota, tX^{-1/2}) \right\}. \end{aligned}$$

By Theorem 5.1 of [11], we have

$$\zeta_{2r+1}(p^{-1} d_0, \iota, tX^{-1/2}) = \frac{1}{\phi_r(p^{-2}) (1 - p^{-2} t^2 X^{-1}) \prod_{i=1}^r (1 - p^{2i-3-2r} t^2 X^{-1})},$$

and

$$\zeta_{2r}(d_0 d, \iota, tX^{-1/2}) = \frac{p^{-1} t X^{-1/2}}{\phi_{r-1}(p^{-2}) (1 - p^{-2} t^2 X^{-1}) \prod_{i=1}^r (1 - p^{2i-3-2r} t^2 X^{-1})}.$$

Thus the assertion can be proved in the same manner as above.

(2) First let $d_0 \in \mathbf{Z}_p^*$. Then by Proposition 4.4.3 (1.1), we have

$$\begin{aligned} K_{n-1}^{(1)}(d_0, \varepsilon, X, t) &= \frac{1}{\phi_{(n-2)/2}(p^{-2})} \\ &\quad + \sum_{r=1}^{(n-2)/2} \sum_{d \in \mathcal{U}} \frac{p^{-r(2r+1)} (t^2 X^{-1})^r \prod_{i=1}^{r-1} (1 - p^{2i-1} X^2)}{2\phi_{(n-2r-2)/2}(p^{-2})} \\ &\quad \times (1 - p^{-1/2} \xi_0 X) (1 + \eta_d p^{r-1/2} X) \xi_0 \eta_d \zeta_{2r}(d_0 d, \varepsilon, tX^{-1/2}) \\ &\quad + \sum_{r=0}^{(n-2)/2} \frac{p^{-(2r+1)(r+1)} (t^2 X^{-1})^{r+1/2} \prod_{i=1}^r (1 - p^{2i-1} X^2)}{\phi_{(n-2r-2)/2}(p^{-2})} \\ &\quad \times (1 - p^{-1/2} \xi_0 X) \zeta_{2r+1}(pd_0, \varepsilon, tX^{-1/2}). \end{aligned}$$

By Theorem 5.2 of [11],

$$\zeta_{2r}(d_0 d, \varepsilon, tX^{-1/2}) = \frac{1 + \xi_0 \eta_d p^{-r}}{\phi_r(p^{-2}) \prod_{i=1}^r (1 - p^{-2i} t^2 X^{-1})},$$

and

$$\zeta_{2r+1}(pd_0, \varepsilon, tX^{-1/2}) = \frac{p^{-r-1} t X^{-1/2}}{\phi_r(p^{-2}) \prod_{i=1}^{r+1} (1 - p^{-2i} t^2 X^{-1})}.$$

Hence $K_{n-1}^{(1)}(d_0, \varepsilon, X, t)$ can be expressed as

$$K_{n-1}^{(1)}(d_0, \varepsilon, X, t) = \frac{T(d_0, \varepsilon, X, t)}{\phi_{(n-2)/2}(p^{-2}) \prod_{i=1}^{n/2} (1 - p^{-2i} t^2 X^{-1})},$$

where $T(d_0, \iota, X, t)$ is a polynomial in t of degree n , and expressed as

$$(C) \quad T(d_0, \iota, X, t) = (p^{-n}X^{-1}t^2)^{n/2}(1 - \xi_0 p^{-1/2}X) \prod_{i=1}^{(n-2)/2} (1 - p^{2i-1}X^2) \\ + (1 - p^{-n}t^2X^{-1})V(X, t),$$

with a polynomial $V(X, t)$ in X, X^{-1} and t . On the other hand, by using the same argument as (1), we can show that

$$(D) \quad T(d_0, \varepsilon, X, t) = \prod_{i=1}^{(n-2)/2} (1 - p^{-2i-1}t^2X)(1 + b_1(X)t^2)$$

with $b_1(X)$ a polynomial in $X + X^{-1}$. Thus, by substituting $p^{n/2}X^{1/2}$ for t in (C) and (D), and comparing them we prove the assertion.

Next let $d_0 \in p\mathbf{Z}_p^*$. Then by Proposition 4.4.3 (1.2), we have

$$K_{n-1}^{(1)}(d_0, \varepsilon, X, t) = X^{1/2} \sum_{r=0}^{(n-2)/2} \frac{p^{-(2r+1)(r+1)-r}(t^2X^{-1})^{r+1/2} \prod_{i=1}^r (1 - p^{2i-1}X^2)}{\phi_{(n-2r-2)/2}(p^{-2})} \\ \times \zeta_{2r+1}(p^{-1}d_0, \varepsilon, tX^{-1/2}).$$

By Theorem 5.2 of [11],

$$\zeta_{2r+1}(p^{-1}d_0, \varepsilon, tX^{-1/2}) = \frac{1}{\phi_r(p^{-2}) \prod_{i=1}^r (1 - p^{-2i}t^2X^{-1})}.$$

Hence

$$K_{n-1}^{(1)}(d_0, \varepsilon, X, t) = p^{-1}t \sum_{r=0}^{(n-2)/2} \frac{p^{-(2r+1)r}(p^{-2}t^2X^{-1})^r \prod_{i=1}^r (1 - p^{2i-1}X^2)}{\phi_{(n-2r-2)/2}(p^{-2})} \\ \times \frac{1}{\phi_r(p^{-2}) \prod_{i=1}^r (1 - p^{-2i}t^2X^{-1})}.$$

Thus the assertion can be proved in the same way as above. \square

Proof of Theorem 4.4.1 in case $p = 2$. The assertion can also be proved by Proposition 4.4.3 (2) in the same way as above. \square

Proposition 4.4.4. *Let k and n be positive even integers. Given a Hecke eigenform $h \in \mathfrak{S}_{k-n/2+1/2}^+(\Gamma_0(4))$, let $f \in \mathfrak{S}_{2k-n}(\Gamma^{(1)})$ be the primitive form as in Section 2. Then*

$$L(s, h) = L(2s, f) \sum_{d_0 \in \mathcal{F}^{(-1)n/2}} c_h(|d_0|) |d_0|^{-s} L(2s - k + n/2 + 1, \left(\frac{d_0}{*}\right))^{-1},$$

where $L(s, \left(\frac{d_0}{*}\right))$ is Dirichlet's L -function for the character $\left(\frac{d_0}{*}\right)$.

Proof. The assertion can immediately be proved by remarking the fact that

$$\sum_{m=1}^{\infty} c_h(|d_0|m^2)m^{-2s} = c_h(|d_0|)L(2s - k + n/2 + 1, \left(\frac{d_0}{*}\right)^{-1}L(2s, f)$$

for $d_0 \in \mathcal{F}^{(-1)^{n/2}}$. □

Proof of Theorem 2.1. By Theorem 4.4.1, we have

$$\prod_p P_{n-1,p}^{(1)}(d_0, \iota_p, \alpha_p, p^{-s+k/2+n/4-1/4}) = |d_0|^{-s+k/2+n/4-5/4} \\ \times \prod_{i=1}^{(n-2)/2} \zeta(2i)L(2s - k - n/2 + 3, \left(\frac{d_0}{*}\right)^{-1} \prod_{i=1}^{(n-2)/2} L(2s - n + 2i + 1, f),$$

and

$$\prod_p P_{n-1,p}^{(1)}(d_0, \varepsilon_p, \alpha_p, p^{-s+k/2+n/4-1/4}) = |d_0|^{-s+k/2+n/4-5/4} \\ \times \prod_{i=1}^{(n-2)/2} \zeta(2i)L(2s - k + n/2 + 1, \left(\frac{d_0}{*}\right)^{-1} \prod_{i=1}^{(n-2)/2} L(2s - n + 2i, f).$$

Thus the assertion follows from Theorem 3.2 and Proposition 4.4.4. □

Remark. Let m be a nonnegative integer, and let k be a positive integer such that $k > m + 2$. Let $E_k^{(m+1)}$ be the Siegel Eisenstein series of weight k and of degree $m + 1$. (For the definition of the Siegel Eisenstein series, see, for example, [6].) Suppose that $m > 0$ and let $e_{k,1}^{(m+1)}$ be the first Fourier-Jacobi coefficient of $E_k^{(m+1)}$. Then $e_{k,1}^{(m+1)}$ belongs to $J_{k,1}(\Gamma_J^{(m)})$. In [6], Hayashida defined the generalized Cohen Eisenstein series $E_{k-1/2}^{(m)}$ as $E_{k-1/2}^{(m)} = \sigma_m(e_{k,1}^{(m+1)})$, where σ_m is the Ibukiyama isomorphism. It turns out that $E_{k-1/2}^{(m)}$ belongs to $\mathfrak{M}_{k-1/2}^+(\Gamma_0^{(m)}(4))$, and in particular, $E_{k-1/2}^{(1)}$ coincides with the Cohen Eisenstein series defined in [5]. Let k and n be positive even integers such that $k > n + 1$. Then, $E_{2k-n}^{(1)}$ is the Hecke eigenform corresponding to $E_{k-n/2+1/2}^{(1)}$ under the Shimura correspondence, and $E_k^{(n)}$ can be regarded as a non-cuspidal version of the Duke-Imamoglu-Ikeda lift of $E_{k-n/2+1/2}^{(1)}$. Therefore, by using the same method as in the proof of Theorem 2.1, we can express the Koecher-Maass series of $E_{k-1/2}^{(n-1)}$ explicitly in terms of $L(s, E_{k-n/2+1/2}^{(1)})$ and $L(s, E_{2k-n}^{(1)})$.

Acknowledgements. The authors thank the referee for suggesting them the remark after the proof of Theorem 2.1. The first named author was partly supported by Grant-in-Aid for Scientific Research, JSPS (No. 21540004), and the second named author was partially supported by Grants-in-Aid for Scientific Research, JSPS (No. 23224001 and No. 22340001).

REFERENCES

- [1] T. Arakawa, *Dirichlet series corresponding to Siegel's modular forms*, Math. Ann. **238** (1978), 157–173.
- [2] T. Arakawa, *Dirichlet series corresponding to Siegel's modular forms with level N* , Tohoku Math. J. **42** (1990), 261–286.
- [3] T. Arakawa, *Köcher-Maass Dirichlet series corresponding to Jacobi forms and Cohen Eisenstein series*, Comment. Math. Univ. St.Paul. **47** (1998), 93–122.
- [4] S. Böcherer and F. Sato, *Rationality of certain formal power series related to local densities*, Comment. Math. Univ. St.Paul. **36** (1987), 53–86.
- [5] H. Cohen, *Sums involving the values at negative integers of L -functions of quadratic characters*, Math. Ann. **217** (1975), 171–185.
- [6] S. Hayashida, *Fourier-Jacobi expansion and the Ikeda lift*, Abh. Math. Sem. Hamburg **81** (2011), 1–17.
- [7] T. Ibukiyama, *On Jacobi forms and Siegel modular forms of half integral weights*, Comment. Math. Univ. St.Paul. **41** (1992), no. 2, 109–124.
- [8] T. Ibukiyama and H. Katsurada, *An explicit formula for Koecher-Maaß Dirichlet series for Eisenstein series of Klingen type*, J. Number Theory **102** (2003), 223–256.
- [9] T. Ibukiyama and H. Katsurada, *An explicit formula for Koecher-Maaß Dirichlet series for the Ikeda lifting*, Abh. Math. Sem. Hamburg **74** (2004), 101–121.
- [10] T. Ibukiyama and H. Katsurada, *Koecher-Maaß series for real analytic Siegel Eisenstein series, Automorphic forms and zeta functions*, 170–197, World Sci. Publ., Hackensack, NJ, 2006.
- [11] T. Ibukiyama and H. Saito, *On zeta functions associated to symmetric matrices. I. An explicit form of zeta functions*, Amer. J. Math. **117** (1995), 1097–1155.
- [12] T. Ikeda, *On the lifting of elliptic modular forms to Siegel cusp forms of degree $2n$* , Ann. of Math. **154** (2001), no. 3, 641–681.
- [13] T. Ikeda, *Pullback of the lifting of elliptic cusp forms and Miyawaki's conjecture*, Duke Math. J. **131** (2006), no. 3, 469–497.
- [14] H. Katsurada, *An explicit formula for Siegel series*, Amer. J. Math. **121** (1999), 415–452.
- [15] H. Katsurada and H. Kawamura, *Ikeda's conjecture on the period of the Duke-Imamoglu-Ikeda lift*, Preprint (2009).

- [16] Y. Kitaoka, *Dirichlet series in the theory of Siegel modular forms*, Nagoya Math. J. **95** (1984), 73–84.
- [17] Y. Kitaoka, *Arithmetic of quadratic forms*, Cambridge Tracts in Math., vol.106, Cambridge Univ. Press, Cambridge, 1993
- [18] W. Kohnen, *Modular forms of half-integral weight on $\Gamma_0(4)$* , Math. Ann. **248** (1980), 249–266.
- [19] H. Maass, *Siegel's modular forms and Dirichlet series*, Lecture Notes in Math., vol. 216, Springer-Verlag, Berlin-New York, 1971.
- [20] G. L. Watson, *The 2-adic density of a quadratic form*, Mathematica **23** (1976), no.1, 94–106.

MURORAN INSTITUTE OF TECHNOLOGY, MIZUMOTO 27-1 MURORAN, 050-8585,
JAPAN

E-mail address: `hidenori@mmm.muroran-it.ac.jp`

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, KITA 10, NISHI 8, KITA-
KU, SAPPORO, 060-0810, JAPAN

E-mail address: `hkawamura@sci.hokudai.ac.jp`