



Title	KOECHER-MAASS SERIES OF THE IKEDA LIFT FOR $U(m,m)$
Author(s)	Katsurada, Hidenori
Citation	Hokkaido University Preprint Series in Mathematics, 1049, 1-21
Issue Date	2014-3-11
DOI	10.14943/84193
Doc URL	http://hdl.handle.net/2115/69853
Type	bulletin (article)
File Information	pre1049.pdf



[Instructions for use](#)

KOECHER-MAASS SERIES OF THE IKEDA LIFT FOR $U(m, m)$

HIDENORI KATSURADA

In memory of Professor Hiroshi Saito
Appearing in *Kyoto Journal of Mathematics*

ABSTRACT. Let $K = \mathbf{Q}(\sqrt{-D})$ be an imaginary quadratic field with discriminant $-D$, and χ the Dirichlet character corresponding to the extension K/\mathbf{Q} . Let $m = 2n$ or $2n + 1$ with n a positive integer. Let f be a primitive form of weight $2k + 1$ and character χ for $\Gamma_0(D)$, or a primitive form of weight $2k$ for $SL_2(\mathbf{Z})$ according as $m = 2n$, or $m = 2n + 1$. For such an f let $I_m(f)$ be the lift of f to the space of Hermitian modular forms constructed by Ikeda. We then give an explicit formula of the Koecher-Maass series $L(s, I_m(f))$ of $I_m(f)$. This is a generalization of Mizuno [Mi06].

1. INTRODUCTION

In [Mi06], Mizuno gave explicit formulas of the Koecher-Maass series of the Hermitian Eisenstein series of degree two and of the Hermitian Maass lift. In this paper, we give an explicit formula of the Koecher-Maass series of the Hermitian Ikeda lift. Let $K = \mathbf{Q}(\sqrt{-D})$ be an imaginary quadratic field with discriminant $-D$. Let \mathcal{O} be the ring of integers in K , and χ the Kronecker character corresponding to the extension K/\mathbf{Q} . For a non-degenerate Hermitian matrix or alternating matrix T with entries in K , let \mathcal{U}_T be the unitary group defined over \mathbf{Q} , whose group $\mathcal{U}_T(R)$ of R -valued points is given by

$$\mathcal{U}_T(R) = \{g \in GL_m(R \otimes K) \mid {}^t \bar{g} T g = T\}$$

for any \mathbf{Q} -algebra R , where \bar{g} denotes the automorphism of $M_n(R \otimes K)$ induced by the non-trivial automorphism of K over \mathbf{Q} . We also define the special unitary group \mathcal{SU}_T over \mathbf{Q}_p by $\mathcal{SU}_T = \mathcal{U}_T \cap R_{K/\mathbf{Q}}(SL_m)$, where $R_{K/\mathbf{Q}}$ is the Weil restriction. In particular we write \mathcal{U}_T as $\mathcal{U}^{(m)}$ or $U(m, m)$ if $T = \begin{pmatrix} \mathcal{O} & -1_m \\ 1_m & \mathcal{O} \end{pmatrix}$. For a more precise description of $\mathcal{U}^{(m)}$ see Section 2. Put $\Gamma_K^{(m)} = U(m, m)(\mathbf{Q}) \cap GL_{2m}(\mathcal{O})$. For a modular form F of weight $2l$ and character ψ for $\Gamma_K^{(m)}$ we define the Koecher-Maass series $L(s, F)$ of F by

$$L(s, F) = \sum_T \frac{c_F(T)}{e^*(T)(\det T)^s},$$

where T runs over all $SL_m(\mathcal{O})$ -equivalence classes of positive definite semi-integral Hermitian matrices of degree m , $c_F(T)$ denotes the T -th Fourier coefficient of F , and $e^*(T) = \#(\mathcal{SU}_T(\mathbf{Q}) \cap SL_m(\mathcal{O}))$.

Date: 2014, February 17.

2010 Mathematics Subject Classification Primary 11F55, 11F67, .

Let k be a non-negative integer. Then for a primitive form $f \in \mathfrak{S}_{2k+1}(T_0(D), \chi)$ Ikeda [Ike08] constructed a lift $I_{2n}(f)$ of f to the space of modular forms of weight $2k+2n$ and a character \det^{-k-n} for $\Gamma_K^{(2n)}$. This is a generalization of the Maass lift considered by Kojima [Koj82], Gritsenko [Gri90], Krieg [Kri91] and Sugano [Su95]. Similarly for a primitive form $f \in \mathfrak{S}_{2k}(SL_2(\mathbf{Z}))$ he constructed a lift $I_{2n+1}(f)$ of f to the space of modular forms of weight $2k+2n$ and a character \det^{k+n} for $\Gamma_K^{(2n+1)}$. For the rest of this section, let $m = 2n$ or $m = 2n+1$. We then call $I_m(f)$ the Ikeda lift of f for $U(m, m)$ or the Hermitian Ikeda lift of degree m . Ikeda also showed that the automorphic form $Lift^{(m)}(f)$ on the adèle group $\mathcal{U}^{(m)}(\mathbf{A})$ associated with $I_m(f)$ is a cuspidal Hecke eigenform whose standard L -function coincides with

$$\prod_{i=1}^m L(s+k+n-i+1/2, f) L(s+k+n-i+1/2, f, \chi),$$

where $L(s+k+n-i+1/2, f)$ is the Hecke L -function of f and $L(s+k+n-i+1/2, f, \chi)$ is its "modified twist" by χ . For the precise definition of $L(s+k+n-i+1/2, f, \chi)$ see Section 2. We also call $Lift^{(m)}(f)$ the adelic Ikeda lift of f for $U(m, m)$. Then we express the Kocher-Maass series of $I_m(f)$ in terms of the L -functions related to f . This result was already obtained in the case $m = 2$ by Mizuno [Mi06].

The method we use is similar to that in the proof of the main result of [IK04] or [IK06]. We explain it more precisely. In Section 3, we reduce our computation to a computation of certain formal power series $\hat{P}_{m,p}(d; X, t)$ in t associated with local Siegel series similarly to [IK04] (cf. Theorem 3.4 and Section 5).

Section 4 is devoted to the computation of them. This computation is similar to that in [IK04], but we should be careful in dealing with the case where p is ramified in K . After such an elaborate computation, we can get explicit formulas of $\hat{P}_{m,p}(d; X, t)$ for all prime numbers p (cf. Theorems 4.3.1, 4.3.2, and 4.3.6). In Section 5, by using explicit formulas for $\hat{P}_{m,p}(d; X, t)$, we immediately get an explicit formula of $L(s, I_m(f))$.

Using the same argument as in the proof our main result, we can give an explicit formula of the Koecher-Maass series of the Hermitian Eisenstein series of any degree, which can be regarded as a zeta function of a certain prehomogeneous vector space. We also note that the method used in this paper is useful for giving an explicit formula for the Rankin-Selberg series of the Hermitian Ikeda lift, and as a result we can prove the period relation of the Hermitian Ikeda lift, which was conjectured by Ikeda [Ike08]. We will discuss these topics in subsequent papers [Kat13] and [Kat14].

Acknowledgments. The author thanks Professor T. Watanabe and Professor R. Schulze-Pillot for giving him many crucial comments on the mass formula for the unitary group. He also thanks Professor Y. Mizuno, Professor T. Ikeda, and A. Yukie for useful discussions. The author was partly supported by JSPS KAKENHI Grant Number 24540005.

Notation. Let R be a commutative ring. We denote by R^\times and R^* the semigroup of non-zero elements of R and the unit group of R , respectively. For a subset S of R we denote by $M_{mn}(S)$ the set of (m, n) -matrices with entries in S . In particular put $M_n(S) = M_{nn}(S)$. Put $GL_m(R) = \{A \in M_m(R) \mid \det A \in R^*\}$, where $\det A$ denotes the determinant of a square matrix A . Let K_0 be a field, and

K a quadratic extension of K_0 , or $K = K_0 \oplus K_0$. In the latter case, we regard K_0 as a subring of K via the diagonal embedding. We also identify $M_{mn}(K)$ with $M_{mn}(K_0) \oplus M_{mn}(K_0)$ in this case. If K is a quadratic extension of K_0 , let ρ be the non-trivial automorphism of K over K_0 , and if $K = K_0 \oplus K_0$, let ρ be the automorphism of K defined by $\rho(a, b) = (b, a)$ for $(a, b) \in K_0$. We sometimes write \bar{x} instead of $\rho(x)$ for $x \in K$ in both cases. Let R be a subring of K . For an (m, n) -matrix $X = (x_{ij})_{m \times n}$ write $X^* = (\bar{x}_{ji})_{n \times m}$, and for an (m, m) -matrix A , we write $A[X] = X^*AX$. Let $\text{Her}_n(R)$ denote the set of Hermitian matrices of degree n with entries in R , that is the subset of $M_n(R)$ consisting of matrices X such that $X^* = X$. Then a Hermitian matrix A of degree n with entries in K is said to be semi-integral over R if $\text{tr}(AB) \in K_0 \cap R$ for any $B \in \text{Her}_n(R)$, where tr denotes the trace of a matrix. We denote by $\widehat{\text{Her}}_n(R)$ the set of semi-integral matrices of degree n over R .

For a subset S of $M_n(R)$ we denote by S^\times the subset of S consisting of non-degenerate matrices. If S is a subset of $\text{Her}_n(\mathbf{C})$ with \mathbf{C} the field of complex numbers, we denote by S^+ the subset of S consisting of positive definite matrices. The group $GL_n(R)$ acts on the set $\text{Her}_n(R)$ in the following way:

$$GL_n(R) \times \text{Her}_n(R) \ni (g, A) \longrightarrow g^*Ag \in \text{Her}_n(R).$$

Let G be a subgroup of $GL_n(R)$. For a G -stable subset \mathcal{B} of $\text{Her}_n(R)$ we denote by \mathcal{B}/G the set of equivalence classes of \mathcal{B} under the action of G . We sometimes identify \mathcal{B}/G with a complete set of representatives of \mathcal{B}/G . We abbreviate $\mathcal{B}/GL_n(R)$ as \mathcal{B}/\sim if there is no fear of confusion. Two Hermitian matrices A and A' with entries in R are said to be G -equivalent and write $A \sim_G A'$ if there is an element X of G such that $A' = A[X]$. For square matrices X and Y we write $X \perp Y = \begin{pmatrix} X & O \\ O & Y \end{pmatrix}$.

We put $\mathbf{e}(x) = \exp(2\pi\sqrt{-1}x)$ for $x \in \mathbf{C}$, and for a prime number p we denote by $\mathbf{e}_p(*)$ the continuous additive character of \mathbf{Q}_p such that $\mathbf{e}_p(x) = \mathbf{e}(x)$ for $x \in \mathbf{Z}[p^{-1}]$.

For a prime number p we denote by $\text{ord}_p(*)$ the additive valuation of \mathbf{Q}_p normalized so that $\text{ord}_p(p) = 1$, and put $|x|_p = p^{-\text{ord}_p(x)}$. Moreover we denote by $|x|_\infty$ the absolute value of $x \in \mathbf{C}$. Let K be an imaginary quadratic field, and \mathcal{O} the ring of integers in K . For a prime number p put $K_p = K \otimes \mathbf{Q}_p$, and $\mathcal{O}_p = \mathcal{O} \otimes \mathbf{Z}_p$. Then K_p is a quadratic extension of \mathbf{Q}_p or $K_p \cong \mathbf{Q}_p \oplus \mathbf{Q}_p$. In the former case, for $x \in K_p$, we denote by \bar{x} the conjugate of x over \mathbf{Q}_p . In the latter case, we identify K_p with $\mathbf{Q}_p \oplus \mathbf{Q}_p$, and for $x = (x_1, x_2)$ with $x_i \in \mathbf{Q}_p$, we put $\bar{x} = (x_2, x_1)$. For $x \in K_p$ we define the norm $N_{K_p/\mathbf{Q}_p}(x)$ by $N_{K_p/\mathbf{Q}_p}(x) = x\bar{x}$, and put $\nu_{K_p}(x) = \text{ord}_p(N_{K_p/\mathbf{Q}_p}(x))$, and $|x|_{K_p} = |N_{K_p/\mathbf{Q}_p}(x)|_p$. Moreover put $|x|_{K_\infty} = |x\bar{x}|_\infty$ for $x \in \mathbf{C}$.

2. MAIN RESULTS

For a positive integer N let

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z}) \mid c \equiv 0 \pmod{N} \right\},$$

and for a Dirichlet character $\psi \pmod{N}$, we denote by $\mathfrak{M}_l(\Gamma_0(N), \psi)$ the space of modular forms of weight l for $\Gamma_0(N)$ and nebentype ψ , and by $\mathfrak{S}_l(\Gamma_0(N), \psi)$ its subspace consisting of cusp forms. We simply write $\mathfrak{M}_l(\Gamma_0(N), \psi)$ (resp. $\mathfrak{S}_l(\Gamma_0(N), \psi)$) as $\mathfrak{M}_l(\Gamma_0(N))$ (resp. as $\mathfrak{S}_l(\Gamma_0(N))$) if ψ is the trivial character.

Throughout the paper, we fix an imaginary quadratic extension K of \mathbf{Q} with discriminant $-D$, and denote by \mathcal{O} the ring of integers in K . For such a K let $\mathcal{U}^{(m)} =$

$U(m, m)$ be the unitary group defined in Section 1. Put $J_m = \begin{pmatrix} O_m & -1_m \\ 1_m & O_m \end{pmatrix}$, where 1_m denotes the unit matrix of degree m . Then

$$\mathcal{U}^{(m)}(\mathbf{Q}) = \{M \in GL_{2m}(K) \mid J_m[M] = J_m\}.$$

Put

$$\Gamma^{(m)} = \Gamma_K^{(m)} = \mathcal{U}^{(m)}(\mathbf{Q}) \cap GL_{2m}(\mathcal{O}).$$

Let \mathfrak{H}_m be the Hermitian upper half-space defined by

$$\mathfrak{H}_m = \{Z \in M_m(\mathbf{C}) \mid \frac{1}{2\sqrt{-1}}(Z - Z^*) \text{ is positive definite}\}.$$

The group $\mathcal{U}^{(m)}(\mathbf{R})$ acts on \mathfrak{H}_m by

$$g\langle Z \rangle = (AZ + B)(CZ + D)^{-1} \text{ for } g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathcal{U}^{(m)}(\mathbf{R}), Z \in \mathfrak{H}_m.$$

We also put $j(g, Z) = \det(CZ + D)$ for such Z and g . Let l be an integer. For a subgroup Γ of $\mathcal{U}^{(m)}(\mathbf{Q})$ commensurable with $\Gamma^{(m)}$ and a character ψ of Γ , we denote by $\mathfrak{M}_l(\Gamma, \psi)$ the space of holomorphic modular forms of weight l with character ψ for Γ . We denote by $\mathfrak{S}_l(\Gamma, \psi)$ the subspace of $\mathfrak{M}_l(\Gamma, \psi)$ consisting of cusp forms. In particular, if ψ is the character of Γ defined by $\psi(\gamma) = (\det \gamma)^{-l}$ for $\gamma \in \Gamma$, we write $\mathfrak{M}_{2l}(\Gamma, \psi)$ as $\mathfrak{M}_{2l}(\Gamma, \det^{-l})$, and so on. Let $F(z)$ be an element of $\mathfrak{M}_{2l}(\Gamma^{(m)}, \det^{-l})$. We then define the Koecher-Maass series $L(s, F)$ for F by

$$L(s, F) = \sum_{T \in \widehat{\text{Her}}_m(\mathcal{O})^+ / SL_m(\mathcal{O})} \frac{c_F(T)}{(\det T)^s e^*(T)},$$

where $c_F(T)$ denotes the T -th Fourier coefficient of F , and $e^*(T) = \#(\mathcal{S}U_T(\mathbf{Q}) \cap SL_m(\mathcal{O}))$.

Now we consider the adelic modular form. Let \mathbf{A} be the adèle ring of \mathbf{Q} , and \mathbf{A}_f the non-archimedean factor of \mathbf{A} . Let $h = h_K$ be a class number of K . Let $G^{(m)} = \text{Res}_{K/\mathbf{Q}}(GL_m)$, and $G^{(m)}(\mathbf{A})$ be the adélization of $G^{(m)}$. Moreover put $\mathcal{C}^{(m)} = \prod_p GL_m(\mathcal{O}_p)$. Let $\mathcal{U}^{(m)}(\mathbf{A})$ be the adélization of $\mathcal{U}^{(m)}$. We define the compact subgroup $\mathcal{K}_0^{(m)}$ of $\mathcal{U}^{(m)}(\mathbf{A}_f)$ by $\mathcal{U}^{(m)}(\mathbf{A}) \cap \prod_p GL_{2m}(\mathcal{O}_p)$, where p runs over all rational primes. Then we have

$$\mathcal{U}^{(m)}(\mathbf{A}) = \bigsqcup_{i=1}^h \mathcal{U}^{(m)}(\mathbf{Q}) \gamma_i \mathcal{K}_0^{(m)} \mathcal{U}^{(m)}(\mathbf{R})$$

with some subset $\{\gamma_1, \dots, \gamma_h\}$ of $\mathcal{U}^{(m)}(\mathbf{A}_f)$. We can take γ_i as

$$\gamma_i = \begin{pmatrix} t_i & 0 \\ 0 & t_i^{*-1} \end{pmatrix},$$

where $\{t_i\}_{i=1}^h = \{(t_{i,p})\}_{i=1}^h$ is a certain subset of $G^{(m)}(\mathbf{A}_f)$ such that $t_1 = 1$, and

$$G^{(m)}(\mathbf{A}) = \bigsqcup_{i=1}^h G^{(m)}(\mathbf{Q}) t_i G^{(m)}(\mathbf{R}) \mathcal{C}^{(m)}.$$

Put $\Gamma_i = \mathcal{U}^{(m)}(\mathbf{Q}) \cap \gamma_i \mathcal{K}_0 \gamma_i^{-1} \mathcal{U}^{(m)}(\mathbf{R})$. Then for an element $(F_1, \dots, F_h) \in \bigoplus_{i=1}^h \mathfrak{M}_{2l}(\Gamma_i, \det^{-l})$, we define $(F_1, \dots, F_h)^\sharp$ by

$$(F_1, \dots, F_h)^\sharp(g) = F_i(x(\mathbf{i})) j(x, \mathbf{i})^{-2l} (\det x)^l$$

for $g = u\gamma_i x\kappa$ with $u \in \mathcal{U}^{(m)}(\mathbf{Q})$, $x \in \mathcal{U}^{(m)}(\mathbf{R})$, $\kappa \in \mathcal{K}_0$. We denote by $\mathcal{M}_{2l}(\mathcal{U}^{(m)}(\mathbf{Q}) \backslash \mathcal{U}^{(m)}(\mathbf{A}), \det^{-l})$ the space of automorphic forms obtained in this way. We also put

$$\mathcal{S}_{2l}(\mathcal{U}^{(m)}(\mathbf{Q}) \backslash \mathcal{U}^{(m)}(\mathbf{A}), \det^{-l}) = \{(F_1, \dots, F_h)^\sharp \mid F_i \in \mathfrak{S}_{2l}(\Gamma_i, \det^{-l})\}.$$

We can define the Hecke operators which act on the space

$\mathcal{M}_{2l}(\mathcal{U}^{(m)}(\mathbf{Q}) \backslash \mathcal{U}^{(m)}(\mathbf{A}), \det^{-l})$. For the precise definition of them, see [Ike08].

Let $\widehat{\text{Her}}_m(\mathcal{O})$ be the set of semi-integral Hermitian matrices over \mathcal{O} of degree m as in the Notation. We note that A belongs to $\widehat{\text{Her}}_m(\mathcal{O})$ if and only if its diagonal components are rational integers and $\sqrt{-D}A \in \text{Her}_m(\mathcal{O})$. For a non-degenerate Hermitian matrix B with entries in K of degree m , put $\gamma(B) = (-D)^{[m/2]} \det B$.

Let $\widehat{\text{Her}}_m(\mathcal{O}_p)$ be the set of semi-integral matrices over \mathcal{O}_p of degree m as in the Notation. We put $\xi_p = 1, -1$, or 0 according as $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$, K_p is an unramified quadratic extension of \mathbf{Q}_p , or K_p is a ramified quadratic extension of \mathbf{Q}_p . For $T \in \widehat{\text{Her}}_m(\mathcal{O}_p)^\times$ we define the local Siegel series $b_p(T, s)$ by

$$b_p(T, s) = \sum_{R \in \text{Her}_n(K_p)/\text{Her}_n(\mathcal{O}_p)} \mathbf{e}_p(\text{tr}(TR)) p^{-\text{ord}_p(\mu_p(R))s},$$

where $\mu_p(R) = [R\mathcal{O}_p^m + \mathcal{O}_p^m : \mathcal{O}_p^m]$. We remark that there exists a unique polynomial $F_p(T, X)$ in X such that

$$b_p(T, s) = F_p(T, p^{-s}) \prod_{i=0}^{[(m-1)/2]} (1 - p^{2i-s}) \prod_{i=1}^{[m/2]} (1 - \xi_p p^{2i-1-s})$$

(cf. Shimura [Sh97]). We then define a Laurent polynomial $\widetilde{F}_p(T, X)$ as

$$\widetilde{F}_p(T, X) = X^{-\text{ord}_p(\gamma(T))} F_p(T, p^{-m} X^2).$$

We remark that we have

$$\begin{aligned} \widetilde{F}_p(T, X^{-1}) &= (-D, \gamma(T))_p \widetilde{F}_p(T, X) && \text{if } m \text{ is even,} \\ \widetilde{F}_p(T, \xi_p X^{-1}) &= \widetilde{F}_p(T, X) && \text{if } m \text{ is even and } p \nmid D, \end{aligned}$$

and

$$\widetilde{F}_p(T, X^{-1}) = \widetilde{F}_p(T, X) \quad \text{if } m \text{ is odd}$$

(cf. [Ike08]). Here $(a, b)_p$ is the Hilbert symbol of $a, b \in \mathbf{Q}_p^\times$. Hence we have

$$\widetilde{F}_p(T, X) = (-D, \gamma(B))_p^{m-1} X^{\text{ord}_p(\gamma(T))} F_p(T, p^{-m} X^{-2}).$$

Now we put

$$\widehat{\text{Her}}_m(\mathcal{O})_i^+ = \{T \in \text{Her}_m(K)^+ \mid t_{i,p}^* T t_{i,p} \in \widehat{\text{Her}}_m(\mathcal{O}_p) \text{ for any } p\}.$$

First let k be a non-negative integer, and $m = 2n$ a positive even integer. Let

$$f(z) = \sum_{N=1}^{\infty} a(N) \mathbf{e}(Nz)$$

be a primitive form in $\mathfrak{S}_{2k+1}(\Gamma_0(D), \chi)$. For a prime number p not dividing D let $\alpha_p \in \mathbf{C}$ such that $\alpha_p + \chi(p)\alpha_p^{-1} = p^{-k}a(p)$, and for $p \mid D$ put $\alpha_p = p^{-k}a(p)$. We note that $\alpha_p \neq 0$ even if $p \mid D$. Then for the Kronecker character χ we define Hecke's L -function $L(s, f, \chi^i)$ twisted by χ^i as

$$L(s, f, \chi^i) = \prod_{p \nmid D} \{(1 - \alpha_p p^{-s+k} \chi(p)^i)(1 - \alpha_p^{-1} p^{-s+k} \chi(p)^{i+1})\}^{-1}$$

$$\times \begin{cases} \prod_{p|D} (1 - \alpha_p p^{-s+k})^{-1} & \text{if } i \text{ is even} \\ \prod_{p|D} (1 - \alpha_p^{-1} p^{-s+k})^{-1} & \text{if } i \text{ is odd.} \end{cases}$$

In particular, if i is even, we sometimes write $L(s, f, \chi^i)$ as $L(s, f)$ as usual. Moreover for $i = 1, \dots, h$ we define a Fourier series

$$I_m(f)_i(Z) = \sum_{T \in \widehat{\text{Her}}_m(\mathcal{O})_i^+} a_{I_m(f)_i}(T) \mathbf{e}(\text{tr}(TZ)),$$

where

$$a_{I_{2n}(f)_i}(T) = |\gamma(T)|^k \prod_p |\det(t_{i,p}) \det(\overline{t_{i,p}})|_p^n \widetilde{F}_p(t_{i,p}^* T t_{i,p}, \alpha_p^{-1}).$$

Next let k be a positive integer and $m = 2n + 1$ a positive odd integer. Let

$$f(z) = \sum_{N=1}^{\infty} a(N) \mathbf{e}(Nz)$$

be a primitive form in $\mathfrak{S}_{2k}(SL_2(\mathbf{Z}))$. For a prime number p let $\alpha_p \in \mathbf{C}$ such that $\alpha_p + \alpha_p^{-1} = p^{-k+1/2} a(p)$. Then we define Hecke's L -function $L(s, f, \chi^i)$ twisted by χ^i as

$$L(s, f, \chi^i) = \prod_p \{(1 - \alpha_p p^{-s+k-1/2} \chi(p)^i)(1 - \alpha_p^{-1} p^{-s+k-1/2} \chi(p)^i)\}^{-1}.$$

In particular, if i is even we write $L(s, f, \chi^i)$ as $L(s, f)$ as usual. Moreover for $i = 1, \dots, h$ we define a Fourier series

$$I_{2n+1}(f)_i(Z) = \sum_{T \in \widehat{\text{Her}}_{2n+1}(\mathcal{O})_i^+} a_{I_{2n+1}(f)_i}(T) \mathbf{e}(\text{tr}(TZ)),$$

where

$$a_{I_{2n+1}(f)_i}(T) = |\gamma(T)|^{k-1/2} \prod_p |\det(t_{i,p}) \det(\overline{t_{i,p}})|_p^{n+1/2} \widetilde{F}_p(t_{i,p}^* T t_{i,p}, \alpha_p^{-1}).$$

Remark. In [Ike08], Ikeda defined $\widetilde{F}_p(T, X)$ as

$$\widetilde{F}_p(T, X) = X^{\text{ord}_p(\gamma(T))} F_p(T, p^{-m} X^{-2}),$$

and we define it by replacing X with X^{-1} in this paper. This change does not affect the results.

Then Ikeda [Ike08] showed the following:

Theorem 2.1. *Let $m = 2n$ or $2n + 1$. Let f be a primitive form in $\mathfrak{S}_{2k+1}(\Gamma_0(D), \chi)$ or in $\mathfrak{S}_{2k}(SL_2(\mathbf{Z}))$ according as $m = 2n$ or $m = 2n + 1$. Moreover let Γ_i be the subgroup of $\mathcal{U}^{(m)}$ defined as above. Then $I_m(f)_i(Z)$ is an element of $\mathfrak{S}_{2k+2n}(\Gamma_i, \det^{-k-n})$ for any i . In particular, $I_m(f) := I_m(f)_1$ is an element of $\mathfrak{S}_{2k+2n}(\Gamma^{(m)}, \det^{-k-n})$.*

This is a Hermitian analogue of the lifting constructed in [Ike01]. We call $I_m(f)$ the Ikeda lift of f for $\mathcal{U}^{(m)}$.

It follows from Theorem 2.1 that we can define an element $(I_m(f)_1, \dots, I_m(f)_h)^\sharp$ of $\mathfrak{S}_{2k+2n}(\mathcal{U}^{(m)}(\mathbf{Q}) \backslash \mathcal{U}^{(m)}(\mathbf{A}), \det^{-k-n})$, which we write $\text{Lift}^{(m)}(f)$.

Theorem 2.2. *Let $m = 2n$ or $2n + 1$. Suppose that $Lift^{(m)}(f)$ is not identically zero. Then $Lift^{(m)}(f)$ is a Hecke eigenform in $\mathcal{S}_{2k+2n}(\mathcal{U}^{(m)}(\mathbf{Q}) \backslash \mathcal{U}^{(m)}(\mathbf{A}), \det^{-k-n})$ and its standard L -function $L(s, Lift^{(m)}(f), \text{st})$ coincides with*

$$\prod_{i=1}^m L(s + k + n - i + 1/2, f) L(s + k + n - i + 1/2, f, \chi)$$

up to bad Euler factors.

We call $Lift^{(m)}(f)$ the adelic Ikeda lift of f for $\mathcal{U}^{(m)}$.

Let Q_D be the set of prime divisors of D . For each prime $q \in Q_D$, put $D_q = q^{\text{ord}_q(D)}$. We define a Dirichlet character χ_q by

$$\chi_q(a) = \begin{cases} \chi(a') & \text{if } (a, q) = 1 \\ 0 & \text{if } q|a \end{cases},$$

where a' is an integer such that

$$a' \equiv a \pmod{D_q} \quad \text{and} \quad a' \equiv 1 \pmod{DD_q^{-1}}.$$

For a subset Q of Q_D put $\chi_Q = \prod_{q \in Q} \chi_q$ and $\chi'_Q = \prod_{q \in Q_D, q \notin Q} \chi_q$. Here we make the convention that $\chi_Q = 1$ and $\chi'_Q = \chi$ if Q is the empty set. Let

$$f(z) = \sum_{N=1}^{\infty} c_f(N) \mathbf{e}(Nz)$$

be a primitive form in $\mathfrak{S}_{2k+1}(\Gamma_0(D), \chi)$. Then there exists a primitive form

$$f_Q(z) = \sum_{N=1}^{\infty} c_{f_Q}(N) \mathbf{e}(Nz)$$

such that

$$c_{f_Q}(p) = \chi_Q(p) c_f(p) \quad \text{for } p \notin Q$$

and

$$c_{f_Q}(p) = \chi'_Q(p) \overline{c_f(p)} \quad \text{for } p \in Q.$$

Let $L(s, \chi^i) = \zeta(s)$ or $L(s, \chi)$ according as i is even or odd, where $\zeta(s)$ and $L(s, \chi)$ are Riemann's zeta function, and the Dirichlet L -function for χ , respectively. Moreover we define $\tilde{\Lambda}(s, \chi^i)$ by

$$\tilde{\Lambda}(s, \chi^i) = 2(2\pi)^{-s} \Gamma(s) L(s, \chi^i)$$

with $\Gamma(s)$ the Gamma function.

Then our main results in this paper are as follows:

Theorem 2.3. *Let k be a nonnegative integer and n a positive integer. Let f be a primitive form in $\mathfrak{S}_{2k+1}(\Gamma_0(D), \chi)$. Then, we have*

$$L(s, I_{2n}(f)) = D^{ns+n^2-n/2-1/2} 2^{-2n+1} \times \prod_{i=2}^{2n} \tilde{\Lambda}(i, \chi^i) \sum_{Q \subset Q_D} \chi_Q((-1)^n) \prod_{j=1}^{2n} L(s - 2n + j, f_Q, \chi^{j-1}).$$

Theorem 2.4. *Let k be a positive integer and n a non-negative integer. Let f be a primitive form in $\mathfrak{S}_{2k}(SL_2(\mathbf{Z}))$. Then, we have*

$$L(s, I_{2n+1}(f)) = D^{ns+n^2+3n/2} 2^{-2n} \prod_{i=2}^{2n+1} \tilde{\Lambda}(i, \chi^i) \prod_{j=1}^{2n+1} L(s-2n-1+j, f, \chi^{j-1}).$$

Remark. We note that $L(s, I_{2n+1}(f))$ has an Euler product.

3. REDUCTION TO LOCAL COMPUTATIONS

To prove our main result, we reduce the problem to local computations. Let $K_p = K \otimes \mathbf{Q}_p$ and $\mathcal{O}_p = \mathcal{O} \otimes \mathbf{Z}_p$ as in Notation. Then K_p is a quadratic extension of \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. In the former case let f_p the exponent of the conductor of K_p/\mathbf{Q}_p . If K_p is ramified over \mathbf{Q}_p , put $e_p = f_p - \delta_{2,p}$, where $\delta_{2,p}$ is Kronecker's delta. If K_p is unramified over \mathbf{Q}_p , put $e_p = f_p = 0$. In the latter case, put $e_p = f_p = 0$. Let K_p be a quadratic extension of \mathbf{Q}_p , and $\varpi = \varpi_p$ and $\pi = \pi_p$ be prime elements of K_p and \mathbf{Q}_p , respectively. If K_p is unramified over \mathbf{Q}_p , we take $\varpi = \pi = p$. If K_p is ramified over \mathbf{Q}_p , we take π so that $\pi = N_{K_p/\mathbf{Q}_p}(\varpi)$. Let $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then put $\varpi = \pi = p$. Let χ_{K_p} be the quadratic character of \mathbf{Q}_p^\times corresponding to the quadratic extension K_p/\mathbf{Q}_p . We note that we have $\chi_{K_p}(a) = (-D_0, a)_p$ for $a \in \mathbf{Q}_p^\times$ if $K_p = \mathbf{Q}_p(\sqrt{-D_0})$ with $D_0 \in \mathbf{Z}_p$. Moreover put $\widetilde{\text{Her}}_m(\mathcal{O}_p) = p^{e_p} \widetilde{\text{Her}}_m(\mathcal{O}_p)$. We note that $\widetilde{\text{Her}}_m(\mathcal{O}_p) = \text{Her}_m(\mathcal{O}_p)$ if K_p is not ramified over \mathbf{Q}_p . Let K be an imaginary quadratic extension of \mathbf{Q} with discriminant $-D$. We then put $\tilde{D} = \prod_{p|D} p^{e_p}$, and $\widetilde{\text{Her}}_m(\mathcal{O}) = \tilde{D} \widetilde{\text{Her}}_m(\mathcal{O})$. An element $X \in M_{ml}(\mathcal{O}_p)$ with $m \geq l$ is said to be primitive if there is an element Y of $M_{m, m-l}(\mathcal{O}_p)$ such that $(X \ Y) \in GL_m(\mathcal{O}_p)$. If K_p is a field, this is equivalent to saying that $\text{rank}_{\mathcal{O}_p/\varpi\mathcal{O}_p} X = l$. If $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$, and $X = (X_1, X_2) \in M_{ml}(\mathbf{Z}_p) \oplus M_{ml}(\mathbf{Z}_p)$, this is equivalent to saying that $\text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} X_1 = \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} X_2 = l$. Now let m and l be positive integers such that $m \geq l$. Then for an integer a and $A \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$, $B \in \widetilde{\text{Her}}_l(\mathcal{O}_p)$ put

$$\mathcal{A}_a(A, B) = \{X \in M_{ml}(\mathcal{O}_p)/p^a M_{ml}(\mathcal{O}_p) \mid A[X] - B \in p^a \widetilde{\text{Her}}_l(\mathcal{O}_p)\},$$

and

$$\mathcal{B}_a(A, B) = \{X \in \mathcal{A}_a(A, B) \mid X \text{ is primitive}\}.$$

Suppose that A and B are non-degenerate. Then the number $p^{a(-2ml+l^2)} \#\mathcal{A}_a(A, B)$ is independent of a if a is sufficiently large. Hence we define the local density $\alpha_p(A, B)$ representing B by A as

$$\alpha_p(A, B) = \lim_{a \rightarrow \infty} p^{a(-2ml+l^2)} \#\mathcal{A}_a(A, B).$$

Similarly we can define the primitive local density $\beta_p(A, B)$ as

$$\beta_p(A, B) = \lim_{a \rightarrow \infty} p^{a(-2ml+l^2)} \#\mathcal{B}_a(A, B)$$

if A is non-degenerate. We remark that the primitive local density $\beta_p(A, B)$ can be defined even if B is not non-degenerate. In particular we write $\alpha_p(A) = \alpha_p(A, A)$. We also define $v_p(A)$ for $A \in \text{Her}_m(\mathcal{O}_p)^\times$ as

$$v_p(A) = \lim_{a \rightarrow \infty} p^{-am^2} \#(\Upsilon_a(A)),$$

where

$$\Upsilon_a(A) = \{X \in M_m(\mathcal{O}_p)/p^a M_m(\mathcal{O}_p) \mid A[X] - A \in p^a \text{Her}_m(\mathcal{O}_p)\}.$$

The relation between $\alpha_p(A)$ and $v_p(A)$ is as follows:

Lemma 3.1. *Let $T \in \widetilde{\text{Her}}_m(\mathcal{O}_p)^\times$. Suppose that K_p is ramified over \mathbf{Q}_p . Then we have*

$$\alpha_p(T) = p^{-m(m+1)f_p/2+m^2\delta_{2,p}} v_p(T).$$

Otherwise, $\alpha_p(T) = v_p(T)$.

Proof. The proof is similar to that in [Kitaoka [Kit93], Lemma 5.6.5], and we here give an outline of the proof. The last assertion is trivial. Suppose that K_p is ramified over \mathbf{Q}_p . Let $\{T_i\}_{i=1}^l$ be a complete set of representatives of $\text{Her}_m(\mathcal{O}_p)/p^{r+e_p} \text{Her}_m(\mathcal{O}_p)$ such that $T_i \equiv T \pmod{p^r \widetilde{\text{Her}}_m(\mathcal{O}_p)}$. Then it is easily seen that

$$l = [p^r \widetilde{\text{Her}}_m(\mathcal{O}_p) : p^{r+e_p} \text{Her}_m(\mathcal{O}_p)] = p^{m(m-1)f_p/2}.$$

Define a mapping

$$\phi : \bigsqcup_{i=1}^l \Upsilon_{r+e_p}(T_i) \longrightarrow \mathcal{A}_r(T, T)$$

by $\phi(X) = X \pmod{p^r}$. For $X \in \mathcal{A}_r(T, T)$ and $Y \in M_m(\mathcal{O}_p)$ we have

$$T[X + p^r Y] \equiv T[X] \pmod{p^r \widetilde{\text{Her}}_m(\mathcal{O}_p)}.$$

Namely, $X + p^r Y$ belongs to $\Upsilon_{r+e_p}(T_i)$ for some i and therefore ϕ is surjective.

Moreover for $X \in \mathcal{A}_r(T, T)$ we have $\#(\phi^{-1}(X)) = p^{2m^2 e_p}$. For a sufficiently large integer r we have $\#\Upsilon_{r+e_p}(T_i) = \#\Upsilon_{r+e_p}(T)$ for any i . Hence

$$\begin{aligned} p^{m(m-1)f_p/2} \#\Upsilon_{r+e_p}(T) &= \sum_{i=1}^l \#\Upsilon_{r+e_p}(T_i) \\ &= p^{2m^2 e_p} \#\mathcal{A}_r(T, T) = p^{m^2 e_p} \#\mathcal{A}_{r+e_p}(T, T). \end{aligned}$$

Recall that $e_p = f_p - \delta_{2,p}$. Hence

$$\#\Upsilon_{r+e_p}(T) = p^{m(m+1)f_p/2-m^2\delta_{2,p}} \#\mathcal{A}_{r+e_p}(T, T).$$

This proves the assertion. \square

For $T \in \text{Her}_m(K)^+$, let $\mathcal{G}(T)$ denote the set of $SL_m(\mathcal{O})$ -equivalence classes of positive definite Hermitian matrices T' such that T' is $SL_m(\mathcal{O}_p)$ -equivalent to T for any prime number p . Moreover put

$$M^*(T) = \sum_{T' \in \mathcal{G}(T)} \frac{1}{e^*(T')}$$

for a positive definite Hermitian matrix T of degree m with entries in \mathcal{O} .

Let \mathcal{U}_1 be the unitary group defined in Section 1. Namely let

$$\mathcal{U}_1 = \{u \in R_{K/\mathbf{Q}}(GL_1) \mid \bar{u}u = 1\}.$$

For an element $T \in \text{Her}_m(\mathcal{O}_p)$, let

$$\widetilde{\mathcal{U}}_{p,T} = \{\det X \mid X \in \mathcal{U}_T(K_p) \cap GL_m(\mathcal{O}_p)\},$$

and put $U_{1,p} = \mathcal{U}_1(K_p) \cap \mathcal{O}_p^*$. Then $\widetilde{U}_{p,T}$ is a subgroup of $U_{1,p}$ of finite index. We then put $l_{p,T} = [U_{1,p} : \widetilde{U}_{p,T}]$. We also put

$$u_p = \begin{cases} (1+p^{-1})^{-1} & \text{if } K_p/\mathbf{Q}_p \text{ is unramified} \\ (1-p^{-1})^{-1} & \text{if } K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p \\ 2^{-1} & \text{if } K_p/\mathbf{Q}_p \text{ is ramified.} \end{cases}$$

To state the Mass formula for SU_T , put $\Gamma_{\mathbf{C}}(s) = 2(2\pi)^{-s}\Gamma(s)$.

Proposition 3.2. *Let $T \in \text{Her}_m(\mathcal{O})^+$. Then*

$$M^*(T) = \frac{(\det T)^m \prod_{i=2}^m D^{i/2} \Gamma_{\mathbf{C}}(i)}{2^{m-1} \prod_p l_{p,T} u_p v_p(T)}.$$

Proof. The assertion is more or less well known (cf. [Re71].) But for the sake of completeness we here give an outline of the proof. Let $SU_T(\mathbf{A})$ be the adelization of SU_T and let $\{x_i\}_{i=1}^H$ be a subset of $SU_T(\mathbf{A})$ such that

$$SU_T(\mathbf{A}) = \bigsqcup_{i=1}^H \mathcal{Q} x_i SU_T(\mathbf{Q}),$$

where $\mathcal{Q} = SU_T(\mathbf{R}) \prod_{p < \infty} (SU_T(K_p) \cap SL_m(\mathcal{O}_p))$. We note that the strong approximation theorem holds for SL_m . Hence, by using the standard method we can prove that

$$M^*(T) = \sum_{i=1}^H \frac{1}{\#(x_i^{-1} \mathcal{Q} x_i \cap SU_T(\mathbf{Q}))}.$$

We recall that the Tamagawa number of SU_T is 1 (cf. Weil [We82]). Hence, by [[Re71], (1.1) and (4.5)], we have

$$M^*(T) = \frac{(\det T)^m \prod_{i=2}^m D^{i/2} \Gamma_{\mathbf{C}}(i)}{2^{m-1} \prod_p l_{p,T}} \frac{v_p(1)}{v_p(T)}.$$

We can easily show that $v_p(1) = u_p^{-1}$. This completes the assertion. \square

Corollary. *Let $T \in \widetilde{\text{Her}}_m(\mathcal{O})^+$. Then*

$$M^*(T) = \frac{2^{c_D m^2} (\det T)^m \prod_{i=2}^m \Gamma_{\mathbf{C}}(i)}{2^{m-1} D^{m(m+1)/4+1/2} \prod_p u_p l_{p,T} \alpha_p(T)},$$

where $c_D = 1$ or 0 according as 2 divides D or not.

For a subset \mathcal{T} of \mathcal{O}_p put

$$\text{Her}_m(\mathcal{T}) = \text{Her}_m(\mathcal{O}_p) \cap M_m(\mathcal{T}),$$

and for a subset \mathcal{S} of \mathcal{O}_p put

$$\text{Her}_m(\mathcal{S}, \mathcal{T}) = \{A \in \text{Her}_m(\mathcal{T}) \mid \det A \in \mathcal{S}\},$$

and $\widetilde{\text{Her}}_m(\mathcal{S}, \mathcal{T}) = \text{Her}_m(\mathcal{S}, \mathcal{T}) \cap \widetilde{\text{Her}}_m(\mathcal{O}_p)$. In particular if \mathcal{S} consists of a single element d we write $\text{Her}_m(\mathcal{S}, \mathcal{T})$ as $\text{Her}_m(d, \mathcal{T})$, and so on. For $d \in \mathbf{Z}_{>0}$ we also define the set $\text{Her}_m(d, \mathcal{O})^+$ in a similar way. For each $T \in \widetilde{\text{Her}}_m(\mathcal{O}_p)^\times$ put

$$F_p^{(0)}(T, X) = F_p(p^{-e_p} T, X)$$

and

$$\tilde{F}_p^{(0)}(T, X) = \tilde{F}_p(p^{-e_p}T, X).$$

We remark that

$$\tilde{F}_p^{(0)}(T, X) = X^{-\text{ord}_p(\det T)} X^{e_p m - f_p[m/2]} F_p^{(0)}(T, p^{-m} X^2).$$

For $d \in \mathbf{Z}_p^\times$ put

$$\lambda_{m,p}(d, X) = \sum_{A \in \widetilde{\text{Her}}_m(d, \mathcal{O}_p)/SL_m(\mathcal{O}_p)} \frac{\tilde{F}_p^{(0)}(A, X)}{u_p l_{p,A} \alpha_p(A)}.$$

An explicit formula for $\lambda_{m,p}(p^i d_0, X)$ will be given in the next section for $d_0 \in \mathbf{Z}_p^*$ and $i \geq 0$.

Now let $\widetilde{\text{Her}}_m = \prod_p (\widetilde{\text{Her}}_m(\mathcal{O}_p)/SL_m(\mathcal{O}_p))$. Then the diagonal embedding induces a mapping

$$\phi : \widetilde{\text{Her}}_m(\mathcal{O}) / \prod_p SL_m(\mathcal{O}_p) \longrightarrow \widetilde{\text{Her}}_m.$$

Proposition 3.3. *In addition to the above notation and the assumption, for a positive integer d let*

$$\widetilde{\text{Her}}_m(d) = \prod_p (\widetilde{\text{Her}}_m(d, \mathcal{O}_p)/SL_m(\mathcal{O}_p)).$$

Then the mapping ϕ induces a bijection from $\widetilde{\text{Her}}_m(d, \mathcal{O}) / \prod_p SL_m(\mathcal{O}_p)$ to $\widetilde{\text{Her}}_m(d)$, which will be denoted also by ϕ .

Proof. The proof is similar to that of [[IS95], Proposition 2.1], but it is a little bit more complex because the class number of K is not necessarily one. It is easily seen that ϕ is injective. Let $(x_p) \in \widetilde{\text{Her}}_m(d)$. Then by Theorem 6.9 of [Sch85], there exists an element y in $\text{Her}_m(K)^+$ such that $\det y \in dN_{K/\mathbf{Q}}(K^\times)$. Then we have $\det y \in \det x_p N_{K_p/\mathbf{Q}_p}(K_p^\times)$ for any p . Thus by [[Jac62], Theorem 3.1] we have $x_p = g_p^* y g_p$ with some $g_p \in GL_m(K_p)$ for any prime number p . For p not dividing Dd we may suppose $g_p \in GL_m(\mathcal{O}_p)$. Hence (g_p) defines an element of $R_{K/\mathbf{Q}}(GL_m)(\mathbf{A}_f)$. Since we have $d^{-1} \det y \in \mathbf{Q}^\times \cap \prod_p N_{K_p/\mathbf{Q}_p}(K_p)$, we see that $d^{-1} \det y = N_{K/\mathbf{Q}}(u)$ with some $u \in K^\times$. Thus, by replacing y with $\begin{pmatrix} 1_{m-1} & \mathcal{O} \\ \mathcal{O} & u^{-1} \end{pmatrix} y \begin{pmatrix} 1_{m-1} & \mathcal{O} \\ \mathcal{O} & u^{-1} \end{pmatrix}$, we may suppose that $\det y = d$. Then we have $N_{K_p/\mathbf{Q}_p}(\det g_p) = 1$. It is easily seen that there exists an element $\delta_p \in GL_m(K_p)$ such that $\det \delta_p = \det g_p^{-1}$ and $\delta_p^* x_p \delta_p = x_p$. Thus we have $g_p \delta_p \in SL_m(K_p)$ and

$$x_p = (g_p \delta_p)^* y g_p \delta_p.$$

By the strong approximation theorem for SL_m there exists an element $\gamma \in SL_m(K)$, $\gamma_\infty \in SL_m(\mathbf{C})$, and $(\gamma_p) \in \prod_p SL_m(\mathcal{O}_p)$ such that

$$(g_p \delta_p) = \gamma \gamma_\infty (\gamma_p).$$

Put $x = \gamma^* y \gamma$. Then x belongs to $\widetilde{\text{Her}}_m(d, \mathcal{O})^+$, and $\phi(x) = (x_p)$. This proves the surjectivity of ϕ . \square

Theorem 3.4. *Let f be a primitive form in $\mathfrak{S}_{2k+1}(\Gamma_0(D), \chi)$ or in $\mathfrak{S}_{2k}(SL_2(\mathbf{Z}))$ according as $m = 2n$ or $2n + 1$. For such an f and a positive integer d_0 put*

$$b_m(f; d_0) = \prod_p \lambda_{m,p}(d_0, \alpha_p^{-1}),$$

where α_p is the Satake p -parameter of f . Moreover put

$$\begin{aligned} \mu_{m,k,D} &= D^{m(s-k+l_0/2)+(k-l_0/2)[m/2]-m(m+1)/4-1/2} \\ &\times 2^{-c_D m(s-k-2n-l_0/2)-m+1} \prod_{i=2}^m \Gamma_{\mathbf{C}}(i), \end{aligned}$$

where $l_0 = 0$ or 1 according as m is even or odd. Then for $\text{Re}(s) \gg 0$, we have

$$L(s, I_m(f)) = \mu_{m,k,D} \sum_{d_0=1}^{\infty} b_m(f; d_0) d_0^{-s+k+2n+l_0/2}.$$

Proof. We note that $L(s, I_m(f))$ can be rewritten as

$$L(s, I_m(f)) = \tilde{D}^{ms} \sum_{T \in \widetilde{\text{Her}}_m(\mathcal{O})^+ / SL_m(\mathcal{O})} \frac{a_{I_m(f)}(\tilde{D}^{-1}T)}{e^*(T)(\det T)^s}.$$

For $T \in \widetilde{\text{Her}}_m(\mathcal{O})^+$ the Fourier coefficient $a_{I_m(f)}(\tilde{D}^{-1}T)$ of $I_m(f)$ is uniquely determined by the genus to which T belongs, and can be expressed as

$$a_{I_m(f)}(\tilde{D}^{-1}T) = (D^{[m/2]} \tilde{D}^{-m} \det T)^{k-l_0/2} \prod_p \tilde{F}_p^{(0)}(T, \alpha_p^{-1}).$$

Thus the assertion follows from Corollary to Proposition 3.2 and Proposition 3.3 similarly to [IS95]. \square

4. FORMAL POWER SERIES ASSOCIATED WITH LOCAL SIEGEL SERIES

For $d_0 \in \mathbf{Z}_p^\times$ put

$$\hat{P}_{m,p}(d_0, X, t) = \sum_{i=0}^{\infty} \lambda_{m,p}^*(p^i d_0, X) t^i,$$

where for $d \in \mathbf{Z}_p^\times$ we define $\lambda_{m,p}^*(d, X)$ as

$$\lambda_{m,p}^*(d, X) = \sum_{A \in \widetilde{\text{Her}}_m(dN_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*), \mathcal{O}_p) / GL_m(\mathcal{O}_p)} \frac{\tilde{F}_p^{(0)}(A, X)}{\alpha_p(A)}.$$

We note that

$$\sum_{A \in \widetilde{\text{Her}}_m(dN_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*), \mathcal{O}_p) / GL_m(\mathcal{O}_p)} \frac{\tilde{F}_p^{(0)}(A, X^{-1})}{\alpha_p(A)}$$

is $\chi_{K_p}((-1)^{m/2}d)\lambda_{m,p}^*(d, X)$ or $\lambda_{m,p}^*(d, X)$ according as m is even and K_p is a field, or not. In Proposition 4.3.7 we will show that we have

$$\lambda_{m,p}^*(d, X) = u_p \lambda_{m,p}(d, X)$$

for $d \in \mathbf{Z}_p^\times$ and therefore

$$\hat{P}_{m,p}(d_0, X, t) = u_p \sum_{i=0}^{\infty} \lambda_{m,p}(p^i d_0, X) t^i.$$

We also define $P_{m,p}(d_0, X, t)$ as

$$P_{m,p}(d_0, X, t) = \sum_{i=0}^{\infty} \lambda_{m,p}^*(\pi_p^i d_0, X) t^i.$$

We note that $P_{m,p}(d_0, X, t) = \hat{P}_{m,p}(d_0, X, t)$ if K_p is unramified over \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$, but it is not necessarily the case if K_p is ramified over \mathbf{Q}_p . In this section, we give explicit formulas of $P_{m,p}(d_0, X, t)$ for all prime numbers p (cf. Theorems 4.3.1 and 4.3.2), and therefore explicit formulas for $\hat{P}_{m,p}(d_0, X, t)$ (cf. Theorem 4.3.6.)

From now on we fix a prime number p . Throughout this section we simply write ord_p as ord and so on if the prime number p is clear from the context. We also write ν_{K_p} as ν . We also simply write $\widetilde{\text{Her}}_{m,p}$ instead of $\widetilde{\text{Her}}_m(\mathcal{O}_p)$, and so on.

4.1. Preliminaries.

Let m be a positive integer. For a non-negative integer $i \leq m$ let

$$\mathcal{D}_{m,i} = GL_m(\mathcal{O}_p) \begin{pmatrix} 1_{m-i} & 0 \\ 0 & \varpi 1_i \end{pmatrix} GL_m(\mathcal{O}_p),$$

and for $W \in \mathcal{D}_{m,i}$, put $\Pi_p(W) = (-1)^i p^{i(i-1)a/2}$, where $a = 2$ or 1 according as K_p is unramified over \mathbf{Q}_p or not. Let $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then for a pair $i = (i_1, i_2)$ of non-negative integers such that $i_1, i_2 \leq m$, let

$$\mathcal{D}_{m,i} = GL_m(\mathcal{O}_p) \left(\begin{pmatrix} 1_{m-i_1} & 0 \\ 0 & p 1_{i_1} \end{pmatrix}, \begin{pmatrix} 1_{m-i_2} & 0 \\ 0 & p 1_{i_2} \end{pmatrix} \right) GL_m(\mathcal{O}_p),$$

and for $W \in \mathcal{D}_{m,i}$ put $\Pi_p(W) = (-1)^{i_1+i_2} p^{i_1(i_1-1)/2+i_2(i_2-1)/2}$. In either case K_p is a quadratic extension of \mathbf{Q}_p , or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$, we put $\Pi_p(W) = 0$ for $W \in M_n(\mathcal{O}_p^\times) \setminus \bigcup_{i=0}^m \mathcal{D}_{m,i}$.

First we remark the following lemma, which can easily be proved by the usual Newton approximation method in \mathcal{O}_p :

Lemma 4.1.1. *Let $A, B \in \widetilde{\text{Her}}_m(\mathcal{O}_p)^\times$. Let e be an integer such that $p^e A^{-1} \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$. Suppose that $A \equiv B \pmod{p^{e+1} \widetilde{\text{Her}}_m(\mathcal{O}_p)}$. Then there exists a matrix $U \in GL_m(\mathcal{O}_p)$ such that $B = A[U]$.*

Lemma 4.1.2. *Let $S \in \widetilde{\text{Her}}_m(\mathcal{O}_p)^\times$ and $T \in \widetilde{\text{Her}}_n(\mathcal{O}_p)^\times$ with $m \geq n$. Then*

$$\alpha_p(S, T) = \sum_{W \in GL_n(\mathcal{O}_p) \setminus M_n(\mathcal{O}_p)^\times} p^{(n-m)\nu(\det W)} \beta_p(S, T[W^{-1}]).$$

Proof. The assertion can be proved by using the same argument as in the proof of [[Kit93], Theorem 5.6.1]. We here give an outline of the proof. For each $W \in M_n(\mathcal{O}_p)$, put

$$\mathcal{B}_e(S, T; W) = \{X \in \mathcal{A}_e(S, T) \mid XW^{-1} \text{ is primitive}\}.$$

Then we have

$$\mathcal{A}_e(S, T) = \bigsqcup_{W \in GL_n(\mathcal{O}_p) \setminus M_n(\mathcal{O}_p)^\times} \mathcal{B}_e(S, T; W).$$

Take a sufficiently large integer e , and for an element W of $M_n(\mathcal{O}_p)$, let $\{R_i\}_{i=1}^r$ be a complete set of representatives of $p^e \widetilde{\text{Her}}_m(\mathcal{O}_p)[W^{-1}]/p^e \widetilde{\text{Her}}_m(\mathcal{O}_p)$. Then we have $r = p^{\nu(\det W)n}$. Put

$$\begin{aligned} \widetilde{\mathcal{B}}_e(S, T; W) = \{X \in M_{mn}(\mathcal{O}_p)/p^e M_{mn}(\mathcal{O}_p)W \mid S[X] \equiv T \pmod{p^e \widetilde{\text{Her}}_m(\mathcal{O}_p)} \\ \text{and } XW^{-1} \text{ is primitive}\}. \end{aligned}$$

Then

$$\#(\widetilde{\mathcal{B}}_e(S, T; W)) = p^{\nu(\det W)m} \#(\mathcal{B}_e(S, T; W)).$$

It is easily seen that

$$S[XW^{-1}] \equiv T[W^{-1}] + R_i \pmod{p^e \widetilde{\text{Her}}_m(\mathcal{O}_p)}$$

for some i . Hence the mapping $X \mapsto XW^{-1}$ induces a bijection from $\widetilde{\mathcal{B}}_e(S, T; W)$ to $\bigsqcup_{i=1}^r \mathcal{B}_e(S, T[W^{-1}] + R_i)$. Recall that $\nu(W) \leq \text{ord}(\det T)$. Hence

$$R_i \equiv O \pmod{p^{\lfloor e/2 \rfloor} \widetilde{\text{Her}}_m(\mathcal{O}_p)},$$

and therefore by Lemma 4.1.1,

$$T[W^{-1}] + R_i = T[W^{-1}][G]$$

for some $G \in GL_n(\mathcal{O}_p)$. Hence

$$\#(\widetilde{\mathcal{B}}_e(S, T; W)) = p^{\nu(\det W)n} \#(\mathcal{B}_e(S, T[W^{-1}])).$$

Hence

$$\begin{aligned} \alpha_p(S, T) &= p^{-2mne+n^2e} \#(\mathcal{A}_e(S, T)) \\ &= p^{-2mne+n^2e} \sum_{W \in GL_n(\mathcal{O}_p) \setminus M_n(\mathcal{O}_p)^\times} p^{\nu(\det W)(-m+n)} \#(\mathcal{B}_e(S, T[W^{-1}])). \end{aligned}$$

This proves the assertion. □

Now by using the same argument as in the proof of [[Kit83], Theorem 1], we obtain

Corollary. *Under the same notation as above, we have*

$$\beta_p(S, T) = \sum_{W \in GL_n(\mathcal{O}_p) \setminus M_n(\mathcal{O}_p)^\times} p^{(n-m)\nu(\det W)} \Pi_p(W) \alpha_p(S, T[W^{-1}]).$$

For two elements $A, A' \in \text{Her}_m(\mathcal{O}_p)$ we simply write $A \sim_{GL_m(\mathcal{O}_p)} A'$ as $A \sim A'$ if there is no fear of confusion. For a variables U and q put

$$(U, q)_m = \prod_{i=1}^m (1 - q^{i-1}U), \quad \phi_m(q) = (q, q)_m.$$

We note that $\phi_m(q) = \prod_{i=1}^m (1 - q^i)$. Moreover for a prime number p put

$$\phi_{m,p}(q) = \begin{cases} \phi_m(q^2) & \text{if } K_p/\mathbf{Q}_p \text{ is unramified} \\ \phi_m(q)^2 & \text{if } K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p \\ \phi_m(q) & \text{if } K_p/\mathbf{Q}_p \text{ is ramified} \end{cases}$$

Lemma 4.1.3. (1) Let $\Omega(S, T) = \{w \in M_m(\mathcal{O}_p) \mid S[w] \sim T\}$. Then we have

$$\frac{\alpha_p(S, T)}{\alpha_p(T)} = \#(\Omega(S, T)/GL_m(\mathcal{O}_p))p^{-m(\text{ord}(\det T) - \text{ord}(\det S))}.$$

(2) Let $\tilde{\Omega}(S, T) = \{w \in M_m(\mathbf{Z}) \mid S \sim T[w^{-1}]\}$. Then we have

$$\frac{\alpha_p(S, T)}{\alpha_p(S)} = \#(GL_m(\mathcal{O}_p) \backslash \tilde{\Omega}(S, T)).$$

Proof. (1) The proof is similar to that of Lemma 2.2 of [BS87]. First we prove

$$\int_{\Omega(S, T)} |dx| = \phi_{m,p}(p^{-1}) \frac{\alpha_p(S, T)}{\alpha_p(T)},$$

where $|dx|$ is the Haar measure on $M_m(K_p)$ normalized so that

$$\int_{M_m(\mathcal{O}_p)} |dx| = 1.$$

To prove this, for a positive integer e let T_1, \dots, T_l be a complete set of representatives of $\{T[\gamma] \bmod p^e \mid \gamma \in GL_m(\mathcal{O}_p)\}$. Then it is easy to see that

$$\int_{\Omega(S, T)} |dx| = p^{-2m^2e} \sum_{i=1}^l \#(\mathcal{A}_e(S, T_i))$$

and, by Lemma 4.1.1, T_i is $GL_m(\mathcal{O}_p)$ -equivalent to T if e is sufficiently large. Hence we have

$$\#(\mathcal{A}_e(S, T_i)) = \#(\mathcal{A}_e(S, T))$$

for any i . Moreover we have

$$l = \#(GL_m(\mathcal{O}_p/p^e\mathcal{O}_p))/\#(\mathcal{A}_e(T, T)) = p^{m^2e} \phi_{m,p}(p^{-1})/\alpha_p(T).$$

Hence

$$\int_{\Omega(S, T)} |dx| = lp^{-2m^2e} \#(\mathcal{A}_e(S, T)) = \phi_{m,p}(p^{-1}) \frac{\alpha_p(S, T)}{\alpha_p(T)},$$

which proves the above equality. Now we have

$$\int_{\Omega(S, T)} |dx| = \sum_{W \in \Omega(S, T)/GL_m(\mathcal{O}_p)} |\det W|_{K_p}^m = \sum_{W \in \Omega(S, T)/GL_m(\mathcal{O}_p)} |\det W \overline{\det W}|_p^m.$$

Remark that for any $W \in \Omega(S, T)/GL_m(\mathcal{O}_p)$ we have $|\det W \overline{\det W}|_p = p^{-m(\text{ord}(\det T) - \text{ord}(\det S))}$. Thus the assertion has been proved.

(2) By Lemma 4.1.2 we have

$$\alpha_p(S, T) = \sum_{W \in GL_m(\mathcal{O}_p) \setminus M_m(\mathcal{O}_p)^\times} \beta_p(S, T[W^{-1}]).$$

Then we have $\beta_p(S, T[W^{-1}]) = \alpha_p(S)$ or 0 according as $S \sim T[W^{-1}]$ or not. Thus the assertion (2) holds. \square

For a subset \mathcal{T} of \mathcal{O}_p , we put

$$\text{Her}_m(\mathcal{T})_k = \{A = (a_{ij}) \in \text{Her}_m(\mathcal{T}) \mid a_{ii} \in \pi^k \mathbf{Z}_p\}.$$

From now on put

$$\text{Her}_{m,*}(\mathcal{O}_p) = \begin{cases} \text{Her}_m(\mathcal{O}_p)_1 & \text{if } p = 2 \text{ and } f_p = 3, \\ \text{Her}_m(\varpi \mathcal{O}_p)_1 & \text{if } p = 2 \text{ and } f_p = 2 \\ \text{Her}_m(\mathcal{O}_p) & \text{otherwise,} \end{cases}$$

where ϖ is a prime element of K_p . Moreover put $i_p = 0$, or 1 according as $p = 2$ and $f_2 = 2$, or not. Suppose that K_p/\mathbf{Q}_p is unramified or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then an element B of $\widetilde{\text{Her}}_m(\mathcal{O}_p)$ can be expressed as $B \sim_{GL_m(\mathcal{O}_p)} 1_r \perp B_2$ with some integer r and $B_2 \in \text{Her}_{m-r,*}(\mathcal{O}_p)$. Suppose that K_p/\mathbf{Q}_p is ramified. For an even positive integer r define Θ_r by

$$\Theta_r = \overbrace{\left(\begin{array}{cc} 0 & \varpi^{i_p} \\ \bar{\varpi}^{i_p} & 0 \end{array} \right) \perp \dots \perp \left(\begin{array}{cc} 0 & \varpi^{i_p} \\ \bar{\varpi}^{i_p} & 0 \end{array} \right)}^{r/2},$$

where $\bar{\varpi}$ is the conjugate of ϖ over \mathbf{Q}_p . Then an element B of $\widetilde{\text{Her}}_m(\mathcal{O}_p)$ is expressed as $B \sim_{GL_m(\mathcal{O}_p)} \Theta_r \perp \pi^{i_p} B_2$ with some even integer r and $B_2 \in \text{Her}_{m-r,*}(\mathcal{O}_p)$. For these results, see Jacobowitz [Jac62].

A non-degenerate square matrix $W = (d_{ij})_{m \times m}$ with entries in \mathcal{O}_p is called reduced if W satisfies the following conditions:

$d_{ii} = p^{e_i}$ with e_i a non-negative integer, d_{ij} is a non-negative integer $\leq p^{e_j} - 1$ for $i < j$ and $d_{ij} = 0$ for $i > j$. It is well known that we can take the set of all reduced matrices as a complete set of representatives of $GL_m(\mathcal{O}_p) \setminus M_m(\mathcal{O}_p)^\times$. Let m be an integer. For $B \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$ put

$$\widetilde{\Omega}(B) = \{W \in GL_m(K_p) \cap M_m(\mathcal{O}_p) \mid B[W^{-1}] \in \widetilde{\text{Her}}_m(\mathcal{O}_p)\}.$$

Let $r \leq m$, and $\psi_{r,m}$ be the mapping from $GL_r(K_p)$ into $GL_m(K_p)$ defined by $\psi_{r,m}(W) = 1_{m-r} \perp W$.

Lemma 4.1.4. (1) Assume that K_p is unramified over \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let $B_1 \in \text{Her}_{m-n_0}(\mathcal{O}_p)$. Then $\psi_{m-n_0,m}$ induces a bijection from $GL_{m-n_0}(\mathcal{O}_p) \setminus \widetilde{\Omega}(B_1)$ to $GL_m(\mathcal{O}_p) \setminus \widetilde{\Omega}(1_{n_0} \perp B_1)$, which will be also denoted by $\psi_{m-n_0,m}$.

(2) Assume that K_p is ramified over \mathbf{Q}_p and that n_0 is even. Let $B_1 \in \widetilde{\text{Her}}_{m-n_0}(\mathcal{O}_p)$. Then $\psi_{m-n_0,m}$ induces a bijection from $GL_{m-n_0}(\mathcal{O}_p) \setminus \widetilde{\Omega}(B_1)$ to $GL_m(\mathcal{O}_p) \setminus \widetilde{\Omega}(\Theta_{n_0} \perp B_1)$, which will be also denoted by $\psi_{m-n_0,m}$. Here i_p is the integer defined above.

Proof. (1) Clearly $\psi_{m-n_0, m}$ is injective. To prove the surjectivity, take a representative W of an element of $GL_m(\mathcal{O}_p) \backslash \widetilde{\Omega}(1_{n_0} \perp B_1)$. Without loss of generality we may assume that W is a reduced matrix. Since we have $(1_{n_0} \perp B_1)[W^{-1}] \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$, we have $W = \begin{pmatrix} 1_{n_0} & 0 \\ 0 & W_1 \end{pmatrix}$ with $W_1 \in \widetilde{\Omega}(B_1)$. This proves the assertion.

(2) The assertion can be proved in the same manner as (1). \square

Lemma 4.1.5. *Let $B \in \widetilde{\text{Her}}_m(\mathcal{O}_p)^\times$. Then we have*

$$\alpha_p(\pi^r dB) = p^{rm^2} \alpha_p(B)$$

for any non-negative integer r and $d \in \mathbf{Z}_p^*$.

Proof. The assertion can be proved by using the same argument as in the proof of (a) of Theorem 5.6.4 of Kitaoka [Ki2]. \square

Now we prove induction formulas for local densities different from Lemma 4.1.2 (cf. Lemmas 4.1.6, 4.1.7, and 4.1.8.) For technical reason, we formulate and prove them in terms of Hermitian modules. Let M be \mathcal{O}_p free module, and let b be a mapping from $M \times M$ to K_p such that

$$b(\lambda_1 u + \lambda_2 u_2, v) = \lambda_1 b(u_1, v) + \lambda_2 b(u_2, v)$$

for $u, v \in M$ and $\lambda_1, \lambda_2 \in \mathcal{O}_p$, and

$$b(u, v) = \overline{b(v, u)} \text{ for } u, v \in M.$$

We call such an M a Hermitian module with a Hermitian inner product b . We set $q(u) = b(u, u)$ for $u \in M$. Take an \mathcal{O}_p -basis $\{u_i\}_{i=1}^m$ of M , and put $T_M = (b(u_i, u_j))_{1 \leq i, j \leq m}$. Then T_M is a Hermitian matrix, and its determinant is uniquely determined, up to $N_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*)$, by M . We say M is non-degenerate if $\det T_M \neq 0$. Conversely for a Hermitian matrix T of degree m , we can define a Hermitian module M_T so that

$$M_T = \mathcal{O}_p u_1 + \mathcal{O}_p u_2 \cdots + \mathcal{O}_p u_m$$

with $(b(u_i, u_j))_{1 \leq i, j \leq m} = T$. Let M_1 and M_2 be submodules of M . We then write $M = M_1 \perp M_2$ if $M = M_1 + M_2$, and $b(u, v) = 0$ for any $u \in M_1, v \in M_2$. Let M and N be Hermitian modules. Then a homomorphism $\sigma : N \rightarrow M$ is said to be an isometry if σ is injective and $b(\sigma(u), \sigma(v)) = b(u, v)$ for any $u, v \in N$. In particular M is said to be isometric to N if σ is an isomorphism. We denote by \widetilde{U}'_M the group of isometries of M to M itself. From now on we assume that $T_M \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$ for a Hermitian module M of rank m . For Hermitian modules M and N over \mathcal{O}_p of rank m and n respectively, put

$$\mathcal{A}'_a(N, M) = \{\sigma : N \rightarrow M/p^a M \mid q(\sigma(u)) \equiv q(u) \pmod{p^{e_p+a}}\},$$

and

$$\mathcal{B}'_a(N, M) := \{\sigma \in \mathcal{A}'_a(N, M) \mid \sigma \text{ is primitive}\}.$$

Here a homomorphism $\sigma : N \rightarrow M$ is said to be primitive if ϕ induces an injective mapping from $N/\varpi N$ to $M/\varpi M$. Then we can define the local density $\alpha'_p(N, M)$ as

$$\alpha'_p(N, M) = \lim_{a \rightarrow \infty} p^{-a(2mn-n^2)} \#(\mathcal{A}'_a(N, M))$$

if M and N are non-degenerate, and the primitive local density $\beta'_p(N, M)$ as

$$\beta'_p(N, M) = \lim_{a \rightarrow \infty} p^{-a(2mn-n^2)} \#(\mathcal{B}'_a(N, M))$$

if M is non-degenerate as in the matrix case. It is easily seen that

$$\alpha_p(S, T) = \alpha'_p(M_T, M_S),$$

and

$$\beta_p(S, T) = \beta'_p(M_T, M_S).$$

Let N_1 be a submodule of N . For each $\phi_1 \in \mathcal{B}'_a(N_1, M)$, put

$$\mathcal{B}'_a(N, M; \phi_1) = \{\phi \in \mathcal{B}'_a(N, M) \mid \phi|_{N_1} = \phi_1\}.$$

We note that we have

$$\mathcal{B}'_a(N, M) = \bigsqcup_{\phi_1 \in \mathcal{B}'_a(N_1, M)} \mathcal{B}'_a(N, M; \phi_1).$$

Suppose that K_p is unramified over \mathbf{Q}_p . Then put $\Xi_m = 1_m$. Suppose that K_p is ramified over \mathbf{Q}_p , and that m is even. Then put $\Xi_m = \Theta_m$.

Lemma 4.1.6. *Let m_1, m_2, n_1 , and n_2 be non-negative integers such that $m_1 \geq n_1$ and $m_1 + m_2 \geq n_1 + n_2$. Moreover suppose that m_1 and n_1 are even if K_p is ramified over \mathbf{Q}_p . Let $A_2 \in \widehat{\text{Her}}_{m_2}(\mathcal{O}_p)$ and $B_2 \in \widehat{\text{Her}}_{n_2}(\mathcal{O}_p)$. Then we have*

$$\beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1} \perp B_2) = \beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1}) \beta_p(\Xi_{m_1-n_1} \perp A_2, B_2),$$

and in particular we have

$$\beta_p(\Xi_{n_1} \perp A_2, \Xi_{n_1} \perp B_2) = \beta_p(\Xi_{n_1} \perp A_2, \Xi_{n_1}) \beta_p(A_2, B_2),$$

Proof. Let $M = M_{\Xi_{m_1} \perp A_2}$, $N_1 = M_{\Xi_{n_1}}$, $N_2 = M_{B_2}$, and $N = N_1 \perp N_2$. Let a be a sufficiently large positive integer. Let $\widetilde{N}_1 = \mathcal{O}_p v_1 \oplus \cdots \oplus \mathcal{O}_p v_{n_1}$ and $\widetilde{N}_2 = \mathcal{O}_p v_{n_1+1} \oplus \cdots \oplus \mathcal{O}_p v_{n_1+n_2}$. For each $\phi_1 \in \mathcal{B}'_a(N_1, M)$, put $u_i = \phi_1(v_i)$ for $i = 1, \dots, n_1$. Then we can take elements $u_{n_1+1}, \dots, u_{m_1+m_2} \in M$ such that

$$(u_i, u_j) = 0 \quad (i = 1, \dots, n_1, j = n_1 + 1, \dots, m_1 + m_2),$$

and

$$((u_i, u_j))_{n_1+1 \leq i, j \leq m_1+m_2} = \Xi_{m_1-n_1} \perp A_2.$$

Put $N'_1 = \mathcal{O}_p u_1 \oplus \cdots \oplus \mathcal{O}_p u_{n_1}$. Then we have $N'_1 = M_{\Xi_{n_1}}$. For $\phi \in \mathcal{B}'_a(N_1, M; \phi_1)$ and $i = 1, \dots, n_2$ we have

$$\phi(v_{n_1+i}) = \sum_{j=1}^{m_1+m_2} a_{n_1+i, j} u_j$$

with $a_{n_1+i, j} \in \mathcal{O}_p$. Put $\Xi_{n_1} = (b_{ij})_{1 \leq i, j \leq n_1}$. Then we have

$$(\phi(v_j), \phi(v_{n_1+i})) = \sum_{\gamma=1}^{n_1} \overline{a_{n_1+i, \gamma}} b_{j\gamma} = 0$$

for $i = 1, \dots, n_2$ and $j = 1, \dots, n_1$. Hence we have $a_{n_1+i, \gamma} = 0$ for $i = 1, \dots, n_2$ and $\gamma = 1, \dots, n_1$. This implies that $\phi|_{N_2} \in \mathcal{B}'_a(N_2, M_{A_2 \perp \Xi_{m_1-n_1}})$. Then the mapping

$$\mathcal{B}'_a(N_1, M; \phi_1) \ni \phi \mapsto \phi|_{N_2} \in \mathcal{B}'_a(N_2, M_{A_2 \perp \Xi_{m_1-n_1}})$$

is bijective. Thus we have

$$\#\mathcal{B}'_a(N, M) = \#\mathcal{B}'_a(N_1, M) \#\mathcal{B}'_a(N_2, M_{\Xi_{m-n_1} \perp A_2}).$$

This implies that

$$\beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1} \perp B_2) = \beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1}) \beta_p(\Xi_{m_1-n_1} \perp A, B_2).$$

□

Lemma 4.1.7. *In addition to the notation and the assumption in Lemma 4.1.6, suppose that A_1 and A_2 are non-degenerate. Then*

$$\alpha_p(\Xi_{m_1} \perp A_2, \Xi_{n_1}) = \beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1}),$$

and we have

$$\alpha_p(\Xi_{m_1} \perp A_2, \Xi_{n_1} \perp B_2) = \alpha_p(\Xi_{m_1} \perp A_2, \Xi_{n_1}) \alpha_p(\Xi_{m_1-n_1} \perp A_2, B_2),$$

and in particular we have

$$\alpha_p(\Xi_{n_1} \perp A_2, \Xi_{n_1} \perp B_2) = \alpha_p(\Xi_{n_1} \perp A_2, \Xi_{n_1}) \alpha_p(A_2, B_2),$$

Proof. The first assertion can easily be proved. By Lemmas 4.1.2 and 4.1.4, we have

$$\begin{aligned} & \alpha_p(\Xi_{m_1} \perp A_2, \Xi_{n_1} \perp B_2) \\ = & \sum_{W \in \widetilde{GL}_{n_1+n_2}(\mathcal{O}_p) \setminus \widetilde{\Omega}(\Xi_{n_1} \perp B_2)} p^{(n_1+n_2-(m_1+m_2))\nu(\det W)} \beta_p(\Xi_{m_1} \perp A_2, (\Xi_{n_1} \perp B_2)[W^{-1}]) \\ = & \sum_{X \in \widetilde{GL}_{n_2}(\mathcal{O}_p) \setminus \widetilde{\Omega}(B_2)} p^{(n_2-(m_1-n_1+m_2))\nu(\det X)} \beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1} \perp B_2[X^{-1}]). \end{aligned}$$

By Lemma 4.1.6 and the first assertion, we have

$$\beta_p(\Xi_{m_1} \perp A_2, \Xi_{n_1} \perp B_2[X^{-1}]) = \alpha_p(\Xi_{m_1} \perp A_2, \Xi_{n_1}) \beta_p(\Xi_{m_1-n_1} \perp A_2, B_2[X^{-1}]).$$

Hence again by Lemma 4.1.2, we prove the second assertion. □

Lemma 4.1.8. (1) *Suppose that K_p is unramified over \mathbf{Q}_p . Let $A \in \text{Her}_l(\mathcal{O}_p)$, $B_1 \in \text{Her}_{n_1}(\mathcal{O}_p)$ and $B_2 \in \text{Her}_{n_2}(\mathcal{O}_p)$ with $m \geq 2n_1$.*

Then we have

$$\beta_p(1_m \perp A, B_1 \perp B_2) = \beta_p(1_m \perp A, B_1) \beta_p((-B_1) \perp 1_{m-2n_1} \perp A, B_2)$$

(2) *Suppose that K_p is ramified over \mathbf{Q}_p . Let $A \in \widetilde{\text{Her}}_l(\mathcal{O}_p)$, $B_1 \in \widetilde{\text{Her}}_{n_1}(\mathcal{O}_p)$, and $B_2 \in \widetilde{\text{Her}}_{n_2}(\mathcal{O}_p)$ with $m \geq n_1$. Then we have*

$$\beta_p(\Theta_{2m} \perp A, B_1 \perp B_2) = \beta_p(\Theta_{2m} \perp A, B_1) \beta_p((-B_1) \perp \Theta_{2m-2n_1} \perp A, B_2).$$

Proof. First suppose that K_p is ramified over \mathbf{Q}_p . Let $M = M_{\Theta_{2m} \perp A}$, $N_1 = M_{B_1}$, $N_2 = M_{B_2}$, and $N = N_1 \perp N_2$. Let a be a sufficiently large positive integer. Let $N_1 = \mathcal{O}_p v_1 \oplus \cdots \oplus \mathcal{O}_p v_{n_1}$ and $N_2 = \mathcal{O}_p v_{n_1+1} \oplus \cdots \oplus \mathcal{O}_p v_{n_1+n_2}$. For each $\phi_1 \in \mathcal{B}'_a(N_1, M)$, put $u_i = \phi_1(v_i)$ for $i = 1, \dots, n_1$. Then we can take elements $u_{n_1+1}, \dots, u_{2m+l} \in M$ such that

$$\begin{aligned} (u_i, u_{n_1+j}) &= \delta_{ij} \varpi^{ip}, \quad (u_{n_1+i}, u_{n_1+j}) = 0 \quad (i, j = 1, \dots, n_1), \\ (u_i, u_j) &= 0 \quad (i = 1, \dots, 2n_1, j = 2n_1+1, \dots, 2m+l), \end{aligned}$$

and

$$((u_i, u_j))_{2n_1+1 \leq i, j \leq 2m+l} = \Theta_{2m-2n_1} \perp A,$$

where δ_{ij} is Kronecker's delta. Let $B_1 = (b_{ij})_{1 \leq i, j \leq n_1}$, and put

$$u'_j = u_j - \bar{\omega}^{-i_p} \sum_{\gamma=1}^{n_1} \bar{b}_{\gamma j} u_{n_1+\gamma}$$

for $j = 1, \dots, n_1$, and $M' = \mathcal{O}_p u'_1 \oplus \dots \oplus \mathcal{O}_p u'_{n_1}$. Then we have $(u'_i, u'_j) = -b_{ij}$ and hence we have $M' = M_{(-B_1)}$. For $\phi \in \mathcal{B}'_a(N_1, M; \phi_1)$ and $i = 1, \dots, n_2$ we have

$$\phi(v_{n_1+i}) = \sum_{j=1}^{2m+l} a_{n_1+i, j} u_j$$

with $a_{n_1+i, j} \in \mathcal{O}_p$. Then we have

$$(\phi(v_j), \phi(v_{n_1+i})) = \sum_{\gamma=1}^{n_1} \overline{a_{n_1+i, \gamma} b_{j\gamma}} + \overline{a_{n_1+i, n_1+j}} \bar{\omega}^{i_p} = 0$$

for $i = 1, \dots, n_2$ and $j = 1, \dots, n_1$. Hence we have

$$\phi(v_{n_1+i}) = \sum_{j=1}^{n_1} a_{n_1+i, j} u'_j + \sum_{j=2n_1+1}^{2m+l} a_{n_1+i, j} u_j.$$

This implies that $\phi|_{N_2} \in \mathcal{B}'_a(N_2, M_{(-B_1)} \perp M_{A \perp \Theta_{2m-2n_1}})$. Then the mapping

$$\mathcal{B}'_a(N_1, M; \phi_1) \ni \phi \mapsto \phi|_{N_2} \in \mathcal{B}'_a(N_2, M_{(-B_1)} \perp M_{A \perp \Theta_{2m-2n_1}})$$

is bijective. Thus we have

$$\#\mathcal{B}'_a(N, M) = \#\mathcal{B}'_a(N_1, M) \#\mathcal{B}'_a(N_2, M_{(-B_1)} \perp M_{\Theta_{2m-2n_1} \perp A}).$$

This implies that

$$\beta_p(\Theta_{2m} \perp A, B_1 \perp B_2) = \beta_p(\Theta_{2m} \perp A, B_1) \beta_p((-B_1) \perp \Theta_{2m-2n_1} \perp A, B_2).$$

This proves (2). Next suppose that K_p is unramified over \mathbf{Q}_p . For an even positive integer r define Θ_r by

$$\Theta_r = \overbrace{\left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right) \perp \dots \perp \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right)}^{r/2}.$$

Then we have $\Theta_r \sim 1_r$. By using the same argument as above we can prove that

$$\beta_p(\Theta_m \perp A, B_1 \perp B_2) = \beta_p(\Theta_m \perp A, B_1) \beta_p((-B_1) \perp \Theta_{m-2n_1} \perp A, B_2)$$

or

$$\beta_p(\Theta_{m-1} \perp 1 \perp A, B_1 \perp B_2) = \beta_p(\Theta_{m-1} \perp 1 \perp A, B_1) \beta_p((-B_1) \perp \Theta_{m-2n_1} \perp 1 \perp A, B_2)$$

according as m is even or not. Thus we prove the assertion (1). \square

Lemma 4.1.9. *Let k be a positive integer.*

(1) *Suppose that K_p is unramified over \mathbf{Q}_p .*

(1.1) *Let $b \in \mathbf{Z}_p$. Then we have*

$$\beta_p(1_{2k}, pb) = (1 - p^{-2k})(1 + p^{-2k+1}).$$

(1.2) Let $b \in \mathbf{Z}_p^*$. Then we have

$$\alpha_p(1_{2k}, b) = \beta_p(1_{2k}, b) = 1 - p^{-2k},$$

and

$$\alpha_p(1_{2k-1}, b) = \beta_p(1_{2k-1}, b) = 1 + p^{-2k+1}.$$

(2) Suppose that K_p is ramified over \mathbf{Q}_p .

(2.1) Let $B \in \text{Her}_{m,*}(\mathcal{O}_p)$ with $m \leq 2$. Then we have

$$\beta_p(\Theta_{2k}, \pi^{i_p} B) = \prod_{i=0}^{m-1} (1 - p^{-2k+2i}).$$

(2.2) Let $B = \begin{pmatrix} 0 & \varpi \\ \varpi & 0 \end{pmatrix}$. Then we have

$$\alpha_p(\Theta_{2k}, B) = \beta_p(\Theta_{2k}, B) = 1 - p^{-2k}.$$

Proof. (1) Put $B = (b)$. Let $p \neq 2$. Then we have $K_p = \mathbf{Q}_p(\sqrt{\varepsilon})$ with $\varepsilon \in \mathbf{Z}_p^*$ such that $(\varepsilon, p)_p = -1$. Then we have

$$\begin{aligned} \#\mathcal{B}_a(1_{2k}, B) &= \#\{(x_i) \in M_{4k,1}(\mathbf{Z}_p)/p^a M_{4k,1}(\mathbf{Z}_p) \mid (x_i) \not\equiv 0 \pmod{p}, \\ &\quad \sum_{i=1}^{2k} (x_{2i-1}^2 - \varepsilon x_{2i}^2) \equiv pb \pmod{p^a}\}. \end{aligned}$$

Let $p = 2$. Then we have $K_2 = \mathbf{Q}_2(\sqrt{-3})$ and

$$\begin{aligned} \#\mathcal{B}_a(1_{2k}, B) &= \#\{(x_i) \in M_{4k,1}(\mathbf{Z}_2)/2^a M_{4k,1}(\mathbf{Z}_2) \mid (x_i) \not\equiv 0 \pmod{2}, \\ &\quad \sum_{i=1}^{2k} (x_{2i-1}^2 + x_{2i-1}x_{2i} + x_{2i}^2) \equiv 2b \pmod{2^a}\}. \end{aligned}$$

In any case, by Lemma 9 of [Kit84], we have

$$\#\mathcal{B}_a(1_{2k}, B) = p^{(4k-1)a} (1 - p^{-2k})(1 + p^{-2k+1}).$$

This proves the assertion (1.1). Similarly the assertion (1.2) holds.

(2) First let $m = 1$, and put $B = (b)$ with $b \in 2\mathbf{Z}_p$. Then $2^{-1}b \in \mathbf{Z}_p$. Let $p \neq 2$, or $p = 2$ and $f_2 = 3$. Then we have $K_p = \mathbf{Q}_p(\varpi)$ with ϖ a prime element of K_p such that $\bar{\varpi} = -\varpi$. Then an element $\mathbf{x} = (x_{2i-1} + \varpi x_{2i})_{1 \leq i \leq 2k}$ of $M_{2k,1}(\mathcal{O}_p)/p^a M_{2k,1}(\mathcal{O}_p)$ is primitive if and only if $(x_{2i-1})_{1 \leq i \leq 2k} \not\equiv 0 \pmod{p}$. Moreover we have

$$\Theta_{2k}[\mathbf{x}] = 2 \sum_{1 \leq i \leq 2k} (x_{2i}x_{2i+1} - x_{2i-1}x_{2i+2})\pi.$$

Hence we have

$$\begin{aligned} \#\mathcal{B}_a(1_{2k}, B) &= \#\{(x_i) \in M_{4k,1}(\mathbf{Z}_p)/p^a M_{4k,1}(\mathbf{Z}_p) \mid (x_{2i-1})_{1 \leq i \leq 2k} \not\equiv 0 \pmod{p} \\ &\quad \sum_{i=1}^{2k} (x_{2i}x_{2i+1} - x_{2i-1}x_{2i+2}) \equiv 2^{-1}b \pmod{p^a}\}. \end{aligned}$$

Let $p = 2$ and $f_2 = 2$. Then we have $K_2 = \mathbf{Q}_2(\varpi)$ with ϖ a prime element of K_2 such that $\eta := 2^{-1}(\varpi + \bar{\varpi}) \in \mathbf{Z}_2^*$. Then we have

$$\begin{aligned} \#\mathcal{B}_a(1_{2k}, B) &= \#\{(x_i) \in M_{4k,1}(\mathbf{Z}_2)/2^a M_{4k,1}(\mathbf{Z}_2) \mid (x_{2i-1})_{1 \leq i \leq 2k} \not\equiv 0 \pmod{2}, \\ &\quad \sum_{i=1}^{2k} \{\eta(x_{2i}x_{2i+1} + x_{2i-1}x_{2i+2}) + x_{2i-1}x_{2i+1} + \pi x_{2i}x_{2i+2}\} \equiv 2^{-1}b \pmod{2^a}\}. \end{aligned}$$

Thus, in any case, by a simple computation we have

$$\#\mathcal{B}_a(1_{2k}, B) = p^{(2k-1)a}(p^{2ka} - p^{2k(a-1)}).$$

Thus the assertion (2.1) has been proved for $m = 1$. Next let $\pi^{i_p}B = (b_{ij})_{1 \leq i, j \leq 2} \in \text{Her}_{2,*}(\mathcal{O}_p)$. Let $M = M_{\Theta_{2k}}$, $N_1 = M_{\pi^{i_p}b_{11}}$, and $N = M_B$. Let a be a sufficiently large positive integer. For each $\phi_1 \in \mathcal{B}'_a(N_1, M)$, put

$$\mathcal{B}'_a(N, M; \phi_1) = \{\phi \in \mathcal{B}'_a(N, M) \mid \phi|_{N_1} = \phi_1\}.$$

Let $N = \mathcal{O}_p v_1 \oplus \mathcal{O}_p v_2$, and put $u_1 = \phi_1(v_1)$. Then we can take elements $u_2, \dots, u_{2k} \in M$ such that

$$M = \mathcal{O}_p u_1 \oplus \mathcal{O}_p u_2 \oplus \dots \oplus \mathcal{O}_p u_{2k}$$

and

$$(u_1, u_2) = \varpi, (u_2, u_2) = 0, (u_i, u_j) = 0 \text{ for } i = 1, 2, j = 3, \dots, 2k, \text{ and } (u_i, u_j)_{3 \leq i, j \leq 2k} = \Theta_{2k-2}.$$

Then by the same argument as in the proof of Lemma 4.1.8, we can prove that

$$\begin{aligned} \mathcal{B}'_a(N, M; \phi_1) &= \{(x_i)_{1 \leq i \leq 2k-1} \in M_{2k-1,1}(\mathcal{O}_p)/p^a M_{2k-1,1}(\mathcal{O}_p) \mid (x_i)_{2 \leq i \leq 2k-2} \not\equiv 0 \pmod{\varpi}, \\ &\quad -x_1 \bar{x}_1 b_{11} - x_1 b_{12} - \bar{x}_1 \bar{b}_{12} + \Theta_{2k-2}[(x_i)_{2 \leq i \leq 2k-2}] \equiv b_{22} \pmod{p^a}\}. \end{aligned}$$

Hence by the assertion for $m = 1$, we have

$$\begin{aligned} \beta_p(\Theta_{2k}, B) &= \beta_p(\Theta_{2k}, b_{11})p^{-a} \sum_{x_1 \in \mathcal{O}_p/\varpi^a \mathcal{O}_p} \beta_p(\Theta_{2k-2}, b_{22} + b_{11}x_1 \bar{x}_1 + x_1 b_{12} + \bar{x}_1 \bar{b}_{12}) \\ &= (1 - p^{-2k})(1 - p^{-2k+2}). \end{aligned}$$

Thus the assertion (2.1) has been proved for $m = 2$. The assertion (2.2) can be proved by using the same argument as above. \square

Lemma 4.1.10. *Let k and m be integers with $k \geq m$.*

(1) *Suppose that K_p is unramified over \mathbf{Q}_p . Let $A \in \text{Her}_l(\mathcal{O}_p)$ and $B \in \text{Her}_m(\mathcal{O}_p)$. Then we have*

$$\beta_p(pA \perp 1_{2k}, pB) = \beta_p(1_{2k}, pB) = \prod_{i=0}^{2m-1} (1 - (-1)^i p^{-2k+i})$$

(2) *Let $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let l be an integer. Let $B \in \text{Her}_m(\mathcal{O}_p)$. Then we have*

$$\beta_p(1_{2k}, pB) = \prod_{i=0}^{2m-1} (1 - p^{-2k+i})$$

(3) *Suppose that K_p is ramified over \mathbf{Q}_p . Let $A \in \text{Her}_{l,*}(\mathcal{O}_p)$ and $B \in \text{Her}_{m,*}(\mathcal{O}_p)$. Then we have*

$$\beta_p(\pi^{i_p} A \perp \Theta_{2k}, \pi^{i_p} B) = \beta_p(\Theta_{2k}, \pi^{i_p} B) = \prod_{i=0}^{m-1} (1 - p^{-2k+2i}).$$

Proof. (1) Suppose that K_p is unramified over \mathbf{Q}_p . We prove the assertion by induction on m . Let $\deg B = 1$, and a be a sufficiently large integer. Then, by Lemma 4.1.9, we have

$$\beta_p(pA \perp 1_{2k}, pB) = p^{-al} \sum_{\mathbf{x} \in M_{11}(\mathcal{O}_p)/p^a M_{11}(\mathcal{O}_p)} \beta_p(1_{2k}, pB - pA[\mathbf{x}])$$

$$= (1 - p^{-2k})(1 + p^{-2k+1}).$$

This proves the assertion for $m = 1$. Let $m > 1$ and suppose that the assertion holds for $m - 1$. Then B can be expressed as $B \sim_{GL_m(\mathcal{O}_p)} B_1 \perp B_2$ with $B_1 \in \text{Her}_1(\mathcal{O}_p)$ and $B_2 \in \text{Her}_{m-1}(\mathcal{O}_p)$. Then by Lemma 4.1.8, we have

$$\beta_p(pA \perp 1_{2k}, pB_1 \perp pB_2) = \beta_p(pA \perp 1_{2k}, pB_1) \beta_p(pA \perp (-pB_1) \perp 1_{2k-2}, pB_2).$$

Thus the assertion holds by the induction assumption.

(2) Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then we easily see that we have

$$\beta_p(1_{2k}, pB) = p^{(-4km+m^2)} \# \mathcal{B}_1(1_{2k}, O_m).$$

We have

$$\begin{aligned} & \mathcal{B}_1(1_{2k}, O_m) \\ &= \{(X, Y) \in M_{2k,l}(\mathbf{Z}_p)/pM_{2k,l}(\mathbf{Z}_p) \oplus M_{2k,l}(\mathbf{Z}_p)/pM_{2k,l}(\mathbf{Z}_p) \mid \\ & \quad {}^t Y X \equiv O_m \pmod{pM_m(\mathbf{Z}_p)} \text{ and } \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} X = \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} Y = m\}. \end{aligned}$$

For each $X \in M_{2k,l}(\mathbf{Z}_p)/pM_{2k,l}(\mathbf{Z}_p)$ such that $\text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} X = m$, put

$$\# \mathcal{B}_1(1_{2k}, O_m; X)$$

$$= \{Y \in M_{2k,l}(\mathbf{Z}_p)/pM_{2k,l}(\mathbf{Z}_p) \mid {}^t Y X \equiv O_m \pmod{pM_m(\mathbf{Z}_p)} \text{ and } \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} Y = m\}.$$

By a simple computation we have

$$\#\{X \in M_{2k,l}(\mathbf{Z}_p)/pM_{2k,l}(\mathbf{Z}_p) \mid \text{rank}_{\mathbf{Z}_p/p\mathbf{Z}_p} X = m\} = \prod_{i=0}^{m-1} (p^{2k} - p^i),$$

and

$$\# \mathcal{B}_1(1_{2k}, O_m; X) = \prod_{i=0}^{m-1} (p^{2k-m} - p^i).$$

This proves the assertion.

(3) Suppose that K_p is ramified over \mathbf{Q}_p . We prove the assertion by induction on m . Let $\deg B = 1$, and a be a sufficiently large integer. Then, by Lemma 4.1.9, we have

$$\beta_p(\pi^{i_p} A \perp \Theta_{2k}, \pi^{i_p} B) = p^{-al} \sum_{\mathbf{x} \in M_{11}(\mathcal{O}_p)/p^a M_{11}(\mathcal{O}_p)} \beta_p(\Theta_{2k}, \pi^{i_p} B - \pi^{i_p} A[\mathbf{x}]) = 1 - p^{-2k}.$$

Let $\deg B = 2$. Then by Lemma 4.1.9, we have

$$\begin{aligned} \beta_p(\pi^{i_p} A \perp \Theta_{2k}, \pi^{i_p} B) &= p^{-2la} \sum_{\mathbf{x} \in M_{12}(\mathcal{O}_p)/p^a M_{12}(\mathcal{O}_p)} \beta_p(\Theta_{2k}, \pi^{i_p} B - \pi^{i_p} A[\mathbf{x}]) \\ &= (1 - p^{-2k})(1 - p^{-2k+2}). \end{aligned}$$

Let $m \geq 3$. Then B can be expressed as $B \sim_{GL_m(\mathcal{O}_p)} B_1 \perp B_2$ with $\deg B_1 \leq 2$. Then the assertion for m holds by Lemma 4.1.8, the induction hypothesis, and Lemma 4.1.9. \square

Lemma 4.1.11. (1) *Suppose that K_p is unramified over \mathbf{Q}_p . Let l and m be an integers with $l \geq m$. Then we have*

$$\alpha_p(1_l, 1_m) = \beta_p(1_l, 1_m) = \prod_{i=0}^{m-1} (1 - (-p)^{-l+i})$$

(2) Let $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let l and m be integers with $l \geq m$. Then we have

$$\alpha_p(1_l, 1_m) = \beta_p(1_l, 1_m) = \prod_{i=0}^{m-1} (1 - p^{-l+i})$$

(3) Suppose that K_p is ramified over \mathbf{Q}_p . Let k and m be even integers with $k \geq m$. Then we have

$$\alpha_p(\Theta_{2k}, \Theta_{2m}) = \beta_p(\Theta_{2k}, \Theta_{2m}) = \prod_{i=0}^{m-1} (1 - p^{-2k+2i}).$$

Proof. In any case, we easily see that the local density coincides with the primitive local density. Suppose that K_p is unramified over \mathbf{Q}_p . Then, by Lemma 4.1.7, we have

$$\alpha_p(1_l, 1_m) = \alpha_p(1_l, 1) \alpha_p(1_{l-1}, 1_{m-1}).$$

We easily see that we have

$$\alpha_p(1_l, 1) = 1 - (-1)^l p^{-l}.$$

This proves the assertion (1). Suppose that K_p is ramified over \mathbf{Q}_p . Then by Lemma 4.1.7, we have

$$\alpha_p(\Theta_{2k}, \Theta_m) = \alpha_p(\Theta_{2k}, \Theta_2) \alpha_p(\Theta_{2k-2}, \Theta_{2m-2}).$$

Moreover by Lemma 4.1.9, we have

$$\alpha_p(\Theta_{2k}, \Theta_2) = 1 - p^{-2k}.$$

This proves the assertion (3). Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then the assertion can be proved similarly to (2) of Lemma 4.1.10. \square

4.2. Primitive densities.

For an element $T \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$, we define a polynomial $G_p(T, X)$ in X by

$$G_p(T, X) = \sum_{i=0}^m \sum_{W \in GL_m(\mathcal{O}_p) \setminus \mathcal{D}_{m,i}} (Xp^m)^{\nu(\det W)} \Pi_p(W) F_p^{(0)}(T[W^{-1}], X).$$

Lemma 4.2.1. (1) Suppose that K_p is unramified over \mathbf{Q}_p . Let $B_1 \in \text{Her}_{m-n_0}(\mathcal{O}_p)$. Then we have

$$\alpha_p(1_{n_0} \perp pB_1) = \prod_{i=1}^{n_0} (1 - (-p)^{-i}) \alpha_p(pB_1)$$

(2) Let $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let $B_1 \in \text{Her}_{m-n_0}(\mathcal{O}_p)$. Then we have

$$\alpha_p(1_{n_0} \perp pB_1) = \prod_{i=1}^{n_0} (1 - p^{-i}) \alpha_p(pB_1)$$

(3) Suppose that K_p is ramified over \mathbf{Q}_p . Let n_0 be even. Let $B_1 \in \text{Her}_{m-n_0, *}(O_p)$. Then we have

$$\alpha_p(\Theta_{n_0} \perp \pi^{i_p} B_1) = \prod_{i=1}^{n_0/2} (1 - p^{-2i}) \alpha_p(\pi^{i_p} B_1).$$

Proof. Suppose that K_p is unramified over \mathbf{Q}_p . By Lemma 4.1.7, we have

$$\alpha_p(1_{n_0} \perp pB_1) = \alpha_p(1_{n_0} \perp pB_1, 1_{n_0}) \alpha_p(pB_1).$$

By using the same argument as in the proof of Lemma 4.1.10, we can prove that we have

$$\alpha_p(1_{n_0} \perp pB_1, 1_{n_0}) = \alpha_p(1_{n_0}),$$

and hence by Lemma 4.1.11, we have

$$\alpha_p(1_{n_0} \perp pB_1) = \prod_{i=1}^{n_0} (1 - (-p)^{-i}) \alpha_p(pB_1).$$

This proves the assertion (1). Similarly the assertions (2) and (3) can be proved. \square

Lemma 4.2.2. Let m be a positive integer and r a non-negative integer such that $r \leq m$.

(1) Suppose that K_p is unramified over \mathbf{Q}_p . Let $T = 1_{m-r} \perp pB_1$ with $B_1 \in \text{Her}_r(O_p)$. Then

$$\beta_p(1_{2k}, T) = \prod_{i=0}^{m+r-1} (1 - p^{-2k+i} (-1)^i).$$

(2) Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let $T = 1_{m-r} \perp pB_1$ with $B_1 \in \text{Her}_r(O_p)$. Then

$$\beta_p(1_{2k}, T) = \prod_{i=0}^{m+r-1} (1 - p^{-2k+i}).$$

(3) Suppose that K_p is ramified over \mathbf{Q}_p and that $m-r$ is even. Let $T = \Theta_{m-r} \perp \pi^{i_p} B_1$ with $B_1 \in \text{Her}_{r, *}(O_p)$. Then

$$\beta_p(\Theta_{2k}, T) = \prod_{i=0}^{(m+r-2)/2} (1 - p^{-2k+2i}).$$

Proof. Suppose that K_p is unramified over \mathbf{Q}_p . By Lemma 4.1.8, we have

$$\beta_p(1_{2k}, T) = \beta_p(1_{2k}, pB_1) \beta_p((-pB_1) \perp 1_{2k-2r}, 1_{m-r}).$$

By using the same argument as in the proof of Lemma 4.1.11, we can prove that we have $\beta_p((-pB_1) \perp 1_{2k-2r}, 1_{m-r}) = \beta_p(1_{2k-2r}, 1_{m-r})$. Hence the assertion follows from Lemmas 4.1.10 and 4.1.11. Similarly the assertions (2) and (3) can be proved. \square

Corollary. (1) Suppose that K_p is unramified over \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let $T = 1_{m-r} \perp pB_1$ with $B_1 \in \text{Her}_r(O_p)$. Then we have

$$G_p(T, Y) = \prod_{i=0}^{r-1} (1 - (\xi_p p)^{m+i} Y).$$

(2) Suppose that K_p is ramified over \mathbf{Q}_p and that $m - r$ is even. Let $T = \Theta_{m-r} \perp \pi^{i_p} B_1$ with $B_1 \in \text{Her}_{r,*}(\mathcal{O}_p)$. Then

$$G_p(T, Y) = \prod_{i=0}^{[(r-2)/2]} (1 - p^{2i+2[(m+1)/2]} Y).$$

Proof. Let k be a positive integer such that $k \geq m$. Put $\Xi_{2k} = \Theta_{2k}$ or 1_{2k} according as K_p is ramified over \mathbf{Q}_p or not. Then it follows from Lemma 14.8 of [Sh97] that we have

$$b_p(B, 2k) = \alpha_p(\Xi_{2k}, B).$$

Hence, by the definition of $G_p(T, X)$ and Corollary to Lemma 4.1.2, we have

$$\beta_p(\Xi_{2k}, T) = G_p(T, p^{-2k}) \prod_{i=0}^{[(m-1)/2]} (1 - p^{2i-2k}) \prod_{i=1}^{[m/2]} (1 - \xi_p p^{2i-1-2k}).$$

Suppose that K_p is unramified over \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then by Lemma 4.2.2, we have

$$G_p(T, p^{-2k}) = \prod_{i=0}^{r-1} (1 - (\xi_p p)^{m+i} p^{-2k}).$$

This equality holds for infinitely many positive integer k , and the both hand sides of it are polynomials in p^{-2k} . Thus the assertion (1) holds. Similarly the assertion (2) holds. \square

Lemma 4.2.3. *Let $B \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$. Then we have*

$$F_p^{(0)}(B, X) = \sum_{W \in GL_m(\mathcal{O}_p) \setminus \widetilde{\Omega}(B)} G_p(B[W^{-1}], X) (p^m X)^{\nu(\det W)}.$$

Proof. Let k be a positive integer such that $k \geq m$. By Lemma 4.1.2, we have

$$\alpha_p(\Xi_{2k}, B) = \sum_{W \in GL_m(\mathcal{O}_p) \setminus \widetilde{\Omega}(B)} \beta_p(\Xi_{2k}, B[W^{-1}]) p^{(-2k+m)\nu(\det W)}.$$

Then the assertion can be proved by using the same argument as in the proof of Corollary to Lemma 4.2.2. \square

Corollary. *Let $B \in \widetilde{\text{Her}}_m(\mathcal{O}_p)$. Then we have*

$$\begin{aligned} \widetilde{F}^{(0)}(B, X) &= X^{e_p m - f_p [m/2]} \sum_{B' \in \widetilde{\text{Her}}_m(\mathcal{O}_p) / GL_m(\mathcal{O}_p)} X^{-\text{ord}(\det B')} \frac{\alpha_p(B', B)}{\alpha_p(B')} \\ &\quad \times G_p(B', p^{-m} X^2) X^{\text{ord}(\det B) - \text{ord}(\det B')}. \end{aligned}$$

Proof. We have

$$\begin{aligned} \widetilde{F}^{(0)}(B, X) &= X^{e_p m - f_p [m/2]} X^{-\text{ord}(\det B)} F^{(0)}(B, p^{-m} X^2) \\ &= X^{e_p m - f_p [m/2]} \sum_{W \in GL_m(\mathcal{O}_p) \setminus \widetilde{\Omega}(B)} X^{-\text{ord}(\det B)} G_p(B[W^{-1}], p^{-m} X^2) (X^2)^{\nu(\det W)} \\ &= X^{e_p m - f_p [m/2]} \\ &\times \sum_{B' \in \widetilde{\text{Her}}_m(\mathcal{O}_p) / GL_m(\mathcal{O}_p)} \sum_{W \in GL_m(\mathcal{O}_p) \setminus \widetilde{\Omega}(B', B)} X^{-\text{ord}(\det B)} G_p(B', p^{-m} X^2) (X^2)^{\nu(\det W)} \end{aligned}$$

$$\begin{aligned}
&= X^{e_p m - f_p \lfloor m/2 \rfloor} \sum_{B' \in \widetilde{\text{Her}}_m(\mathcal{O}_p)/GL_m(\mathcal{O}_p)} X^{-\text{ord}(\det B')} \#(GL_m(\mathcal{O}_p) \backslash \widetilde{\Omega}(B', B)) \\
&\quad \times G_p(B', p^{-m} X^2) X^{\text{ord}(\det B) - \text{ord}(\det B')}.
\end{aligned}$$

Thus the assertion follows from (2) of Lemma 4.1.3. \square

Let

$$\mathcal{F}_{m,p,*}(d_0) = \bigcup_{i=0}^{\infty} (\widetilde{\text{Her}}_m(\pi^i d_0 N_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*), \mathcal{O}_p) \cap \text{Her}_{m,*}(\mathcal{O}_p)).$$

First suppose that K_p is unramified over \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Let H_m be a function on $\text{Her}_m(\mathcal{O}_p)^\times$ satisfying the following condition:

$$H_m(1_{m-r} \perp pB) = H_r(pB) \text{ for any } B \in \text{Her}_r(\mathcal{O}_p).$$

Let $d_0 \in \mathbf{Z}_p^*$. Then we put

$$Q(d_0, H_m, r, t) = \sum_{B \in p^{-1} \mathcal{F}_{r,p}(d_0) \cap \text{Her}_r(\mathcal{O}_p)} \frac{H_m(1_{m-r} \perp pB)}{\alpha_p(1_{m-r} \perp pB)} t^{\text{ord}(\det(pB))}.$$

Next suppose that K_p is ramified over \mathbf{Q}_p . Let H_m be a function on $\text{Her}_m(\mathcal{O}_p)^\times$ satisfying the following condition:

$$H_m(\Theta_{m-r} \perp \pi^{i_p} B) = H_r(\pi^{i_p} B) \text{ for any } B \in \text{Her}_{r,*}(\mathcal{O}_p) \text{ if } m-r \text{ is even.}$$

Let $d_0 \in \mathbf{Z}_p^*$ and $m-r$ be even. Then we put

$$Q(d_0, H_m, r, t) = \sum_{B \in \pi^{-i_p} \widetilde{\mathcal{F}}_{r,p}(d_0) \cap \text{Her}_{r,*}(\mathcal{O}_p)} \frac{H_m(\Theta_{m-r} \perp \pi^{i_p} B)}{\alpha_p(\Theta_{m-r} \perp \pi^{i_p} B)} t^{\text{ord}(\det(\pi^{i_p} B))}.$$

Then by Lemma 4.2.1 we easily obtain the following.

Proposition 4.2.4. (1) *Suppose that K_p is unramified over \mathbf{Q}_p or $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then for any $d_0 \in \mathbf{Z}_p^*$ and a non-negative integer r we have*

$$Q(d_0, H_m, r, t) = \frac{Q(d_0, H_r, r, t)}{\phi_{m-r}(\xi_p p^{-1})}.$$

(2) *Suppose that K_p is ramified over \mathbf{Q}_p . Then for any $d_0 \in \mathbf{Z}_p^*$ and a non-negative integer r such that $m-r$ is even, we have*

$$Q(d_0, H_m, r, t) = \frac{Q(d_0, H_r, r, t)}{\phi_{(m-r)/2}(p^{-2})}.$$

4.3. Explicit formulas of formal power series of Koecher-Maass type.

In this section we give an explicit formula for $P_m(d_0, X, t)$.

Theorem 4.3.1. *Let m be even, and $d_0 \in \mathbf{Z}_p^*$.*

(1) *Suppose that K_p is unramified over \mathbf{Q}_p . Then*

$$P_m(d_0, X, t) = \frac{1}{\phi_m(-p^{-1}) \prod_{i=1}^m (1 - t(-p)^{-i} X)(1 + t(-p)^{-i} X^{-1})}.$$

(2) *Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then*

$$P_m(d_0, X, t) = \frac{1}{\phi_m(p^{-1}) \prod_{i=1}^m (1 - tp^{-i} X)(1 - tp^{-i} X^{-1})}.$$

(3) *Suppose that K_p is ramified over \mathbf{Q}_p . Then*

$$P_m(d_0, X, t) = \frac{t^{mi_p/2}}{2\phi_{m/2}(p^{-2})} \times \left\{ \frac{1}{\prod_{i=1}^{m/2} (1 - tp^{-2i+1} X^{-1})(1 - tp^{-2i} X)} + \frac{\chi_{K_p}((-1)^{m/2} d_0)}{\prod_{i=1}^{m/2} (1 - tp^{-2i} X^{-1})(1 - tp^{-2i+1} X)} \right\}.$$

Theorem 4.3.2. *Let m be odd, and $d_0 \in \mathbf{Z}_p^*$.*

(1) *Suppose that K_p is unramified over \mathbf{Q}_p . Then*

$$P_m(d_0, X, t) = \frac{1}{\phi_m(-p^{-1}) \prod_{i=1}^m (1 + t(-p)^{-i} X)(1 + t(-p)^{-i} X^{-1})}.$$

(2) *Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then*

$$P_m(d_0, X, t) = \frac{1}{\phi_m(p^{-1}) \prod_{i=1}^m (1 - tp^{-i} X)(1 - tp^{-i} X^{-1})}.$$

(3) *Suppose that K_p is ramified over \mathbf{Q}_p . Then*

$$P_m(d_0, X, t) = \frac{t^{(m+1)i_p/2 + \delta_{2p}}}{2\phi_{(m-1)/2}(p^{-2}) \prod_{i=1}^{(m+1)/2} (1 - tp^{-2i+1} X)(1 - tp^{-2i+1} X^{-1})}.$$

To prove Theorems 4.3.1 and 4.3.2, put

$$K_m(d_0, X, t) = \sum_{B' \in \widetilde{\mathcal{F}}_{r,p}(d_0)} \frac{G_p(B', p^{-m} X^2)}{\alpha_p(B')} (tX^{-1})^{\text{ord}(\det B')}.$$

Proposition 4.3.3. *Let m and d_0 be as above. Then we have*

$$P_m(d_0, X, t) = X^{me_p - [m/2]f_p} K_m(d_0, X, t) \times \begin{cases} \prod_{i=1}^m (1 - t^2 X^2 p^{2i-2-2m})^{-1} & \text{if } K_p/\mathbf{Q}_p \text{ is unramified} \\ \prod_{i=1}^m (1 - tXp^{i-1-m})^{-2} & \text{if } K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p \\ \prod_{i=1}^m (1 - tXp^{i-1-m})^{-1} & \text{if } K_p/\mathbf{Q}_p \text{ is ramified.} \end{cases}$$

Proof. We note that B' belongs to $\widetilde{\text{Her}}_{m,p}(d_0)$ if B belongs to $\widetilde{\text{Her}}_{m-l,p}(d_0)$ and $\alpha_p(B', B) \neq 0$. Hence by Corollary to Lemma 4.2.3 we have

$$\begin{aligned} & P_m(d_0, X, t) \\ &= X^{me_p - [m/2]f_p} \sum_{B \in \widetilde{\mathcal{F}}_{m,p}(d_0)} \frac{1}{\alpha_p(B)} \sum_{B'} \frac{G_p(B', p^{-m} X^2) X^{-\text{ord}(B')} \alpha_p(B', B)}{\alpha_p(B')} \end{aligned}$$

$$\begin{aligned}
& \times X^{\text{ord}(\det B) - \text{ord}(\det B')} t^{\text{ord}(\det B)} \\
& = X^{me_p - [m/2]f_p} \sum_{B' \in \tilde{\mathcal{F}}_{m,p}(d_0)} \frac{G_p(B', p^{-m}X^2)}{\alpha_p(B')} (tX^{-1})^{\text{ord}(B')} \\
& \quad \times \sum_{B \in \tilde{\mathcal{F}}_{m,p}(d_0)} \frac{\alpha_p(B', B)}{\alpha_p(B)} (tX)^{\text{ord}(\det B) - \text{ord}(\det B')}.
\end{aligned}$$

Hence by using the same argument as in the proof of [[BS87], Theorem 5], and by (1) of Lemma 4.1.3, we have

$$\begin{aligned}
& \sum_{B \in \tilde{\mathcal{F}}_{m,p}(d_0)} \frac{\alpha_p(B', B)}{\alpha_p(B)} (tX)^{\text{ord}(\det B) - \text{ord}(\det B')} \\
& = \sum_{W \in M_m(\mathcal{O}_p)^\times / GL_m(\mathcal{O}_p)} (tXp^{-m})^{\nu(\det W)} \\
& = \begin{cases} \prod_{i=1}^m (1 - t^2 X^2 p^{2i-2-2m})^{-1} & \text{if } K_p/\mathbf{Q}_p \text{ is unramified} \\ \prod_{i=1}^m (1 - tXp^{i-1-m})^{-2} & \text{if } K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p \\ \prod_{i=1}^m (1 - tXp^{i-1-m})^{-1} & \text{if } K_p/\mathbf{Q}_p \text{ is ramified.} \end{cases}
\end{aligned}$$

Thus the assertion holds. \square

In order to prove Theorems 4.3.1 and 4.3.2, we introduce some notation. For a positive integer r and $d_0 \in \mathbf{Z}_p^\times$ let

$$\zeta_m(d_0, t) = \sum_{T \in \mathcal{F}_{m,p,*}(d_0)} \frac{1}{\alpha_p(T)} t^{\text{ord}(\det T)}.$$

We make the convention that $\zeta_0(d_0, t) = 1$ or 0 according as $d_0 \in \mathbf{Z}_p^*$ or not. To obtain an explicit formula of $\zeta_m(d_0, t)$ let $Z_m(u, d)$ be the integral defined as

$$Z_{m,*}(u, d) = \int_{\mathcal{F}_{m,p,*}(d_0)} |\det x|^{s-m} |dx|,$$

where $u = p^{-s}$, and $|dx|$ is the measure on $\text{Her}_m(K_p)$ so that the volume of $\text{Her}_m(\mathcal{O}_p)$ is 1. Then by Theorem 4.2 of [Sa97] we obtain:

Proposition 4.3.4. *Let $d_0 \in \mathbf{Z}_p^*$.*

(1) *Suppose that K_p is unramified over \mathbf{Q}_p . Then*

$$Z_{m,*}(u, d_0) = \frac{(p^{-1}, p^{-2})_{[(m+1)/2]} (-p^{-2}, p^{-2})_{[m/2]}}{\prod_{i=1}^m (1 - (-1)^{m+i} p^{i-1} u)}.$$

(2) *Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then*

$$Z_{m,*}(u, d_0) = \frac{\phi_m(p^{-1})}{\prod_{i=1}^m (1 - p^{i-1} u)}.$$

(3) *Suppose that K_p is ramified over \mathbf{Q}_p .*

(3.1) *Let $p \neq 2$. Then*

$$Z_{m,*}(u, d_0) = \frac{1}{2} (p^{-1}, p^{-2})_{[(m+1)/2]}$$

$$(3.2) \quad \times \begin{cases} \frac{1}{\prod_{i=1}^{(m+1)/2} (1-p^{2i-2}u)} & \text{if } m \text{ is odd,} \\ \left(\frac{1}{\prod_{i=1}^{m/2} (1-p^{2i-1}u)} + \frac{\chi_{K_p}((-1)^{m/2}d_0)p^{-m/2}}{\prod_{i=1}^{m/2} (1-p^{2i-2}u)} \right) & \text{if } m \text{ is even.} \end{cases}$$

Let $p = 2$ and $f_2 = 2$. Then

$$Z_{m,*}(u, d_0) = \frac{1}{2}(p^{-1}, p^{-2})_{[(m+1)/2]}$$

$$(3.3) \quad \times \begin{cases} \frac{u^{(m+1)/2}}{\prod_{i=1}^{(m+1)/2} (1-p^{2i-2}u)} & \text{if } m \text{ is odd,} \\ u^{m/2}p^{-m/2} \left(\frac{1}{\prod_{i=1}^{m/2} (1-p^{2i-1}u)} + \frac{\chi_{K_p}((-1)^{m/2}d_0)p^{-m/2}}{\prod_{i=1}^{m/2} (1-p^{2i-2}u)} \right) & \text{if } m \text{ is even.} \end{cases}$$

Let $p = 2$ and $f_2 = 3$. Then

$$Z_{m,*}(u, d_0) = \frac{1}{2}(p^{-1}, p^{-2})_{[(m+1)/2]}$$

$$\times \begin{cases} \frac{u}{\prod_{i=1}^{(m+1)/2} (1-p^{2i-2}u)} & \text{if } m \text{ is odd,} \\ p^{-m} \left(\frac{1}{\prod_{i=1}^{m/2} (1-p^{2i-1}u)} + \frac{\chi_{K_p}((-1)^{m/2}d_0)p^{-m/2}}{\prod_{i=1}^{m/2} (1-p^{2i-2}u)} \right) & \text{if } m \text{ is even.} \end{cases}$$

Proof. First suppose that K_p is unramified over \mathbf{Q}_p , $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$, or K_p is ramified over \mathbf{Q}_p and $p \neq 2$. Then $Z_{m,*}(u, d_0)$ coincides with $Z_m(u, d_0)$ in [[Sa97], Theorem 4.2]. Hence the assertion follows from (1) and (2) and the former half of (3) of [loc. cit]. Next suppose that $p = 2$ and $f_2 = 2$. Then $Z_{m,*}(u, d_0)$ is not treated in [loc. cit], but we can prove the assertion (3.2) using the same argument as in the proof of the latter half of (3) of [loc. cit]. Similarly we can prove (3.3) by using the same argument as in the proof of the former half of (3) of [loc. cit]. \square

Corollary. Let $d_0 \in \mathbf{Z}_p^*$.

(1) Suppose that K_p is unramified over \mathbf{Q}_p . Then

$$\zeta_m(d_0, t) = \frac{1}{\phi_m(-p^{-1})} \frac{1}{\prod_{i=1}^m (1 + (-1)^i p^{-i}t)}.$$

(2) Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then

$$\zeta_m(d_0, t) = \frac{1}{\phi_m(p^{-1})} \frac{1}{\prod_{i=1}^m (1 - p^{-i}t)}.$$

(3) Suppose that K_p is ramified over \mathbf{Q}_p .

(3.1) Let m be even. Then

$$\zeta_m(d_0, t) = \frac{p^{m(m+1)f_p/2 - m^2\delta_{2,p}} \kappa_p(t)}{2\phi_{m/2}(p^{-2})}$$

$$\times \left\{ \frac{1}{\prod_{i=1}^{m/2} (1 - p^{-2i+1}t)} + \frac{\chi_{K_p}((-1)^{m/2}d_0)p^{-i_p m/2}}{\prod_{i=1}^{m/2} (1 - p^{-2i}t)} \right\},$$

where $i_p = 0$, or 1 according as $p = 2$ and $f_p = 2$, or not, and

$$\kappa_p(t) = \begin{cases} 1 & \text{if } p \neq 2 \\ t^{m/2}p^{-m(m+1)/2} & \text{if } p = 2 \text{ and } f_2 = 2 \\ p^{-m} & \text{if } p = 2 \text{ and } f_2 = 3 \end{cases}$$

(3.2) Let m be odd. Then

$$\zeta_m(d_0, t) = \frac{p^{m(m+1)f_p/2-m^2\delta_{2,p}}\kappa_p(t)}{2\phi_{(m-1)/2}(p^{-2})} \frac{1}{\prod_{i=1}^{(m+1)/2}(1-p^{-2i+1}t)},$$

where

$$\kappa_p(t) = \begin{cases} 1 & \text{if } p \neq 2 \\ t^{(m+1)/2}p^{-m(m+1)/2} & \text{if } p = 2 \text{ and } f_2 = 2 \\ tp^{-m} & \text{if } p = 2 \text{ and } f_2 = 3 \end{cases}$$

Proof. First suppose that K_p is unramified over \mathbf{Q}_p . Then by a simple computation we have

$$\zeta_m(d_0, t) = \frac{Z_{m,*}(p^{-m}t, d_0)}{\phi_m(p^{-2})}.$$

Next suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then similarly to above

$$\zeta_m(d_0, t) = \frac{Z_{m,*}(p^{-m}t, d_0)}{\phi_m(p^{-1})^2}.$$

Finally suppose that K_p is ramified over \mathbf{Q}_p . Then by a simple computation and Lemma 3.1

$$\zeta_m(d_0, t) = \frac{p^{m(m+1)f_p/2-m^2\delta_{2,p}}Z_{m,*}(p^{-m}t, d_0)}{\phi_m(p^{-1})}.$$

Thus the assertions follow from Proposition 4.3.4. \square

Proposition 4.3.5. Let $d_0 \in \mathbf{Z}_p^*$.

(1) Suppose that K_p is unramified over \mathbf{Q}_p . Then

$$\begin{aligned} & K_m(d_0, X, t) \\ &= \sum_{r=0}^m \frac{p^{-r^2}(tX^{-1})^r \prod_{i=0}^{r-1}(1-(-1)^m(-p)^i X^2)}{\phi_{m-r}(-p^{-1})} \zeta_r(d_0, tX^{-1}). \end{aligned}$$

(2) Suppose that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then

$$\begin{aligned} & K_m(d_0, X, t) \\ &= \sum_{r=0}^m \frac{p^{-r^2}(tX^{-1})^r \prod_{i=0}^{r-1}(1-p^i X^2)}{\phi_{m-r}(p^{-1})} \zeta_r(d_0, tX^{-1}). \end{aligned}$$

(3) Suppose that K_p is ramified over \mathbf{Q}_p . Then

$$\begin{aligned} & K_m(d_0, X, t) \\ &= \sum_{r=0}^{m/2} \frac{p^{-4i_p r^2}(tX^{-1})^{(m/2+r)i_p} \prod_{i=0}^{r-1}(1-p^{2i} X^2)}{\phi_{(m-2r)/2}(p^{-2})} \zeta_{2r}((-1)^{m/2-r} d_0, tX^{-1}) \end{aligned}$$

if m is even, and

$$\begin{aligned} & K_m(d_0, X, t) \\ &= \sum_{r=0}^{(m-1)/2} \frac{p^{-(2r+1)^2 i_p} (tX^{-1})^{((m+1)/2+r)i_p} \prod_{i=0}^{r-1}(1-p^{2i+1} X^2)}{\phi_{(m-2r-1)/2}(p^{-2})} \zeta_{2r+1}((-1)^{(m-2r-1)/2} d_0, tX^{-1}) \end{aligned}$$

if m is odd.

Proof. The assertions can be proved by using Corollary to Lemma 4.2.2 and Proposition 4.2.4 (cf. [[IK06], Proposition 3.1]). \square

It is well known that $\#(\mathbf{Z}_p^*/N_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*)) = 2$ if K_p/\mathbf{Q}_p is ramified. Hence we can take a complete set \mathcal{N}_p of representatives of $\mathbf{Z}_p^*/N_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*)$ so that $\mathcal{N}_p = \{1, \xi_0\}$ with $\chi_{K_p}(\xi_0) = -1$.

Proof of Theorem 4.3.1. (1) By Corollary to Proposition 4.3.4 and Proposition 4.3.5, we have

$$K_m(d_0, X, t) = \frac{1}{\phi_m(-p^{-1})} \frac{L_m(d_0, X, t)}{\prod_{i=1}^m (1 + (-1)^i p^{-i} X^{-1} t)},$$

where $L_m(d_0, X, t)$ is a polynomial in t of degree m . Hence

$$P_m(d_0, X, t) = \frac{1}{\phi_m(-p^{-1})} \frac{L_m(d_0, X, t)}{\prod_{i=1}^m (1 + (-1)^i p^{-i} X^{-1} t) \prod_{i=1}^m (1 - p^{-2i} X^2 t^2)}.$$

We have

$$\tilde{F}(B, -X^{-1}) = \tilde{F}(B, X)$$

for any $B \in \tilde{F}_p^{(0)}(B, X)$. Hence we have

$$P_m(d_0, -X^{-1}, t) = P_m(d_0, X, t),$$

and therefore the denominator of the rational function $P_m(d_0, X, t)$ in t is at most

$$\prod_{i=1}^m (1 + (-1)^i p^{-i} X^{-1} t) \prod_{i=1}^m (1 - (-1)^i p^{-i} X t).$$

Thus

$$P_m(d_0, X, t) = \frac{a}{\phi_m(-p^{-1}) \prod_{i=1}^m (1 + (-1)^i p^{-i} X^{-1} t) \prod_{i=1}^m (1 - (-1)^i p^{-i} X t)},$$

with some constant a . It is easily seen that we have $a = 1$. This proves the assertion.

(2) The assertion can be proved by using the same argument as above.

(3) By Corollary to Proposition 4.3.4 and Proposition 4.3.5, we have

$$\begin{aligned} & K_m(d, X, t) \\ &= \frac{1}{2} \left\{ \frac{L^{(0)}(X, t)}{\prod_{i=1}^{m/2} (1 - p^{-2i+1} X^{-1} t)} + \frac{\chi_{K_p}((-1)^{m/2} d_0) L^{(1)}(X, t)}{\prod_{i=1}^{m/2} (1 - p^{-2i} X^{-1} t)} \right\} \end{aligned}$$

with some polynomials $L^{(0)}(X, t)$ and $L^{(1)}(X, t)$ in t of degrees at most m . Thus we have

$$P_m(d, X, t) = \frac{1}{2} \left\{ \frac{L^{(0)}(X, t)}{\prod_{i=1}^{m/2} (1 - p^{-2i+1} X^{-1} t) \prod_{i=1}^m (1 - p^{-i} X t)} + \frac{\chi_{K_p}((-1)^{m/2} d_0) L^{(1)}(X, t)}{\prod_{i=1}^{m/2} (1 - p^{-2i} X^{-1} t) \prod_{i=1}^m (1 - p^{-i} X t)} \right\}.$$

For $l = 0, 1$ put

$$P_m^{(l)}(X, t) = \frac{1}{2} \sum_{d \in \mathcal{N}_p} \chi_{K_p}((-1)^{m/2} d)^l P_m(d, X, t).$$

Then

$$P_m^{(0)}(X, t) = \frac{L^{(0)}(X, t)}{2\phi_{m/2}(p^{-2})} \frac{1}{\prod_{i=1}^{m/2} (1 - p^{-2i+1} X^{-1} t) \prod_{i=1}^m (1 - p^{-i} X t)},$$

and

$$P_m^{(1)}(X, t) = \frac{L^{(1)}(X, t)}{2\phi_{m/2}(p^{-2})} \frac{1}{\prod_{i=1}^{m/2} (1 - p^{-2i} X^{-1} t) \prod_{i=1}^m (1 - p^{-i} X t)}.$$

Then by the functional equation of Siegel series we have

$$P_m(d, X^{-1}, t) = \chi_{K_p}((-1)^{m/2}d)P_m(d, X, t)$$

for any $d \in \mathcal{N}_p$. Hence we have

$$P_m^{(0)}(X^{-1}, t) = P_m^{(1)}(X, t).$$

Hence the reduced denominator of the rational function $P_m^{(0)}(X, t)$ in t is at most

$$\prod_{i=1}^{m/2} (1 - p^{-2i+1}X^{-1}t) \prod_{i=1}^{m/2} (1 - p^{-2i}Xt),$$

and similarly to (1) we have

$$P_m^{(0)}(X, t) = \frac{1}{2\phi_{m/2}(p^{-2}) \prod_{i=1}^{m/2} (1 - p^{-2i+1}X^{-1}t) \prod_{i=1}^{m/2} (1 - p^{-2i}Xt)}.$$

Similarly

$$P_m^{(1)}(X, t) = \frac{1}{2\phi_{m/2}(p^{-2}) \prod_{i=1}^{m/2} (1 - p^{-2i}X^{-1}t) \prod_{i=1}^{m/2} (1 - p^{-2i+1}Xt)}.$$

We have

$$P_m(d_0, X, t) = P_m^{(0)}(X, t) + \chi_{K_p}((-1)^{m/2}d_0)P_m^{(1)}(X, t).$$

This proves the assertion. \square

Proof of Theorem 4.3.2. The assertion can also be proved by using the same argument as above. \square

Theorem 4.3.6. Let $d_0 \in \mathbf{Z}_p^*$.

(1) Suppose that K_p is unramified over \mathbf{Q}_p or that $K_p = \mathbf{Q}_p \oplus \mathbf{Q}_p$. Then

$$\hat{P}_m(d_0, X, t) = P_m(d_0, X, t)$$

for any $m > 0$.

(2) Suppose that K_p is ramified over \mathbf{Q}_p . Then

$$\hat{P}_{2n+1}(d_0, X, t) = P_{2n+1}(d_0, X, t)$$

and

$$\hat{P}_{2n}(d_0, X, t) = \frac{1}{2\phi_n(p^{-2})} \times \left\{ \frac{t^{ni_p}}{\prod_{i=1}^n (1 - tp^{-2i+1}X^{-1})(1 - tp^{-2i}X)} + \frac{\chi_{K_p}((-1)^n d_0)(t\chi_{K_p}(p))^{ni_p}}{\prod_{i=1}^n (1 - tp^{-2i}\chi_{K_p}(p)X^{-1})(1 - tp^{-2i+1}\chi_{K_p}(p)X)} \right\}.$$

Proof. The assertion (1) is clear from the definition. We note that $P_m(d_0, X, t)$ does not depend on the choice of π . Suppose that K_p is ramified over \mathbf{Q}_p . If $m = 2n + 1$, then it follows from (3) of Theorem 4.3.2 that we have

$$\lambda_{m,p}^*(\pi^i d, X) = \lambda_{m,p}^*(\pi^i, X)$$

for any $d \in \mathbf{Z}_p^*$, and in particular we have

$$\lambda_{m,p}^*(p^i d_0, X) = \lambda_{m,p}^*(\pi^i, X).$$

This proves the assertion. Suppose that $m = 2n$. Write $\hat{P}_{2n}(d_0, X, t)$ as

$$\hat{P}_{2n}(d_0, X, t) = \hat{P}_{2n}(d_0, X, t)_{\text{even}} + \hat{P}_{2n}(d_0, X, t)_{\text{odd}},$$

where

$$\hat{P}_{2n}(d_0, X, t)_{\text{even}} = \frac{1}{2} \{ \hat{P}_{2n}(d_0, X, t) + \hat{P}_{2n}(d_0, X, -t) \},$$

and

$$\hat{P}_{2n}(d_0, X, t)_{\text{odd}} = \frac{1}{2} \{ \hat{P}_{2n}(d_0, X, t) - \hat{P}_{2n}(d_0, X, -t) \}.$$

We have

$$\hat{P}_{2n}(d_0, X, t)_{\text{even}} = \sum_{i=0}^{\infty} \lambda_{2n,p}^* (p^{2i} d_0, X, Y) t^{2i} = \sum_{i=0}^{\infty} \lambda_{2n,p}^* (\pi^{2i} d_0, X, Y) t^{2i}$$

and

$$\hat{P}_{2n}(d_0, X, t)_{\text{odd}} = \sum_{i=0}^{\infty} \lambda_{2n,p}^* (p^{2i+1} d_0, X) t^{2i+1} = \sum_{i=0}^{\infty} \lambda_{2n,p}^* (\pi^{2i+1} d_0 \pi p^{-1}, X) t^{2i+1}.$$

Hence we have

$$\hat{P}_{2n}(d_0, X, t)_{\text{even}} = \frac{1}{2} \{ P_{2n}(d_0, X, t) + P_{2n}(d_0, X, -t) \},$$

and

$$\hat{P}_{2n}(d_0, X, Y, t)_{\text{odd}} = \frac{1}{2} \{ P_{2n}(d_0 \pi p^{-1}, X, t) - P_{2n}(d_0 \pi p^{-1}, X, -t) \},$$

and hence we have

$$\begin{aligned} \hat{P}_{2n}(d_0, X, t) &= P_{2n}^{(0)}(d_0, X, t) + \frac{1}{2} (1 + \chi_{K_p}(\pi p^{-1})) \chi_{K_p}((-1)^n d_0) P_{2n}^{(1)}(d_0, X, t) \\ &\quad + \frac{1}{2} (1 - \chi_{K_p}(\pi p^{-1})) \chi_{K_p}((-1)^n d_0) P_{2n}^{(1)}(d_0, X, -t). \end{aligned}$$

Assume that $\chi_{K_p}(\pi p^{-1}) = 1$. Then $\chi(d_0 \pi p^{-1}) = \chi(d_0)$, and we have

$$\hat{P}_{2n}(d_0, X, t) = P_{2n}(d_0, X, t).$$

Suppose that $\chi_{K_p}(\pi p^{-1}) = -1$. Then $\chi(d_0 \pi p^{-1}) = -\chi(d_0)$, and we have

$$\hat{P}_{2n}(d_0, X, t) = P_{2n}^{(0)}(d_0, X, t) + \chi_{K_p}((-1)^n d_0) P_{2n}^{(1)}(d_0, X, -t)$$

Since $\pi \in N_{K_p/\mathbf{Q}_p}(K_p^\times)$, we have $\chi_{K_p}(\pi p^{-1}) = \chi_{K_p}(p)$. This proves the assertion. \square

Corollary. *Let $m = 2n$ be even. Suppose that K_p is ramified over \mathbf{Q}_p . For $l = 0, 1$ put*

$$\hat{P}_{2n}^{(l)}(X, t) = \frac{1}{2} \sum_{d \in \mathcal{N}_p} \chi_{K_p}((-1)^n d)^l \hat{P}_{2n}(d, X, t).$$

Then we have

$$\hat{P}_{2n}(d, X, t) = \frac{1}{2} (\hat{P}_{2n}^{(0)}(X, t) + \chi_{K_p}((-1)^n d) \hat{P}_{2n}^{(1)}(X, t)),$$

and

$$\hat{P}_{2n}^{(0)}(X, t) = P_{2n}^{(0)}(X, t),$$

and

$$\hat{P}_{2n}^{(1)}(X, t) = P_{2n}^{(1)}(X, \chi_{K_p}(p)t),$$

The following result will be used to prove Theorems 2.3 and 2.4.

Proposition 4.3.7. *Let $d \in \mathbf{Z}_p^\times$. Then we have*

$$\lambda_{m,p}^*(d, X) = u_p \lambda_{m,p}(d, X).$$

Proof. Let I be the left-hand side of the above equation. Let

$$GL_m(\mathcal{O}_p)_1 = \{U \in GL_m(\mathcal{O}_p) \mid \overline{\det U} \det U = 1\}.$$

Then there exists a bijection from $\widetilde{\text{Her}}_m(d, \mathcal{O}_p)/GL_m(\mathcal{O}_p)_1$ to $\widetilde{\text{Her}}_m(dN_{K_p/\mathbf{Q}_p}(\mathcal{O}_p^*), \mathcal{O}_p)/GL_m(\mathcal{O}_p)$. Hence

$$I = \sum_{A \in \widetilde{\text{Her}}_m(d, \mathcal{O}_p)/GL_m(\mathcal{O}_p)_1} \frac{\widetilde{F}_p^{(0)}(A, X)}{\alpha_p(A)}.$$

Now for $T \in \widetilde{\text{Her}}_m(d, \mathcal{O}_p)$, let l be the number of $SL_m(\mathcal{O}_p)$ -equivalence classes in $\widetilde{\text{Her}}_m(d, \mathcal{O}_p)$ which are $GL_m(\mathcal{O}_p)$ -equivalent to T . Then it can easily be shown that $l = l_{p,T}$. Hence the assertion holds. \square

5. PROOF OF THE MAIN THEOREM

Proof of Theorem 2.3. For a while put $\lambda_p^*(d) = \lambda_{m,p}^*(d, \alpha_p^{-1})$. Then by Theorem 3.4 and Proposition 4.3.7, we have

$$L(s, I_{2n}(f)) = \mu_{2n,k,D} \sum_d \prod_p (u_p^{-1} \lambda_p^*(d)) d^{-s+k+2n}.$$

Then by (1) and (2) of Theorem 4.3.1, and (1) of Theorem 4.3.6, $\lambda_p^*(d)$ depends only on $p^{\text{ord}_p(d)}$ if $p \nmid D$. Hence we write $\lambda_p^*(d)$ as $\widetilde{\lambda}_p(p^{\text{ord}_p(d)})$. On the other hand, if $p \mid D$, by (3) of Theorem 4.3. and (2) of Theorem 4.3.6, $\lambda_p^*(d)$ can be expressed as

$$\lambda_p^*(d) = \lambda_p^{(0)}(d) + \chi_{K_p}((-1)^n d p^{-\text{ord}_p(d)}) \lambda_p^{(1)}(d),$$

where $\lambda_p^{(l)}(d)$ is a rational number depending only on $p^{\text{ord}_p(d)}$ for $l = 0, 1$. Hence we write $\lambda_p^{(l)}(d)$ as $\widetilde{\lambda}_p^{(l)}(p^{\text{ord}_p(d)})$. Then we have

$$\begin{aligned} b_m(f; d) &= \sum_{Q \subset Q_D} \prod_{p \mid d, p \nmid D} \left(u_p^{-1} \widetilde{\lambda}_p(p^{\text{ord}_p(d)}) \prod_{q \in Q} \chi_{K_q}(p^{\text{ord}_p(d)}) \right) \\ &\quad \times \prod_{p \mid d, p \mid D, p \neq q} \left(u_p^{-1} \widetilde{\lambda}_p^{(0)}(p^{\text{ord}_p(d)}) \prod_{q \in Q} \chi_{K_q}(p^{\text{ord}_p(d)}) \right) \\ &\quad \times \prod_{p \mid d, p \in Q} \left(u_p^{-1} \widetilde{\lambda}_p^{(1)}(p^{\text{ord}_p(d)}) \prod_{q \in Q, q \neq p} \chi_{K_q}(p^{\text{ord}_p(d)}) \right) \prod_{q \in Q} \chi_{K_q}((-1)^n) \end{aligned}$$

for a positive integer d . We note that for a subset Q of Q_D we have

$$\chi_Q(m) = \prod_{q \in Q} \chi_{K_q}(m)$$

for an integer m coprime to any $q \in Q$, and

$$\chi'_Q(p) = \chi_{K_p}(p) \prod_{q \in Q, q \neq p} \chi_{K_q}(p)$$

for any $p \in Q$. Hence, by Theorems 4.3.1 and 4.3.6, and Corollary to Theorem 4.3.6, we have

$$\begin{aligned}
L(s, I_{2n}(f)) &= \mu_{2n,k,D} \sum_{Q \subset Q_D} \prod_{p \nmid D} \sum_{i=0}^{\infty} u_p^{-1} \tilde{\lambda}_p(p^i) \chi_Q(p^i) p^{(-s+k+2n)i} \\
&\quad \times \prod_{p|D, p \notin Q} \sum_{i=0}^{\infty} u_p^{-1} \tilde{\lambda}_p^{(0)}(p^i) \chi_Q(p^i) p^{(-s+k+2n)i} \chi_Q((-1)^n) \\
&\quad \times \prod_{p \in Q} \sum_{i=0}^{\infty} u_p^{-1} \tilde{\lambda}_p^{(1)}(p^i) \left(\prod_{q \in Q, q \neq p} \chi_{K_q}(p^i) \right) p^{(-s+k+2n)i}. \\
&= \mu_{2n,k,D} \sum_{Q \subset Q_D} \chi_Q((-1)^n) \prod_{p \nmid D} (u_p^{-1} P_{2n,p}(1, \alpha_p^{-1}, \chi_Q(p) p^{-s+k+2n})) \\
&\quad \times \prod_{p|D, p \notin Q} (u_p^{-1} P_{2n,p}^{(0)}(\alpha_p^{-1}, \chi_Q(p) p^{-s+k+2n})) \prod_{p \in Q} (u_p^{-1} P_{2n,p}^{(1)}(\alpha_p^{-1}, \chi'_Q(p) p^{-s+k+2n})).
\end{aligned}$$

Now for $l = 0, 1$ write $P_{2n,p}^{(l)}(X, t)$ as

$$P_{2n,p}^{(l)}(X, t) = t^{ni_p} \tilde{P}_{2n,p}^{(l)}(X, t),$$

where $i_p = 0$ or 1 according as $4 \nmid D$ or not. Notice that $u_p = (1 - \chi(p)p^{-1})^{-1}$ if $p \nmid D$ and $u_p = 2^{-1}$ if $p|D$. Hence we have

$$\begin{aligned}
L(s, I_{2n}(f)) &= \mu_{2n,k,D} \sum_{Q \subset Q_D} \chi_Q((-1)^n) \\
&\quad \times \prod_{p \in Q'_D} p^{(-s+k+2n)n} \left(\prod_{p \in Q'_D, p \notin Q} \chi_Q(p) \prod_{p \in Q'_D \cap Q} \chi'_Q(p) \right)^n \\
&\quad \times \prod_{p \nmid D} ((1 - \chi(p)p^{-1}) P_{2n,p}(1, \alpha_p^{-1}, \chi_Q(p) p^{-s+k+2n})) \\
&\quad \times \prod_{p|D, p \notin Q} (2\tilde{P}_{2n,p}^{(0)}(\alpha_p^{-1}, \chi_Q(p) p^{-s+k+2n})) \prod_{p \in Q} (2\tilde{P}_{2n,p}^{(1)}(\alpha_p^{-1}, \chi'_Q(p) p^{-s+k+2n})),
\end{aligned}$$

where $Q'_D = Q_D \setminus \{2\}$ or Q_D according as $4 \nmid D$ or not. Note that

$$2^{c_D n(-s+k+2n)} \prod_{p \in Q'_D} p^{(-s+k+2n)n} = D^{(-s+k+2n)n},$$

and

$$\prod_{p \in Q'_D, p \notin Q} \chi_Q(p) \prod_{p \in Q'_D \cap Q} \chi'_Q(p) = 1.$$

Thus the assertion follows from Theorem 4.3.1.

Proof of Theorem 2.4. The assertion follows directly from Theorems 3.4 and 4.3.2.

REFERENCES

- [BS87] S. Böcherer and F. Sato, *Rationality of certain formal power series related to local densities*, Comment. Math. Univ. St. Paul. **36** (1987), 53–86.
- [Gri90] V. A. Gritsenko, *The Maass space for $SU(2, 2)$. The Hecke ring, and zeta functions* (Russian), Translated in Proc. Steklov Inst. Math. 1991, no. 4, 75–86. *Galois theory, rings, algebraic groups and their applications* (Russian), Trudy Mat. Inst. Steklov. 183 (1990), 68–78, 223–225.
- [IK04] T. Ibukiyama and H. Katsurada, *An explicit formula for Koecher-Maaß Dirichlet series for the Ikeda lifting*, Abh. Math. Sem. Hamburg **74** (2004), 101–121.
- [IK06] ———, *Koecher-Maaß series for real analytic Siegel Eisenstein series*, Automorphic forms and zeta functions, 170–197, World Sci. Publ., Hackensack, NJ, 2006.
- [IS95] T. Ibukiyama and H. Saito, *On zeta functions associated to symmetric matrices. I. An explicit form of zeta functions*, Amer. J. Math. **117** (1995), 1097–1155.
- [Ike01] T. Ikeda, *On the lifting of elliptic modular forms to Siegel cusp forms of degree $2n$* , Ann. of Math. **154** (2001), no. 3, 641–681.
- [Ike08] ———, *On the lifting of hermitian modular forms*, Compositio Math. **144**(2008) 1107–1154.
- [Jac62] R. Jacobowitz, *Hermitian forms over local fields*, Amer. J. Math. **84**(1962), 441–465.
- [Kat13] H. Katsurada, *On the period of the Ikeda lift for $U(m, m)$* , Preprint 2013.
- [Kat14] ———, *Koecher-Maass series of the adelic Eisenstein series and the adelic Ikeda lift for $U(m, m)$* , Preprint 2014.
- [Kit83] Y. Kitaoka, *A note on local densities of quadratic forms*, Nagoya Math. J. **92**(1983), 145–152.
- [Kit84] ———, *Dirichlet series in the theory of Siegel modular forms*, Nagoya Math. J. **95** (1984), 73–84.
- [Kit93] ———, *Arithmetic of quadratic forms*, Cambridge Tracts in Mathematics, 106. Cambridge University Press, Cambridge, 1993
- [Koj82] H. Kojima, *An arithmetic of Hermitian modular forms of degree two*, Invent. Math. **69**(1982), 217–227.
- [Kri91] A. Krieg, *The Maass spaces on the Hermitian half-space of degree 2*, Math. Ann. **289**(1991), 663–681.
- [Mi06] Y. Mizuno, *An explicit formula of the Koecher-Maass series associated with hermitian modular forms belonging to the Maass space*, Manuscripta Math. **119**(2006), 159–181.
- [Re71] U. Rehmann, *Klassenzahlen einiger totaldefiniter klassischer Gruppen ueber Zahlkoerpern*, Dissertation, Göttingen, 1971
- [Sa97] H. Saito, *Explicit formula of orbital p -adic zeta functions associated to symmetric and Hermitian matrices*, Comment. Math. Univ. St. Pauli **46** (1997), 175–216.
- [Sch85] W. Scharlau, *Quadratic forms and Hermitian forms*, Springer -Verlag, Berlin Heidelberg New York Tokyo, 1985.
- [Sh97] G. Shimura, *Euler products and Eisenstein series*, CBMS Regional Conference Series in Math. 93(1997), Amer. Math. Soc.
- [Sh00] ———, *Arithmeticity in the theory of automorphic forms*, Mathematical Surveys and Monographs, 82, Amer. Math. Soc., 2000.
- [Su95] T. Sugano, *Jacobi forms and the theta lifting*, Comment Math. Univ. St. Pauli **44**(1995), 1–58.
- [We82] A. Weil, *Adeles and algebraic groups*, Birkhäuser, Boston, 1982.

Hidenori KATSURADA

Muroran Institute of Technology 27-1 Mizumoto, Muroran, 050-8585, Japan

E-mail: hidenori@mmm.muroran-it.ac.jp