



Title	CONTACT GEOMETRY OF SECOND ORDER I
Author(s)	Yamaguchi,K.
Citation	Hokkaido University Preprint Series in Mathematics, 927, 1-39
Issue Date	2008-10-8
DOI	10.14943/84421
Doc URL	http://hdl.handle.net/2115/70235
Type	bulletin (article)
File Information	pre927.pdf



[Instructions for use](#)

CONTACT GEOMETRY OF SECOND ORDER I

KEIZO YAMAGUCHI

1. INTRODUCTION

In [C1] and [C2], E.Cartan studied involutive systems of second order partial differential equations for a scalar function with 2 or 3 independent variables, following the tradition of the geometric theory of partial differential equations developed by Monge, Jacobi, Lie, Darboux, Goursat and others. In fact he investigated the contact equivalence and the integration problems of such involutive systems of second order. In this course, he found out the link between the contact equivalence of involutive systems of second order and the geometry of differential systems (Pffafian systems) on five dimensional spaces.

The main purpose of the present paper is to reformulate his study as the **Contact Geometry of Second Order**. As is well known, the classical theory of systems of the first order partial differential equations for a scalar function can be rephrased as the submanifold theory of contact manifolds. In this spirit, we formulate the submanifold theory of second order contact manifolds as the geometry of PD manifolds of second order ([Y1], [Y6] see §4).

By Bäcklund Theorem (see §2.2), the symbols of second order equations become the first invariants under contact transformations. In fact, in [C2], E.Cartan first classified involutive symbols algebraically and wrote the structure equations of such involutive systems of second order with 3 independent variables. To capture good classes of second order equations, we cannot pursue this line in general (see the discussion in §3.3). Our guiding principle in this paper is to utilize Parabolic Geometries, directly or combined with the reduction procedures, to find good classes of PD manifolds of second order. Here the **Parabolic Geometry** is a geometry modeled after the homogeneous space G/G' , where G is a (semi-)simple Lie group and G' is a parabolic subgroup of G (cf. [Bai]). Precisely, in this paper, we mean, by a Parabolic Geometry, the **Geometry associated with the Simple Graded Lie Algebra** in the sense of N.Tanaka ([T4]). As for the reduction procedures, we will establish the First Reduction Theorem for PD manifolds admitting non-trivial Chauchy characteristic systems in §4 and will treat the second reduction procedures (two step reductions) as Part II in the sequel to this paper.

Now let us proceed to describe the contents of each sections. In §2, we will recall the geometric (Grassmannian) construction of Jet spaces and prepare basics of differential systems, especially the Tanaka Theory of (linear) differential systems. In §3, we will discuss the symbols of second order equations as the first invariants in the Contact Geometry of Second Order. We will formulate the submanifold theory of second order contact manifolds in §4 as the geometry of PD manifolds of second order and establish the First Reduction Theorem. In §5, we will first prepare the notation for the simple graded Lie algebras and state the Prolongation Theorem. Then we will exhibit Parabolic Geometries, which directly correspond to the geometry of PD manifolds of second order. Finally, in §6, we will discuss one of the Typical classes of involutive systems of second order as the application of the First Reduction Theorem and will show several examples

of Parabolic Geometries, which are linked to the geometry of PD manifolds of second order through the First Reduction Theorem.

This paper constitutes the extended version of our previous paper [Y6].

2. GEOMETRY OF JET SPACES.

We will recall the geometric construction of Jet spaces and fix our notations for the basic notion for differential systems, following [Y5] and [Y7].

2.1. Spaces of Contact Elements. Let us start with the construction of the space $J(M, n)$ of contact elements to M : Let M be a (real or complex) manifold of dimension $m + n$. Fixing the number n , we consider the space of n -dimensional **contact elements** to M , i.e., the **Grassmannian bundle** over M consisting of all n -dimensional contact elements to M ;

$$J(M, n) = \bigcup_{x \in M} J_x \xrightarrow{\pi} M,$$

where $J_x = \text{Gr}(T_x(M), n)$ is the Grassmann manifold of all n -dimensional subspaces of the tangent space $T_x(M)$ to M at x . Each element $u \in J(M, n)$ is a linear subspace of $T_x(M)$ of codimension m , where $x = \pi(u)$. Hence we have a differential system C of codimension m on $J(M, n)$ by putting:

$$C(u) = \pi_*^{-1}(u) \subset T_u(J(M, n)) \xrightarrow{\pi_*} T_x(M).$$

for each $u \in J(M, n)$. C is called the **Canonical System** on $J(M, n)$. Introducing the *inhomogeneous Grassmann coordinate* $(x_1, \dots, x_n, z^1, \dots, z^m, p_1^1, \dots, p_n^m)$ of $J(M, n)$ around $u_o \in J(M, n)$, C is defined by;

$$C = \{\varpi^1 = \dots = \varpi^m = 0\},$$

where

$$\varpi^\alpha = dz^\alpha - \sum_{i=1}^n p_i^\alpha dx_i, \quad (\alpha = 1, \dots, m).$$

Here $(x_1, \dots, x_n, z^1, \dots, z^m)$ is a coordinate system of M around $x_o = \pi(u_o)$ such that $dx_1 \wedge \dots \wedge dx_n|_{u_o} \neq 0$. Coordinate functions p_i^α are introduced by

$$dz^\alpha|_u = \sum_{i=1}^n p_i^\alpha(u) dx_i|_u.$$

$(J(M, n), C)$ is the (geometric) 1-jet space for n -dimensional submanifolds in M . Let M, \hat{M} be manifolds (of dimension $m + n$) and $\varphi : M \rightarrow \hat{M}$ be a diffeomorphism between them. Then φ induces the **isomorphism** $\varphi_* : (J(M, n), C) \rightarrow (J(\hat{M}, n), \hat{C})$, i.e., the differential map $\varphi_* : J(M, n) \rightarrow J(\hat{M}, n)$ is a diffeomorphism sending C onto \hat{C} .

2.2. Second Order Contact Manifolds. Let J be a manifold and C be a (linear) differential system on J of codimension 1. Namely C is a subbundle of $T(J)$ of codimension 1. Thus, locally at each point u of J , there exists a 1-form ϖ defined around $u \in J$ such that

$$C = \{\varpi = 0\}.$$

Then (J, C) is called a **contact manifold** if $\varpi \wedge (d\varpi)^n$ forms a volume element of J . This condition is equivalent to the following conditions (1), (2) or (3);

- (1) The restriction $d\varpi|_C$ of $d\varpi$ to $C(u)$ is non-degenerate at each point $u \in J$.

- (2) There exists a coframe $\{\varpi, \omega_1, \dots, \omega_n, \pi_1, \dots, \pi_n\}$ defined around $u \in J$ such that the following holds;

$$d\varpi \equiv \omega_1 \wedge \pi_1 + \dots + \omega_n \wedge \pi_n \quad (\text{mod } \varpi)$$

- (3) The Cauchy characteristic system $\text{Ch}(C)$ of C is trivial (see §2.3 below).

By the **Darboux Theorem**, a contact manifold (J, C) of dimension $2n + 1$ can be regarded locally as a space of 1-jets for one unknown function. Namely, at each point of (J, C) , there exists a canonical coordinate system $(x_1, \dots, x_n, z, p_1, \dots, p_n)$ such that

$$C = \left\{ dz - \sum_{i=1}^n p_i dx_i = 0 \right\}.$$

Starting from a contact manifold (J, C) , we can construct the geometric second order jet space $(L(J), E)$ as follows: We consider the **Lagrange-Grassmann bundle** $L(J)$ over J consisting of all n -dimensional integral elements of (J, C) ;

$$L(J) = \bigcup_{u \in J} L_u \xrightarrow{\pi} J,$$

where L_u is the Grassmann manifolds of all lagrangian (or **legendrian**) subspaces of the symplectic vector space $(C(u), d\varpi)$. Here ϖ is a local contact form on J . Then the canonical system E on $L(J)$ is defined by

$$E(v) = \pi_*^{-1}(v) \subset T_v(L(J)) \xrightarrow{\pi_*} T_u(J), \quad \text{for } v \in L(J).$$

Let us fix a point $v_o \in L(J)$. Starting from a canonical coordinate system $(x_1, \dots, x_n, z, p_1, \dots, p_n)$ of (J, C) around $u_o = \pi(v_o)$ such that $dx_1 \wedge \dots \wedge dx_n|_{v_o} \neq 0$, we can introduce a coordinate system (x_i, z, p_i, p_{ij}) ($1 \leq i \leq j \leq n$) by defining coordinate functions p_{ij} as follows;

$$dp_i|_v = \sum_{i=1}^n p_{ij}(v) dx_j|_v.$$

Then, since $v \in C(u)$, we have $dz|_v = \sum_{i=1}^n p_i(u) dx_i|_v$ and, since $d\varpi|_v = 0$, we get $p_{ij} = p_{ji}$.

Thus E is defined on this canonical coordinate system by

$$E = \{\varpi = \varpi_1 = \dots = \varpi_n = 0\},$$

where

$$\varpi = dz - \sum_{i=1}^n p_i dx_i, \quad \text{and} \quad \varpi_i = dp_i - \sum_{j=1}^n p_{ij} dx_j \quad \text{for } i = 1, \dots, n.$$

Let (J, C) , (\hat{J}, \hat{C}) be contact manifolds of dimension $2n + 1$ and $\varphi : (J, C) \rightarrow (\hat{J}, \hat{C})$ be a contact diffeomorphism between them. Then φ induces an isomorphism $\varphi_* : (L(J), E) \rightarrow (L(\hat{J}), \hat{E})$. Conversely we have (cf. Theorem 3.2 [Y1])

Theorem 2.1. (Bäcklund) *Let (J, C) and (\hat{J}, \hat{C}) be contact manifolds of dimension $2n + 1$. Then, for an isomorphism $\Phi : (L(J), E) \rightarrow (L(\hat{J}), \hat{E})$, there exists a contact diffeomorphism $\varphi : (J, C) \rightarrow (\hat{J}, \hat{C})$ such that $\Phi = \varphi_*$.*

2.3. Derived Systems and Cauchy Characteristic Systems. Now we prepare basic notions for **(linear) differential systems** (or Pfaffian systems). By a (linear) differential system (M, D) , we mean a subbundle D of the tangent bundle $T(M)$ of a manifold M of dimension d . Locally D is defined by 1-forms $\omega_1, \dots, \omega_{d-r}$ such that $\omega_1 \wedge \dots \wedge \omega_{d-r} \neq 0$ at each point, where r is the rank of D ;

$$D = \{ \omega_1 = \dots = \omega_{d-r} = 0 \}.$$

For two differential systems (M, D) and (\hat{M}, \hat{D}) , a diffeomorphism φ of M onto \hat{M} is called an **isomorphism** of (M, D) onto (\hat{M}, \hat{D}) if the differential map φ_* of φ sends D onto \hat{D} .

For a non-integrable differential system D , we consider the **Derived System** ∂D of D , which is defined, in terms of sections, by

$$\partial \mathcal{D} = \mathcal{D} + [\mathcal{D}, \mathcal{D}].$$

where $\mathcal{D} = \Gamma(D)$ denotes the space of sections of D .

Furthermore the **Cauchy Characteristic System** $\text{Ch}(D)$ of (M, D) is defined at each point $x \in M$ by

$$\text{Ch}(D)(x) = \{ X \in D(x) \mid X \lrcorner d\omega_i \equiv 0 \pmod{\omega_1, \dots, \omega_s} \text{ for } i = 1, \dots, s \},$$

where \lrcorner denotes the interior multiplication, i.e., $X \lrcorner d\omega(Y) = d\omega(X, Y)$ and $s = d - r$. When $\text{Ch}(D)$ is a differential system (i.e., has constant rank), it is always completely integrable.

Moreover **Higher Derived Systems** $\partial^k D$ are usually defined successively (cf. [BCG₃]) by

$$\partial^k D = \partial(\partial^{k-1} D),$$

where we put $\partial^0 D = D$ for convention.

On the other hand we define the k -th **Weak Derived System** $\partial^{(k)} D$ of D inductively by

$$\partial^{(k)} \mathcal{D} = \partial^{(k-1)} \mathcal{D} + [\mathcal{D}, \partial^{(k-1)} \mathcal{D}],$$

where $\partial^{(0)} D = D$ and $\partial^{(k)} \mathcal{D}$ denotes the space of sections of $\partial^{(k)} D$.

2.4. Review of Tanaka Theory. A differential system (M, D) is called **regular**, if $D^{-(k+1)} = \partial^{(k)} D$ are subbundles of $T(M)$ for every integer $k \geq 1$. For a regular differential system (M, D) , we have ([T2], Proposition 1.1)

(S1) *There exists a unique integer $\mu > 0$ such that, for all $k \geq \mu$,*

$$D^{-k} = \dots = D^{-\mu} \supsetneq D^{-\mu+1} \supsetneq \dots \supsetneq D^{-2} \supsetneq D^{-1} = D,$$

(S2) $[\mathcal{D}^p, \mathcal{D}^q] \subset \mathcal{D}^{p+q}$ for all $p, q < 0$.

where \mathcal{D}^p denotes the space of sections of D^p . (S2) implies subbundles D^p define a filtration on M .

Let (M, D) be a regular differential system such that $T(M) = D^{-\mu}$. As a first invariant for non-integrable differential systems, we now define the **symbol algebra** $\mathfrak{m}(x)$ associated with a differential system (M, D) at $x \in M$, which was introduced by N. Tanaka [T2].

We put $\mathfrak{g}_{-1}(x) = D^{-1}(x)$, $\mathfrak{g}_p(x) = D^p(x)/D^{p+1}(x)$ ($p < -1$) and

$$\mathfrak{m}(x) = \bigoplus_{p=-1}^{-\mu} \mathfrak{g}_p(x).$$

Let π_p be the projection of $D^p(x)$ onto $\mathfrak{g}_p(x)$. Then, for $X \in \mathfrak{g}_p(x)$ and $Y \in \mathfrak{g}_q(x)$, the bracket product $[X, Y] \in \mathfrak{g}_{p+q}(x)$ is defined by

$$[X, Y] = \pi_{p+q}([\tilde{X}, \tilde{Y}]_x),$$

where \tilde{X} and \tilde{Y} are any element of \mathcal{D}^p and \mathcal{D}^q respectively such that $\pi_p(\tilde{X}_x) = X$ and $\pi_q(\tilde{Y}_x) = Y$.

Endowed with this bracket operation, by (S2) above, $\mathfrak{m}(x)$ becomes a nilpotent graded Lie algebra such that $\dim \mathfrak{m}(x) = \dim M$ and satisfies

$$\mathfrak{g}_p(x) = [\mathfrak{g}_{p+1}(x), \mathfrak{g}_{-1}(x)] \quad \text{for } p < -1.$$

We call $\mathfrak{m}(x)$ the **Symbol Algebra of** (M, D) at $x \in M$.

Furthermore, let \mathfrak{m} be a FGLA (fundamental graded Lie algebra) of μ -th kind, that is,

$$\mathfrak{m} = \bigoplus_{p=-1}^{-\mu} \mathfrak{g}_p$$

is a nilpotent graded Lie algebra such that

$$\mathfrak{g}_p = [\mathfrak{g}_{p+1}, \mathfrak{g}_{-1}] \quad \text{for } p < -1.$$

Then (M, D) is called of type \mathfrak{m} if the symbol algebra $\mathfrak{m}(x)$ is isomorphic to \mathfrak{m} at each $x \in M$.

Conversely, given a FGLA $\mathfrak{m} = \bigoplus_{p=-1}^{-\mu} \mathfrak{g}_p$, we can construct a model differential system of type \mathfrak{m} as follows: Let $M(\mathfrak{m})$ be the simply connected Lie group with Lie algebra \mathfrak{m} . Identifying \mathfrak{m} with the Lie algebra of left invariant vector fields on $M(\mathfrak{m})$, \mathfrak{g}_{-1} defines a left invariant subbundle $D_{\mathfrak{m}}$ of $T(M(\mathfrak{m}))$. By definition of symbol algebras, it is easy to see that $(M(\mathfrak{m}), D_{\mathfrak{m}})$ is a regular differential system of type \mathfrak{m} . $(M(\mathfrak{m}), D_{\mathfrak{m}})$ is called the **Standard Differential System of Type \mathfrak{m}** . The Lie algebra of all infinitesimal automorphisms of $(M(\mathfrak{m}), D_{\mathfrak{m}})$ can be calculated algebraically as the **Prolongation $\mathfrak{g}(\mathfrak{m})$ of \mathfrak{m}** ([T1], cf. [Y5]).

In fact, let $\mathfrak{m} = \bigoplus_{p < 0} \mathfrak{g}_p$ be a fundamental graded Lie algebra of μ -th kind defined over a field \mathbb{K} . Here \mathbb{K} denotes the field of real numbers \mathbb{R} or that of complex numbers \mathbb{C} . We put

$$\mathfrak{g}(\mathfrak{m}) = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p(\mathfrak{m}),$$

where $\mathfrak{g}_p(\mathfrak{m}) = \mathfrak{g}_p$ for $p < 0$, $\mathfrak{g}_0(\mathfrak{m})$ is the Lie algebra of all (gradation preserving) derivations of graded Lie algebra \mathfrak{m} and $\mathfrak{g}_k(\mathfrak{m})$ is defined inductively by the following for $k \geq 1$;

$$\mathfrak{g}_k(\mathfrak{m}) = \left\{ u \in \bigoplus_{p < 0} \mathfrak{g}_{p+k} \otimes \mathfrak{g}_p^* \mid u([Y, Z]) = [u(Y), Z] - [u(Z), Y] \right\}.$$

Thus, as a vector space over \mathbb{K} , $\mathfrak{g}_k(\mathfrak{m})$ is a linear subspace of $\text{End}(\mathfrak{m}, \mathfrak{m}^k) = \mathfrak{m}^k \otimes \mathfrak{m}^*$, where $\mathfrak{m}^k = \mathfrak{m} \oplus \mathfrak{g}_0(\mathfrak{m}) \oplus \cdots \oplus \mathfrak{g}_{k-1}(\mathfrak{m})$. The bracket operation of $\mathfrak{g}(\mathfrak{m})$ is given accordingly (see [T1], [Y5] for detail).

The structure of the Lie algebra $\mathcal{A}(M(\mathfrak{m}), D_{\mathfrak{m}})$ of all infinitesimal automorphisms of $(M(\mathfrak{m}), D_{\mathfrak{m}})$ can be described by $\mathfrak{g}(\mathfrak{m})$. Especially $\mathcal{A}(M(\mathfrak{m}), D_{\mathfrak{m}})$ is isomorphic to $\mathfrak{g}(\mathfrak{m})$, when $\mathfrak{g}(\mathfrak{m})$ is finite dimensional.

Let \mathfrak{g}_0 be a subalgebra of $\mathfrak{g}_0(\mathfrak{m})$. We define a subspace \mathfrak{g}_k of $\mathfrak{g}_k(\mathfrak{m})$ for $k \geq 1$ inductively by

$$\mathfrak{g}_k = \left\{ u \in \mathfrak{g}_k(\mathfrak{m}) \mid [u, \mathfrak{g}_{-1}] \subset \mathfrak{g}_{k-1} \right\}.$$

Then, putting

$$\mathfrak{g}(\mathfrak{m}, \mathfrak{g}_0) = \mathfrak{m} \oplus \bigoplus_{k \geq 0} \mathfrak{g}_k,$$

we see, with the generating condition of \mathfrak{m} , that $\mathfrak{g}(\mathfrak{m}, \mathfrak{g}_0)$ is a graded subalgebra of $\mathfrak{g}(\mathfrak{m})$. $\mathfrak{g}(\mathfrak{m}, \mathfrak{g}_0)$ is called the prolongation of $(\mathfrak{m}, \mathfrak{g}_0)$.

We will recall in §5.1 when $\mathfrak{g}(\mathfrak{m})$ or $\mathfrak{g}(\mathfrak{m}, \mathfrak{g}_0)$ becomes finite dimensional and simple.

2.5. Symbol Algebra of $(L(J), E)$. As an example to calculate symbol algebras, let us show that $(L(J), E)$ is a regular differential system of type $\mathfrak{c}^2(n)$:

$$\mathfrak{c}^2(n) = \mathfrak{c}_{-3} \oplus \mathfrak{c}_{-2} \oplus \mathfrak{c}_{-1},$$

where $\mathfrak{c}_{-3} = \mathbb{R}$, $\mathfrak{c}_{-2} = V^*$ and $\mathfrak{c}_{-1} = V \oplus S^2(V^*)$. Here V is a vector space of dimension n and the bracket product of $\mathfrak{c}^2(n)$ is defined accordingly through the pairing between V and V^* such that V and $S^2(V^*)$ are both abelian subspaces of \mathfrak{c}_{-1} . This fact can be checked as follows: Let us take a canonical coordinate system (x_i, z, p_i, p_{ij}) ($1 \leq i \leq j \leq n$) of $(L(J), E)$. Then we have a coframe $\{\varpi, \varpi_i, dx_i, dp_{ij}\}$ ($1 \leq i \leq j \leq n$) at each point in this coordinate neighborhood, where $\varpi = dz - \sum_{i=1}^n p_i dx_i$, $\varpi_i = dp_i - \sum_{j=1}^n p_{ij} dx_j$ ($i = 1, \dots, n$). Now take the dual frame $\{\frac{\partial}{\partial z}, \frac{\partial}{\partial p_i}, \frac{d}{dx_i}, \frac{\partial}{\partial p_{ij}}\}$, of this coframe, where

$$\frac{d}{dx_i} = \frac{\partial}{\partial x_i} + p_i \frac{\partial}{\partial z} + \sum_{j=1}^n p_{ij} \frac{\partial}{\partial p_j}$$

is the classical notation. Notice that $\{\frac{d}{dx_i}, \frac{\partial}{\partial p_{ij}}\}$ ($1 \leq i \leq j \leq n$) forms a free basis of $\Gamma(E)$. Then we have

$$\begin{aligned} \left[\frac{\partial}{\partial p_{ii}}, \frac{d}{dx_j} \right] &= \delta_j^i \frac{\partial}{\partial p_i}, & \left[\frac{\partial}{\partial p_{ij}}, \frac{d}{dx_k} \right] &= \delta_k^i \frac{\partial}{\partial p_i} + \delta_k^j \frac{\partial}{\partial p_j} \quad \text{for } i \neq j, \\ \left[\frac{\partial}{\partial p_i}, \frac{d}{dx_j} \right] &= \delta_j^i \frac{\partial}{\partial z}, & \left[\frac{d}{dx_i}, \frac{d}{dx_j} \right] &= 0. \end{aligned}$$

It follows that $T(L(J)) = \partial^{(2)}E$ and the derived system ∂E of E satisfies the following :

$$\partial E = \{\varpi = 0\} = \pi_*^{-1}C, \quad \text{Ch}(\partial E) = \text{Ker } \pi_*.$$

These facts provide the proof of Theorem 1 (cf. Theorem 3.2 [Y1]).

Moreover, in terms of the defining 1-forms of E and ∂E around $v \in L(J)$, the structure of the symbol algebra $\mathfrak{c}^2(n)$ can be described by the following Structure Equation of E.Cartan([C1],[C2]);

$$\begin{cases} d\varpi \equiv \omega_1 \wedge \varpi_1 + \dots + \omega_n \wedge \varpi_n \quad (\text{mod } \varpi, \varpi_i \wedge \varpi_j (1 \leq i \leq j \leq n)) \\ \left\{ \begin{array}{l} d\varpi_1 \equiv \omega_1 \wedge \pi_{11} + \dots + \omega_n \wedge \pi_{1n} \\ \dots \\ d\varpi_n \equiv \omega_1 \wedge \pi_{n1} + \dots + \omega_n \wedge \pi_{nn} \end{array} \right. \quad (\text{mod } \varpi, \varpi_1, \dots, \varpi_n) \end{cases}$$

where $\partial E = \{\varpi = 0\}$, $E = \{\varpi = \varpi_1 = \dots = \varpi_n = 0\}$ and $\{\varpi, \varpi_i, \omega_i, \pi_{ij} \quad (1 \leq i \leq j \leq n)\}$ forms a coframe around $v \in L(J)$. Here we understand that $\pi_{ij} = \pi_{ji}$.

Similarly we see that $(J(M, n), C)$ is a regular differential system of type $\mathfrak{c}^1(n, m)$:

$$\mathfrak{c}^1(n, m) = \mathfrak{c}_{-2} \oplus \mathfrak{c}_{-1},$$

where $\mathfrak{c}_{-2} = W$ and $\mathfrak{c}_{-1} = V \oplus W \otimes V^*$ for vector spaces V and W of dimension n and m respectively, and the bracket product of $\mathfrak{c}^1(n, m)$ is defined accordingly through the pairing between V and V^* such that V and $W \otimes V^*$ are both abelian subspaces of \mathfrak{c}_{-1} .

3. SYMBOLS OF SECOND ORDER EQUATIONS.

In view of the Bäcklund Theorem, we will discuss the symbols of second order equations as the first invariants in the Contact Geometry of Second Order.

3.1. Symbol Algebras. Let R be a submanifold of $L(J)$ satisfying the following condition:

$$(R.0) \quad p : R \rightarrow J ; \text{ submersion,}$$

where $p = \pi|_R$ and $\pi : L(J) \rightarrow J$ is the projection. This condition implies that the system of equations R of second order contains no equations of first order. We have two differential systems $C^1 = \partial E$ and $C^2 = E$ on $L(J)$. We denote by D^1 and D^2 those differential systems on R obtained by restricting C^1 and C^2 to R . Moreover we denote by the same symbols those 1-forms obtained by restricting the defining 1-forms $\{\varpi, \varpi_1, \dots, \varpi_n\}$ of the canonical system E to R , where $\varpi = dz - \sum_{i=1}^n p_i dx_i$, and $\varpi_i = dp_i - \sum_{j=1}^n p_{ij} dx_j$ ($i = 1, \dots, n$). Then it follows from (R.0) that these 1-forms are independent at each point on R and that

$$D^1 = \{\varpi = 0\}, \quad D^2 = \{\varpi = \varpi_1 = \dots = \varpi_n = 0\}.$$

Thus D^1 and D^2 are subbundles of $T(R)$ such that $\partial D^2 \subset D^1$. Hence subbundles D^2, D^1 and $T(R)$ define a filtration on R . Namely, putting $D^{-1} = D^2, D^{-2} = D^1, D^p = T(R)$ for $p \leq -3$, we have

$$[D^p, D^q] \subset D^{p+q} \quad \text{for } p, q < 0,$$

where $D^p = \Gamma(D^p)$.

Now we define the **Symbol Algebra** $\mathfrak{s}(v)$ of R at $v \in R$ by

$$\mathfrak{s}(v) = \mathfrak{s}_{-3}(v) \oplus \mathfrak{s}_{-2}(v) \oplus \mathfrak{s}_{-1}(v),$$

where $\mathfrak{s}_{-3}(v) = T_v(R)/D^1(v)$, $\mathfrak{s}_{-2}(v) = D^1(v)/D^2(v)$ and $\mathfrak{s}_{-1}(v) = D^2(v)$. The bracket operation in $\mathfrak{s}(v)$ is defined, similarly as in §2.3, as follows: For $X \in \mathfrak{s}_p(v)$ and $Y \in \mathfrak{s}_q(v)$, let us take $\tilde{X} \in D^p$ and $\tilde{Y} \in D^q$ such that $X = \pi_p((\tilde{X})_v)$ and $Y = \pi_q((\tilde{Y})_v)$, where $\pi_p : D^p(v) \rightarrow \mathfrak{s}_p(v)$ is the projection. Then the bracket product is defined by

$$[X, Y] = \pi_{p+q}([\tilde{X}, \tilde{Y}]_v) \in \mathfrak{s}_{p+q}(v).$$

The bracket product $[X, Y]$ is well-defined for $X \in \mathfrak{s}_p(v)$ and $Y \in \mathfrak{s}_q(v)$, i.e., is independent of the choice of \tilde{X} and \tilde{Y} . In fact, in our case, this can be shown as follows: The defining 1-forms $\varpi, \varpi_1, \dots, \varpi_n$ for D^1 and D^2 actually define a basis $\{A\}$ of $\mathfrak{s}_{-3}(v)$ and $\{B_1, \dots, B_n\}$ of $\mathfrak{s}_{-2}(v)$ such that $\varpi(\tilde{A}) = 1, \pi_{-3}(\tilde{A}) = A, \varpi_i(\tilde{B}_j) = \delta_j^i, \pi_{-2}(\tilde{B}_i) = B_i$ and $\tilde{B}_i \in D^1(v)$. Then, for $X_1, X_2 \in \mathfrak{s}_{-1}(v) = D^2(v)$, we calculate

$$d\varpi_i(X_1, X_2) = \tilde{X}_1(\varpi_i(\tilde{X}_2)) - \tilde{X}_2(\varpi_i(\tilde{X}_1)) - \varpi_i([\tilde{X}_1, \tilde{X}_2]) = -\varpi_i([\tilde{X}_1, \tilde{X}_2]).$$

Thus, putting $\beta_i = -d\varpi_i(X_1, X_2)$, we get

$$[X_1, X_2] = \beta_1 B_1 + \dots + \beta_n B_n \in \mathfrak{s}_{-2}(v).$$

For $X \in \mathfrak{s}_{-1}(v)$ and $Y \in \mathfrak{s}_{-2}(v)$, we calculate

$$d\varpi(X, \tilde{Y}_v) = \tilde{X}_v(\varpi(\tilde{Y})) - \tilde{Y}_v(\varpi(\tilde{X})) - \varpi([\tilde{X}, \tilde{Y}]_v) = -\varpi([\tilde{X}, \tilde{Y}]_v).$$

Similarly we have $d\varpi(X_1, X_2) = 0$ for $X_1, X_2 \in \mathfrak{s}_{-1}(v)$. Thus $d\varpi(X, \tilde{Y}_v)$ depends only on $X \in \mathfrak{s}_{-1}(v)$ and $Y \in \mathfrak{s}_{-2}(v)$. Hence, putting $\alpha = -d\varpi(X, \tilde{Y}_v)$, we have

$$[X, Y] = \alpha A \in \mathfrak{s}_{-3}(v).$$

Moreover it follows that, for $X \in \mathfrak{s}_{-1}(v)$,

$$X \lrcorner d\varpi(Y) = 0 \quad \text{for } \forall Y \in D^1(v) \quad \text{if and only if } [X, \mathfrak{s}_{-2}(v)] = 0.$$

Hence, from $\text{Ch}(D^1) = \text{Ker } p_* \subset D^2$, we have, putting $\mathfrak{f}(v) = \text{Ch}(D^1)(v)$,

$$\mathfrak{f}(v) = \{X \in \mathfrak{s}_{-1}(v) \mid [X, \mathfrak{s}_{-2}(v)] = 0\}.$$

$\mathfrak{f}(v)$ is a subspace of $\mathfrak{s}_{-1}(v)$ of codimension n .

By the description of the bracket operation in $\mathfrak{s}(v)$ above, since ϖ and $\varpi_1, \dots, \varpi_n$ are the restriction of defining 1-forms of $C^1 = \partial E$ and $C^2 = E$ on $L(J)$, we immediately see that ϖ and $\varpi_1, \dots, \varpi_n$, at the same time, define bases of $\mathfrak{g}_{-3}(v)$ and $\mathfrak{g}_{-2}(v)$ of the symbol algebra $\mathfrak{m}(v) = \mathfrak{g}_{-3}(v) \oplus \mathfrak{g}_{-2}(v) \oplus \mathfrak{g}_{-1}(v) (\cong \mathfrak{c}^2(n))$ of $(L(J), E)$ at $v \in L(J)$ so that $\mathfrak{s}(v)$ is a graded subalgebra of $\mathfrak{m}(v)$ satisfying $\mathfrak{s}_{-3}(v) = \mathfrak{g}_{-3}(v)$, $\mathfrak{s}_{-2}(v) = \mathfrak{g}_{-2}(v)$ and $\mathfrak{f}(v) = T_v(R) \cap \text{Ch}(C^1)(v)$.

Now we consider the following compatibility condition for R :

$$(C) \quad p^{(1)} : R^{(1)} \rightarrow R \text{ is onto.}$$

where $R^{(1)}$ is the first prolongation of R . Namely we assume that there exists an n -dimensional integral element V of (R, D^2) at each $v \in R$ such that

$$\mathfrak{s}_{-1}(v) = V \oplus \mathfrak{f}(v).$$

V is an abelian subalgebra in $\mathfrak{s}(v)$. By fixing a basis of $\mathfrak{s}_{-3}(v)$, $\mathfrak{s}_{-3}(v)$ is identified with \mathbb{R} and, through $[\cdot, \cdot] : \mathfrak{s}_{-2}(v) \times \mathfrak{s}_{-1}(v) \rightarrow \mathfrak{s}_{-3}(v) \cong \mathbb{R}$, $\mathfrak{s}_{-2}(v)$ is identified with V^* , since $V \cap \mathfrak{f}(v) = \{0\}$ and $\mathfrak{f}(v) = \{X \in \mathfrak{s}_{-1}(v) \mid [X, \mathfrak{s}_{-2}(v)] = 0\}$. Moreover we have a map $\mu : \mathfrak{f}(v) \rightarrow S^2(V^*)$ defined by

$$\mu(f)(v_1, v_2) = [[f, v_1], v_2] \in \mathfrak{s}_{-3}(v) \cong \mathbb{R} \quad \text{for } f \in \mathfrak{f}(v).$$

Here $\mu(f)(v_1, v_2) = \mu(f)(v_2, v_1)$ follows from $[v_1, v_2] = 0$ and the Jacobi identity of $\mathfrak{s}(v)$. We can check the injectivity of μ as follows: If $\mu(f) = 0$, we have $[f, v_1] = 0$ for $\forall v_1 \in V$ by $\mathfrak{s}_{-2}(v) \cong V^*$. Then we have $[f, \mathfrak{s}_{-1}(v)] = 0$, since $\mathfrak{f}(v)$ is abelian, which implies $f \in \mathfrak{f}(v) \cap \text{Ch}(D^2)(v)$. From $\text{Ch}(D^1)(v) \cap \text{Ch}(D^2)(v) = 0$ (see §4.1), we obtain $f = 0$.

Hence, by fixing a basis of $\mathfrak{s}_{-3}(v)$ and the brackets in $\mathfrak{s}(v)$, we obtain

$$\mathfrak{s}_{-3}(v) \cong \mathbb{R}, \quad \mathfrak{s}_{-2}(v) \cong V^*, \quad \mathfrak{s}_{-1}(v) = V \oplus \mathfrak{f}(v) \quad \text{and} \quad \mathfrak{f}(v) \subset S^2(V^*).$$

Thus $\mathfrak{f}(v) \subset S^2(V^*)$ is the first invariant of R under contact transformation. We will first examine $\mathfrak{f}(v) \subset S^2(V^*)$ in the case $\dim V = 2$ in the next section.

3.2. Case $n = 2$. When $\dim V = 2$, we have $\dim S^2(V^*) = 3$. Through the natural pairing of $S^2(V)$ and $S^2(V^*)$ as subspaces of $V \otimes V$ and $V^* \otimes V^*$, we identify $S^2(V)$ with the dual space of $S^2(V^*)$. For a basis $\{e_1, e_2\}$ of V , we have a basis $\{e_1^* \otimes e_1^*, 2e_1^* \otimes e_2^*, e_2^* \otimes e_2^*\}$ of $S^2(V^*)$ and its dual basis $\{e_1 \otimes e_1, e_1 \otimes e_2, e_2 \otimes e_2\}$ of $S^2(V)$, where $\{e_1^*, e_2^*\}$ is the dual basis of $\{e_1, e_2\}$ and $e_i \otimes e_j = \frac{1}{2}(e_i \otimes e_j + e_j \otimes e_i)$. For a subspace \mathfrak{f} of $S^2(V^*)$, we denote by \mathfrak{f}^\perp the annihilator of \mathfrak{f} in $S^2(V)$.

Now let us classify subspaces \mathfrak{f} of $S^2(V^*)$ under the action of $GL(V)$.

(1) $\text{codim } \mathfrak{f} = 1$

In this case $\dim \mathfrak{f}^\perp = 1$. Hence we can classify a generator f of \mathfrak{f}^\perp as a quadratic form and obtain the following classification into three cases, i.e., there exists a basis of V such that

$$\mathfrak{f}^\perp = \begin{cases} \langle \{e_1 \otimes e_1\} \rangle, \\ \langle \{e_1 \otimes e_2\} \rangle, \\ \langle \{e_1 \otimes e_1 + e_2 \otimes e_2\} \rangle \end{cases}$$

The third case occurs when we classify over \mathbb{R} . Here f is of rank 1, rank 2 (indefinite) and rank 2 (definite) respectively.

(2) $\text{codim } \mathfrak{f} = 2$

In this case $\dim \mathfrak{f} = 1$. Hence, similarly as above, we have the following classification into three cases, i.e., there exists a basis of V such that

$$\mathfrak{f} = \begin{cases} \langle \{e_2^* \otimes e_2^*\} \rangle, \\ \langle \{e_1^* \otimes e_2^*\} \rangle, \\ \langle \{e_1^* \otimes e_1^* + e_2^* \otimes e_2^*\} \rangle, \end{cases}$$

Thus, dually, we have

$$\mathfrak{f}^\perp = \begin{cases} \langle \{e_1 \otimes e_1, e_1 \otimes e_2\} \rangle, \\ \langle \{e_1 \otimes e_1, e_2 \otimes e_2\} \rangle, \\ \langle \{e_1 \otimes e_2, e_1 \otimes e_1 - e_2 \otimes e_2\} \rangle \end{cases}$$

The third case occurs when we classify over \mathbb{R} . We note here that, for the prolongation $\mathfrak{f}^{(1)} = \mathfrak{f} \otimes V^* \cap S^3(V^*)$, we have $(\mathfrak{f}^{(1)})^\perp = \langle \{e_1 \otimes e_1 \otimes e_1, e_1 \otimes e_1 \otimes e_2, e_1 \otimes e_2 \otimes e_2, e_2 \otimes e_2 \otimes e_2\} \rangle$ for the second and third cases (see §3.3). Namely $\mathfrak{f}^{(1)} = \{0\}$ for the second and third cases, whereas the first case is involutive.

We can classify the symbol algebra $\mathfrak{s}(v)$ of R at $v \in R$ according to the above classification for $\mathfrak{f}(v) \subset S^2(V^*)$ under the condition (C). In the case $\text{codim } R = 1$, R is called **parabolic**, **hyperbolic** and **elliptic** at v according as f is of rank 1, rank 2 (indefinite) and rank 2 (definite) respectively, where f is a generator of $(\mathfrak{f}(v))^\perp \subset S^2(V)$ (see §3.3).

Now we assume the regularity for the symbol algebras. Namely assume that symbol algebras $\mathfrak{s}(v)$ of R are locally isomorphic to the fixed symbol $\mathfrak{s} = \mathfrak{s}_{-3} \oplus \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1}$ where $\mathfrak{s}_{-3} = \mathbb{R}$, $\mathfrak{s}_{-2} = V^*$ and $\mathfrak{s}_{-1} = V \oplus \mathfrak{f}$ for the fixed $\mathfrak{f} \subset S^2(V^*)$. Then, for example, the Structure Equation reads as follows:

(i) $\mathfrak{f}^\perp = \langle \{e_1 \otimes e_2\} \rangle$

$$\begin{cases} d\varpi \equiv \omega_1 \wedge \varpi_1 + \omega_2 \wedge \varpi_2 & (\text{mod } \varpi) \\ d\varpi_1 \equiv \omega_1 \wedge \pi_{11} & (\text{mod } \varpi, \varpi_1, \varpi_2) \\ d\varpi_2 \equiv \omega_2 \wedge \pi_{22} & (\text{mod } \varpi, \varpi_1, \varpi_2) \end{cases}$$

(ii) $\mathfrak{f}^\perp = \langle \{e_1 \otimes e_1, e_1 \otimes e_2\} \rangle$

$$\begin{cases} d\varpi \equiv \omega_1 \wedge \varpi_1 + \omega_2 \wedge \varpi_2 & (\text{mod } \varpi) \\ d\varpi_1 \equiv 0 & (\text{mod } \varpi, \varpi_1, \varpi_2) \\ d\varpi_2 \equiv \omega_2 \wedge \pi_{22} & (\text{mod } \varpi, \varpi_1, \varpi_2) \end{cases}$$

(iii) $\mathfrak{f}^\perp = \langle \{e_1 \otimes e_1, e_2 \otimes e_2\} \rangle$

$$\begin{cases} d\varpi \equiv \omega_1 \wedge \varpi_1 + \omega_2 \wedge \varpi_2 & (\text{mod } \varpi) \\ d\varpi_1 \equiv \omega_2 \wedge \pi_{12} & (\text{mod } \varpi, \varpi_1, \varpi_2) \\ d\varpi_2 \equiv \omega_1 \wedge \pi_{21} & (\text{mod } \varpi, \varpi_1, \varpi_2) \end{cases}$$

Thus we see, from (ii), that R admits a 1-dimensional Cauchy characteristic system in case $\mathfrak{f}^\perp = \langle \{e_1 \otimes e_1, e_1 \otimes e_2\} \rangle$, i.e, in case R is of codimension 2 and \mathfrak{f} is involutive. In fact, in [C1], E.Cartan characterized overdetermined involutive system R by the condition that R admits a 1-dimensional Cauchy characteristic system.

We will discuss the case when R admits a non-trivial Cauchy characteristic system in §4.2 in general. We will encounter the case (iii) in §5.3 as an example of Parabolic Geometry associated with $\mathfrak{sl}(4)$.

3.3. Involutive Symbols. In general we will consider the case when $\dim V = n$. We identify $S^2(V)$ with the dual space of $S^2(V^*)$ through the natural pairing of $S^2(V)$ and $S^2(V^*)$ as subspaces of $V \otimes V$ and $V^* \otimes V^*$. Then, for a basis $\{e_1, \dots, e_n\}$ of V , we have a basis $\{e_1^* \otimes e_1^*, \dots, e_n^* \otimes e_n^*, 2e_i^* \otimes e_j^* (1 \leq i < j \leq n)\}$ of $S^2(V^*)$ and its dual basis $\{e_i \otimes e_j (1 \leq i \leq j \leq n)\}$ of $S^2(V)$, where $\{e_1^*, \dots, e_n^*\}$ is the dual basis of V^* .

Here we note the adjoint map $\sigma^*(v) : S^k(V) \rightarrow S^{k+1}(V)$ of the interior multiplication $\sigma(v) : S^{k+1}(V^*) \rightarrow S^k(V^*)$ by a vector $v \in V$; $\sigma(v)(f) = v \lrcorner f$, i.e., $\sigma(v)(f)(v_1, \dots, v_k) = f(v, v_1, \dots, v_k)$ for $f \in S^{k+1}(V^*)$, is given by

$$\sigma^*(v)(a) = v \otimes a \quad \text{for } a \in S^k(V).$$

Hence, for a subspace $\mathfrak{f} \subset S^2(V^*)$ such that $\mathfrak{f}^\perp = \langle \{f_1, \dots, f_s\} \rangle$, the first prolongation $\mathfrak{f}^{(1)} = \mathfrak{f} \otimes V^* \cap S^3(V^*)$ is given by $(\mathfrak{f}^{(1)})^\perp = \langle \{e_i \otimes f_j \mid 1 \leq i \leq n, 1 \leq j \leq s\} \rangle$.

As in §3.2, we can classify codimension 1 subspace $\mathfrak{f} \subset S^2(V^*)$ as follows; In this case, we can classify a generator f of \mathfrak{f}^\perp as a quadratic form and obtain

$$f = e_1 \otimes e_1 \pm e_2 \otimes e_2 \pm \dots \pm e_r \otimes e_r,$$

for a basis $\{e_1, \dots, e_n\}$ of V , where r is the rank of f and we have an index when we classify over \mathbb{R} . In each case, $\{e_n, \dots, e_1\}$ forms a regular basis for \mathfrak{f} and the Cartan characters are given by $s_i = n - i + 1$ for $i = 1, \dots, n - 1$ and $s_n = 0$. \mathfrak{f} is always involutive.

For a single equation of second order

$$R = \{F(x_1, \dots, x_n, z, p_1, \dots, p_n, p_{11}, \dots, p_{nn}) = 0\} \subset L(J),$$

we observe the following: From §2.5, we calculate

$$\left[\left[\frac{\partial}{\partial p_{ii}}, X \right], X \right] = v_i^2 \frac{\partial}{\partial z}, \quad \left[\left[\frac{\partial}{\partial p_{ij}}, X \right], X \right] = 2v_i \cdot v_j \frac{\partial}{\partial z} \quad (i \neq j),$$

for $X = \sum_{i=1}^n v_i \frac{d}{dx_i}$. Thus $\frac{\partial}{\partial p_{ii}}$ is identified with $e_i^* \otimes e_i^*$ in $S^2(V^*)$ and $\frac{\partial}{\partial p_{ij}}$ is identified with $2e_i^* \otimes e_j^*$ in $S^2(V^*)$, where $\{e_i = \frac{d}{dx_i} (i = 1, \dots, n)\}$ forms a basis of V . Then, from $\mathfrak{f}(v) = T_v(R) \cap \text{Ch}(C^2)(v)$ at $v \in R$ and

$$dF \equiv \sum_{1 \leq i \leq j \leq n} \frac{\partial F}{\partial p_{ij}} dp_{ij} \pmod{\varpi, \varpi_1, \dots, \varpi_n, dx_1, \dots, dx_n},$$

we see that $(\mathfrak{f}(v))^\perp$ is generated by

$$f = \sum_{1 \leq i \leq j \leq n} \frac{\partial F}{\partial p_{ij}}(v) e_i \otimes e_j.$$

Next we consider the case when $\text{codim } \mathfrak{f} = 2$. In this case the involutiveness becomes rather a restrictive condition and, in fact, we have (cf. [C2] and [Y2])

Proposition 3.1. *Let \mathfrak{f} be a subspace of $S^2(V^*)$ of codimension 2. Then \mathfrak{f} is involutive if and only if there exists a basis $\{e_1, \dots, e_n\}$ of V such that the annihilator \mathfrak{f}^\perp of \mathfrak{f} in $S^2(V)$ is generated by $e_1 \otimes e_2$ and $e_1 \otimes e_3$ or by $e_1 \otimes e_1$ and $e_1 \otimes e_2$.*

Proof. First we observe that $S^2(V^*)$ is an involutive subspace of $V^* \otimes V^*$ with the Cartan characters $\sigma_i = n - i + 1$ for $i = 1, \dots, n$. Let \mathfrak{f} be a subspace of $S^2(V^*)$ of codimension 2 and let s_1, \dots, s_n be the Cartan characters of \mathfrak{f} . Then we have $s_i \leq \sigma_i$ for $i = 1, \dots, n$ and

$$\dim \mathfrak{f} = s_1 + \dots + s_n = \sigma_1 + \dots + \sigma_n - 2.$$

Since the Cartan characters have the property $s_1 \geq \dots \geq s_n$, if $s_{i_0} = \sigma_{i_0} - 2$ for some $i_0 \in \{1, \dots, n\}$, it follows $s_{i_0+1} = \sigma_{i_0+1} = n - i_0 > s_{i_0} = \sigma_{i_0} - 2 = n - i_0 - 1$, which is a contradiction. Hence there exist j and k ($1 \leq j < k \leq n$) such that $s_j = \sigma_j - 1$ and $s_k = \sigma_k - 1$.

Now assume that \mathfrak{f} is involutive. Then, from $\dim \mathfrak{f}^{(1)} = s_1 + 2s_2 + \dots + ns_n$, we obtain

$$\text{codim } \mathfrak{f}^{(1)} = \sum_{i=1}^n i(\sigma_i - s_i) = j + k \leq (n-1) + n = 2n - 1.$$

This implies that the generator $\{e_1 \otimes f, \dots, e_n \otimes f, e_1 \otimes g, \dots, e_n \otimes g\}$ of $(\mathfrak{f}^{(1)})^\perp$ in $S^3(V)$ are linearly dependent, where f and $g \in S^2(V)$ are the generator of \mathfrak{f}^\perp . Namely there exist v_1 and $v_2 \in V$ such that

$$v_1 \otimes f - v_2 \otimes g = 0.$$

Here v_1 and v_2 are linearly independent since f and g are independent. Hence there exists $v \in V$ such that $f = v_2 \otimes v$ and $g = v_1 \otimes v$. In case $\{v_1, v_2, v\}$ are independent, there exists a basis $\{e_1, \dots, e_n\}$ of V such that $f = e_1 \otimes e_2$ and $g = e_1 \otimes e_3$. In case $\{v_1, v_2, v\}$ are dependent, there exists a basis of V such that \mathfrak{f}^\perp is generated by $e_1 \otimes e_1$ and $e_1 \otimes e_2$. Consequently, in these cases, we have $s_i = n - i + 1$ for $1 \leq i \leq n - 2$, $s_{n-1} = 1$ and $s_n = 0$. \square

In [C2], E.Cartan, in fact, first classified involutive subspaces $\mathfrak{f} \subset S^2(V^*)$ when $\dim V = 3$ and immediately wrote the Structure Equation for each involutive system in this case.

However we cannot pursue this line in general by the following facts. By counting the dimensions, we see that the dimension of $\text{Gr}(S^2(V^*), r)$ exceeds the dimension of $GL(V)$ for $n \geq 4$ and $r, s \geq 2$, where $r + s = \dim \text{Gr}(S^2(V^*), r)$. Hence we will have a functional moduli if we try to classify r dimensional subspaces \mathfrak{f} in $S^2(V^*)$ for $n \geq 4$ and $r, s \geq 2$. We suspect this phenomena even if we assume the involutiveness of \mathfrak{f} .

Thus we need other guide lines to proceed. In this paper, after preparing the structure theory for submanifolds in $L(J)$ in §4, we will utilize Parabolic Geometries to find good classes of second order equations in §5 and §6.

3.4. Typical Symbols. We exhibit here typical examples of involutive symbols in $S^2(V^*)$, which are the only invariants of the corresponding involutive systems of second order, which were found in [Y4]. Namely we describe here the involutive subspaces $\mathfrak{f}^1(r)$, $\mathfrak{f}^2(r)$ and $\mathfrak{f}^3(r)$ of $S^2(V^*)$ which have the following property: Let R be an involutive systems of second order, which is regular of type \mathfrak{f} , i.e., R satisfies the condition (C) such that the symbol $\mathfrak{f}(v)$ at each point $v \in R$ is isomorphic to $\mathfrak{f} \subset S^2(V^*)$, where \mathfrak{f} is one of $\mathfrak{f}^1(r)$, $\mathfrak{f}^2(r)$ or $\mathfrak{f}^3(r)$. Then, as in the case of the system of first order partial differential equations of one dependent variable, R can be transformed to the model linear equation by a contact transformation(cf. [Y4]).

$$(1) \quad \mathfrak{f}^1(r) \subset S^2(V^*) \quad (2 \leq r \leq n-2) \quad V = V_r \oplus V_s$$

$$(\mathfrak{f}^1(r))^\perp = \langle \{e_i \otimes e_\alpha \mid 1 \leq i \leq r, r+1 \leq \alpha \leq n\} \rangle = V_r \otimes_S V_s,$$

When $\mathfrak{f} = \mathfrak{f}^1(r)$ (see [Y4] §2), there exists a canonical coordinate system (x_i, z, p_i, p_{ij}) ($1 \leq i \leq j \leq n$) of $L(J)$ such that

$$R = \left\{ \frac{\partial^2 z}{\partial x_i \partial x_\alpha} = 0 \quad (1 \leq i \leq r, r+1 \leq \alpha \leq n) \right\}.$$

$$(2) \quad \mathfrak{f}^2(r) \subset S^2(V^*) \quad (r \geq 2)$$

$$(\mathfrak{f}^2(r))^\perp = \langle \{e_i \otimes e_j \mid 1 \leq i \leq j \leq r\} \rangle = S^2(V_r),$$

When $\mathfrak{f} = \mathfrak{f}^2(r)$ (see [Y4] §3), there exists a canonical coordinate system (x_i, z, p_i, p_{ij}) ($1 \leq i \leq j \leq n$) of $L(J)$ such that

$$R = \left\{ \frac{\partial^2 z}{\partial x_i \partial x_j} = 0 \quad (1 \leq i \leq j \leq r) \right\}.$$

$$(3) \quad \mathfrak{f}^3(r) \subset S^2(V^*) \quad (r \leq n-2)$$

$$(\mathfrak{f}^3(r))^\perp = \langle \{e_i \otimes e_a \mid 1 \leq i \leq r, 1 \leq a \leq n\} \rangle = V_r \otimes_S V,$$

When $\mathfrak{f} = \mathfrak{f}^3(r)$ (see [Y4] §4), there exists a canonical coordinate system (x_i, z, p_i, p_{ij}) ($1 \leq i \leq j \leq n$) of $L(J)$ such that

$$R = \left\{ \frac{\partial^2 z}{\partial x_i \partial x_a} = 0 \quad (1 \leq i \leq r, 1 \leq a \leq n) \right\}.$$

Here $\{e_1, \dots, e_n\}$ is a basis of V , $V_r = \langle \{e_1, \dots, e_r\} \rangle$ and $V_s = \langle \{e_{r+1}, \dots, e_n\} \rangle$.

We need Reduction Theorems to explain why second order equations with these symbols have the property that their symbols are the only invariants under contact transformations. We will explain this fact for the type $\mathfrak{f}^3(r)$ in §6.1 by utilizing the First Reduction Theorem in §4. The other cases will be explained by utilizing the two step reduction procedure in Part II.

4. PD MANIFOLDS OF SECOND ORDER.

We will here formulate the submanifold theory for $(L(J), E)$ as the geometry of *PD* manifolds of second order ([Y1]) and discuss the First Reduction Theorem.

4.1. Realization Theorem. Let R be a submanifold of $L(J)$ satisfying the following condition:

$$(R.0) \quad p : R \rightarrow J ; \text{ submersion,}$$

where $p = \pi|_R$ and $\pi : L(J) \rightarrow J$ is the projection. Let D^1 and D^2 be differential systems on R obtained by restricting $C^1 = \partial E$ and $C^2 = E$ to R . Moreover we denote by the same symbols those 1-forms obtained by restricting the defining 1-forms $\{\varpi, \varpi_1, \dots, \varpi_n\}$ of the canonical system E to R . Then it follows from (R.0) that these 1-forms are independent at each point on R and that

$$D^1 = \{\varpi = 0\}, \quad D^2 = \{\varpi = \varpi_1 = \dots = \varpi_n = 0\}.$$

In fact $(R; D^1, D^2)$ further satisfies the following conditions:

$$(R.1) \quad D^1 \text{ and } D^2 \text{ are differential systems of codimension 1 and } n+1 \text{ respectively.}$$

$$(R.2) \quad \partial D^2 \subset D^1.$$

$$(R.3) \quad \text{Ch}(D^1) \text{ is a subbundle of } D^2 \text{ of codimension } n.$$

$$(R.4) \quad \text{Ch}(D^1)(v) \cap \text{Ch}(D^2)(v) = \{0\} \quad \text{at each } v \in R.$$

Here (R.2) follows from $d\varpi \equiv 0 \pmod{\varpi, \varpi_1, \dots, \varpi_n}$. (R.3) follows from $\text{Ch}(D^1) = \text{Ker } p_* = \{ dz = dx_1 = \dots = dx_n = dp_1 = \dots = dp_n = 0 \}$. Moreover the last condition follows easily from the Realization Lemma below.

Conversely these four conditions characterize submanifolds in $L(J)$ satisfying (R.0). To see this, we first recall the following **Realization Lemma**, which characterizes submanifolds of $(J(M, n), C)$.

Realization Lemma. *Let R and M be manifolds. Assume that the quadruple (R, D, p, M) satisfies the following conditions :*

- (1) p is a map of R into M of constant rank.
- (2) D is a differential system on R such that $F = \text{Ker } p_*$ is a subbundle of D of codimension n .

Then there exists a unique map ψ of R into $J(M, n)$ satisfying $p = \pi \cdot \psi$ and $D = \psi_*^{-1}(C)$, where C is the canonical differential system on $J(M, n)$ and $\pi : J(M, n) \rightarrow M$ is the projection. Furthermore, let v be any point of R . Then ψ is in fact defined by

$$\psi(v) = p_*(D(v)) \quad \text{as a point of } \text{Gr}(T_{p(v)}(M)),$$

and satisfies

$$\text{Ker}(\psi_*)_v = F(v) \cap \text{Ch}(D)(v).$$

where $\text{Ch}(D)$ is the Cauchy Characteristic System of D .

For the proof, see Lemma 1.5 [Y1].

In view of this Lemma, we call the triplet $(R; D^1, D^2)$ of a manifold and two differential systems on it a **PD manifold of second order** if these satisfy the above four conditions (R.1) to (R.4). Here we note, by (R.2), subbundles D^2 , D^1 and $T(R)$ define a filtration on R . Hence we can form the symbol algebra $\mathfrak{s}(v) = \mathfrak{s}_{-3}(v) \oplus \mathfrak{s}_{-2}(v) \oplus \mathfrak{s}_{-1}(v)$ of $(R; D^1, D^2)$ at $v \in R$ as in §3.1.

We have the (local) Realization Theorem for PD manifolds as follows: From conditions (R.1) and (R.3), it follows that the codimension of the foliation defined by the completely integrable system $\text{Ch}(D^1)$ is $2n + 1$. Assume that R is regular with respect to $\text{Ch}(D^1)$, i.e., the space $J = R/\text{Ch}(D^1)$ of leaves of this foliation is a manifold of dimension $2n + 1$ such that each fibre of the projection $p : R \rightarrow J = R/\text{Ch}(D^1)$ is connected and p is a submersion. Then D^1 drops down to J . Namely there exists a differential system C on J of codimension 1 such that $D^1 = p_*^{-1}(C)$. From $\text{Ch}(C) = \{0\}$, (J, C) becomes a contact manifold of dimension $2n + 1$. Conditions (R.1) and (R.2) guarantees that the image of the following map ι is a legendrian subspace of (J, C) :

$$\iota(v) = p_*(D^2(v)) \subset C(u), \quad u = p(v).$$

Finally the condition (R.4) shows that $\iota : R \rightarrow L(J)$ is an immersion by Realization Lemma for (R, D^2, p, J) . Furthermore we have (Corollary 5.4 [Y1])

Theorem 4.1. *Let $(R; D^1, D^2)$ and $(\hat{R}; \hat{D}^1, \hat{D}^2)$ be PD manifolds of second order. Assume that R and \hat{R} are regular with respect to $\text{Ch}(D^1)$ and $\text{Ch}(\hat{D}^1)$ respectively. Let (J, C) and (\hat{J}, \hat{C}) be the associated contact manifolds. Then an isomorphism $\Phi : (R; D^1, D^2) \rightarrow (\hat{R}; \hat{D}^1, \hat{D}^2)$ induces a contact diffeomorphism $\varphi : (J, C) \rightarrow (\hat{J}, \hat{C})$ such that the following commutes;*

$$\begin{array}{ccc}
R & \xrightarrow{\iota} & L(J) \\
\Phi \downarrow & & \downarrow \varphi_* \\
\hat{R} & \xrightarrow{\hat{\iota}} & L(\hat{J}).
\end{array}$$

Proof. Since Φ is an isomorphism of (R, D^1, D^2) onto $(\hat{R}, \hat{D}^1, \hat{D}^2)$, we have $\Phi_*(D^1) = \hat{D}^1$. Hence we get $\Phi_*(\text{Ch}(D^1)) = \text{Ch}(\hat{D}^1)$. Therefore, since $\text{Ch}(D^1) = \text{Ker } p_*$ and each fibre of $p : R \rightarrow J$ is connected, Φ is fibre-preserving and induces a unique diffeomorphism φ of J onto \hat{J} such that $\hat{p} \cdot \Phi = \varphi \cdot p$. By $D^1 = (p_*)^{-1}(C)$ and $\hat{D}^1 = (\hat{p}_*)^{-1}(\hat{C})$, φ is a contact diffeomorphism of J onto \hat{J} . Put $\bar{\iota} = (\varphi_*)^{-1} \cdot \hat{\iota} \cdot \Phi$. Then it is easy to see that $\bar{\iota}$ is a map of R into $L(J)$ satisfying $\pi \cdot \bar{\iota} = p$ and $D^2 = (\bar{\iota}_*)^{-1}(C^2)$. Therefore by the uniqueness of the canonical immersion ι of R into $L(J)$, we obtain $\iota = \bar{\iota}$, i.e., $\hat{\iota} \cdot \Phi = \varphi_* \cdot \iota$. \square

By this theorem, the submanifold theory for $(L(J), E)$ is reformulated as the geometry of PD manifolds of second order.

4.2. First Reduction Theorem. When $D^1 = \partial D^2$ holds for a PD manifold $(R; D^1, D^2)$ of second order, the geometry of $(R; D^1, D^2)$ reduces to that of (R, D^2) and the Tanaka theory is directly applicable to this case. We will treat this case as Parabolic Geometries associated with PD manifolds of second order in §5. Concerning about this situation, we will show the following proposition under the compatibility condition (C) :

$$(C) \quad p^{(1)} : R^{(1)} \rightarrow R \text{ is onto.}$$

where $R^{(1)}$ is the first prolongation of $(R; D^1, D^2)$, i.e.,

$$R^{(1)} = \{n\text{-dim. integral elements of } (R, D^2), \text{ transversal to } F = \text{Ker } p_*\} \subset J(R, n),$$

(cf. Proposition 5.11 [Y1]).

Proposition 4.1. *Let $(R; D^1, D^2)$ be a PD manifold of second order satisfying the condition (C) above. Then the following equality holds at each point v of R :*

$$\dim D^1(v) - \dim \partial D^2(v) = \dim \text{Ch}(D^2)(v).$$

In particular $D^1 = \partial D^2$ holds if and only if $\text{Ch}(D^2) = \{0\}$.

Proof. By the condition (C), there exists an n -dimensional integral element V of (R, D^2) at v such that

$$D^2(v) = \mathfrak{s}_{-1}(v) = V \oplus \mathfrak{f}(v).$$

In terms of the symbol algebra $\mathfrak{s}(v) = \mathfrak{s}_{-3}(v) \oplus \mathfrak{s}_{-2}(v) \oplus \mathfrak{s}_{-1}(v)$, we have

$$\text{Ch}(D^2)(v) = \{X \in \mathfrak{s}_{-1}(v) \mid [X, \mathfrak{s}_{-1}(v)] = 0\}$$

For a vector $X = w_X + f_X \in \text{Ch}(D^2)(v)$, where $w_X \in V$ and $f_X \in \mathfrak{f}(v)$, $[X, V] = 0$ forces $f_X = 0$, that is, $\text{Ch}(D^2)(v) \subset V \subset \mathfrak{s}_{-1}(v)$. We put $\mathcal{C}(v) = \text{Ch}(D^2)(v) \subset V$. Then we have

$$\mathcal{C}(v) = \{w \in V \mid w \rfloor f = 0 \text{ for all } f \in \mathfrak{f}(v)\} \subset V.$$

Thus $\mathcal{C}(v) = E$ is the largest subspace of V such that $\mathfrak{f}(v) \subset S^2(E^\perp)$. On the other hand, let us consider the derived system ∂D^2 in terms of the symbol algebra $\mathfrak{s}(v)$. From $[\mathfrak{s}_{-1}(v), \mathfrak{s}_{-1}(v)] = [V, \mathfrak{f}(v)]$, putting $\mathcal{D}(v) = \pi_2(\partial D^2(v))$, we have

$$\mathcal{D}(v) = \{z \rfloor f \mid z \in V, f \in \mathfrak{f}(v)\} \subset V^*,$$

under the identification $\mathfrak{s}_{-2}(v) \cong V^*$, where $\pi_2 : D^1(v) \rightarrow \mathfrak{s}_{-2}(v)$ is the projection. Thus w belongs to the annihilator of $\mathcal{D}(v)$ iff $\langle w, z \rfloor f \rangle = 0$ for all $z \in V$ and all $f \in \mathfrak{f}(v)$, hence

iff $w \rfloor f = 0$ for all $f \in \mathfrak{f}(v)$. This implies $\mathcal{D}(v) = (\mathcal{C}(v))^\perp$, which completes the proof of Proposition. \square

When a PD manifold $(R; D^1, D^2)$ admits a non-trivial Cauchy characteristics, i.e., when $\text{rank Ch}(D^2) > 0$, the geometry of $(R; D^1, D^2)$ is further reducible to the geometry of a single differential system. Here we will be concerned with the local equivalence of $(R; D^1, D^2)$, hence we may assume that R is regular with respect to $\text{Ch}(D^2)$, i.e., the leaf space $X = R/\text{Ch}(D^2)$ is a manifold such that the projection $\rho : R \rightarrow X$ is a submersion and there exists a differential system D on X satisfying $D^2 = \rho_*^{-1}(D)$. Then the local equivalence of $(R; D^1, D^2)$ is further reducible to that of (X, D) as in the following: We assume that $(R; D^1, D^2)$ satisfies the condition (C) above and $\text{Ch}(D^2)$ is a subbundle of rank r ($0 < r < n$). Then, by Proposition 4.1, ∂D^2 is a subbundle of D^1 of codimension r .

By the information of the symbol algebra $\mathfrak{s}(v)$ of $(R; D^1, D^2)$ at $v \in R$, we can find locally independent 1-forms $\varpi, \varpi_i, \omega_i$ ($i = 1, \dots, n$) around v such that

$$D^1 = \{\varpi = 0\},$$

$$\partial D^2 = \{\varpi = \varpi_1 = \dots = \varpi_r = 0\}, \quad D^2 = \{\varpi = \varpi_1 = \dots = \varpi_n = 0\}$$

$$\begin{cases} d\varpi \equiv \omega_1 \wedge \varpi_1 + \dots + \omega_n \wedge \varpi_n \pmod{\varpi} \\ d\varpi_i \equiv 0 \pmod{\varpi, \varpi_1, \dots, \varpi_n} & i = 1, \dots, r \\ d\varpi_\alpha \not\equiv 0 \pmod{\varpi, \varpi_1, \dots, \varpi_n} & \alpha = r+1, \dots, n \end{cases}$$

Thus we have

$$\begin{cases} d\varpi \equiv \omega_{r+1} \wedge \varpi_{r+1} + \dots + \omega_n \wedge \varpi_n \pmod{\varpi, \varpi_1, \dots, \varpi_r} \\ d\varpi_i \equiv \pi_i^{r+1} \wedge \varpi_{r+1} + \dots + \pi_i^n \wedge \varpi_n \pmod{\varpi, \varpi_1, \dots, \varpi_r} & i = 1, \dots, r \end{cases}$$

for some 1-forms π_i^α . This shows that the Cartan rank of $(X, \partial D)$ (see [BCG₃] II §4) equals to $s = n - r$ at $x = \rho(v) \in X$, which gives us a necessary condition for a differential system (X, D) to be obtained from a PD manifold $(R; D^1, D^2)$ as $X = R/\text{Ch}(D^2)$.

From (X, D) , at least locally, we can reconstruct the PD manifold $(R; D^1, D^2)$ as follows. First let us consider the collection $P(X)$ of hyperplanes v in each tangent space $T_x(X)$ at $x \in X$ which contains the fibre $\partial D(x)$ of the derived system ∂D of D .

$$P(X) = \bigcup_{x \in X} P_x \subset J(X, m-1),$$

$$P_x = \{v \in \text{Gr}(T_x(X), m-1) \mid v \supset \partial D(x)\} \cong \mathbb{P}(T_x(X)/\partial D(x)) = \mathbb{P}^r,$$

where $m = \dim X$ and $r = \text{rank Ch}(D^2)$. Moreover D_X^1 is the canonical system obtained by the Grassmannian construction and D_X^2 is the lift of D . Precisely, D_X^1 and D_X^2 are given by

$$D_X^1(v) = \nu_*^{-1}(v) \supset D_X^2(v) = \nu_*^{-1}(D(x)),$$

for each $v \in P(X)$ and $x = \rho(v)$, where $\nu : P(X) \rightarrow X$ is the projection. Then we have a map κ of R into $P(X)$ given by

$$\kappa(v) = \rho_*(D^1(v)) \subset T_x(X),$$

for each $v \in R$ and $x = \rho(v)$. By Realization Lemma for (R, D^1, ρ, X) , κ is a map of constant rank such that

$$\text{Ker } \kappa_* = \text{Ch}(D^1) \cap \text{Ker } \rho_* = \text{Ch}(D^1) \cap \text{Ch}(D^2) = \{0\}.$$

Thus κ is an immersion and, by a dimension count, in fact, a local diffeomorphism of R into $P(X)$ such that

$$\kappa_*(D^1) = D_X^1 \quad \text{and} \quad \kappa_*(D^2) = D_X^2.$$

Namely $\kappa : (R, D^1, D^2) \rightarrow (P(X), D_X^1, D_X^2)$ is a local isomorphism of PD manifolds of second order. Thus $(R; D^1, D^2)$ is reconstructed from (X, D) , at least locally, as a part of $(P(X); D_X^1, D_X^2)$. (Precisely, in general, $(P(X), D_X^1, D_X^2)$ becomes a PD manifold on an open subset. See Proposition 4.2.) By the construction of $(P(X); D_X^1, D_X^2)$, an isomorphism of (X, D) naturally lifts to an isomorphism of $(P(X); D_X^1, D_X^2)$.

Summarizing the above consideration, we obtain the following First Reduction Theorem for PD manifolds admitting non-trivial Cauchy characteristics.

Theorem 4.2. *Let (R, D^1, D^2) and $(\hat{R}; \hat{D}^1, \hat{D}^2)$ be PD manifolds satisfying the condition (C) such that $Ch(D^2)$ and $Ch(\hat{D}^2)$ are subbundles of rank r ($0 < r < n$). Assume that R and \hat{R} are regular with respect to $Ch(D^2)$ and $Ch(\hat{D}^2)$ respectively. Let (X, D) and (\hat{X}, \hat{D}) be the leaf spaces, where $X = R/Ch(D^2)$ and $\hat{X} = \hat{R}/Ch(\hat{D}^2)$. Let us fix points $v_o \in R$ and $\hat{v}_o \in \hat{R}$ and put $x_o = \rho(v_o)$ and $\hat{x}_o = \hat{\rho}(\hat{v}_o)$. Then a local isomorphism $\psi : (R; D^1, D^2) \rightarrow (\hat{R}; \hat{D}^1, \hat{D}^2)$ such that $\psi(v_o) = \hat{v}_o$ induces a local isomorphism $\varphi : (X, D) \rightarrow (\hat{X}, \hat{D})$ such that $\varphi(x_o) = \hat{x}_o$ and $\varphi_*(\kappa(x_o)) = \hat{\kappa}(\hat{x}_o)$, and vice versa.*

4.3. Construction of $(R(X); D_X^1, D_X^2)$. Now we will characterize differential systems (X, D) , which are obtained by the First Reduction Theorem from PD manifolds $(R; D^1, D^2)$ as $X = R/Ch(D^2)$. We already saw that the necessary condition for (X, D) is that ∂D is of Cartan rank $s = n - r$. We will show that this condition is also sufficient.

Let (X, D) be a differential system satisfying the following conditions;

- (X.1) D is a differential system of codimension $n + 1$ such that $Ch(D)$ is trivial.
- (X.2) ∂D is a differential system of codimension $r + 1$.
- (X.3) ∂D is of Cartan rank $s = n - r$.

Under the conditions (X.1) and (X.2), Cartan rank of ∂D is less than or equal to s (see the proof of Proposition 4.2 below). Thus (X.3) is a nondegeneracy condition for (X, D) .

We form the weak symbol algebra $\mathfrak{t}(x)$ of (X, D) at $x \in X$ as follows: Put $\mathfrak{t}_{-3}(x) = T_x(X)/\partial D(x)$, $\mathfrak{t}_{-2}(x) = \partial D(x)/D(x)$ and $\mathfrak{t}_{-1}(x) = D(x)$. The subbundles D , ∂D and $T(X)$ give a filtration on X . Hence as in the symbol algebra of PD manifolds, we can introduce the Lie brackets in

$$\mathfrak{t}(x) = \mathfrak{t}_{-3}(x) \oplus \mathfrak{t}_{-2}(x) \oplus \mathfrak{t}_{-1}(x),$$

so that $\mathfrak{t}(x)$ becomes a graded Lie algebra.

Now let us consider the collection $P(X)$ of hyperplanes v in each tangent space $T_x(X)$ at $x \in X$ which contains the fibre $\partial D(x)$ of the derived system ∂D of D .

$$P(X) = \bigcup_{x \in X} P_x \subset J(X, m - 1),$$

$$P_x = \{v \in \text{Gr}(T_x(X), m - 1) \mid v \supset \partial D(x)\} \cong \mathbb{P}(T_x(X)/\partial D(x)) = \mathbb{P}^r,$$

where $m = \dim X$ and $r + 1 = \text{codim } \partial D$. Moreover D_X^1 is the canonical system obtained by the Grassmannian construction and D_X^2 is the lift of D . In fact, D_X^1 and D_X^2 are given by

$$D_X^1(v) = \nu_*^{-1}(v) \supset D_X^2(v) = \nu_*^{-1}(D(x)),$$

for each $v \in P(X)$ and $x = \nu(v)$, where $\nu : P(X) \rightarrow X$ is the projection. For a point $v \in P(X)$, we define the **symbol subspace** $\hat{f}(v)$ of $D(x)$ by

$$\hat{f}(v) = \{X \in \mathfrak{t}_{-1}(x) \mid [X, \mathfrak{t}_{-2}(x)] \subset \hat{v}\},$$

where $x = \nu(v) \in X$, $\hat{v} = p_{-3}(v) \subset \mathfrak{t}_{-3}(x)$ and $p_{-3} : T_x(X) \rightarrow \mathfrak{t}_{-3} = T_x(X)/\partial D(x)$ is the projection. We put

$$R(X) = \{v \in P(X) \mid \text{codim } \hat{f}(v) = s\}$$

($R(X)$ is an open subset of $P(X)$ under the condition (X.3). See below). We denote the restrictions of differential systems D_X^1 and D_X^2 of $P(X)$ to $R(X)$ by the same symbols.

Then we have

Proposition 4.2. *($R(X), D_X^1, D_X^2$) is a PD manifold of second order.*

Proof. By (X.1) and the construction of $P(X)$, it follows that D_X^1 and D_X^2 are differential systems on $P(X)$ of codimension 1 and $n+1$ respectively. Moreover $\partial D_X^2 \subset D_X^1$ holds on $P(X)$ by construction. Since $P(X)$ is a submanifold of $J(X, m-1)$, Realization Lemma for $(P(X); D_X^1, \nu, X)$ implies

$$\text{Ker } \bar{\iota}_* = \text{Ker } \nu_* \cap \text{Ch}(D_X^1) = \text{Ch}(D_X^1) \cap \text{Ch}(D_X^2) = \{0\},$$

where $\bar{\iota} : P(X) \rightarrow J(X, m-1)$ is the inclusion. Thus it remains to show that $R(X)$ is an open subset of $P(X)$ under the condition (X.3) and $\text{Ch}(D_X^1)$ is a subbundle of D_X^2 of codimension n on $R(X)$.

For this purpose, let us take a point $v_o \in P(X)$. We can find locally independent 1-forms $\varpi_0, \dots, \varpi_r, \pi_1, \dots, \pi_s$ on a neighborhood U of $x_o = \nu(v_o) \in X$ such that

$$\partial D = \{\varpi_0 = \dots = \varpi_r = 0\}, \quad D = \{\varpi_0 = \dots = \varpi_r = \pi_1 = \dots = \pi_s = 0\}.$$

Here we may assume that $v_o = \{\varpi_0 = 0\} \subset T_{x_o}(X)$. Then we have

$$d\varpi_i \equiv 0 \pmod{\varpi_0, \dots, \varpi_r, \pi_1, \dots, \pi_s} \quad \text{for } i = 0, \dots, r.$$

Hence there exist 1-forms β_i^α such that

$$d\varpi_i \equiv \beta_i^1 \wedge \pi_1 + \dots + \beta_i^s \wedge \pi_s \pmod{\varpi_0, \dots, \varpi_r} \quad \text{for } i = 0, \dots, r.$$

Thus the Cartan rank of ∂D is less than or equal to s .

Now let us consider

$$\varpi = \varpi_0 + \lambda_1 \varpi_1 + \dots + \lambda_r \varpi_r$$

on U . Namely we consider a point $v \in P(X)$ such that $v = \{\varpi = 0\} \subset T_x(X)$, where $x = \nu(v) \in X$. Here $(\lambda_1, \dots, \lambda_r)$ constitutes an inhomogeneous coordinate of the fibres of $\nu : P(X) \rightarrow X$. Denoting the pullback on $P(X)$ of 1-forms on X by the same symbol, we have

$$D_X^1 = \{\varpi = 0\},$$

and

$$d\varpi = d\varpi_0 + \sum_{i=1}^r \lambda_i d\varpi_i + \sum_{i=1}^r d\lambda_i \wedge \varpi_i.$$

on $\nu^{-1}(U)$. From $d\varpi_i \equiv \sum_{\alpha=1}^s \beta_i^\alpha \wedge \pi_\alpha + \sum_{j=1}^r \gamma_i^j \wedge \varpi_j \pmod{\varpi}$ for $i = 0, \dots, r$, we calculate

$$d\varpi \equiv \sum_{\alpha=1}^s (\beta_0^\alpha + \sum_{i=1}^r \lambda_i \beta_i^\alpha) \wedge \pi_\alpha + \sum_{i=1}^r (d\lambda_i + \gamma_0^i + \sum_{j=1}^r \lambda_j \gamma_j^i) \wedge \varpi_i \pmod{\varpi}.$$

For a vector $Y \in \partial D(x)$, we have

$$Y \rfloor d\varpi \equiv - \sum_{\alpha=1}^s \pi_\alpha(Y) (\beta_0^\alpha + \sum_{i=1}^r \lambda_i \beta_i^\alpha) \pmod{\varpi_0, \dots, \varpi_r, \pi_1, \dots, \pi_s}.$$

By the definition of brackets in the weak symbol algebra $\mathfrak{t}(x)$, it follows that

$$\hat{\mathfrak{f}}(v) = \{X \in D(x) \mid (\beta_0^\alpha + \sum_{i=1}^r \lambda_i \beta_i^\alpha)(X) = 0 \text{ for } \alpha = 1, \dots, s\}.$$

Hence we see that $\text{codim } \hat{\mathfrak{f}}(v) = s$ if and only if $\{\beta_0^\alpha + \sum_{i=1}^r \lambda_i \beta_i^\alpha\}_{\alpha=1}^s$ are independent $\pmod{\varpi_0, \dots, \varpi_r, \pi_1, \dots, \pi_s}$ at $x \in X$. Thus $R(X)$ is a non-empty open subset of $P(X)$ under the condition (X.3). Moreover we have, at each $v \in R(X)$,

$$\begin{aligned} \text{Ch}(D_X^1)(v) = \{ \varpi = \pi_\alpha = \beta_0^\alpha + \sum_{i=1}^r \lambda_i \beta_i^\alpha = \varpi_i = d\lambda_i + \gamma_0^i + \sum_{j=1}^r \lambda_j \gamma_j^i = 0 \\ (i = 1, \dots, r, \alpha = 1, \dots, s) \}. \end{aligned}$$

Therefore $\text{Ch}(D_X^1)$ is a subbundle of D_X^2 of codimension n on $R(X)$, which completes the proof of Proposition. \square

4.4. Symbol Subspaces. We will consider the relation between the symbol subspaces $\mathfrak{f}(v) = \text{Ch}(D_X^1)(v)$ and $\hat{\mathfrak{f}}(v) \in D(x)$ for a point $v \in R(X)$ and $x = \nu(v) \in X$.

First observe that there exists an integral element V of $(R(X), D_X^2)$ at v such that

$$\mathfrak{s}_{-1}(v) = V \oplus \mathfrak{f}(v) (= D_X^2(v)),$$

if and only if there exists an integral element W of (X, D) at x such that

$$\mathfrak{t}_{-1}(x) = W \oplus \hat{\mathfrak{f}}(v) (= D(x)).$$

In fact $W = \nu_*(V)$ and $V = \nu_*^{-1}(W)$, where $\nu_* : D_X^2(v) \rightarrow D(x)$ is onto, $\text{Ker } \nu_* = \text{Ch}(D_X^2)(v)$ and $\nu_* : \mathfrak{f}(v) \rightarrow \hat{\mathfrak{f}}(v)$ is a linear isomorphism.

Put $\hat{\mathfrak{t}}_{-3}(v) = \mathfrak{t}_{-3}(x)/\hat{v}$, and let us consider

$$\hat{\mathfrak{t}}(v) = \hat{\mathfrak{t}}_{-3}(v) \oplus \mathfrak{t}_{-2}(x) \oplus \mathfrak{t}_{-1}(x).$$

$\hat{\mathfrak{t}}(v)$ is a quotient algebra of $\mathfrak{t}(x)$ and $\hat{\mathfrak{f}}(v) = \{X \in \mathfrak{t}_{-1}(x) \mid [X, \mathfrak{t}_{-2}(x)] = 0\}$ in $\hat{\mathfrak{t}}(v)$. Let us fix a basis of $\hat{\mathfrak{t}}_{-3}(v)$. Then, as in the proof of Proposition 4.2, the basis of $\mathfrak{s}_{-3}(v)$ is fixed. Hence we have $\mathfrak{s}_{-3}(v) \cong \mathbb{R}$, $\mathfrak{s}_{-2}(v) \cong V^*$ and $\mathfrak{f}(v) \subset S^2(V^*)$. Moreover we have $\hat{\mathfrak{t}}_{-3}(v) \cong \mathbb{R}$, $\mathfrak{t}_{-2}(x) \cong W^*$ and the map $\mu : \hat{\mathfrak{f}}(v) \rightarrow S^2(W^*)$ is defined by

$$\mu(\tilde{f})(w_1, w_2) = [[\tilde{f}, w_1], w_2] \in \mathbb{R} \cong \hat{\mathfrak{t}}_{-3}(v) \quad \text{for } \tilde{f} \in \hat{\mathfrak{f}}(v).$$

Here we note that there exists a unique $f \in \mathfrak{f}(v)$ such that $\tilde{f} = \nu_*(f)$ and we obtain, by $\mathfrak{f}(v) \subset S^2(E^\perp)$ and the definition of the brackets of $\mathfrak{s}(v)$ and $\hat{\mathfrak{t}}(v)$,

$$[f, v_1] = \kappa^*([\tilde{f}, w_1]) \in V^*,$$

where $w_1 = \kappa(v_1)$, $\kappa^* : W^* \rightarrow V^*$, $\kappa = \nu_* : V \rightarrow W$ and $\text{Ker } \kappa = E$, $\kappa^*(W^*) = E^\perp$. Moreover we have

$$[[f, v_1], v_2] = [[\tilde{f}, w_1], w_2] \quad \text{for } w_2 = \kappa(v_2).$$

Namely μ is injective and $\kappa_2^*(\hat{\mathfrak{f}}(v)) = \mathfrak{f}(v) \subset S^2(E^\perp) \subset S^2(V^*)$ where $\kappa_2^* : S^2(W^*) \rightarrow S^2(V^*)$ is induced from $\kappa^* : W^* \rightarrow V^*$.

Next we consider the algebraic prolongation $\mathfrak{f}(v)^{(1)} \subset S^3(V^*)$ of $\mathfrak{f}(v) \subset S^2(V^*)$,

$$\mathfrak{f}(v)^{(1)} = \mathfrak{f}(v) \otimes V^* \cap S^3(V^*).$$

Then, since $\mathfrak{f}(v) \subset S^2(E^\perp)$, we observe $\mathfrak{f}(v)^{(1)} \subset S^3(E^\perp)$. Therefore, for the prolongation $\hat{\mathfrak{f}}(v)^{(1)} = \hat{\mathfrak{f}}(v) \otimes W^* \cap S^3(W^*)$ of $\hat{\mathfrak{f}}(v) \subset S^2(W^*)$, we get

$$\kappa_3^*(\hat{\mathfrak{f}}(v)^{(1)}) = \mathfrak{f}(v)^{(1)} \subset S^3(E^\perp).$$

where $\kappa_3^* : S^3(W^*) \rightarrow S^3(V^*)$ is induced from $\kappa^* : W^* \rightarrow V^*$. Repeatedly we obtain $\kappa_{k+2}^*(\hat{\mathfrak{f}}(v)^{(k)}) = \mathfrak{f}(v)^{(k)} \subset S^{k+2}(E^\perp)$ for the higher prolongations.

Moreover, from $\mathfrak{f}(v) \subset S^2(E^\perp)$, we also observe that, for a regular basis $\{w_1, \dots, w_s\}$ of W for $\hat{\mathfrak{f}}(v) \subset S^2(W^*)$, we obtain the regular basis $\{v_1, \dots, v_s, v_{s+1}, \dots, v_n\}$ of V for $\mathfrak{f}(v) \subset S^2(V^*)$ by taking $v_i \in V$ such that $\kappa(v_i) = w_i$ ($i = 1, \dots, s$) and adding a basis $\{v_{s+1}, \dots, v_n\}$ of $E = \text{Ker } \kappa$. In fact, in this case, we have

$$\kappa_2^*(\hat{\mathfrak{f}}_k) = \mathfrak{f}_k \quad \text{for } k = 1, \dots, s \quad \text{and} \quad \mathfrak{f}_k = 0 \quad \text{for } k = s+1, \dots, n.$$

where $\hat{\mathfrak{f}}_k = \{\tilde{f} \in \hat{\mathfrak{f}}(v) \mid w_1 \lrcorner \tilde{f} = \dots = w_k \lrcorner \tilde{f} = 0\}$ and $\mathfrak{f}_k = \{f \in \mathfrak{f}(v) \mid v_1 \lrcorner f = \dots = v_k \lrcorner f = 0\}$.

Summarizing the above discussion, we obtain

Proposition 4.3. *Notations being as above:*

- (1) $\mathfrak{f}(v) \subset S^2(V^*)$ is involutive if and only if $\hat{\mathfrak{f}}(v) \subset S^2(W^*)$ is involutive.
- (2) $\mathfrak{f}(v) \subset S^2(V^*)$ is of finite type if and only if $\hat{\mathfrak{f}}(v) \subset S^2(W^*)$ is of finite type.

5. PARABOLIC GEOMETRIES ASSOCIATED WITH PD MANIFOLDS OF SECOND ORDER.

We will here exhibit Parabolic Geometries which directly correspond to the geometry of PD manifolds of second order, following [Y5] and [YY2].

5.1. Differential Systems associated with Simple Graded Lie Algebras (Parabolic Geometries). We first recall basic materials for simple graded Lie algebras over \mathbb{C} and state the Prolongation Theorem. We will work mainly over \mathbb{C} in this section for the sake of simplicity.

Let \mathfrak{g} be a finite dimensional simple Lie algebra over \mathbb{C} . Let us fix a Cartan subalgebra \mathfrak{h} of \mathfrak{g} and choose a simple root system $\Delta = \{\alpha_1, \dots, \alpha_\ell\}$ of the root system Φ of \mathfrak{g} relative to \mathfrak{h} . Then every $\alpha \in \Phi$ is an (all non-negative or all non-positive) integer coefficient linear combination of elements of Δ and we have the root space decomposition of \mathfrak{g} ;

$$\mathfrak{g} = \bigoplus_{\alpha \in \Phi^+} \mathfrak{g}_\alpha \oplus \mathfrak{h} \oplus \bigoplus_{\alpha \in \Phi^-} \mathfrak{g}_{-\alpha},$$

where $\mathfrak{g}_\alpha = \{X \in \mathfrak{g} \mid [h, X] = \alpha(h)X \text{ for } h \in \mathfrak{h}\}$ is (1-dimensional) root space (corresponding to $\alpha \in \Phi$) and Φ^+ denotes the set of positive roots.

Now let us take a nonempty subset Δ_1 of Δ . Then Δ_1 defines the partition of Φ^+ as in the following and induces the gradation of $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ as follows:

$$\begin{aligned} \Phi^+ &= \bigcup_{p \geq 0} \Phi_p^+, & \Phi_p^+ &= \left\{ \alpha = \sum_{i=1}^{\ell} n_i \alpha_i \mid \sum_{\alpha_i \in \Delta_1} n_i = p \right\}, \\ \mathfrak{g}_p &= \bigoplus_{\alpha \in \Phi_p^+} \mathfrak{g}_\alpha, & \mathfrak{g}_0 &= \bigoplus_{\alpha \in \Phi_0^+} \mathfrak{g}_\alpha \oplus \mathfrak{h} \oplus \bigoplus_{\alpha \in \Phi_0^+} \mathfrak{g}_{-\alpha}, & \mathfrak{g}_{-p} &= \bigoplus_{\alpha \in \Phi_p^+} \mathfrak{g}_{-\alpha}, \end{aligned}$$

$$[\mathfrak{g}_p, \mathfrak{g}_q] \subset \mathfrak{g}_{p+q} \quad \text{for } p, q \in \mathbb{Z}.$$

Moreover the negative part $\mathfrak{m} = \bigoplus_{p < 0} \mathfrak{g}_p$ satisfies the following generating condition :

$$\mathfrak{g}_p = [\mathfrak{g}_{p+1}, \mathfrak{g}_{-1}] \quad \text{for } p < -1$$

We denote the simple graded Lie algebra $\mathfrak{g} = \bigoplus_{p=-\mu}^{\mu} \mathfrak{g}_p$ obtained from Δ_1 in this manner by (X_ℓ, Δ_1) , when \mathfrak{g} is a simple Lie algebra of type X_ℓ . Here X_ℓ stands for the Dynkin diagram of \mathfrak{g} representing Δ and Δ_1 is a subset of vertices of X_ℓ . Moreover we have

$$\mu = \sum_{\alpha_i \in \Delta_1} n_i(\theta),$$

where $\theta = \sum_{i=1}^{\ell} n_i(\theta) \alpha_i$ is the highest root of Φ^+ .

Conversely we have (Theorem 3.12 [Y5])

Theorem A. *Let $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ be a simple graded Lie algebra over \mathbb{C} satisfying the generating condition. Let X_ℓ be the Dynkin diagram of \mathfrak{g} . Then $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ is isomorphic to a graded Lie algebra (X_ℓ, Δ_1) for some $\Delta_1 \subset \Delta$. Moreover (X_ℓ, Δ_1) and (X_ℓ, Δ'_1) are isomorphic if and only if there exists a diagram automorphism ϕ of X_ℓ such that $\phi(\Delta_1) = \Delta'_1$.*

In the real case, we can utilize the Satake diagram of \mathfrak{g} to describe gradations of \mathfrak{g} (Theorem 3.12 [Y5]).

By Theorem A, the classification of the gradation $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ of \mathfrak{g} satisfying the generating condition coincides with that of parabolic subalgebras $\mathfrak{g}' = \bigoplus_{p \geq 0} \mathfrak{g}_p$ of \mathfrak{g} . Accordingly, to each simple graded Lie algebra (X_ℓ, Δ_1) , there corresponds a unique R -space $M_{\mathfrak{g}} = G/G'$ (compact simply connected homogeneous complex manifold) (see [Y5] §4.1 for detail). Furthermore, when $\mu \geq 2$, there exists the G -invariant differential system $D_{\mathfrak{g}}$ on $M_{\mathfrak{g}}$, which is induced from \mathfrak{g}_{-1} , and the standard differential system $(M(\mathfrak{m}), D_{\mathfrak{m}})$ of type \mathfrak{m} becomes an open submanifold of $(M_{\mathfrak{g}}, D_{\mathfrak{g}})$. For the Lie algebras of all infinitesimal automorphisms of $(M_{\mathfrak{g}}, D_{\mathfrak{g}})$, hence of $(M(\mathfrak{m}), D_{\mathfrak{m}})$, we have the following **Prolongation Theorem** (Theorem 5.2 [Y5]).

Prolongation Theorem. *Let $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ be a simple graded Lie algebra over \mathbb{C} satisfying the generating condition. Then $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ is the prolongation of $\mathfrak{m} = \bigoplus_{p < 0} \mathfrak{g}_p$ except for the following three cases.*

- (1) $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is of depth 1.
- (2) $\mathfrak{g} = \bigoplus_{p=-2}^2 \mathfrak{g}_p$ is a contact gradation.
- (3) $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ is isomorphic to $(A_\ell, \{\alpha_1, \alpha_i\})$ ($1 < i < \ell$) or $(C_\ell, \{\alpha_1, \alpha_\ell\})$.

Furthermore $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ is the prolongation of $(\mathfrak{m}, \mathfrak{g}_0)$ except when $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ is isomorphic to $(A_\ell, \{\alpha_1\})$ or $(C_\ell, \{\alpha_1\})$.

Here R -spaces corresponding to the above exceptions (1), (2) and (3) are as follows: (1) correspond to compact irreducible hermitian symmetric spaces. (2) correspond to contact manifolds of Boothby type (Standard contact manifolds), which exist uniquely for each simple Lie algebra other than $\mathfrak{sl}(2, \mathbb{C})$ (see §5.2 below). In case of (3), $(J(\mathbb{P}^\ell, i), C)$ corresponds to $(A_\ell, \{\alpha_1, \alpha_i\})$ and $(L(\mathbb{P}^{2\ell-1}), E)$ corresponds to $(C_\ell, \{\alpha_1, \alpha_\ell\})$ ($1 < i < \ell$), where \mathbb{P}^ℓ denotes the ℓ -dimensional complex projective space and $\mathbb{P}^{2\ell-1}$ is the Standard contact manifold of type C_ℓ corresponding to $(C_\ell, \{\alpha_1\})$. Here we note that R -spaces corresponding to (2) and (3) are all Jet spaces of the first or second order.

For the real version of this theorem, we refer the reader to Theorem 5.3 [Y5].

Now the **Parabolic Geometry** is a geometry modeled after the homogeneous space G/G' , where G is a (semi-)simple Lie group and G' is a parabolic subgroup of G (cf. [Bai]). Precisely, in this paper, we mean, by a Parabolic Geometry, the **Geometry associated with the Simple Graded Lie Algebra** in the sense of N.Tanaka ([T4]).

In fact, let $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ be a simple graded Lie algebra over \mathbb{R} satisfying the generating condition. Let M be a manifold with a G_0^\sharp -structure of type \mathfrak{m} in the sense of [T4] (for the precise definition, see §2 of [T4]). In [T4], under the assumption that \mathfrak{g} is the prolongation of $(\mathfrak{m}, \mathfrak{g}_0)$, N. Tanaka constructed the **Normal Cartan Connection** (P, ω) of **Type \mathfrak{g}** over M , which settles the equivalence problem for the G_0^\sharp -structure of type \mathfrak{m} in the following sense: Let M and \hat{M} be two manifolds with G_0^\sharp -structures of type \mathfrak{m} . Let (P, ω) and $(\hat{P}, \hat{\omega})$ be the normal connections of type \mathfrak{g} over M and \hat{M} respectively. Then a diffeomorphism φ of M onto \hat{M} preserving the G_0^\sharp -structures lifts uniquely to an isomorphism φ^\sharp of (P, ω) onto $(\hat{P}, \hat{\omega})$ and vice versa ([T4], Theorem 2.7).

Here we note that, if \mathfrak{g} is the prolongation of \mathfrak{m} , a G_0^\sharp -structure on M is nothing but a regular differential system of type \mathfrak{m} (see [T4, §2.2]). Thus a Parabolic Geometry modeled after G/G' is the geometry of PD manifold of second order with the symbol algebra $\mathfrak{s} = \mathfrak{s}_{-3} \oplus \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1}$, if \mathfrak{g} is the prolongation of \mathfrak{m} and \mathfrak{m} is isomorphic to \mathfrak{s} .

Hence, among simple graded Lie algebras $\mathfrak{g} = \bigoplus_{p=-\mu}^{\mu} \mathfrak{g}_p \cong (X_\ell, \Delta_1)$, we will seek those algebras such that $\mathfrak{m} = \bigoplus_{p<0} \mathfrak{g}_p$ is isomorphic to the symbol algebra of PD manifolds of second order. Thus a necessary condition for this is $\mu = 3$ and $\dim \mathfrak{g}_{-3} = 1$. Then, by the above construction of (X_ℓ, Δ_1) , \mathfrak{g}_3 should be the highest root space. This forces $\Delta_\theta \subset \Delta_1$ where (X_ℓ, Δ_θ) is the (standard) contact gradation of \mathfrak{g} (see §5.2 below). These two conditions confine the possibility of (X_ℓ, Δ_1) . In fact, a simple graded Lie algebra (X_ℓ, Δ_1) , which satisfies both $\mu = 3$ and $\Delta_\theta \subset \Delta_1$ is one of the following: $(A_\ell, \{\alpha_1, \alpha_i, \alpha_\ell\})$ ($1 < i \leq \lfloor \frac{\ell+1}{2} \rfloor$), $(B_\ell, \{\alpha_1, \alpha_2\})$, $(C_\ell, \{\alpha_1, \alpha_\ell\})$, $(D_\ell, \{\alpha_1, \alpha_2\})$, $(D_\ell, \{\alpha_2, \alpha_\ell\})$, $(E_6, \{\alpha_1, \alpha_2\})$ and $(E_7, \{\alpha_1, \alpha_7\})$ up to conjugacy.

In fact, as we will see in §5.3 and §5.4, these simple graded Lie algebras (X_ℓ, Δ_1) represent the Parabolic Geometries of PD manifolds of second order (of finite type), except for $(C_\ell, \{\alpha_1, \alpha_\ell\})$, which is one of the exception in Prolongation Theorem and represents the Parabolic Geometry of third order equations of finite type (cf. [YY2] §3 Case (4)).

Furthermore, in §6.2 and §6.3, by utilizing the First Reduction Theorem, we will see more examples of Parabolic Geometry, which is associated with the geometry of (X, D) in §4.3.

5.2. Standard Contact Manifolds. Each simple Lie algebra \mathfrak{g} over \mathbb{C} has the highest root θ . Let Δ_θ denote the subset of Δ consisting of all vertices which are connected to $-\theta$ in the Extended Dynkin diagram of X_ℓ ($\ell \geq 2$). This subset Δ_θ of Δ , by the construction in §5.1, defines a gradation (or a partition of Φ^+), which distinguishes the highest root θ . Then, this gradation (X_ℓ, Δ_θ) turns out to be a contact gradation, which is unique up to conjugacy (Theorem 4.1 [Y5]). Explicitly we have $\Delta_\theta = \{\alpha_1, \alpha_\ell\}$ for A_ℓ type and $\Delta_\theta = \{\alpha_\theta\}$ for other types. Here $\alpha_\theta = \alpha_2, \alpha_1, \alpha_2$ for B_ℓ, C_ℓ, D_ℓ types respectively and $\alpha_\theta = \alpha_2, \alpha_1, \alpha_8, \alpha_1, \alpha_2$ for E_6, E_7, E_8, F_4, G_2 types respectively.


Moreover we have the adjoint (or equivalently coadjoint) representation, which has θ as the highest weight. The R -space $J_\mathfrak{g}$ corresponding to (X_ℓ, Δ_θ) can be obtained as the projectiviation of the (co-)adjoint orbit of G passing through the root vector of θ . By this construction, $J_\mathfrak{g}$ has the natural contact structure $C_\mathfrak{g}$ induced from the symplectic

structure as the coadjoint orbit, which corresponds to the contact gradation (X_ℓ, Δ_θ) (cf. [Y5], §4). Standard contact manifolds $(J_{\mathfrak{g}}, C_{\mathfrak{g}})$ were first found by Boothby ([Bo]) as compact simply connected homogeneous complex contact manifolds.


For the explicit description of the standard contact manifolds of the classical type, we refer the reader to §4.3 [Y5].

In §5.3 and §5.4, the model equation $(R_{\mathfrak{g}}, D_{\mathfrak{g}}^2)$ can be realized as a R -space orbit in $L(J_{\mathfrak{g}})$.


Extended Dynkin Diagrams with the coefficient of Highest Root (cf. [Bu])




$A_\ell (\ell > 1)$




$B_\ell (\ell > 2)$




$C_\ell (\ell > 1)$




$D_\ell (\ell > 3)$




E_6




F_4



E_7



G_2



E_8

5.3. Classical Type Examples. We will describe here the symbol algebra $\mathfrak{m} = \bigoplus_{p < 0} \mathfrak{g}_p$ of each (X_ℓ, Δ_1) of the classical type and give the model equations of second order. We refer the reader to §3 of [YY2] for the detailed description of the symbol algebras in matrices form. In this and next subsections, we will discuss in the complex analytic or the real C^∞ category depending on whether $\mathbb{K} = \mathbb{C}$ or \mathbb{R} .

(1) **Case of $(A_\ell, \{\alpha_1, \alpha_{i+1}, \alpha_\ell\})$** ($1 < i + 1 \leq \lfloor \frac{\ell+1}{2} \rfloor$).

We have the following matrix representation of $(A_\ell, \{\alpha_1, \alpha_{i+1}, \alpha_\ell\})$:

$$\mathfrak{sl}(\ell + 1, \mathbb{K}) = \mathfrak{g}_{-3} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2 \oplus \mathfrak{g}_3.$$

where the gradation is given by subdividing matrices as follows;

$$\begin{aligned} \mathfrak{g}_{-3} &= \left\{ \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 \end{array} \right) \middle| a \in \mathbb{K} \right\} \cong \mathbb{K}, \\ \mathfrak{g}_{-2} &= \left\{ \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\xi_2 & 0 & 0 & 0 \\ 0 & {}^t\xi_1 & 0 & 0 \end{array} \right) \middle| \xi_1 \in \mathbb{K}^i \quad \xi_2 \in \mathbb{K}^j \right\}, \\ \mathfrak{g}_{-1} &= V \oplus \mathfrak{f}, \\ \mathfrak{f} &= \left\{ \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & A & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \middle| A \in M(j, i) \right\}, \\ V &= \left\{ \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & {}^tx_2 & 0 \end{array} \right) \middle| x_1 \in \mathbb{K}^i, \quad x_2 \in \mathbb{K}^j \right\} \cong \mathbb{K}^{\ell-1}, \\ \mathfrak{g}_0 &= \left\{ \left(\begin{array}{cccc} b & 0 & 0 & 0 \\ 0 & B & 0 & 0 \\ 0 & 0 & C & 0 \\ 0 & 0 & 0 & c \end{array} \right) \middle| \begin{array}{l} b, c \in \mathbb{K}, B \in \mathfrak{gl}(i, \mathbb{K}), C \in \mathfrak{gl}(j, \mathbb{K}), \\ b + c + \text{tr}B + \text{tr}C = 0 \end{array} \right\} \\ \mathfrak{g}_k &= \{ {}^tX \mid X \in \mathfrak{g}_{-k} \}, (k = 1, 2, 3), \end{aligned}$$

where $i + j = \ell - 1$. Then we have

$$\begin{aligned} \mathfrak{m} &= \mathfrak{g}_{-3} \oplus \mathfrak{g}_{-2} \oplus (V \oplus \mathfrak{f}) \\ &= \left\{ \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 \\ -\xi_2 & A & 0 & 0 \\ a & {}^t\xi_1 & {}^tx_2 & 0 \end{array} \right) = \hat{a} + \check{\xi} + \hat{x} + \hat{A} \middle| \begin{array}{l} \xi = \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix}, x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{K}^{\ell-1}, \\ a \in \mathbb{K}, A \in M(j, i) \end{array} \right\}. \end{aligned}$$

By calculating $[\check{\xi}, \hat{x}]$ and $[[\hat{A}, \hat{x}], \hat{x}]$, we have

$$[\check{\xi}, \hat{x}] = \widehat{({}^t\xi x)}, \quad [[\hat{A}, \hat{x}], \hat{x}] = \widehat{(-2{}^tx_2 Ax_1)}, \quad 2{}^tx_2 Ax_1 = ({}^tx_1, {}^tx_2) \begin{pmatrix} 0 & {}^tA \\ A & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

Thus we have $\mathfrak{g}_{-2} \cong V^*$ and $\mathfrak{f} = \langle \{e_k^* \otimes e_\alpha^* \mid 1 \leq k \leq i, i+1 \leq \alpha \leq \ell-1\} \rangle \subset S^2(V^*)$ for a basis $\{e_1, \dots, e_{\ell-1}\}$ of V . This implies that the model equation of the second order is given by

$$\frac{\partial^2 z}{\partial x_k \partial x_l} = \frac{\partial^2 z}{\partial x_\alpha \partial x_\beta} = 0 \quad \text{for } 1 \leq k, l \leq i \quad \text{and } i+1 \leq \alpha, \beta \leq \ell-1,$$

where z is dependent variable and $x_1, \dots, x_{\ell-1}$ are independent variables.

(2) **Case of** $(B_\ell, \{\alpha_1, \alpha_2\})$ ($\ell \geq 3$), $(D_\ell, \{\alpha_1, \alpha_2\})$ ($\ell \geq 4$).

Similarly in this case, we have $\dim \mathfrak{f} = 1$ and $\mathfrak{f} = \langle \{e_1^* \odot e_1^* + \cdots + e_n^* \odot e_n^*\} \rangle$ for a basis $\{e_1, \dots, e_n\}$ of V . Thus we have the following model equation of the second order:

$$\frac{\partial^2 z}{\partial x_p \partial x_q} = \delta_{pq} \frac{\partial^2 z}{\partial^2 x_1} \quad \text{for } 1 \leq p, q \leq n,$$

where z is dependent variable and x_1, \dots, x_n are independent variables. This equation is the embedding equation as a hypersurface for the quadric Q^n . For the explicit matrix description of the gradation, we refer the reader to Case (5) in §3 of [YY2].

(3) **Case of** $(D_\ell, \{\alpha_2, \alpha_\ell\})$ ($\ell \geq 4$).

Similarly in this case, we have $\dim \mathfrak{f} = \frac{1}{2}(\ell - 1)(\ell - 2)$ and

$$\mathfrak{f} = \langle \left\{ \sum_{p,q=1}^{\ell-1} a_{pq} (e_p^1)^* \odot (e_q^2)^* \mid A = (a_{pq}) \in \mathfrak{o}(\ell - 1) \right\} \rangle$$

for a basis $\{e_1^1, \dots, e_{\ell-1}^1, e_1^2, \dots, e_{\ell-1}^2\}$ of V . Thus we have the following model equation of the second order:

$$\frac{\partial^2 z}{\partial x_p^i \partial x_q^j} + \frac{\partial^2 z}{\partial x_q^i \partial x_p^j} = 0 \quad \text{for } 1 \leq i, j \leq 2, \quad 1 \leq p < q \leq \ell - 1,$$

where z is dependent variable and $x_1^1, \dots, x_{\ell-1}^1, x_1^2, \dots, x_{\ell-1}^2$ are independent variables. This equation is the Plücker embedding equation for the Grassman manifold $\text{Gr}(\ell + 1, 2)$ (see [SYY] §3). For the explicit matrix description of the gradation, we refer the reader to Case (10) in §3 of [YY2].

5.4. Exceptional Type Examples. We here only describe the model equation $(R_{\mathfrak{g}}, D_{\mathfrak{g}}^2)$ of second order in the form of the standard differential system of type \mathfrak{m} . We refer the reader to §4 of [YY2] for the detailed description of the symbol algebras by the use of the Chevalley basis of \mathfrak{g} .

(1) **Case of** $(E_6, \{\alpha_1, \alpha_2\})$.

The symbol algebra $\mathfrak{m} = \mathfrak{g}_{-3} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$ is described as follows:

$$\mathfrak{g}_{-3} \cong \mathbb{K}, \quad \mathfrak{g}_{-2} \cong V^*, \quad \mathfrak{g}_{-1} = V \oplus \mathfrak{f}, \quad \mathfrak{f} \subset S^2(V^*) \quad \text{and} \quad \dim V = 10, \quad \dim \mathfrak{f} = 5.$$

Here the standard differential system $(M(\mathfrak{m}), D_{\mathfrak{m}})$ of type \mathfrak{m} in this case is given by

$$D_{\mathfrak{m}} = \{ \varpi = \varpi_1 = \varpi_2 = \cdots = \varpi_{10} = 0 \},$$

where

$$\begin{aligned} \varpi &= dz - p_1 dx_1 - \cdots - p_{10} dx_{10}, \\ \varpi_1 &= dp_1 + q_5 dx_8 + q_4 dx_9 + q_3 dx_{10}, & \varpi_2 &= dp_2 - q_5 dx_5 - q_4 dx_7 + q_2 dx_{10}, \\ \varpi_3 &= dp_3 + q_5 dx_4 + q_4 dx_6 + q_1 dx_{10}, & \varpi_4 &= dp_4 + q_5 dx_3 - q_3 dx_7 - q_2 dx_9, \\ \varpi_5 &= dp_5 - q_5 dx_2 + q_3 dx_6 - q_1 dx_9, & \varpi_6 &= dp_6 + q_4 dx_3 + q_3 dx_5 + q_2 dx_8, \\ \varpi_7 &= dp_7 - q_4 dx_2 - q_3 dx_4 + q_1 dx_8, & \varpi_8 &= dp_8 + q_5 dx_1 + q_2 dx_6 + q_1 dx_7, \\ \varpi_9 &= dp_9 + q_4 dx_1 - q_2 dx_4 - q_1 dx_5 & \varpi_{10} &= dp_{10} + q_3 dx_1 + q_2 dx_2 + q_1 dx_3. \end{aligned}$$

Here $(x_1, \dots, x_{10}, z, p_1, \dots, p_{10}, q_1, \dots, q_5)$ is a coordinate system of $M(\mathfrak{m}) \cong \mathbb{K}^{26}$.

(2) **Case of** $(E_7, \{\alpha_1, \alpha_7\})$.

The symbol algebra $\mathfrak{m} = \mathfrak{g}_{-3} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$ is described as follows:

$$\mathfrak{g}_{-3} \cong \mathbb{K}, \quad \mathfrak{g}_{-2} \cong V^*, \quad \mathfrak{g}_{-1} = V \oplus \mathfrak{f}, \quad \mathfrak{f} \subset S^2(V^*) \quad \text{and} \quad \dim V = 16, \quad \dim \mathfrak{f} = 10.$$

Here the standard differential system $(M(\mathfrak{m}), D_{\mathfrak{m}})$ of type \mathfrak{m} in this case is given by

$$D_{\mathfrak{m}} = \{ \varpi = \varpi_1 = \varpi_2 = \cdots = \varpi_{16} = 0 \},$$

where

$$\begin{aligned} \varpi &= dz - p_1 dx_1 - \cdots - p_{16} dx_{16}, \\ \varpi_1 &= dp_1 + q_{10} dx_{11} + q_9 dx_{12} + q_8 dx_{14} + q_7 dx_{15} + q_5 dx_{16}, \\ \varpi_2 &= dp_2 - q_{10} dx_9 - q_9 dx_{10} - q_8 dx_{13} + q_6 dx_{15} + q_4 dx_{16}, \\ \varpi_3 &= dp_3 + q_{10} dx_6 + q_9 dx_8 - q_7 dx_{13} - q_6 dx_{14} + q_3 dx_{16}, \\ \varpi_4 &= dp_4 - q_{10} dx_5 - q_9 dx_7 - q_5 dx_{13} - q_4 dx_{14} - q_3 dx_{15}, \\ \varpi_5 &= dp_5 - q_{10} dx_4 + q_8 dx_8 + q_7 dx_{10} + q_6 dx_{12} + q_2 dx_{16}, \\ \varpi_6 &= dp_6 + q_{10} dx_3 - q_8 dx_7 + q_5 dx_{10} + q_4 dx_{12} - q_2 dx_{15}, \\ \varpi_7 &= dp_7 - q_9 dx_4 - q_8 dx_6 - q_7 dx_9 - q_6 dx_{11} + q_1 dx_{16}, \\ \varpi_8 &= dp_8 + q_9 dx_3 + q_8 dx_5 - q_5 dx_9 - q_4 dx_{11} - q_1 dx_{15}, \\ \varpi_9 &= dp_9 - q_{10} dx_2 - q_7 dx_7 - q_5 dx_8 + q_3 dx_{12} + q_2 dx_{14}, \\ \varpi_{10} &= dp_{10} - q_9 dx_2 + q_7 dx_5 + q_5 dx_6 - q_3 dx_{11} + q_1 dx_{14}, \\ \varpi_{11} &= dp_{11} + q_{10} dx_1 - q_6 dx_7 - q_4 dx_8 - q_3 dx_{10} - q_2 dx_{13}, \\ \varpi_{12} &= dp_{12} + q_9 dx_1 + q_6 dx_5 + q_4 dx_6 + q_3 dx_9 - q_1 dx_{13}, \\ \varpi_{13} &= dp_{13} - q_8 dx_2 - q_7 dx_3 - q_5 dx_4 - q_2 dx_{11} - q_1 dx_{12}, \\ \varpi_{14} &= dp_{14} + q_8 dx_1 - q_6 dx_3 - q_4 dx_4 + q_2 dx_9 + q_1 dx_{10}, \\ \varpi_{15} &= dp_{15} + q_7 dx_1 + q_6 dx_2 - q_3 dx_4 - q_2 dx_6 - q_1 dx_8, \\ \varpi_{16} &= dp_{16} + q_5 dx_1 + q_4 dx_2 + q_3 dx_3 + q_2 dx_5 + q_1 dx_7. \end{aligned}$$

Here $(x_1, \dots, x_{16}, z, p_1, \dots, p_{16}, q_1, \dots, q_{10})$ is a coordinate system of $M(\mathfrak{m}) \cong \mathbb{K}^{43}$.

The model linear equations in §5.3 (2), (3) and §5.4 (1) and (2) appeared as the embedding equations of the corresponding symmetric spaces into the projective spaces in [SYY] and [HY] (see [SYY] §1). Except for §5.3 (2), these equations have rigidity properties. As is pointed out in §5 of [YY2], by the vanishing of the second cohomology (cf. Theorem 2.7 and 2.9 [T4], Proposition 5.5 [Y5]), we observe that Parabolic Geometries associated with $(D_\ell, \{\alpha_2, \alpha_\ell\})$, $(E_6, \{\alpha_1, \alpha_2\})$ and $(E_7, \{\alpha_1, \alpha_7\})$ have no local invariant. Thus the model second order equations of these cases is solely characterized by their symbols $\mathfrak{f} \subset S^2(V^*)$ under the condition (C), as in the case of Typical involutive symbols in §2.4.

6. EXAMPLES OF FIRST REDUCTION THEOREM.

Utilizing the First Reduction Theorem, we will discuss the Typical class of type $\mathfrak{f}^3(r)$ and exhibit several examples of PD manifolds of second order given through Parabolic Geometries on (X, D) .

6.1. Typical Class of Type $\mathfrak{f}^3(r)$. Let $(R; D^1, D^2)$ be a PD manifold of second order satisfying the condition (C), which is regular of type $\mathfrak{f}^3(r)$ ($r \leq n-2$). Namely $(R; D^1, D^2)$ is a PD manifold of second order such that symbol algebra $\mathfrak{s}(v)$ at each point $v \in R$ is isomorphic to $\mathfrak{s} = \mathfrak{s}_{-3} \oplus \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1}$ where

$$\mathfrak{s}_{-3} = \mathbb{R}, \quad \mathfrak{s}_{-2} = V^* \quad \text{and} \quad \mathfrak{s}_{-1} = V \oplus \mathfrak{f}^3(r).$$

Then, by Lemma 1.2 [Y4], there exists a coframe $\{\varpi, \varpi_a, \omega_a, \pi_{\alpha\beta} (1 \leq a \leq n, r+1 \leq \alpha, \beta \leq n)\}$ on a neighborhood U of each point $v \in R$ such that $D^1 = \{\varpi = 0\}$, $D^2 = \{\varpi = \varpi_1 = \cdots = \varpi_n = 0\}$ and that the following equalities hold:

$$\begin{aligned} d\varpi &\equiv \omega_1 \wedge \varpi_1 + \cdots + \omega_n \wedge \varpi_n \pmod{\varpi}, \\ \begin{cases} d\varpi_i &\equiv 0 \pmod{\varpi, \varpi_1, \dots, \varpi_n}, \\ d\varpi_\alpha &\equiv \omega_{r+1} \wedge \pi_{\alpha r+1} + \cdots + \omega_n \wedge \pi_{\alpha n} \pmod{\varpi, \varpi_1, \dots, \varpi_n}. \end{cases} \end{aligned}$$

for $1 \leq i \leq r$ and $r+1 \leq \alpha \leq n$ and $\pi_{\alpha\beta} = \pi_{\beta\alpha}$. Thus we see that $\text{Ch}(D^2)$ is a subbundle of D^2 of rank r . A coframe $\{\varpi, \varpi_a, \omega_a, \pi_{\alpha\beta} (1 \leq a \leq n, r+1 \leq \alpha, \beta \leq n)\}$ on U satisfying $D^1 = \{\varpi = 0\}$ and $D^2 = \{\varpi = \varpi_1 = \cdots = \varpi_n = 0\}$ is called adapted if it satisfies the above structure equations. Then we have (Lemma 4.1 [Y4])

Lemma A. *Let $(R; D^1, D^2)$ be as above and $r \leq n-2$. Then there exists an adapted coframe on U such that the following equalities hold:*

$$d\varpi_i \equiv 0 \pmod{\varpi_1, \dots, \varpi_r} \text{ for } i = 1, \dots, r.$$

By this lemma, for $\partial D^2 = \{\varpi = \varpi_1 = \cdots = \varpi_r = 0\}$, we have

$$\begin{cases} d\varpi \equiv \omega_{r+1} \wedge \varpi_{r+1} + \cdots + \omega_n \wedge \varpi_n \pmod{\varpi, \varpi_1, \dots, \varpi_r}, \\ d\varpi_i \equiv 0 \pmod{\varpi, \varpi_1, \dots, \varpi_r}. \end{cases}$$

Hence $\partial^2 D^2 = \partial^{(2)} D^2 = \{\varpi_1 = \cdots = \varpi_r = 0\}$ and $B = \partial^2 D^2$ is completely integrable.

Now we assume that R is regular with respect to $\text{Ch}(D^2)$, i.e., the leaf space $X = R/\text{Ch}(D^2)$ is a manifold such that the projection $\rho : R \rightarrow X$ is a submersion and there exists differential system D on X satisfying $D^2 = \rho_*^{-1}(D)$. Then, from the above information, we see that there exists a coframe $\{\varpi, \varpi_1, \dots, \varpi_n, \omega_{r+1}, \dots, \omega_n, \pi_{\alpha\beta} (r+1 \leq \alpha \leq \beta \leq n)\}$ on a neighborhood of each $x \in X$ such that

$$\begin{aligned} D &= \{\varpi = \varpi_1 = \cdots = \varpi_n = 0\}, \partial D = \{\varpi = \varpi_1 = \cdots = \varpi_r = 0\}, \\ \partial^2 D &= \partial^{(2)} D = \{\varpi_1 = \cdots = \varpi_r = 0\}, \end{aligned}$$

and that

$$\begin{cases} d\varpi_i \equiv 0 \pmod{\varpi_1, \dots, \varpi_r} \\ d\varpi \equiv \omega_{r+1} \wedge \varpi_{r+1} + \cdots + \omega_n \wedge \varpi_n \pmod{\varpi, \varpi_1, \dots, \varpi_r}, \\ d\varpi_\alpha \equiv \omega_{r+1} \wedge \pi_{\alpha r+1} + \cdots + \omega_n \wedge \pi_{\alpha n} \pmod{\varpi, \varpi_1, \dots, \varpi_n}. \end{cases}$$

for $1 \leq i \leq r$ and $r+1 \leq \alpha \leq n$ and $\pi_{\alpha\beta} = \pi_{\beta\alpha}$. Thus $B = \partial^2 D$ is completely integrable. Let p_1, \dots, p_r be the independent first integral of B around $x \in X$. Then we obtain

$$\begin{aligned} D &= \{\varpi = \varpi_{r+1} = \cdots = \varpi_n = dp_1 = \cdots = dp_r = 0\}, \\ \begin{cases} d\varpi \equiv \omega_{r+1} \wedge \varpi_{r+1} + \cdots + \omega_n \wedge \varpi_n \pmod{\varpi, dp_1, \dots, dp_r}, \\ d\varpi_\alpha \equiv \omega_{r+1} \wedge \pi_{\alpha r+1} + \cdots + \omega_n \wedge \pi_{\alpha n} \pmod{\varpi, \varpi_{r+1}, \dots, \varpi_n, dp_1, \dots, dp_r}. \end{cases} \end{aligned}$$

for $r+1 \leq \alpha \leq n$.

Namely (X, D) is a **parametrized second order contact manifold**. Hence, by the Darboux theorem, we obtain a coordinate system $(x_\alpha, z, p_i, p_\alpha, p_{\alpha\beta})$ ($1 \leq i \leq r, r+1 \leq \alpha, \beta \leq n$) around $x \in X$ such that (see [Y4] §4.2 and [Y7] §1.4)

$$D = \{\hat{\varpi} = \hat{\varpi}_{r+1} = \cdots = \hat{\varpi}_n = dp_1 = \cdots = dp_r = 0\},$$

where

$$\hat{\omega} = dz - \sum_{\alpha=r+1}^n p_{\alpha} dx_{\alpha}, \quad \hat{\omega}_{\alpha} = dp_{\alpha} - \sum_{\beta=r+1}^n p_{\alpha\beta} dx_{\beta}.$$

We refer the reader to §4.2 [Y4] for the detail to obtain a canonical coordinate system of $(R; D^1, D^2)$.

Moreover we observe that (X, D) satisfies three conditions in §4.3,

$$R(X) = \{v \in P(X) \mid v \not\propto \partial^2 D(x) \mid x = \nu(v)\}.$$

and $\hat{f}(v) = \text{Ch}(\partial D)(x) (\cong S^2((V_s)^*)) \subset D(x)$ for $v \in R(X)$. Thus, by Proposition 4, R is involutive, because $S^2((V_s)^*)$ is involutive.

6.2. G_2 -Geometry. Let $(X_{\ell}, \Delta_{\theta})$ be the (standard) contact gradation. Then we have $\Delta_{\theta} = \{\alpha_{\theta}\}$ except for A_{ℓ} type (see §5.2). As we observed in §6.3 in [Y6], for the exceptional simple Lie algebras, there exists, without exception, a unique simple root α_G next to α_{θ} such that the coefficient of α_G in the highest root is 3. We will consider simple graded Lie algebras $(X_{\ell}, \{\alpha_G\})$ of depth 3 and will show that regular differential systems of these types satisfy the conditions (X.1) to (X.3) in §4.3.

Explicitly we will here consider the following simple graded Lie algebras of depth 3: $(G_2, \{\alpha_1\})$, $(F_4, \{\alpha_2\})$, $(E_6, \{\alpha_4\})$, $(E_7, \{\alpha_3\})$, $(E_8, \{\alpha_7\})$, $(B_{\ell}, \{\alpha_1, \alpha_3\})$ ($\ell \geq 3$), $(D_{\ell}, \{\alpha_1, \alpha_3\})$ ($\ell \geq 5$) and $(D_4, \{\alpha_1, \alpha_3, \alpha_4\})$. These graded Lie algebras have the common feature with $(G_2, \{\alpha_1\})$ as follows: In these cases, $\mathfrak{m} = \mathfrak{g}_{-3} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$ satisfies $\dim \mathfrak{g}_{-3} = 2$ and $\dim \mathfrak{g}_{-1} = 2 \dim \mathfrak{g}_{-2}$. Moreover, in the description of the gradation in terms of the root space decomposition in §5.1, we have $\Phi_3^+ = \{\theta, \theta - \alpha_{\theta}\}$ such that the coefficient of α_{θ} in each $\beta \in \Phi_2^+$ is 1 and Φ_1^+ consists of roots $\theta - \beta$, $\theta - \alpha_{\theta} - \beta$ for each $\beta \in \Phi_2^+$. Hence, ignoring the bracket product in \mathfrak{g}_{-1} , we can describe the bracket products of other part of \mathfrak{m} , in terms of paring, by

$$\mathfrak{g}_{-3} = W, \quad \mathfrak{g}_{-2} = V \quad \text{and} \quad \mathfrak{g}_{-1} = W \otimes V^*,$$

where $\dim W = 2$.

Thus let (X, D) be a regular differential system of type \mathfrak{m} , where \mathfrak{m} is the negative part of one of the above graded Lie algebras. Then $(X, \partial D)$ is a regular differential system of type $\mathfrak{c}^1(s, 2)$. Namely, there exists a coframe $\{\varpi_1, \varpi_2, \pi_1, \dots, \pi_s, \pi_1^1, \dots, \pi_1^s, \pi_2^1, \dots, \pi_2^s\}$ around $x \in X$ such that

$$\partial D = \{\varpi_1 = \varpi_2 = 0\},$$

and

$$\begin{cases} d\varpi_1 \equiv \pi_1^1 \wedge \pi_1 + \dots + \pi_1^s \wedge \pi_s \pmod{\varpi_1, \varpi_2} \\ d\varpi_2 \equiv \pi_2^1 \wedge \pi_1 + \dots + \pi_2^s \wedge \pi_s \pmod{\varpi_1, \varpi_2} \end{cases}$$

Thus (X, D) satisfies the conditions (X.1) to (X.3) in §4.3.

Now, putting $\varpi = \lambda_1 \varpi_1 + \lambda_2 \varpi_2$, we consider a point $v \in P(X)$ such that $v = \{\varpi = 0\} \subset T_x(X)$, where $x = \nu(v)$. Then, for $(\lambda_1, \lambda_2) \neq (0, 0)$, $\{\lambda_1 \pi_1^i + \lambda_2 \pi_2^i (i = 1, \dots, s)\}$ are linearly independent (mod $\varpi_1, \varpi_2, \pi_1, \dots, \pi_s$). Thus $\hat{f}(v) \subset D(x)$ is of codimension s at each $v \in P(X)$ (see the proof of Proposition 3). Hence we obtain $R(X) = P(X)$ in this case, i.e., $R(X)$ is a \mathbb{P}^1 -bundle over X .

In fact, when (X, D) is the model space $(M_{\mathfrak{g}}, D_{\mathfrak{g}})$ of type $(X_{\ell}, \{\alpha_G\})$, $R(X)$ can be identified with the model space $(R_{\mathfrak{g}}, E_{\mathfrak{g}})$ of type $(X_{\ell}, \{\alpha_{\theta}, \alpha_G\})$ as follows (here, we understand α_G denotes two simple roots α_1 and α_3 in case of BD_{ℓ} types and three simple

roots α_1, α_3 and α_4 in case of D_4): Let $(J_{\mathfrak{g}}, C_{\mathfrak{g}})$ be the standard contact manifold of type $(X_{\ell}, \{\alpha_{\theta}\})$. Then we have the double fibration;

$$\begin{array}{ccc} R_{\mathfrak{g}} & \xrightarrow{\pi_c} & J_{\mathfrak{g}} \\ \pi_g \downarrow & & \\ & & M_{\mathfrak{g}} \end{array}$$

Here $(X_{\ell}, \{\alpha_{\theta}, \alpha_G\})$ is a graded Lie algebra of depth 5 and satisfies the following: $\dim \check{\mathfrak{g}}_{-5} = \dim \check{\mathfrak{g}}_{-4} = 1$, $\dim \check{\mathfrak{g}}_{-3} = \dim \check{\mathfrak{g}}_{-2} = s$ and $\dim \check{\mathfrak{g}}_{-1} = s + 1$. In fact, comparing with the gradation of $(X_{\ell}, \{\alpha_G\})$, we have $\check{\Phi}_5^+ = \{\theta\}$, $\check{\Phi}_4^+ = \{\theta - \alpha_{\theta}\}$, $\check{\Phi}_3^+ = \Phi_2^+$, $\check{\Phi}_2^+$ consists of roots $\theta - \beta$ for each $\beta \in \check{\Phi}_3^+$ and $\check{\Phi}_1^+$ consists of roots α_{θ} and $\theta - \alpha_{\theta} - \beta$ for each $\beta \in \check{\Phi}_3^+$. Thus we see that $\partial^{(3)}E_{\mathfrak{g}} = (\pi_c)_*^{-1}(C_{\mathfrak{g}})$, $\partial^{(2)}E_{\mathfrak{g}} = (\pi_g)_*^{-1}(\partial D_{\mathfrak{g}})$ and $\partial E_{\mathfrak{g}} = (\pi_g)_*^{-1}(D_g)$. We put $D^1 = \partial^{(3)}E_{\mathfrak{g}}$ and $D^2 = \partial E_{\mathfrak{g}}$. Then $(R_{\mathfrak{g}}; D^1, D^2)$ is a PD manifold of second order. In fact, we have an isomorphism of $(R_{\mathfrak{g}}; D^1, D^2)$ onto $(R(M_{\mathfrak{g}}); D_{M_{\mathfrak{g}}}^1, D_{M_{\mathfrak{g}}}^2)$ by the Realization Lemma for $(R_{\mathfrak{g}}, D^1, \pi_g, M_{\mathfrak{g}})$ and an embedding of $R_{\mathfrak{g}}$ into $L(J_{\mathfrak{g}})$ by the Realization Lemma for $(R_{\mathfrak{g}}, D^2, \pi_c, J_{\mathfrak{g}})$. Thus $R_{\mathfrak{g}}$ is identified with a R -space orbit in $L(J_{\mathfrak{g}})$.

Now we will calculate the symbol of $(R(X); D_X^1, D_X^2)$ by utilizing the model PD manifold $(R_{\mathfrak{g}}; D^1, D^2)$ of second order, especially when $(X_{\ell}, \{\alpha_{\theta}, \alpha_G\})$ is of BD_{ℓ} types. Let us describe the gradation of $(BD_{\ell}, \{\alpha_1, \alpha_2, \alpha_3\})$ or $(D_4, \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\})$ in matrices form as follows: First we describe

$$\mathfrak{o}(k+6) = \{X \in \mathfrak{gl}(k+6, \mathbb{K}) \mid {}^t X J + J X = 0\},$$

where

$$J = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & I_k & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \in \mathfrak{gl}(k+6, \mathbb{K}), \quad I_k = (\delta_{ij}) \in \mathfrak{gl}(k, \mathbb{K}).$$

Here $I_k \in \mathfrak{gl}(k, \mathbb{K})$ is the unit matrix and the gradation is given again by subdividing matrices as follows;

$$\check{\mathfrak{g}}_{-5} = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ y & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -y & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \right\}, \quad \check{\mathfrak{g}}_{-4} = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \xi_0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\xi_0 & 0 & 0 & 0 & 0 \end{pmatrix} \right\}, \quad y, \xi_0 \in \mathbb{K}$$

$$\check{\mathfrak{g}}_{-3} = \left\{ \left(\begin{array}{ccccccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \xi & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \xi_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\xi_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -{}^t\xi & 0 & 0 & 0 \end{array} \right) \mid \xi \in \mathbb{K}^k, \xi_1 \in \mathbb{K} \right\},$$

$$\check{\mathfrak{g}}_{-2} = \left\{ \left(\begin{array}{ccccccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & x & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -{}^tx & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_1 & 0 & 0 \end{array} \right) = \hat{x} + \hat{x}_1 \mid x \in \mathbb{K}^k, x_1 \in \mathbb{K} \right\},$$

$$\check{\mathfrak{g}}_{-1} = \left\{ \left(\begin{array}{ccccccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & x_0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -{}^ta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -a_1 & 0 \end{array} \right) = \hat{x}_0 + \hat{a}_1 + \hat{a} \mid \begin{array}{l} x_0, a_1 \in \mathbb{K}, \\ a \in \mathbb{K}^k \end{array} \right\},$$

$$\check{\mathfrak{g}}_0 = \left\{ \left(\begin{array}{ccccccc} b & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & c & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & e & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & B & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -e & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -c & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -b \end{array} \right) \mid b, c, e \in \mathbb{K}, B \in \mathfrak{o}(k) \right\}$$

$$\check{\mathfrak{g}}_\ell = \{ {}^tX \mid X \in \mathfrak{g}_{-\ell} \}, (\ell = 1, 2, 3, 4, 5),$$

Then, for $X = \hat{x} + \hat{x}_1 + \hat{x}_0$ and $A = \hat{a} + \hat{a}_1$, we calculate

$$[[A, X], X] = (2x_1 {}^t a x - a_1 {}^t x x) \in \check{\mathfrak{g}}_{-5}$$

Thus we obtain

$$\mathfrak{f} = \langle \{2e_1^* \otimes e_2^*, \dots, 2e_1^* \otimes e_{k+1}^*, e_2^* \otimes e_2^* + \dots + e_{k+1}^* \otimes e_{k+1}^*\} \subset S^2(E^\perp),$$

where $\{e_0, e_1, \dots, e_{k+1}\}$ is a basis of V and $E = \langle \{e_0\} \rangle$ is the Cauchy characteristic direction. In case $k = 1$, $\mathfrak{f} = \langle \{2e_1^* \otimes e_2^*, e_2^* \otimes e_2^*\} \rangle$ is an involutive subspace of $S^2(E^\perp)$. Hence $(R_{\mathfrak{g}}; D^1, D^2)$ is involutive when $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ is of type $(B_3, \{\alpha_1, \alpha_2, \alpha_3\})$. In case $k > 1$, we have

$$\mathfrak{f}^\perp = \langle \{e_1 \otimes e_1, e_i \otimes e_j (2 \leq i < j \leq k+1),$$

$$e_2 \otimes e_2 - e_{k+1} \otimes e_{k+1}, \dots, e_k \otimes e_k - e_{k+1} \otimes e_{k+1}\} \subset S^2(W),$$

where $W = \langle \{e_1, \dots, e_{k+1}\} \rangle$. Then we see that $(\mathfrak{f}^{(1)})^\perp$ contains every $e_i \otimes e_i \otimes e_i$ for $i = 1, \dots, k+1$, which implies \mathfrak{f} is of finite type. We can also check that exceptional cases other than G_2 are of finite type by utilizing R -space orbit $(R_{\mathfrak{g}}; D^1, D^2)$, whereas

$\mathfrak{f} = S^2(W^*)$ is involutive in case of G_2 , where $\dim W = 1$. We will discuss these cases in a uniform way in other occasion.

6.3. Other Examples. We exhibit here two other examples of simple graded Lie algebras $\mathfrak{g} = \bigoplus_{p \in \mathbb{Z}} \mathfrak{g}_p$ of depth 3, such that regular differential systems (X, D) of type \mathfrak{m} satisfy the conditions (X.1) to (X.3) in §4.3, where $\mathfrak{m} = \bigoplus_{p < 0} \mathfrak{g}_p$.

The first example is of type $(C_\ell, \{\alpha_2, \alpha_\ell\})$. Here we have $\dim \mathfrak{g}_{-3} = 3$, $\dim \mathfrak{g}_{-2} = 2(\ell - 2)$ and $\dim \mathfrak{g}_{-1} = 2(\ell - 2) + \frac{1}{2}(\ell - 2)(\ell - 1)$. Utilizing the calculation in Case (3) of §3 in [YY2], we have the following description of the standard differential system $(M_{\mathfrak{m}}, D_{\mathfrak{m}})$ of type \mathfrak{m} in this case:

$$D_{\mathfrak{m}} = \{\varpi_0 = \varpi_1 = \varpi_2 = \pi_1^1 = \cdots = \pi_1^{\ell-2} = \pi_2^1 = \cdots = \pi_2^{\ell-2} = 0\},$$

where

$$\left\{ \begin{array}{l} \varpi_0 = d y_0 - \sum_{\alpha=1}^{\ell-2} \xi_1^\alpha dx_2^\alpha - \sum_{\alpha=1}^{\ell-2} \xi_2^\alpha dx_1^\alpha, \\ \varpi_1 = d y_1 - 2 \sum_{\alpha=1}^{\ell-2} \xi_1^\alpha dx_1^\alpha, \quad \varpi_2 = d y_2 - 2 \sum_{\alpha=1}^{\ell-2} \xi_2^\alpha dx_2^\alpha, \\ \pi_p^\alpha = d \xi_p^\alpha - \sum_{\beta=1}^{\ell-2} a_{\alpha\beta} dx_p^\beta \quad (p = 1, 2 \quad \alpha = 1, \dots, \ell - 2), \end{array} \right.$$

Here we put $y_{12} = y_0$, $y_{11} = y_1$, $y_{22} = y_2$ and $a_{\alpha\beta} = a_{\beta\alpha}$ for $1 \leq \alpha, \beta \leq \ell - 2$. Let (X, D) be a regular differential system of type \mathfrak{m} . Then we have the structure equation of $(X, \partial D)$ as follows;

$$\left\{ \begin{array}{l} d\varpi_0 \equiv \omega_2^1 \wedge \pi_1^1 + \cdots + \omega_2^{\ell-2} \wedge \pi_1^{\ell-2} + \omega_1^1 \wedge \pi_2^1 + \cdots + \omega_1^{\ell-2} \wedge \pi_2^{\ell-2} \\ d\varpi_1 \equiv 2 \omega_1^1 \wedge \pi_1^1 + \cdots + 2 \omega_1^{\ell-2} \wedge \pi_1^{\ell-2} \\ d\varpi_2 \equiv 2 \omega_2^1 \wedge \pi_2^1 + \cdots + 2 \omega_2^{\ell-2} \wedge \pi_2^{\ell-2} \end{array} \pmod{\varpi_0, \varpi_1, \varpi_2} \right.$$

Now, putting $\varpi = \lambda_0 \varpi_0 + \lambda_1 \varpi_1 + \lambda_2 \varpi_2$, we consider a point $v \in P(X)$ such that $v = \{\varpi = 0\} \subset T_x(X)$, where $x = \nu(v)$. Then, from

$$d\varpi \equiv \sum_{\alpha=1}^{\ell-2} (\lambda_0 \omega_2^\alpha + 2 \lambda_1 \omega_1^\alpha) \wedge \pi_1^\alpha + \sum_{\alpha=1}^{\ell-2} (\lambda_0 \omega_1^\alpha + 2 \lambda_2 \omega_2^\alpha) \wedge \pi_2^\alpha, \pmod{\varpi_0, \varpi_1, \varpi_2},$$

and

$$(\lambda_0 \omega_2^\alpha + 2 \lambda_1 \omega_1^\alpha) \wedge (\lambda_0 \omega_1^\alpha + 2 \lambda_2 \omega_2^\alpha) = (4\lambda_1 \lambda_2 - \lambda_0^2) \omega_1^\alpha \wedge \omega_2^\alpha,$$

we see that (X, D) satisfies the condition (X.1) to (X.3) in §4.3 and we obtain

$$R(X) = \{v \in P(X) \mid 4\lambda_1 \lambda_2 - \lambda_0^2 \neq 0\},$$

where $(\lambda_0, \lambda_1, \lambda_2)$ is the homogeneous coordinate of the fibre $\nu : P(X) \rightarrow X$.

By the calculation in Case (3) of §3 in [YY2], we have $y_{pq} = 2 \sum_{\alpha, \beta=1}^{\ell-2} a_{\alpha\beta} x_p^\alpha x_q^\beta$ for $1 \leq p \leq q \leq 2$ and $a_{\alpha\beta} = a_{\beta\alpha}$ ($1 \leq \alpha, \beta \leq \ell - 2$) so that

$$\lambda_0 y_0 + \lambda_1 y_1 + \lambda_2 y_2 = 2 \sum_{\alpha, \beta=1}^{\ell-2} a_{\alpha\beta} (\lambda_0 x_1^\alpha x_2^\beta + \lambda_1 x_1^\alpha x_1^\beta + \lambda_2 x_2^\alpha x_2^\beta).$$

Hence, from

$$\begin{aligned} \sum_{\alpha, \beta=1}^{\ell-2} a_{\alpha\beta}(\lambda_0 x_1^\alpha x_2^\beta + \lambda_1 x_1^\alpha x_1^\beta + \lambda_2 x_2^\alpha x_2^\beta) &= \sum_{\alpha=1}^{\ell-2} a_{\alpha\alpha}(\lambda_0 x_1^\alpha x_2^\alpha + \lambda_1 x_1^\alpha x_1^\alpha + \lambda_2 x_2^\alpha x_2^\alpha) \\ &+ \sum_{\alpha < \beta} a_{\alpha\beta}(\lambda_0(x_1^\alpha x_2^\beta + x_1^\beta x_2^\alpha) + 2\lambda_1 x_1^\alpha x_1^\beta + 2\lambda_2 x_2^\alpha x_2^\beta), \end{aligned}$$

we have

$$\begin{aligned} \hat{f}(v) &= \langle \{\lambda_0(e_1^\alpha)^* \odot (e_2^\alpha)^* + \lambda_1(e_1^\alpha)^* \odot (e_1^\alpha)^* + \lambda_2(e_2^\alpha)^* \odot (e_2^\alpha)^* (1 \leq \alpha \leq \ell - 2), \\ &\lambda_0((e_1^\alpha)^* \odot (e_2^\beta)^* + (e_1^\beta)^* \odot (e_2^\alpha)^*) + 2\lambda_1(e_1^\alpha)^* \odot (e_1^\beta)^* + 2\lambda_2(e_2^\alpha)^* \odot (e_2^\beta)^* \\ &(1 \leq \alpha < \beta \leq \ell - 2)\} \rangle \subset S^2(W^*), \end{aligned}$$

where $W = \langle \{e_1^1, \dots, e_1^{\ell-2}, e_2^1, \dots, e_2^{\ell-2}\} \rangle$. Thus, assuming $\lambda_0 \neq 0$, we get

$$\begin{aligned} \hat{f}(v)^\perp &= \langle \{\lambda_0 e_1^\alpha \odot e_1^\alpha - 2\lambda_1 e_1^\alpha \odot e_2^\alpha, \lambda_0 e_2^\alpha \odot e_2^\alpha - 2\lambda_2 e_1^\alpha \odot e_2^\alpha (1 \leq \alpha \leq \ell - 2), \\ &e_1^\alpha \odot e_2^\beta - e_1^\beta \odot e_2^\alpha, \lambda_0 e_1^\alpha \odot e_1^\beta - 2\lambda_1 e_1^\alpha \odot e_2^\beta, \lambda_0 e_2^\alpha \odot e_2^\beta - 2\lambda_2 e_1^\alpha \odot e_2^\beta, \\ &(1 \leq \alpha < \beta \leq \ell - 2)\} \rangle \subset S^2(W), \end{aligned}$$

Then, from the first two generator of $\hat{f}(v)^\perp$, we see that $\hat{f}^{(1)}(v)^\perp$ contains $e_1^\alpha \odot e_1^\alpha \odot e_2^\alpha$ and $e_1^\alpha \odot e_2^\alpha \odot e_2^\alpha$. Moreover it follows that $\hat{f}^{(1)}(v)^\perp$ contains every $e_1^\alpha \odot e_1^\alpha \odot e_1^\alpha$ and $e_2^\alpha \odot e_2^\alpha \odot e_2^\alpha$ for $\alpha = 1, \dots, \ell - 2$, which implies that $\hat{f}(v)$ is of finite type.

Now let us construct the model equation of second order from the coordinate description of the standard differential system (M_m, D_m) . We calculate

$$\begin{aligned} \varpi &= \varpi_0 + \lambda_1 \varpi_1 + \lambda_2 \varpi_2 \\ &= dy_0 + \lambda_1 dy_1 + \lambda_2 dy_2 - \sum_{\alpha=1}^{\ell-2} (\xi_2^\alpha + 2\lambda_1 \xi_1^\alpha) dx_1^\alpha - \sum_{\alpha=1}^{\ell-2} (\xi_1^\alpha + 2\lambda_2 \xi_2^\alpha) dx_2^\alpha \\ &= d(y_0 + \lambda_1 y_1 + \lambda_2 y_2) - y_1 d\lambda_1 - y_2 d\lambda_2 \\ &\quad - \sum_{\alpha=1}^{\ell-2} (\xi_2^\alpha + 2\lambda_1 \xi_1^\alpha) dx_1^\alpha - \sum_{\alpha=1}^{\ell-2} (\xi_1^\alpha + 2\lambda_2 \xi_2^\alpha) dx_2^\alpha \end{aligned}$$

Thus we put

$$\begin{cases} z = y_0 + \lambda_1 y_1 + \lambda_2 y_2, & x_1^0 = \lambda_1, & x_2^0 = \lambda_2, & p_1^0 = y_1, & p_2^0 = y_2, \\ p_1^\alpha = \xi_2^\alpha + 2\lambda_1 \xi_1^\alpha, & p_2^\alpha = \xi_1^\alpha + 2\lambda_2 \xi_2^\alpha & (\alpha = 1, \dots, \ell - 2). \end{cases}$$

Then we have

$$\xi_1^\alpha = \frac{p_2^\alpha - 2x_2^0 p_1^\alpha}{1 - 4x_1^0 x_2^0}, \quad \xi_2^\alpha = \frac{p_1^\alpha - 2x_1^0 p_2^\alpha}{1 - 4x_1^0 x_2^0},$$

and

$$\begin{aligned} \varpi &= dz - \sum_{\alpha=0}^{\ell-2} p_1^\alpha dx_1^\alpha - \sum_{\alpha=0}^{\ell-2} p_2^\alpha dx_2^\alpha, \\ \pi_1^0 &= dp_1^0 - 2 \sum_{\alpha=1}^{\ell-2} \xi_1^\alpha dx_1^\alpha, & \pi_2^0 &= dp_2^0 - 2 \sum_{\alpha=1}^{\ell-2} \xi_2^\alpha dx_2^\alpha, \end{aligned}$$

$$\begin{aligned}\pi_1^\alpha &= dp_1^\alpha - 2\xi_1^\alpha dx_1^0 - 2\sum_{\beta=1}^{\ell-2}\lambda_1 a_{\alpha\beta} dx_1^\beta - \sum_{\beta=1}^{\ell-2} a_{\alpha\beta} dx_2^\beta, \\ \pi_2^\alpha &= dp_2^\alpha - 2\xi_2^\alpha dx_2^0 - \sum_{\beta=1}^{\ell-2} a_{\alpha\beta} dx_1^\beta - 2\sum_{\beta=1}^{\ell-2}\lambda_2 a_{\alpha\beta} dx_2^\beta,\end{aligned}$$

where $\pi_1^0 = \varpi_1$, $\pi_2^0 = \varpi_2$, $\pi_1^\alpha = \varpi_2^\alpha + 2\lambda_1\varpi_1^\alpha$ and $\pi_2^\alpha = \varpi_1^\alpha + 2\lambda_2\varpi_2^\alpha$. Hence we obtain the following model equation of second order :

$$\begin{aligned}\frac{\partial^2 z}{\partial x_1^0 \partial x_1^0} &= \frac{\partial^2 z}{\partial x_2^0 \partial x_2^0} = \frac{\partial^2 z}{\partial x_1^0 \partial x_2^0} = 0, & \frac{\partial^2 z}{\partial x_1^0 \partial x_2^\alpha} &= \frac{\partial^2 z}{\partial x_2^0 \partial x_1^\alpha} = 0, \\ \frac{\partial^2 z}{\partial x_1^0 \partial x_1^\alpha} &= \frac{2}{1 - 4x_1^0 x_2^0} \left(\frac{\partial z}{\partial x_2^\alpha} - 2x_2^0 \frac{\partial z}{\partial x_1^\alpha} \right) \quad (= 2\xi_1^\alpha), \\ \frac{\partial^2 z}{\partial x_2^0 \partial x_2^\alpha} &= \frac{2}{1 - 4x_1^0 x_2^0} \left(\frac{\partial z}{\partial x_1^\alpha} - 2x_1^0 \frac{\partial z}{\partial x_2^\alpha} \right) \quad (= 2\xi_2^\alpha), \\ \frac{\partial^2 z}{\partial x_1^\alpha \partial x_1^\beta} &= 2x_1^0 \frac{\partial^2 z}{\partial x_2^\alpha \partial x_1^\beta}, & \frac{\partial^2 z}{\partial x_2^\alpha \partial x_2^\beta} &= 2x_2^0 \frac{\partial^2 z}{\partial x_2^\alpha \partial x_1^\beta}, \\ \frac{\partial^2 z}{\partial x_2^\alpha \partial x_1^\beta} &= \frac{\partial^2 z}{\partial x_2^\beta \partial x_1^\alpha} \quad (= a_{\alpha\beta}) & (1 \leq \alpha \leq \beta \leq \ell - 2).\end{aligned}$$

Our second example is of type $(E_7, \{\alpha_6, \alpha_7\})$. Here we have $\dim \mathfrak{g}_{-3} = 10$, $\dim \mathfrak{g}_{-2} = 16$ and $\dim \mathfrak{g}_{-1} = 16 + 1$. Utilizing the calculation in Case (4) of §4 in [YY2], we have the following description of the standard differential system $(M_{\mathfrak{m}}, D_{\mathfrak{m}})$ of type \mathfrak{m} in this case:

$$D_{\mathfrak{m}} = \{\varpi_1 = \cdots = \varpi_{10} = \pi_1 = \cdots = \pi_{16} = 0\},$$

where

$$\begin{aligned}\varpi_1 &= dy_1 - p_{11}dx_1 + p_9dx_2 - p_6dx_3 + p_5dx_4 + p_4dx_5 - p_3dx_6 + p_2dx_9 - p_1dx_{11}, \\ \varpi_2 &= dy_2 - p_{13}dx_1 + p_{10}dx_2 - p_8dx_3 + p_7dx_4 + p_4dx_7 - p_3dx_8 + p_2dx_{10} - p_1dx_{13}, \\ \varpi_3 &= dy_3 - p_{14}dx_1 + p_{12}dx_2 - p_8dx_5 + p_7dx_6 + p_6dx_7 - p_5dx_8 + p_2dx_{12} - p_1dx_{14}, \\ \varpi_4 &= dy_4 - p_{15}dx_1 + p_{12}dx_3 - p_{10}dx_5 + p_9dx_7 + p_7dx_9 - p_5dx_{10} + p_3dx_{12} - p_1dx_{15}, \\ \varpi_5 &= dy_5 - p_{15}dx_2 + p_{14}dx_3 - p_{13}dx_5 + p_{11}dx_7 + p_7dx_{11} - p_5dx_{13} + p_3dx_{14} - p_2dx_{15}, \\ \varpi_6 &= dy_6 - p_{16}dx_1 + p_{12}dx_4 - p_{10}dx_6 + p_9dx_8 + p_8dx_9 - p_6dx_{10} + p_4dx_{12} - p_1dx_{16}, \\ \varpi_7 &= dy_7 - p_{16}dx_2 + p_{14}dx_4 - p_{13}dx_6 + p_{11}dx_8 + p_8dx_{11} - p_6dx_{13} + p_4dx_{14} - p_2dx_{16}, \\ \varpi_8 &= dy_8 - p_{16}dx_3 + p_{15}dx_4 - p_{13}dx_9 + p_{11}dx_{10} + p_{10}dx_{11} - p_9dx_{13} + p_4dx_{15} - p_3dx_{16}, \\ \varpi_9 &= dy_9 - p_{16}dx_5 + p_{15}dx_6 - p_{14}dx_9 + p_{12}dx_{11} + p_{11}dx_{12} - p_9dx_{14} + p_6dx_{15} - p_5dx_{16}, \\ \varpi_{10} &= dy_{10} - p_{16}dx_7 + p_{15}dx_8 - p_{14}dx_{10} + p_{13}dx_{12} + p_{12}dx_{13} - p_{10}dx_{14} + p_8dx_{15} - p_7dx_{16}, \\ \pi_i &= dp_i - adx_i \quad (i = 1, 2, \dots, 16).\end{aligned}$$

Let (X, D) be a regular differential system of type \mathbf{m} . Then we have the structure equation of $(X, \partial D)$ as follows;

$$\begin{aligned}
d\varpi_1 &\equiv \omega_1 \wedge \pi_{11} - \omega_2 \wedge \pi_9 + \omega_3 \wedge \pi_6 - \omega_4 \wedge \pi_5 - \omega_5 \wedge \pi_4 + \omega_6 \wedge \pi_3 - \omega_9 \wedge \pi_2 + \omega_{11} \wedge \pi_1, \\
d\varpi_2 &\equiv \omega_1 \wedge \pi_{13} - \omega_2 \wedge \pi_{10} + \omega_3 \wedge \pi_8 - \omega_4 \wedge \pi_7 - \omega_7 \wedge \pi_4 + \omega_8 \wedge \pi_3 - \omega_{10} \wedge \pi_2 + \omega_{13} \wedge \pi_1, \\
d\varpi_3 &\equiv \omega_1 \wedge \pi_{14} - \omega_2 \wedge \pi_{12} + \omega_5 \wedge \pi_8 - \omega_6 \wedge \pi_7 - \omega_7 \wedge \pi_6 + \omega_8 \wedge \pi_5 - \omega_{12} \wedge \pi_2 + \omega_{14} \wedge \pi_1, \\
d\varpi_4 &\equiv \omega_1 \wedge \pi_{15} - \omega_3 \wedge \pi_{12} + \omega_5 \wedge \pi_{10} - \omega_7 \wedge \pi_9 - \omega_9 \wedge \pi_7 + \omega_{10} \wedge \pi_5 - \omega_{12} \wedge \pi_3 + \omega_{15} \wedge \pi_1, \\
d\varpi_5 &\equiv \omega_2 \wedge \pi_{15} - \omega_3 \wedge \pi_{14} + \omega_5 \wedge \pi_{13} - \omega_7 \wedge \pi_{11} - \omega_{11} \wedge \pi_7 + \omega_{13} \wedge \pi_5 - \omega_{14} \wedge \pi_3 + \omega_{15} \wedge \pi_2, \\
d\varpi_6 &\equiv \omega_1 \wedge \pi_{16} - \omega_4 \wedge \pi_{12} + \omega_6 \wedge \pi_{10} - \omega_8 \wedge \pi_9 - \omega_9 \wedge \pi_8 + \omega_{10} \wedge \pi_6 - \omega_{12} \wedge \pi_4 + \omega_{16} \wedge \pi_1, \\
d\varpi_7 &\equiv \omega_2 \wedge \pi_{16} - \omega_4 \wedge \pi_{14} + \omega_6 \wedge \pi_{13} - \omega_8 \wedge \pi_{11} - \omega_{11} \wedge \pi_8 + \omega_{13} \wedge \pi_6 - \omega_{14} \wedge \pi_4 + \omega_{16} \wedge \pi_2, \\
d\varpi_8 &\equiv \omega_3 \wedge \pi_{16} - \omega_4 \wedge \pi_{15} + \omega_9 \wedge \pi_{13} - \omega_{10} \wedge \pi_{11} - \omega_{11} \wedge \pi_{10} + \omega_{13} \wedge \pi_9 - \omega_{15} \wedge \pi_4 + \omega_{16} \wedge \pi_3, \\
d\varpi_9 &\equiv \omega_5 \wedge \pi_{16} - \omega_6 \wedge \pi_{15} + \omega_9 \wedge \pi_{14} - \omega_{11} \wedge \pi_{12} - \omega_{12} \wedge \pi_{11} + \omega_{14} \wedge \pi_9 - \omega_{15} \wedge \pi_6 + \omega_{16} \wedge \pi_5, \\
d\varpi_{10} &\equiv \omega_7 \wedge \pi_{16} - \omega_8 \wedge \pi_{15} + \omega_{10} \wedge \pi_{14} - \omega_{12} \wedge \pi_{13} - \omega_{13} \wedge \pi_{12} + \omega_{14} \wedge \pi_{10} - \omega_{15} \wedge \pi_8 + \omega_{16} \wedge \pi_7, \\
& \pmod{\varpi_1, \dots, \varpi_{10}}.
\end{aligned}$$

Now, putting $\varpi = \varpi_1 + \sum_{i=2}^{10} \lambda_i \varpi_i$, we consider a point $v \in P(X)$ such that $v = \{\varpi = 0\} \subset T_x(X)$, where $x = \nu(v)$. Then we have

$$d\varpi \equiv \check{\omega}_1 \wedge \pi_1 + \dots + \check{\omega}_{16} \wedge \pi_{16} \pmod{\varpi_1, \dots, \varpi_{10}},$$

where

$$\begin{aligned}
\check{\omega}_1 &= \omega_{11} + \lambda_2 \omega_{13} + \lambda_3 \omega_{14} + \lambda_4 \omega_{15} + \lambda_6 \omega_{16}, & \check{\omega}_2 &= -\omega_9 - \lambda_2 \omega_{10} - \lambda_3 \omega_{12} + \lambda_5 \omega_{15} + \lambda_7 \omega_{16}, \\
\check{\omega}_3 &= \omega_6 + \lambda_2 \omega_8 - \lambda_4 \omega_{12} - \lambda_5 \omega_{14} + \lambda_8 \omega_{16}, & \check{\omega}_4 &= -\omega_5 - \lambda_2 \omega_7 - \lambda_6 \omega_{12} - \lambda_7 \omega_{14} - \lambda_8 \omega_{15}, \\
\check{\omega}_5 &= -\omega_4 + \lambda_3 \omega_8 + \lambda_4 \omega_{10} + \lambda_5 \omega_{13} + \lambda_9 \omega_{16}, & \check{\omega}_6 &= \omega_3 - \lambda_3 \omega_7 + \lambda_6 \omega_{10} + \lambda_7 \omega_{13} - \lambda_9 \omega_{15}, \\
\check{\omega}_7 &= -\lambda_2 \omega_4 - \lambda_3 \omega_6 - \lambda_4 \omega_9 - \lambda_5 \omega_{11} + \lambda_{10} \omega_{16}, & \check{\omega}_8 &= \lambda_2 \omega_3 + \lambda_3 \omega_5 - \lambda_6 \omega_9 - \lambda_7 \omega_{11} - \lambda_{10} \omega_{15}, \\
\check{\omega}_9 &= -\omega_2 - \lambda_4 \omega_7 - \lambda_6 \omega_8 + \lambda_8 \omega_{13} + \lambda_9 \omega_{14}, & \check{\omega}_{10} &= -\lambda_2 \omega_2 + \lambda_4 \omega_5 + \lambda_6 \omega_6 - \lambda_8 \omega_{11} + \lambda_{10} \omega_{14}, \\
\check{\omega}_{11} &= \omega_1 - \lambda_5 \omega_7 - \lambda_7 \omega_8 - \lambda_8 \omega_{10} - \lambda_9 \omega_{12}, & \check{\omega}_{12} &= -\lambda_3 \omega_2 - \lambda_4 \omega_3 - \lambda_6 \omega_4 - \lambda_9 \omega_{11} - \lambda_{10} \omega_{13}, \\
\check{\omega}_{13} &= \lambda_2 \omega_1 + \lambda_5 \omega_5 + \lambda_7 \omega_6 + \lambda_8 \omega_9 - \lambda_{10} \omega_{12}, & \check{\omega}_{14} &= \lambda_3 \omega_1 - \lambda_5 \omega_3 - \lambda_7 \omega_4 + \lambda_9 \omega_9 + \lambda_{10} \omega_{10}, \\
\check{\omega}_{15} &= \lambda_4 \omega_1 + \lambda_5 \omega_2 - \lambda_8 \omega_4 - \lambda_9 \omega_6 - \lambda_{10} \omega_8, & \check{\omega}_{16} &= \lambda_6 \omega_1 + \lambda_7 \omega_2 + \lambda_8 \omega_3 + \lambda_9 \omega_5 + \lambda_{10} \omega_7.
\end{aligned}$$

We calculate

$$\begin{aligned}
\check{\omega}_7 &= -\lambda_5 \check{\omega}_1 + \lambda_4 \check{\omega}_2 - \lambda_3 \check{\omega}_3 + \lambda_2 \check{\omega}_5 + \lambda \omega_{16}, & \check{\omega}_8 &= -\lambda_7 \check{\omega}_1 + \lambda_6 \check{\omega}_2 - \lambda_3 \check{\omega}_4 + \lambda_2 \check{\omega}_6 - \lambda \omega_{15}, \\
\check{\omega}_{10} &= -\lambda_8 \check{\omega}_1 + \lambda_6 \check{\omega}_3 - \lambda_4 \check{\omega}_4 + \lambda_2 \check{\omega}_9 + \lambda \omega_{14}, & \check{\omega}_{12} &= -\lambda_9 \check{\omega}_1 + \lambda_6 \check{\omega}_5 - \lambda_4 \check{\omega}_6 + \lambda_3 \check{\omega}_9 - \lambda \omega_{13}, \\
\check{\omega}_{13} &= -\lambda_8 \check{\omega}_2 + \lambda_7 \check{\omega}_3 - \lambda_5 \check{\omega}_4 + \lambda_2 \check{\omega}_{11} - \lambda \omega_{12}, & \check{\omega}_{14} &= -\lambda_9 \check{\omega}_2 + \lambda_7 \check{\omega}_5 - \lambda_5 \check{\omega}_6 + \lambda_3 \check{\omega}_{11} + \lambda \omega_{10}, \\
\check{\omega}_{15} &= -\lambda_9 \check{\omega}_3 + \lambda_8 \check{\omega}_5 - \lambda_5 \check{\omega}_9 + \lambda_4 \check{\omega}_{11} - \lambda \omega_8, & \check{\omega}_{16} &= -\lambda_9 \check{\omega}_4 + \lambda_8 \check{\omega}_6 - \lambda_7 \check{\omega}_9 + \lambda_6 \check{\omega}_{11} + \lambda \omega_7,
\end{aligned}$$

where $\lambda = \lambda_{10} - \lambda_2 \lambda_9 + \lambda_3 \lambda_8 - \lambda_4 \lambda_7 + \lambda_5 \lambda_6$. Thus we see that (X, D) satisfies the condition (X.1) to (X.3) in §4.3 and we obtain

$$R(X) = \{v \in P(X) \mid \lambda \neq 0\},$$

where $(\lambda_2, \dots, \lambda_{10})$ is the inhomogeneous coordinate of the fibre $\nu : P(X) \rightarrow X$. Moreover we have

$$\hat{f}(v) = \{X \in \mathfrak{g}_{-1}(x) \mid [X, \mathfrak{g}_{-2}(x)] = 0\} \quad \text{for each } v \in R(X), x = \nu(v).$$

and $\dim \hat{f}(v) = 1$. Utilizing the calculation in Case (4) of §4 in [YY2], we have

$$f = (e_1^* \otimes e_{11}^* - e_2^* \otimes e_9^* + e_3^* \otimes e_6^* - e_4^* \otimes e_5^*) + \lambda_2 (e_1^* \otimes e_{13}^* - e_2^* \otimes e_{10}^* + e_3^* \otimes e_8^* - e_4^* \otimes e_7^*)$$

$$\begin{aligned}
& + \lambda_3(e_1^* \otimes e_{14}^* - e_2^* \otimes e_{12}^* + e_5^* \otimes e_8^* - e_6^* \otimes e_7^*) + \lambda_4(e_1^* \otimes e_{15}^* - e_3^* \otimes e_{12}^* + e_5^* \otimes e_{10}^* - e_7^* \otimes e_9^*) \\
& + \lambda_5(e_2^* \otimes e_{15}^* - e_3^* \otimes e_{14}^* + e_5^* \otimes e_{13}^* - e_7^* \otimes e_{11}^*) + \lambda_6(e_1^* \otimes e_{16}^* - e_4^* \otimes e_{12}^* + e_6^* \otimes e_{10}^* - e_8^* \otimes e_9^*) \\
& + \lambda_7(e_2^* \otimes e_{16}^* - e_4^* \otimes e_{14}^* + e_6^* \otimes e_{13}^* - e_8^* \otimes e_{11}^*) + \lambda_8(e_3^* \otimes e_{16}^* - e_4^* \otimes e_{15}^* + e_9^* \otimes e_{13}^* - e_{10}^* \otimes e_{11}^*) \\
& + \lambda_9(e_5^* \otimes e_{16}^* - e_6^* \otimes e_{15}^* + e_9^* \otimes e_{14}^* - e_{11}^* \otimes e_{12}^*) + \lambda_{10}(e_7^* \otimes e_{16}^* - e_8^* \otimes e_{15}^* + e_{10}^* \otimes e_{14}^* - e_{12}^* \otimes e_{13}^*)
\end{aligned}$$

for the generator f of $\hat{\mathfrak{f}}(v)$. Then we calculate

$$\begin{aligned}
f &= \alpha_1 \otimes \alpha_{11} - \alpha_2 \otimes \alpha_9 + \alpha_3 \otimes \alpha_6 - \alpha_4 \otimes \alpha_5 \\
&\quad + \lambda(e_7^* \otimes e_{16}^* - e_8^* \otimes e_{15}^* + e_{10}^* \otimes e_{14}^* - e_{12}^* \otimes e_{13}^*),
\end{aligned}$$

where $\lambda = \lambda_{10} - \lambda_2\lambda_9 + \lambda_3\lambda_8 - \lambda_4\lambda_7 + \lambda_5\lambda_6$ and

$$\begin{aligned}
\alpha_1 &= e_{11}^* + \lambda_2 e_{13}^* + \lambda_3 e_{14}^* + \lambda_4 e_{15}^* + \lambda_6 e_{16}^*, & \alpha_2 &= -e_9^* - \lambda_2 e_{10}^* - \lambda_3 e_{12}^* + \lambda_5 e_{15}^* + \lambda_7 e_{16}^*, \\
\alpha_3 &= e_6^* + \lambda_2 e_8^* - \lambda_4 e_{12}^* - \lambda_5 e_{14}^* + \lambda_8 e_{16}^*, & \alpha_4 &= -e_5^* - \lambda_2 e_7^* - \lambda_6 e_{12}^* - \lambda_7 e_{14}^* - \lambda_8 e_{15}^*, \\
\alpha_5 &= -e_4^* + \lambda_3 e_8^* + \lambda_4 e_{10}^* + \lambda_5 e_{13}^* + \lambda_9 e_{16}^*, & \alpha_6 &= e_3^* - \lambda_3 e_7^* + \lambda_6 e_{10}^* + \lambda_7 e_{13}^* - \lambda_9 e_{15}^*, \\
\alpha_9 &= -e_2^* - \lambda_4 e_7^* - \lambda_6 e_8^* + \lambda_8 e_{13}^* + \lambda_9 e_{14}^*, & \alpha_{11} &= e_1^* - \lambda_5 e_7^* - \lambda_7 e_8^* - \lambda_8 e_{10}^* - \lambda_9 e_{12}^*,
\end{aligned}$$

Thus f is a non-degenerate quadratic form in $S^2(W^*)$, where $W = \langle \{e_1, \dots, e_{16}\} \rangle$. Hence $\hat{\mathfrak{f}}(v)$ is of finite type (see Case (5) of §3 in [YY2]).

Now let us construct the model equation of second order from the coordinate description of the standard differential system (M_m, D_m) . We calculate

$$\begin{aligned}
\varpi &= \varpi_1 + \lambda_2 \varpi_2 + \dots + \lambda_{10} \varpi_{10} \\
&= dy_1 + \lambda_2 dy_2 + \dots + \lambda_{10} dy_{10} - \hat{p}_1 dx_1 - \dots - \hat{p}_{16} dx_{16} \\
&= d(y_1 + \lambda_2 y_2 + \dots + \lambda_{10} y_{10}) - y_2 d\lambda_2 - \dots - y_{10} d\lambda_{10} - \hat{p}_1 dx_1 - \dots - \hat{p}_{16} dx_{16}, \\
&= dz - \hat{p}_1 dx_1 - \dots - \hat{p}_{16} dx_{16} - \hat{p}_{17} dx_{17} - \dots - \hat{p}_{25} dx_{25},
\end{aligned}$$

where we put

$$\begin{aligned}
z &= y_1 + \lambda_2 y_2 + \dots + \lambda_{10} y_{10}, \\
\hat{p}_1 &= p_{11} + \lambda_2 p_{13} + \lambda_3 p_{14} + \lambda_4 p_{15} + \lambda_6 p_{16}, & \hat{p}_2 &= -p_9 - \lambda_2 p_{10} - \lambda_3 p_{12} + \lambda_5 p_{15} + \lambda_7 p_{16}, \\
\hat{p}_3 &= p_6 + \lambda_2 p_8 - \lambda_4 p_{12} - \lambda_5 p_{14} + \lambda_8 p_{16}, & \hat{p}_4 &= -p_5 - \lambda_2 p_7 - \lambda_6 p_{12} - \lambda_7 p_{14} - \lambda_8 p_{15}, \\
\hat{p}_5 &= -p_4 + \lambda_3 p_8 + \lambda_4 p_{10} + \lambda_5 p_{13} + \lambda_9 p_{16}, & \hat{p}_6 &= p_3 - \lambda_3 p_7 + \lambda_6 p_{10} + \lambda_7 p_{13} - \lambda_9 p_{15}, \\
\hat{p}_7 &= -\lambda_2 p_4 - \lambda_3 p_6 - \lambda_4 p_9 - \lambda_5 p_{11} + \lambda_{10} p_{16}, & \hat{p}_8 &= \lambda_2 p_3 + \lambda_3 p_5 - \lambda_6 p_9 - \lambda_7 p_{11} - \lambda_{10} p_{15}, \\
\hat{p}_9 &= -p_2 - \lambda_4 p_7 - \lambda_6 p_8 + \lambda_8 p_{13} + \lambda_9 p_{14}, & \hat{p}_{10} &= -\lambda_2 p_2 + \lambda_4 p_5 + \lambda_6 p_6 - \lambda_8 p_{11} + \lambda_{10} p_{14}, \\
\hat{p}_{11} &= p_1 - \lambda_5 p_7 - \lambda_7 p_8 - \lambda_8 p_{10} - \lambda_9 p_{12}, & \hat{p}_{12} &= -\lambda_3 p_2 - \lambda_4 p_3 - \lambda_6 p_4 - \lambda_9 p_{11} - \lambda_{10} p_{13}, \\
\hat{p}_{13} &= \lambda_2 p_1 + \lambda_5 p_5 + \lambda_7 p_6 + \lambda_8 p_9 - \lambda_{10} p_{12}, & \hat{p}_{14} &= \lambda_3 p_1 - \lambda_5 p_3 - \lambda_7 p_4 + \lambda_9 p_9 + \lambda_{10} p_{10}, \\
\hat{p}_{15} &= \lambda_4 p_1 + \lambda_5 p_2 - \lambda_8 p_4 - \lambda_9 p_6 - \lambda_{10} p_8, & \hat{p}_{16} &= \lambda_6 p_1 + \lambda_7 p_2 + \lambda_8 p_3 + \lambda_9 p_5 + \lambda_{10} p_7, \\
\hat{p}_{17} &= y_2, \dots, \hat{p}_{\alpha+15} = y_\alpha, \dots, \hat{p}_{25} = y_{10}, & x_{17} &= \lambda_2, \dots, x_{\alpha+15} = \lambda_\alpha, \dots, x_{25} = \lambda_{10}.
\end{aligned}$$

Then we have, for $(R(M_m); D_{M_m}^1, D_{M_m}^2)$,

$$D_{M_m}^1 = \{\varpi = 0\}, \quad D_{M_m}^2 = \{\varpi_k = \pi_i = 0 \quad (1 \leq k \leq 10, 1 \leq i \leq 16)\},$$

where we denote the pullback on $R(M_m)$ of 1-forms on M_m by the same symbol. By taking the exterior derivatives of both sides of the above defining equations for \hat{p}_i ($i = 1, \dots, 16$), we put

$$\hat{\pi}_1 = d\hat{p}_1 - a(dx_{11} + x_{17}dx_{13} + x_{18}dx_{14} + x_{19}dx_{15} + x_{21}dx_{16})$$

$$\begin{aligned}
& -p_{13}dx_{17} - p_{14}dx_{18} - p_{15}dx_{19} - p_{16}dx_{21}, \\
\hat{\pi}_2 = d\hat{p}_2 - a(-dx_9 - x_{17}dx_{10} - x_{18}dx_{12} + x_{20}dx_{15} + x_{22}dx_{16}) \\
& \quad + p_{10}dx_{17} + p_{12}dx_{18} - p_{15}dx_{20} - p_{16}dx_{22}, \\
\hat{\pi}_3 = d\hat{p}_3 - a(dx_6 + x_{17}dx_8 - x_{19}dx_{12} - x_{20}dx_{14} + x_{23}dx_{16}) \\
& \quad - p_8dx_{17} + p_{12}dx_{19} + p_{14}dx_{20} - p_{16}dx_{23}, \\
\hat{\pi}_4 = d\hat{p}_4 - a(-dx_5 - x_{17}dx_7 - x_{21}dx_{12} - x_{22}dx_{14} - x_{23}dx_{15}) \\
& \quad + p_7dx_{17} + p_{12}dx_{21} + p_{14}dx_{22} + p_{15}dx_{23}, \\
\hat{\pi}_5 = d\hat{p}_5 - a(-dx_4 + x_{18}dx_8 + x_{19}dx_{10} + x_{20}dx_{13} + x_{24}dx_{16}) \\
& \quad - p_8dx_{18} - p_{10}dx_{19} - p_{13}dx_{20} - p_{16}dx_{24}, \\
\hat{\pi}_6 = d\hat{p}_6 - a(dx_3 - x_{18}dx_7 + x_{21}dx_{10} + x_{22}dx_{13} - x_{24}dx_{15}) \\
& \quad + p_7dx_{18} - p_{10}dx_{21} - p_{13}dx_{22} + p_{15}dx_{24}, \\
\hat{\pi}_7 = d\hat{p}_7 - a(-x_{17}dx_4 - x_{18}dx_6 - x_{19}dx_9 - x_{20}dx_{11} + x_{25}dx_{16}) \\
& \quad + p_4dx_{17} + p_6dx_{18} + p_9dx_{19} + p_{11}dx_{20} - p_{16}dx_{25}, \\
\hat{\pi}_8 = d\hat{p}_8 - a(x_{17}dx_3 + x_{18}dx_5 - x_{21}dx_9 - x_{22}dx_{11} - x_{25}dx_{15}) \\
& \quad - p_3dx_{17} - p_5dx_{18} + p_9dx_{21} + p_{11}dx_{22} + p_{15}dx_{25}, \\
\hat{\pi}_9 = d\hat{p}_9 - a(-dx_2 - x_{19}dx_7 - x_{21}dx_8 + x_{23}dx_{13} + x_{24}dx_{14}) \\
& \quad + p_7dx_{19} + p_8dx_{21} - p_{13}dx_{23} - p_{14}dx_{24}, \\
\hat{\pi}_{10} = d\hat{p}_{10} - a(-x_{17}dx_2 + x_{19}dx_5 + x_{21}dx_6 - x_{23}dx_{11} + x_{25}dx_{14}) \\
& \quad + p_2dx_{17} - p_5dx_{19} - p_6dx_{21} + p_{11}dx_{23} - p_{14}dx_{25}, \\
\hat{\pi}_{11} = d\hat{p}_{11} - a(dx_1 - x_{20}dx_7 - x_{22}dx_8 - x_{23}dx_{10} - x_{24}dx_{12}) \\
& \quad + p_7dx_{20} + p_8dx_{22} + p_{10}dx_{23} + p_{12}dx_{24}, \\
\hat{\pi}_{12} = d\hat{p}_{12} - a(-x_{18}dx_2 - x_{19}dx_3 - x_{21}dx_4 - x_{24}dx_{11} - x_{25}dx_{13}) \\
& \quad + p_2dx_{18} + p_3dx_{19} + p_4dx_{21} + p_{11}dx_{24} + p_{13}dx_{25}, \\
\hat{\pi}_{13} = d\hat{p}_{13} - a(x_{17}dx_1 + x_{20}dx_5 + x_{22}dx_6 + x_{23}dx_9 - x_{25}dx_{12}) \\
& \quad - p_1dx_{17} - p_5dx_{20} - p_6dx_{22} - p_9dx_{23} + p_{12}dx_{25}, \\
\hat{\pi}_{14} = d\hat{p}_{14} - a(x_{18}dx_1 - x_{20}dx_3 - x_{22}dx_4 + x_{24}dx_9 + x_{25}dx_{10}) \\
& \quad - p_1dx_{18} + p_3dx_{20} + p_4dx_{22} - p_9dx_{24} - p_{10}dx_{25}, \\
\hat{\pi}_{15} = d\hat{p}_{15} - a(x_{19}dx_1 + x_{20}dx_2 - x_{23}dx_4 - x_{24}dx_6 - x_{25}dx_8) \\
& \quad - p_1dx_{19} - p_2dx_{20} + p_4dx_{23} + p_6dx_{24} + p_8dx_{25}, \\
\hat{\pi}_{16} = d\hat{p}_{16} - a(x_{21}dx_1 + x_{22}dx_2 + x_{23}dx_3 + x_{24}dx_5 + x_{25}dx_7) \\
& \quad - p_1dx_{21} - p_2dx_{22} - p_3dx_{23} - p_5dx_{24} - p_7dx_{25}.
\end{aligned}$$

Then we see that $\{\hat{\pi}_1, \dots, \hat{\pi}_{16}\}$ can be written as the linear combinations of $\{\pi_1, \dots, \pi_{16}\}$ with the same coefficients (in λ 's) such that $\{\hat{p}_1, \dots, \hat{p}_{16}\}$ are written as the linear combination of $\{p_1, \dots, p_{16}\}$ as in the above equations. Hence we have

$$D_{M_m}^1 = \{\varpi = o\}, \quad D_{M_m}^2 = \{\varpi = \hat{\pi}_1 = \dots = \hat{\pi}_{16} = \hat{\pi}_{17} = \dots = \hat{\pi}_{25} = 0\},$$

where $\hat{\pi}_{15+\alpha} = \varpi_\alpha$ ($2 \leq \alpha \leq 10$) are written as follows:

$$\hat{\pi}_{17} = d\hat{p}_{17} - p_{13}dx_1 + p_{10}dx_2 - p_8dx_3 + p_7dx_4 + p_4dx_7 - p_3dx_8 + p_2dx_{10} - p_1dx_{13},$$

$$\begin{aligned}
\hat{\pi}_{18} &= d\hat{p}_{18} - p_{14}dx_1 + p_{12}dx_2 - p_8dx_5 + p_7dx_6 + p_6dx_7 - p_5dx_8 + p_2dx_{12} - p_1dx_{14}, \\
\hat{\pi}_{19} &= d\hat{p}_{19} - p_{15}dx_1 + p_{12}dx_3 - p_{10}dx_5 + p_9dx_7 + p_7dx_9 - p_5dx_{10} + p_3dx_{12} - p_1dx_{15}, \\
\hat{\pi}_{20} &= d\hat{p}_{20} - p_{15}dx_2 + p_{14}dx_3 - p_{13}dx_5 + p_{11}dx_7 + p_7dx_{11} - p_5dx_{13} + p_3dx_{14} - p_2dx_{15}, \\
\hat{\pi}_{21} &= d\hat{p}_{21} - p_{16}dx_1 + p_{12}dx_4 - p_{10}dx_6 + p_9dx_8 + p_8dx_9 - p_6dx_{10} + p_4dx_{12} - p_1dx_{16}, \\
\hat{\pi}_{22} &= d\hat{p}_{22} - p_{16}dx_2 + p_{14}dx_4 - p_{13}dx_6 + p_{11}dx_8 + p_8dx_{11} - p_6dx_{13} + p_4dx_{14} - p_2dx_{16}, \\
\hat{\pi}_{23} &= d\hat{p}_{23} - p_{16}dx_3 + p_{15}dx_4 - p_{13}dx_9 + p_{11}dx_{10} + p_{10}dx_{11} - p_9dx_{13} + p_4dx_{15} - p_3dx_{16}, \\
\hat{\pi}_{24} &= d\hat{p}_{24} - p_{16}dx_5 + p_{15}dx_6 - p_{14}dx_9 + p_{12}dx_{11} + p_{11}dx_{12} - p_9dx_{14} + p_6dx_{15} - p_5dx_{16}, \\
\hat{\pi}_{25} &= d\hat{p}_{25} - p_{16}dx_7 + p_{15}dx_8 - p_{14}dx_{10} + p_{13}dx_{12} + p_{12}dx_{13} - p_{10}dx_{14} + p_8dx_{15} - p_7dx_{16}.
\end{aligned}$$

Moreover we calculate

$$\begin{aligned}
\lambda p_{16} &= x_{20}\hat{p}_1 - x_{19}\hat{p}_2 + x_{18}\hat{p}_3 - x_{17}\hat{p}_5 + \hat{p}_7, \\
\lambda p_{15} &= -x_{22}\hat{p}_1 + x_{21}\hat{p}_2 - x_{18}\hat{p}_4 + x_{17}\hat{p}_6 - \hat{p}_8, \\
\lambda p_{14} &= x_{23}\hat{p}_1 - x_{21}\hat{p}_3 + x_{19}\hat{p}_4 - x_{17}\hat{p}_9 + \hat{p}_{10}, \\
\lambda p_{13} &= -x_{24}\hat{p}_1 + x_{21}\hat{p}_5 - x_{19}\hat{p}_6 + x_{18}\hat{p}_9 - \hat{p}_{12}, \\
\lambda p_{12} &= -x_{23}\hat{p}_2 + x_{22}\hat{p}_3 - x_{20}\hat{p}_4 + x_{17}\hat{p}_{11} - \hat{p}_{13}, \\
\lambda p_{11} &= x_{25}\hat{p}_1 - x_{21}\hat{p}_7 + x_{19}\hat{p}_8 - x_{18}\hat{p}_{10} + x_{17}\hat{p}_{12}, \\
\lambda p_{10} &= x_{24}\hat{p}_2 - x_{22}\hat{p}_5 + x_{20}\hat{p}_6 - x_{18}\hat{p}_{11} + \hat{p}_{14}, \\
\lambda p_9 &= -x_{25}\hat{p}_2 + x_{22}\hat{p}_7 - x_{20}\hat{p}_8 + x_{18}\hat{p}_{13} - x_{17}\hat{p}_{14}, \\
\lambda p_8 &= -x_{24}\hat{p}_3 + x_{23}\hat{p}_5 - x_{20}\hat{p}_9 + x_{19}\hat{p}_{11} - \hat{p}_{15}, \\
\lambda p_7 &= x_{24}\hat{p}_4 - x_{23}\hat{p}_6 + x_{22}\hat{p}_9 - x_{21}\hat{p}_{11} + \hat{p}_{16}, \\
\lambda p_6 &= x_{25}\hat{p}_3 - x_{23}\hat{p}_7 + x_{20}\hat{p}_{10} - x_{19}\hat{p}_{13} + x_{17}\hat{p}_{15}, \\
\lambda p_5 &= -x_{25}\hat{p}_4 + x_{23}\hat{p}_8 - x_{22}\hat{p}_{10} + x_{21}\hat{p}_{13} - x_{17}\hat{p}_{16}, \\
\lambda p_4 &= -x_{25}\hat{p}_5 + x_{24}\hat{p}_7 - x_{20}\hat{p}_{12} + x_{19}\hat{p}_{14} - x_{18}\hat{p}_{15}, \\
\lambda p_3 &= x_{25}\hat{p}_6 - x_{24}\hat{p}_8 + x_{22}\hat{p}_{12} - x_{21}\hat{p}_{14} + x_{18}\hat{p}_{16}, \\
\lambda p_2 &= -x_{25}\hat{p}_9 + x_{24}\hat{p}_{10} - x_{23}\hat{p}_{12} + x_{21}\hat{p}_{15} - x_{19}\hat{p}_{16}, \\
\lambda p_1 &= x_{25}\hat{p}_{11} - x_{24}\hat{p}_{13} + x_{23}\hat{p}_{14} - x_{22}\hat{p}_{15} + x_{20}\hat{p}_{16},
\end{aligned}$$

where $\lambda = x_{25} - x_{17}x_{24} + x_{18}x_{23} - x_{19}x_{22} + x_{20}x_{21}$. Thus we obtain the following model equation of second order :

$$\begin{aligned}
\frac{\partial^2 z}{\partial x_1 \partial x_{11}} &= -\frac{\partial^2 z}{\partial x_2 \partial x_9} = \frac{\partial^2 z}{\partial x_3 \partial x_6} = -\frac{\partial^2 z}{\partial x_4 \partial x_5} \quad (= a) \\
\frac{\partial^2 z}{\partial x_1 \partial x_{13}} &= -\frac{\partial^2 z}{\partial x_2 \partial x_{10}} = \frac{\partial^2 z}{\partial x_3 \partial x_8} = -\frac{\partial^2 z}{\partial x_4 \partial x_7} = x_{17} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_1 \partial x_{14}} &= -\frac{\partial^2 z}{\partial x_2 \partial x_{12}} = \frac{\partial^2 z}{\partial x_5 \partial x_8} = -\frac{\partial^2 z}{\partial x_6 \partial x_7} = x_{18} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_1 \partial x_{15}} &= -\frac{\partial^2 z}{\partial x_3 \partial x_{12}} = \frac{\partial^2 z}{\partial x_5 \partial x_{10}} = -\frac{\partial^2 z}{\partial x_7 \partial x_9} = x_{19} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_2 \partial x_{15}} &= -\frac{\partial^2 z}{\partial x_3 \partial x_{14}} = \frac{\partial^2 z}{\partial x_5 \partial x_{13}} = -\frac{\partial^2 z}{\partial x_7 \partial x_{11}} = x_{20} \frac{\partial^2 z}{\partial x_1 \partial x_{11}}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial^2 z}{\partial x_1 \partial x_{16}} &= -\frac{\partial^2 z}{\partial x_4 \partial x_{12}} = \frac{\partial^2 z}{\partial x_6 \partial x_{10}} = -\frac{\partial^2 z}{\partial x_8 \partial x_9} = x_{21} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_2 \partial x_{16}} &= -\frac{\partial^2 z}{\partial x_4 \partial x_{14}} = \frac{\partial^2 z}{\partial x_6 \partial x_{13}} = -\frac{\partial^2 z}{\partial x_8 \partial x_{11}} = x_{22} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_3 \partial x_{16}} &= -\frac{\partial^2 z}{\partial x_4 \partial x_{15}} = \frac{\partial^2 z}{\partial x_9 \partial x_{13}} = -\frac{\partial^2 z}{\partial x_{10} \partial x_{11}} = x_{23} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_5 \partial x_{16}} &= -\frac{\partial^2 z}{\partial x_6 \partial x_{15}} = \frac{\partial^2 z}{\partial x_9 \partial x_{14}} = -\frac{\partial^2 z}{\partial x_{11} \partial x_{12}} = x_{24} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_7 \partial x_{16}} &= -\frac{\partial^2 z}{\partial x_8 \partial x_{15}} = \frac{\partial^2 z}{\partial x_{10} \partial x_{14}} = -\frac{\partial^2 z}{\partial x_{12} \partial x_{13}} = x_{25} \frac{\partial^2 z}{\partial x_1 \partial x_{11}} \\
\frac{\partial^2 z}{\partial x_{13} \partial x_{17}} &= \frac{\partial^2 z}{\partial x_{14} \partial x_{18}} = \frac{\partial^2 z}{\partial x_{15} \partial x_{19}} = \frac{\partial^2 z}{\partial x_{16} \partial x_{21}} \quad (= p_1) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_{11}} - x_{24} \frac{\partial z}{\partial x_{13}} + x_{23} \frac{\partial z}{\partial x_{14}} - x_{22} \frac{\partial z}{\partial x_{15}} + x_{20} \frac{\partial z}{\partial x_{16}} \right), \\
\frac{\partial^2 z}{\partial x_{10} \partial x_{17}} &= \frac{\partial^2 z}{\partial x_{12} \partial x_{18}} = -\frac{\partial^2 z}{\partial x_{15} \partial x_{20}} = -\frac{\partial^2 z}{\partial x_{16} \partial x_{22}} \quad (= -p_2) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_9} - x_{24} \frac{\partial z}{\partial x_{10}} + x_{23} \frac{\partial z}{\partial x_{12}} - x_{21} \frac{\partial z}{\partial x_{15}} + x_{19} \frac{\partial z}{\partial x_{16}} \right), \\
\frac{\partial^2 z}{\partial x_8 \partial x_{17}} &= -\frac{\partial^2 z}{\partial x_{12} \partial x_{19}} = -\frac{\partial^2 z}{\partial x_{14} \partial x_{20}} = \frac{\partial^2 z}{\partial x_{16} \partial x_{23}} \quad (= p_3) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_6} - x_{24} \frac{\partial z}{\partial x_8} + x_{22} \frac{\partial z}{\partial x_{12}} - x_{21} \frac{\partial z}{\partial x_{14}} + x_{18} \frac{\partial z}{\partial x_{16}} \right), \\
\frac{\partial^2 z}{\partial x_7 \partial x_{17}} &= \frac{\partial^2 z}{\partial x_{12} \partial x_{21}} = \frac{\partial^2 z}{\partial x_{14} \partial x_{22}} = \frac{\partial^2 z}{\partial x_{15} \partial x_{23}} \quad (= -p_4) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_5} - x_{24} \frac{\partial z}{\partial x_7} + x_{20} \frac{\partial z}{\partial x_{12}} - x_{19} \frac{\partial z}{\partial x_{14}} + x_{18} \frac{\partial z}{\partial x_{15}} \right), \\
\frac{\partial^2 z}{\partial x_8 \partial x_{18}} &= \frac{\partial^2 z}{\partial x_{10} \partial x_{19}} = \frac{\partial^2 z}{\partial x_{13} \partial x_{20}} = \frac{\partial^2 z}{\partial x_{16} \partial x_{24}} \quad (= p_5) \\
&= \lambda^{-1} \left(-x_{25} \frac{\partial z}{\partial x_4} + x_{23} \frac{\partial z}{\partial x_8} - x_{22} \frac{\partial z}{\partial x_{10}} + x_{21} \frac{\partial z}{\partial x_{13}} - x_{17} \frac{\partial z}{\partial x_{16}} \right), \\
-\frac{\partial^2 z}{\partial x_7 \partial x_{18}} &= \frac{\partial^2 z}{\partial x_{10} \partial x_{21}} = \frac{\partial^2 z}{\partial x_{13} \partial x_{22}} = -\frac{\partial^2 z}{\partial x_{15} \partial x_{24}} \quad (= p_6) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_3} - x_{23} \frac{\partial z}{\partial x_7} + x_{20} \frac{\partial z}{\partial x_{10}} - x_{19} \frac{\partial z}{\partial x_{13}} + x_{17} \frac{\partial z}{\partial x_{15}} \right), \\
-\frac{\partial^2 z}{\partial x_4 \partial x_{17}} &= -\frac{\partial^2 z}{\partial x_6 \partial x_{18}} = -\frac{\partial^2 z}{\partial x_9 \partial x_{19}} = -\frac{\partial^2 z}{\partial x_{11} \partial x_{20}} = \frac{\partial^2 z}{\partial x_{16} \partial x_{25}} \quad (= p_7) \\
&= \lambda^{-1} \left(x_{24} \frac{\partial z}{\partial x_4} - x_{23} \frac{\partial z}{\partial x_6} + x_{22} \frac{\partial z}{\partial x_9} - x_{21} \frac{\partial z}{\partial x_{11}} + \frac{\partial z}{\partial x_{16}} \right), \\
-\frac{\partial^2 z}{\partial x_3 \partial x_{17}} &= -\frac{\partial^2 z}{\partial x_5 \partial x_{18}} = \frac{\partial^2 z}{\partial x_9 \partial x_{21}} = \frac{\partial^2 z}{\partial x_{11} \partial x_{22}} = \frac{\partial^2 z}{\partial x_{15} \partial x_{25}} \quad (= -p_8) \\
&= \lambda^{-1} \left(x_{24} \frac{\partial z}{\partial x_3} - x_{23} \frac{\partial z}{\partial x_5} + x_{20} \frac{\partial z}{\partial x_9} - x_{19} \frac{\partial z}{\partial x_{11}} + \frac{\partial z}{\partial x_{15}} \right),
\end{aligned}$$

$$\begin{aligned}
\frac{\partial^2 z}{\partial x_7 \partial x_{19}} &= \frac{\partial^2 z}{\partial x_8 \partial x_{21}} = -\frac{\partial^2 z}{\partial x_{13} \partial x_{23}} = -\frac{\partial^2 z}{\partial x_{14} \partial x_{24}} \quad (= -p_9) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_2} - x_{22} \frac{\partial z}{\partial x_7} + x_{20} \frac{\partial z}{\partial x_8} - x_{18} \frac{\partial z}{\partial x_{13}} + x_{17} \frac{\partial z}{\partial x_{14}} \right), \\
-\frac{\partial^2 z}{\partial x_2 \partial x_{17}} &= \frac{\partial^2 z}{\partial x_5 \partial x_{19}} = \frac{\partial^2 z}{\partial x_6 \partial x_{21}} = -\frac{\partial^2 z}{\partial x_{11} \partial x_{23}} = \frac{\partial^2 z}{\partial x_{14} \partial x_{25}} \quad (= p_{10}) \\
&= \lambda^{-1} \left(x_{24} \frac{\partial z}{\partial x_2} - x_{22} \frac{\partial z}{\partial x_5} + x_{20} \frac{\partial z}{\partial x_6} - x_{18} \frac{\partial z}{\partial x_{11}} + \frac{\partial z}{\partial x_{14}} \right), \\
-\frac{\partial^2 z}{\partial x_7 \partial x_{20}} &= -\frac{\partial^2 z}{\partial x_8 \partial x_{22}} = -\frac{\partial^2 z}{\partial x_{10} \partial x_{23}} = -\frac{\partial^2 z}{\partial x_{12} \partial x_{24}} \quad (= p_{11}) \\
&= \lambda^{-1} \left(x_{25} \frac{\partial z}{\partial x_1} - x_{21} \frac{\partial z}{\partial x_7} + x_{19} \frac{\partial z}{\partial x_8} - x_{18} \frac{\partial z}{\partial x_{10}} + x_{17} \frac{\partial z}{\partial x_{12}} \right), \\
\frac{\partial^2 z}{\partial x_2 \partial x_{18}} &= \frac{\partial^2 z}{\partial x_3 \partial x_{19}} = \frac{\partial^2 z}{\partial x_4 \partial x_{21}} = \frac{\partial^2 z}{\partial x_{11} \partial x_{24}} = \frac{\partial^2 z}{\partial x_{13} \partial x_{25}} \quad (= -p_{12}) \\
&= \lambda^{-1} \left(x_{23} \frac{\partial z}{\partial x_2} - x_{22} \frac{\partial z}{\partial x_3} + x_{20} \frac{\partial z}{\partial x_4} - x_{17} \frac{\partial z}{\partial x_{11}} + \frac{\partial z}{\partial x_{13}} \right), \\
-\frac{\partial^2 z}{\partial x_1 \partial x_{17}} &= -\frac{\partial^2 z}{\partial x_5 \partial x_{20}} = -\frac{\partial^2 z}{\partial x_6 \partial x_{22}} = -\frac{\partial^2 z}{\partial x_9 \partial x_{23}} = \frac{\partial^2 z}{\partial x_{12} \partial x_{25}} \quad (= -p_{13}) \\
&= \lambda^{-1} \left(x_{24} \frac{\partial z}{\partial x_1} - x_{21} \frac{\partial z}{\partial x_5} + x_{19} \frac{\partial z}{\partial x_6} - x_{18} \frac{\partial z}{\partial x_9} + \frac{\partial z}{\partial x_{12}} \right), \\
\frac{\partial^2 z}{\partial x_1 \partial x_{18}} &= -\frac{\partial^2 z}{\partial x_3 \partial x_{20}} = -\frac{\partial^2 z}{\partial x_4 \partial x_{22}} = \frac{\partial^2 z}{\partial x_9 \partial x_{24}} = \frac{\partial^2 z}{\partial x_{10} \partial x_{25}} \quad (= p_{14}) \\
&= \lambda^{-1} \left(x_{23} \frac{\partial z}{\partial x_1} - x_{21} \frac{\partial z}{\partial x_3} + x_{19} \frac{\partial z}{\partial x_4} - x_{17} \frac{\partial z}{\partial x_9} + \frac{\partial z}{\partial x_{10}} \right), \\
-\frac{\partial^2 z}{\partial x_1 \partial x_{19}} &= -\frac{\partial^2 z}{\partial x_2 \partial x_{20}} = \frac{\partial^2 z}{\partial x_4 \partial x_{23}} = \frac{\partial^2 z}{\partial x_6 \partial x_{24}} = \frac{\partial^2 z}{\partial x_8 \partial x_{25}} \quad (= -p_{15}) \\
&= \lambda^{-1} \left(x_{22} \frac{\partial z}{\partial x_1} - x_{21} \frac{\partial z}{\partial x_2} + x_{18} \frac{\partial z}{\partial x_4} - x_{17} \frac{\partial z}{\partial x_6} + \frac{\partial z}{\partial x_8} \right), \\
\frac{\partial^2 z}{\partial x_1 \partial x_{21}} &= \frac{\partial^2 z}{\partial x_2 \partial x_{22}} = \frac{\partial^2 z}{\partial x_3 \partial x_{23}} = \frac{\partial^2 z}{\partial x_5 \partial x_{24}} = \frac{\partial^2 z}{\partial x_7 \partial x_{25}} \quad (= p_{16}) \\
&= \lambda^{-1} \left(x_{20} \frac{\partial z}{\partial x_1} - x_{19} \frac{\partial z}{\partial x_2} + x_{18} \frac{\partial z}{\partial x_3} - x_{17} \frac{\partial z}{\partial x_5} + \frac{\partial z}{\partial x_7} \right), \\
\frac{\partial^2 z}{\partial x_i \partial x_j} &= 0 \quad \text{otherwise.}
\end{aligned}$$

One can check that, among simple graded Lie algebras (of depth 3) of class (D) in §5 of [YY2], regular differential systems (X, D) of type \mathfrak{m} satisfy the condition (X.1) to (X.3) in §4.3 when \mathfrak{m} is the negative part of one of the simple graded Lie algebras $(C_\ell, \{\alpha_i, \alpha_\ell\})$ for $i = 2, \dots, \ell - 1$, $(D_\ell, \{\alpha_i, \alpha_\ell\})$ ($2 < i < \ell - 1$), when i is even, or $(E_7, \{\alpha_6, \alpha_7\})$, whereas the condition (X.3) is not satisfied by the other cases.

REFERENCES

- [Bai] T.N.Baily, *Parabolic Invariant Theory and Geometry* in “The Penrose Transform and Analytic Cohomology in Representation Theory” Contemp. Math. **154**, Amer. Math. Soc., (1993).
- [Bo] W. M. Boothby, *Homogeneous complex contact manifolds*, Proc. Symp. Pure Math., Amer. Math. Soc. **3** (1961), 144–154.
- [Bu] N.Bourbaki, *Groupes et algèbres de Lie, Chapitre 4,5 et 6*, Hermann, Paris (1968).
- [BCG₃] R. Bryant, S. S. Chern, R. B. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Springer-Verlag, New-York (1986).
- [C1] E.Cartan, *Les systèmes de Pfaff à cinq variables et les équations aux dérivées partielles du second ordre*, Ann. Ec. Normale **27** (1910), 109–192.
- [C2] -----, *Sur les systèmes en involution d'équations aux dérivées partielles du second ordre à une fonction inconnue de trois variables indépendantes*, Bull. Soc. Math. France **39** (1911), 352–443.
- [HY] J.Hwang and K. Yamaguchi, *Characterization of hermitian Symmetric Spaces by Fundamental Forms*, Duke Math. J. **120** No.3 (2003), 621–634.
- [Hu] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, New York (1972).
- [Ko] B. Kostant, *Lie algebra cohomology and generalized Borel-Weil theorem*, Ann. Math. **74** (1961), 329–387.
- [Ku] M. Kuranishi, *Lectures on involutive systems of partial differential equations*, Pub.Soc.Mat., São Paulo (1967).
- [SYY] T. Sasaki, K. Yamaguchi and M. Yoshida, *On the Rigidity of Differential Systems modeled on Hermitian Symmetric Spaces and Disproofs of a Conjecture concerning Modular Interpretations of Configuration Spaces*, Advanced Studies in Pure Math. **25** (1997), 318–354.
- [Se] Y. Se-ashi, *On differential invariants of integrable finite type linear differential equations*, Hokkaido Math.J. **17** (1988), 151–195.
- [St] S. Sternberg *Lectures on Differential Geometry*, Prentice-Hall, New Jersey (1964).
- [T1] N. Tanaka *On generalized graded Lie algebras and geometric structures I*, J. Math. Soc. Japan **19** (1967), 215–254.
- [T2] -----, *On differential systems, graded Lie algebras and pseudo-groups*, J. Math. Kyoto Univ.**10** (1970), 1–82.
- [T3] -----, *On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections*, Japan. J. Math.**2** (1976), 131–190.
- [T4] -----, *On the equivalence problems associated with simple graded Lie algebras*, Hokkaido Math. J. **8** (1979), 23–84.
- [T5] -----, *On geometry and integration of systems of second order ordinary differential equations*, Proc. Symposium on Differential Geometry, 1982, 194–205 (in Japanese)
- [Y1] K.Yamaguchi *Contact geometry of higher order*, Japanese J. of Math. **8** (1982), 109–176.
- [Y2] -----, *On involutive systems of second order of codimension 2*, Proc. of Japan Acad. **58**, Ser A, No.7 (1982), 302–305.
- [Y3] -----, *Geometrization of Jet bundles*, Hokkaido Math. J. **12** (1983), 27–40.
- [Y4] -----, *Typical classes in involutive systems of second order*, Japanese J. Math. **11** (1985), 265–291.
- [Y5] -----, *Differential systems associated with simple graded Lie algebras*, Adv. Studies in Pure Math. **22** (1993), 413–494.
- [Y6] -----, *G₂-Geometry of Overdetermined Systems of Second Order*, Trends in Math. (Analysis and Geometry in Several Complex Variables) (1999), Birkhäuser, Boston 289–314.
- [Y7] -----, *Geometry of Linear Differential Systems Towards Contact Geometry of Second Order*, IMA Volumes in Mathematics and its Applications **144** (Symmetries and Overdetermined Systems of Partial Differential Equations) (2007), 151–203.
- [YY1] K. Yamaguchi and T. Yatsui, *Geometry of Higher Order Differential Equations of Finite Type associated with Symmetric Spaces*, Advanced Studies in Pure Mathematics **37** (2002), 397–458.
- [YY2] K.Yamaguchi and T.Yatsui, *Parabolic Geometries associated with Differential Equations of Finite Type*, Progress in Mathematics **252** (From Geometry to Quantum Mechanics: In Honor of Hideki Omori) (2007), 161–209

K.Yamaguchi, DEPARTMENT OF MATHEMATICS,, FACULTY OF SCIENCE,, HOKKAIDO UNIVERSITY,, SAPPORO 060-0810,, JAPAN, E-MAIL yamaguch@math.sci.hokudai.ac.jp