



Title	The Euler characteristic of a generic wave front in a 3-manifold
Author(s)	Izumiya,S.; Marar,W.L.
Citation	Hokkaido University Preprint Series in Mathematics, 137, 2-4
Issue Date	1992-2
DOI	10.14943/84430
Doc URL	<a href="http://hdl.handle.net/2115/70243">http://hdl.handle.net/2115/70243</a>
Type	bulletin (article)
File Information	137.pdf



[Instructions for use](#)

**THE EULER CHARACTERISTIC OF  
A GENERIC WAVE FRONT IN  
A 3-MANIFOLD**

**S. Izumiya and W.L. Marar**

**Series #137. February 1992**

HOKKAIDO UNIVERSITY  
PREPRINT SERIES IN MATHEMATICS

- # 112: K. Matsuda, An analogy of the theorem of Hector and Duminy, 10 pages. 1991.
- # 113: S. Takahashi, On a regularity criterion up to the boundary for weak solutions of the Navier-Stokes equations, 23 pages. 1991.
- # 114: T. Nakazi, Sum of two inner functions and exposed points in  $H^1$ , 18 pages. 1991.
- # 115: A. Arai, De Rham operators, Laplacians, and Dirac operators on topological vector spaces, 27 pages. 1991.
- # 116: T. Nishimori, A note on the classification of non-singular flows with transverse similarity structures, 17 pages. 1991.
- # 117: T. Hibi, A lower bound theorem for Ehrhart polynomials of convex polytopes, 6 pages. 1991.
- # 118: R. Agemi, H. Takamura, The lifespan of classical solutions to nonlinear wave equations in two space dimensions, 30 pages. 1991.
- # 119: S. Altschuler, S. Angenent and Y. Giga, Generalized motion by mean curvature for surfaces of rotation, 15 pages. 1991.
- # 120: T. Nakazi, Invariant subspaces in the bidisc and commutators, 20 pages. 1991.
- # 121: A. Arai, Commutation properties of the partial isometries associated with anticommuting self-adjoint operators, 25 pages. 1991.
- # 122: Y.-G. Chen, Blow-up solutions to a finite difference analogue of  $u_t = \Delta u + u^{1+\alpha}$  in  $N$ -dimensional balls, 31 pages. 1991.
- # 123: A. Arai, Fock-space representations of the relativistic supersymmetry algebra in the two-dimensional spacetime, 13 pages. 1991.
- # 124: S. Izumiya, The theory of Legendrian unfoldings and first order differential equations, 16 pages. 1991.
- # 125: T. Hibi, Face number inequalities for matroid complexes and Cohen-Macaulay types of Stanley-Reisner rings of distributive lattices, 17 pages. 1991.
- # 126: S. Izumiya, Completely integrable holonomic systems of first order differential equations, 35 pages. 1991.
- # 127: G. Ishikawa, S. Izumiya and K. Watanabe, Vector fields near a generic submanifold, 9 pages. 1991.
- # 128: A. Arai, I. Mitoma, Comparison and nuclearity of spaces of differential forms on topological vector spaces, 27 pages. 1991.
- # 129: K. Kubota, Existence of a global solution to a semi-linear wave equation with initial data of non-compact support in low space dimensions, 53 pages. 1991.
- # 130: S. Altschuler, S. Angenent and Y. Giga, Mean curvature flow through singularities for surfaces of rotation, 62 pages. 1991.
- # 131: M. Giga, Y. Giga and H. Sohr,  $L^p$  estimates for the Stokes system, 13 pages. 1991.
- # 132: Y. Okabe, T. Ootsuka, Applications of the theory of  $KM_2O$ -Langevin equations to the non-linear prediction problem for the one-dimensional strictly stationary time series, 27 pages. 1992.
- # 133: Y. Okabe, Applications of the theory of  $KM_2O$ -Langevin equations to the linear prediction problem for the multi-dimensional weakly stationary time series, 22 pages. 1992.
- # 134: P. Aviles, Y. Giga and N. Komuro, Duality formulas and variational integrals, 22 pages. 1992.
- # 135: S. Izumiya, The Clairaut type equation, 6 pages. 1992.
- # 136: S. Izumiya, Singular solutions of first order differential equations, 6 pages. 1992.

# THE EULER CHARACTERISTIC OF A GENERIC WAVE FRONT IN A 3-MANIFOLD

S.IZUMIYA AND W.L.MARAR

**Abstract.** We give a relation between Euler characteristics of a generic closed Legendrian surface and its wavefront.

## INTRODUCTION

In this note we shall compute the Euler characteristic of a generic wave front in a 3-manifold.

Let  $N$  be a  $(2n + 1)$ -dimensional smooth manifold and  $K$  be a contact structure on  $N$  (i.e.  $K$  is a nondegenerate tangent hyperplane field on  $N$ ). An immersion  $i : L \rightarrow N$  is said to be *Legendrian* if  $\dim L = n$  and  $di_x(T_x L) \subset K_x$  for any  $x \in L$ . We say that a smooth fibre bundle  $\pi : E \rightarrow M$  is *Legendrian* if its total space  $E$  is furnished with a contact structure and its fibres are Legendrian submanifolds. For a Legendrian immersion  $i : L \rightarrow E$ ,  $\pi \circ i : L \rightarrow M$  is called a *Legendrian map* and the image of the Legendrian map  $\pi \circ i$  is called *the wave front* of  $i$ . It is denoted by  $W(i)$ .

From now on, we only consider the case of  $n = 2$ . Then it is known that a generic wave front has (semi cubic) cuspidal edges ( $A_2$ ), swallowtails ( $A_3$ ) and points of transversal self intersection ( $A_1A_1$ ,  $A_1A_2$ ,  $A_1A_1A_1$ ) as singularities ([1], see Fig. 1). We shall refer to the  $A_1A_1A_1$ -type point as a *triple point* of  $i$ .




Fig.1

If  $L$  is a closed surface, then the number of swallowtails and triple points are finite. Our main result is the following :

**THEOREM.** Let  $i : L \rightarrow E$  be a generic Legendrian immersion of a closed surface. Then we have

$$\chi(W(i)) = \chi(L) + T(i) + \frac{S(i)}{2},$$

---

1991 Mathematics Subject Classification. Primary 58C27.

Key words and phrases. Euler characteristics, wavefronts.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-T}\mathcal{E}\mathcal{X}$

where  $\chi(X)$  is the Euler characteristic of  $X$ ,  $T(i)$  is the number of triple points on  $W(i)$  and  $S(i)$  is the number of swallowtails.

We remark that the corresponding result for a generic wave front in a 2-manifold is easily verified and that

$$\chi(W(i)) = -d(i),$$

where  $d(i)$  denote the number of double points on  $W(i)$ .

On the other hand, we consider an equidistant surface of a closed surface in 3-dimensional Euclidian space. If the distance is sufficiently small, then the equidistant surface is diffeomorphic to the original surface. However, for some distances, singularities may be appeared in the equidistant surface and it may not be homeomorphic to the original surface. Hence, it is interesting to study the relation between topologies of the equidistant surface and of the original surface. These subjects are studied in the theory of Legendrian singularities. In fact, it is known that singularities of equidistant surfaces are locally diffeomorphic to singularities of wave fronts (see [1]). Then we show the general property of the Euler characteristic of global wave fronts.

In order to prove the theorem, we shall apply the method which has been introduced to compute the Euler characteristic of the image of a stable perturbation of an  $\mathcal{A}$ -finite map germ in [2].

All maps considered here are class  $C^\infty$  unless stated otherwise.

### 1. PROOF OF THE THEOREM

In this section we shall give a proof of the theorem. Let  $i : L \rightarrow E$  be a generic Legendrian immersion of a closed surface. Since the Euler characteristic is a topological invariant, then we can ignore cuspidal edges. We now define the following sets:

$$\begin{aligned} D^2(i) &= cl\{x \in S \mid \#(\pi \circ i)^{-1} \pi \circ i(x) \geq 2\}, \\ D^3(i) &= \{x \in D^2(i) \mid \#(\pi \circ i)^{-1} \pi \circ i(x) = 3\}, \\ D^2(i, (2)) &= \{x \in D^2(i) \mid \#(\pi \circ i)^{-1} \pi \circ i(x) = 1\}, \end{aligned}$$

where  $clX$  is the topological closure of  $X$ . Then we have the following diagram:

$$\begin{array}{ccc} & D^3(i) & \\ & \downarrow h & \\ D^2(i, (2)) & \xrightarrow{j} & D^2(i) \\ & \downarrow k & \\ L & \xrightarrow{\pi \circ i} & W(i) \subset M, \end{array}$$

where  $h, j, k$  are inclusions.

By the characterization of generic wave fronts (see [ ]),  $D^2(i)$  is a union of curves on  $L$  with self-intersection and circles,  $D^3(i)$  is the inverse image of triple points and  $D^2(i, (2))$  is the set of swallowtails of  $\pi \circ i$ . It follows that these are immersed submanifolds of  $L$  with  $\dim D^2(i) = 1$  and  $\dim D^3(i) = \dim D^2(i, (2)) = 0$ .

In order to prove the theorem, we need the following formula.

LEMMA 1.1.  $\chi(W(i)) = \chi(L) - \frac{1}{2}\chi(D^2(i)) + \frac{1}{2}\chi(D^2(i, (2))) - \frac{1}{6}\chi(D^3(i))$ .

PROOF: Consider the equation

$$(*) \quad \chi(W(i)) = \alpha \chi(L) + \beta \chi(D^2(i)) + \gamma \chi(D^2(i, (2))) + \delta \chi(D^3(i)),$$

where  $\alpha, \beta, \gamma$  and  $\delta$  are unknown variables. We solve this by a purely combinatorial method.

We now construct a triangulation  $K_i$  of the stratified set  $W(i)$  as follows: We start to triangulate  $W(i)$  by including the image of  $D^2(i, (2))$  and the image of  $D^3(i)$  among the vertices of  $K_i$ . After this, we build up the one-skeleton  $K_i^{(1)}$  of  $K_i$  so that the image of  $D^2(i)$  is a subcomplex of  $K_i^{(1)}$ . We complete our procedure by constructing the two-skeleton  $K_i^{(2)}$ .

Since  $\pi \circ i$  and its restrictions to  $D^2(i)$ ,  $D^2(i, (2))$  and  $D^3(i)$  are proper and finite-to-one mappings, then we can pull back  $K_i$  to obtain a triangulation for  $L$ ,  $D^2(i)$ ,  $D^2(i, (2))$  and  $D^3(i)$ . Let  $C_j^X$  be the number of  $j$ -cells in  $X$ , where  $X = W(i)$ ,  $L$ ,  $D^2(i)$ ,  $D^2(i, (2))$  or  $D^3(i)$ . Then the equation (\*) can be written by

$$\begin{aligned} \sum_j (-1)^j C_j^{W(i)} &= \alpha \sum_j (-1)^j C_j^L + \beta \sum_j (-1)^j C_j^{D^2(i)} \\ &+ \gamma \sum_j (-1)^j C_j^{D^2(i, (2))} + \delta \sum_j (-1)^j C_j^{D^3(i)}, \end{aligned}$$

where  $C_j^X = 0$  if  $i > \dim X$ . So, if we can find real numbers  $\alpha, \beta, \gamma$  and  $\delta$  such that

$$(**) \quad C_j^{W(i)} = \alpha C_j^L + \beta C_j^{D^2(i)} + \gamma C_j^{D^2(i, (2))} + \delta C_j^{D^3(i)},$$

for any  $j$ , then we have solutions of the equation (\*). By the construction of the triangulation, we may concentrate on solving (\*\*) in the case when  $j = 0$ . We remark that  $\pi \circ i$  is 3 to 1 over the points in the image of  $D^3(i)$ , 1 to 1 over the points in the image of  $D^2(i, (2))$ , 2 to 1 over the points in the image of  $D^2(i) - (D^2(i, (2)) \cup D^3(i))$ , and 1 to 1 over the points in the image of  $S - D^2(i)$ . It follows that the equation

$$C_0^{W(i)} = \alpha C_0^L + \beta C_0^{D^2(i)} + \gamma C_0^{D^2(i, (2))} + \delta C_0^{D^3(i)}$$

is equivalent to the system of linear equations :

$$\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 3 & 3 & 0 & 3 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \\ \delta \end{pmatrix}.$$

We can easily solve this equation, so that  $\alpha = 1$ ,  $\beta = -1/2$ ,  $\gamma = 1/2$  and  $\delta = -1/6$ . This completes the proof.

Then we can prove the theorem.

PROOF OF THE THEOREM: By the definition we have  $\chi(D^2(i, (2))) = S(i)$  and  $\chi(D^3(i)) = 3T(i)$ . Since  $D^2(i)$  is a union of closed curves on the surface  $L$  with  $3T(i)$  crossings, then we can triangulate it with  $3T(i) + n$  0-cells and  $6T(i) + n$  1-cells, where  $n$  is the number of circles in  $D^2(i)$ . It follows that  $\chi(D^2(i)) = -3T(i)$ . If we substitute these on the formula in Lemma 1.1, then we have

$$\chi(W(i)) = \chi(L) + T(i) + \frac{1}{2}S(i).$$

This completes the proof of the theorem.

ACKNOWLEDGEMENT. This work has been done during the authors' stay at the University of Liverpool. The authors would like to thank the department of Pure Mathematics. The first author acknowledges the financial support of JSPS and the second author the financial support of CAPES.

#### REFERENCES

1. V. I. Arnol'd, "Singularities of Caustics and Wave fronts, Mathematics and Its Applications (Soviet Series)," Kluwer Academic Publisher, 1990.
2. W. L. Marar, *The Euler characteristic of the disentanglement of the image of a corank 1 map germ*, in "Springer Lecture Notes in Mathematics 1462," Springer, 1991, p. 212-220.

(S. Izumiya) Department of Mathematics, Faculty of Science, Hokkaido Univeristy, Sapporo 060, Japan  
(W. L. Marar) Instituto de Ciências Matemáticas de São Carlos, Universidade de São Paulo, Caixa Postal 668, 13560-São Carlos (SP)-Brazil