



Title	Laparoscopic proximal gastrectomy with oblique jejunogastrostomy
Author(s)	Tanaka, Kimitaka; Ebihara, Yuma; Kurashima, Yo; Nakanishi, Yoshitsugu; Asano, Toshimichi; Noji, Takehiro; Murakami, Soichi; Nakamura, Toru; Tsuchikawa, Takahiro; Okamura, Keisuke; Shichinohe, Toshiaki; Hirano, Satoshi
Citation	Langenbeck's archives of surgery, 402(6), 995-1002 https://doi.org/10.1007/s00423-017-1587-4
Issue Date	2017-09
Doc URL	http://hdl.handle.net/2115/71398
Rights	The final publication is available at link.springer.com
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	LangenbecksArchSurg402_995.pdf



[Instructions for use](#)

Title: Laparoscopic Proximal Gastrectomy with Oblique Jejunogastrostomy

Type of Submission: How-I-do-it article

Names of the Authors:

Kimitaka Tanaka, MD, PhD¹; Yuma Ebihara, MD, PhD¹; Yo Kurashima, MD, PhD¹; Yoshitsugu Nakanishi, MD, PhD¹; Toshimichi Asano, MD, PhD¹; Takehiro Noji, MD, PhD¹; Soichi Murakami, MD, PhD¹; Toru Nakamura, MD, PhD¹; Takahiro Tsuchikawa, MD, PhD¹; Keisuke Okamura, MD, PhD¹; Toshiaki Shichinohe, MD, PhD¹; and Satoshi Hirano, MD, PhD¹

Affiliation:

Department of Gastroenterological Surgery II, Hokkaido University Graduate School of Medicine
North 15, West 7, Kita-ku, Sapporo 060-8638, Japan
Tel: +81-11-706-7714, Fax: +81-11-706-7158

Corresponding Author:

Yuma Ebihara, MD, PhD
Department of Gastroenterological Surgery II, Hokkaido University Graduate School of Medicine
North 15, West 7, Kita-ku, Sapporo 060-8638, Japan
Tel: +81-11-706-7714, Fax: +81-11-706-7158
E-mail: yuma-ebi@wc4.so-net.ne.jp

Funding: No financial support was provided for this study.

Conflict of Interest: The authors declare that they have no conflict of interest.

Abstract

Background

Proximal early gastric cancer is a good indication for totally laparoscopic proximal gastrectomy (TLPG) with double-tract reconstruction (DTR). However, when most of the dietary intake passes through the escape route of the jejunum, the functional benefits of proximal gastrectomy might be similar to those after total gastrectomy. Our DTR procedure was improved for easy passage through the remnant stomach. The purpose of this study was to present a novel technique for intracorporeal DTR using linear staplers after TLPG and to investigate surgical outcomes.

Methods

DTR was performed using linear staplers only. A side-to-side jejunogastrostomy with twisting of both the remnant stomach and the anal jejunum was performed for the purpose of passing meals through the remnant stomach (an oblique jejunogastrostomy technique). The 10 patients who underwent TLPG with DTR from January 2011 to August 2016 in Hokkaido University Hospital were retrospectively reviewed. Their clinicopathological characteristics and surgical and postoperative outcomes were collected and analyzed.

Results

The median duration of operation was 285 (range 146 - 440) min. No patients required blood transfusions. The number of dissected lymph nodes was 32 (range 22 - 56). There were no intraoperative complications, and no cases were converted to open surgery. All patients were pT1N0M0 stage IA. No anastomotic leakage or complications were detected. Postoperative gastrography after reconstruction showed that contrast medium flowed mainly to the remnant stomach. The average percentage body weight loss was 14.0% ± 7.1% at 10 months. The average percentage decrease in serum hemoglobin was 5.4% ± 10.4% at 12 months.

Conclusions

This novel technique for intracorporeal DTR provided a considerable advantage by the passage of dietary intake to the remnant stomach after LPG.

Keywords: proximal gastrectomy, double-tract reconstruction, gastric cancer, laparoscopic surgery

Author contributions:

Study conception and design: Kimitaka Tanaka, Yuma Ebihara, Yo Kurashima, and Satoshi Hirano

Acquisition of data: Kimitaka Tanaka, Yuma Ebihara, Yo Kurashima

Drafting of manuscript: Kimitaka, Tanaka, Yuma Ebihara, and Satoshi Hirano

Critical revision of manuscript: Yuma Ebihara, Yo Kurashima, Yoshitsugu Nakanishi, Toshimichi Asano, Takehiro Noji, Soichi Murakami, Toru Nakamura, Takahiro Tsuchikawa, Keisuke Okamura, Toshiaki Shichinohe, and Satoshi Hirano

Introduction

Gastric cancer is the most common form of cancer in East Asian countries [1]. The reported incidence of early gastric cancer (EGC) has increased as a result of improved surveillance by the national cancer screening program in Japan [2]. It has also been found that *H. pylori* infection is associated with gastric cancer. *H. pylori* eradication therapy by antibiotic treatment decreases the incidence of distal gastric cancer [3]. On the other hand, the incidence of proximal gastric cancer in Korea has gradually increased from 5.3% to 14.0% [4]. In recent decades, the oncological safety of minimally invasive surgery for the treatment of EGC has been established [5]. Laparoscopic gastrectomy with regional lymph node dissection has been used in the treatment of EGC with low mortality and morbidity and improved patient quality of life [6]. We have performed totally laparoscopic gastrectomy for EGC, including totally laparoscopic distal gastrectomy (TLDG), totally laparoscopic proximal gastrectomy (TLPG), and totally laparoscopic total gastrectomy (TLTG) with intracorporeal anastomosis, using a laparoscopic linear stapler [7].

Proximal gastrectomy for EGC was a significant improvement over total gastrectomy in terms of maintaining physiological function and quality of life, such as weight loss, the necessity for additional meals, dumping [8], and maintaining high hemoglobin levels [9,10]. Proximal EGC is a good indication for TLPG when functional preservation or minimal invasiveness is taken into consideration. However, total gastrectomy has still been widely performed as a standard treatment for early upper-third gastric cancer to achieve a tumor-free resection margin and extended lymph node dissection [11]. The oncological safety and functional benefits of proximal gastrectomy have been reported in several studies. They concluded that the long-term overall survival of patients who underwent proximal gastrectomy for proximal EGC was similar to that of those who underwent total gastrectomy [12-14]. Among the various reconstructions of laparoscopic proximal gastrectomy, such as esophagogastrostomy [15], jejunal interposition [16], and double-tract reconstruction (DTR) [17], we have performed TLPG with DTR for EGC. Aikou et al.[18] first reported DTR after proximal gastrectomy and the original gastrojejunostomy method called the N-shaped double-tract method. One of the reasons why we chose DTR was its technical similarity to the DTR with Roux-en-Y reconstruction for laparoscopic total gastrectomy, which has been established as a standard laparoscopic reconstruction procedure. However, there is some concern that, with DTR, most of the dietary intake might escape into the jejunum. To prevent this disadvantageous phenomenon, we have developed a newly-devised DTR that allows dietary intake to pass easily through the remnant stomach.

The purpose of this study was to present a novel technique for intracorporeal DTR using linear staplers after proximal gastrectomy and to investigate the short- and mid-term outcomes of LPG-DTR.

Patients and methods

Patient selection

Between January 2011 and August 2016, 10 patients diagnosed with EGC preoperatively underwent LPG with DTR at Hokkaido University Hospital, Sapporo, Japan. Preoperative assessments were carried out by endoscopy, computed tomography (CT), and endoscopic ultrasound. The eligibility criteria of the patients were: (1) gastric cancer invaded within the submucosal layer (cT1); (2) suspected to have no lymph node metastases (cN0); and (3) tumor located more than a 5-cm distance from the angular region and the remnant stomach would be more than half the size of the preoperative stomach. Patients who had a previous history of upper gastrointestinal surgery were excluded from DTR. Informed consent was obtained from all patients. Patients' characteristics including age, sex, body mass index (BMI), ASA physical status classification system (ASA), history of abdominal surgery and endoscopic submucosal dissection (ESD), and chronic disease status were recorded. Operation time, blood loss, time to resume a soft diet, postoperative hospital stay, postoperative complications, and pathological findings were also analyzed. Postoperative morbidity was evaluated using the Clavien-Dindo classification [19]. Among early complications, the main complications were investigated, including intraperitoneal or digestive tract hemorrhage, anastomotic leakage, bowel or anastomosis obstruction, and abdominal infection. Specimens were evaluated according to the Japanese Classification of Gastric Carcinoma established by the Japanese Research Society for Gastric Cancer [20]. Follow-up was conducted to evaluate late complications, including reflux esophagitis and anastomotic stenosis, the changes of body weight and hemoglobin concentration, and the rates of recurrence and survival.

Procedures for gastrectomy

Patients are placed in the supine reverse Trendelenburg position with the legs apart under general anesthesia. The operator is positioned between the legs of the patient, the assistant is on the left side, and the scopist is on the opposite side. Five trocars (Xcel[®] Ethicon Endo-Surgery, Cincinnati, OH, USA) are used, and a 12-mm paraumbilical port is subsequently extended to 3.0 cm when extracting specimens. After carbon dioxide pneumoperitoneum is established at a pressure of 10 mmHg, a laparoscope (3CCD Video System SX-2, Olympus, Tokyo, Japan) is introduced through this port, and four other trocars (three 12-mm trocars and one 5-mm trocar) are placed (Fig. 1). We dissect lymph nodes and coagulate vessels using laparoscopic coagulation shears (SONOSURG-X[®], Olympus Medical Systems or Harmonic Ace[®], Ethicon Endo-Surgery) and a vessel sealing system (Ligasure Maryland Jaw[™], Medtronic, Mansfield, MA, USA). The basic extent of lymph node dissection in the present series was D1 + dissection of lymph node Nos. 1, 2, 3a, 4sa, 4sb, 7, 8a, 9, and 11p; lymph node regions and dissection were decided according to the Japanese Classification of Gastric Carcinoma published by the Japanese Gastric Cancer Association [21]. The lymph nodes of Nos. 4d, 5, and 6 with the right gastroepiploic vessels and right gastric vessels are preserved. The bottom of the esophagus and the distal 1/3 to 1/2 stomach are transected by 60-mm endoscopic linear staplers (Powered Echelon 60[®], Ethicon Endo-Surgery). The proximal and distal free margins are more than 2 cm.

Procedures for double-tract reconstruction (DTR)

DTR after TLPG is performed using only linear staplers. The jejunum is transected by a 60-mm endoscopic linear stapler at a point 20-cm distant from the ligament of Treitz. For esophagojejunostomy, we use functional end-to-end technique [22,7]. Thereafter, 10-mm transverse incisions are created at the antimesenteric wall of the jejunum 20 cm below the esophagojejunostomy (Fig. 2A), and 10-mm incisions are also created at the greater curvature side of the remnant stomach (Fig. 2B). Jaws of a 60-mm linear stapler (Powered Echelon 60[®], Ethicon Endo-Surgery) are inserted into the holes. Then, both the stomach and jejunum are twisted posteriorly, and the posterior wall of the remnant stomach and the posterior wall of the jejunum are put together. An oblique side-to-side jejunogastrostomy from the antimesenteric wall to the posterior wall is performed (Fig. 2C, D). The entry hole for this stapler is closed with a running suture (3-0 Vicryl, Ethicon). In this way, the jejunum returns the torsion of the jejunum to the counter-clockwise direction and rides on the remnant stomach (Fig. 2E). Finally, a side-to-side jejunojejunostomy is made between the jejunum 20 cm below the jejunogastrostomy and the proximal jejunum. The jejunojejunostomy is pulled toward the cranial side without sacrificing the small intestine (Fig. 3). Thereby, the distal jejunum from the jejunogastrostomy can easily bend to the ventral side, and the exit route becomes narrow (Online Resource 1). We called this jejunogastrostomy with twisting of both the remnant stomach and the anal jejunum and the jejunojejunostomy pulled toward the cranial side an oblique jejunogastrostomy technique.

Ethics and Consent

The Hokkaido University Hospital Ethics Committee approved this study in 2016 (No. 016-0194). The consent form stated the aim of the study on the web site of our hospital and the participants' right to decline to participate or opt out at any time. Informed consent was obtained from all individual participants included in the study. The ethics committee/IRB approved this consent procedure.

Results

Clinical characteristics of patients

Ten patients (7 males, 3 females; median age 70 (range 55 - 77) y; median BMI 23.7 (range 18.3 - 28.7) kg/m²) were included. Six patients had previously undergone endoscopic submucosal dissection (ESD). Two patients had undergone previous abdominal operations, including 1 appendectomy and 1 colorectal cancer operation. The patients' clinical characteristics are presented in Table 1.

Operative and pathological data

The operative and pathological data are summarized in Table 2. The median duration of operation was 285 (range 146 - 440) min. No patients required blood transfusions. The number of dissected lymph nodes was 32 (range 22-56). There were no intraoperative complications, and no cases were converted to open surgery. Cancer had invaded

the submucosa in all 10 patients. All patients were pT1N0M0 stage IA. Histological types included 5 well-differentiated cases, 3 moderately differentiated cases, and 2 poorly differentiated cases. The free margins of the tumor were no less than 2 cm from the distal end (2.1-8.5 cm).

Postoperative outcomes

The operative and pathological data are summarized in Table 3. The median time to resume a soft diet postoperatively was 3 (range 3-5) days, and the median postoperative hospital stay was 13 (range 9 - 16) days. No patients required second surgery. No anastomotic leakage and no complications were seen. Recurrence was not observed in any patient after LPG over a median follow-up period of 10.7 (range 1 - 37.6) months. Some patients underwent endoscopy follow-up and upper gastrointestinal series examinations. They showed the wide entry hole of the gastrojejunostomy and the narrow route of the distal jejunum. No patients complained of dumping syndrome, while two patients complained of moderate reflux symptoms of heartburn, but they soon recovered with proton-pump inhibitor treatment. The average percentage body weight loss was 14.0% (SD \pm 7.1%). The average percentage decrease in serum hemoglobin was 5.4% (SD \pm 10.4%). The median follow-up periods for these two were 10 months and 12 months, respectively.

Discussion

Three representative reconstruction procedures after proximal gastrectomy have been reported: esophagogastrostomy, jejunal pouch interposition reconstruction, and DTR. DTR was thought to be the best reconstruction procedure with respect to anastomosis-related late complications, especially postoperative reflux esophagitis. Esophagogastrostomy is simpler than the other procedures because it includes only one anastomosis, and it is the most popular and the classical reconstruction [23]. However, a relatively high incidence (27.4% to 67.4%) of postoperative anastomosis-related complications such as reflux esophagitis has been reported [24,25,13,14]. Jejunal (pouch) interposition reconstruction is the second most common reconstruction [23]. Some papers [26-28] reported that proximal gastrectomy with jejunal interposition had a high incidence (10.2%, 9.1%, 31.8%) of anastomotic stricture. Kinoshita et al [27] speculated about the reason for the stricture. They suggested that the small amount of bile reflux to the interposed jejunum and tension to the interposed jejunum cause stenosis. In four studies [17,29-31] of laparoscopic proximal gastrectomy with DTR, three reported postoperative complications. In the literature, there were no anastomosis-related complications. The morbidity rates were 9.5%, 11.6%, and 25%. Symptoms related to reflux esophagitis occurred in 0%, 4.7%, and 4.8%. The present results for postoperative complications were similar to these reports. As to long-term function, changes of serum hemoglobin and body weight after gastrectomy were reported. The present results for body weight loss and decrease in serum hemoglobin were not worse than the body weight loss (14.1 - 30.0%) [32-36] and decrease in serum hemoglobin (9.7%) [33] at 12 months after total gastrectomy. Ahn et al. [17] first reported 43 cases of DTR after LPG, and they

performed a Roux-en-Y esophagojejunostomy by intracorporeal anastomosis with a circular stapler by a mini-laparotomy, and side-to-side gastrojejunostomy was performed in an extracorporeal fashion using 2 linear staplers. Nomura et al. [29] used circular staplers to complete the esophagojejunostomies. In the remaining two reports, linear staplers were used to complete all anastomoses, which is similar to the present procedure. Hong et al. [30] first reported TLPG with DTR with a triangular stapling technique (delta-shaped anastomosis). Linear staplers can make a larger size anastomosis than a circular stapler [37], and thus the occurrence of anastomotic stenosis can be reduced. With extracorporeal anastomosis, it was sometimes difficult to move the intestinal tracts just under the mini-laparotomy, especially in obese patients. We selected DTR to be able to perform intracorporeal totally laparoscopic surgery and to have less anastomosis-related problems, such as reflux esophagitis and anastomotic stenosis.

DTR was first reported by Aikou et al. [18] in terms of gaining the smooth transfer of larger foods through the duodenal route. There are some concerns with DTR that the functional benefits of proximal gastrectomy, such as preserving the antrum, might be lost when most of the dietary intake passes through the escape route of the jejunum. In order to overcome this disadvantage, we modified the intracorporeal anastomosis of jejunogastrostomy in the traditional DTR by narrowing the exit route of the jejunum to induce meals to pass into the remnant stomach. Aikou et al. [18] also reported the original jejunogastrostomy method after proximal gastrectomy called the N-shaped method, due to the flow of larger quantities of food into the antrum of the residual stomach. The residual stomach was twisted 180 degrees anteriorly beforehand, the twisted residual stomach was restored to its usual position, and the jejunum was then shaped like the letter "N" around the gastro-jejunostomy. Seeing this anastomotic technique, we modified it to be suitable for intracorporeal laparoscopic reconstruction. The anastomotic method of jejunogastrostomy with twisting of only the stomach created an anastomotic site in which anastomosis ischemia was prevented. On the other hand, with twisting of both the stomach and the jejunum, the dietary intake could easily flow into the remnant stomach because the anastomotic site of the jejunogastrostomy was made at the posterior wall by the jejunum riding on the stomach and faced slightly more vertically against the flow of the meal, returning the torsion of the jejunum to the counter-clockwise direction. There appear to be two reasons for this: 1) the twisted jejunum returns to the original position because the rim of the jejunum was fixed by the mesentery and the remnant stomach was relatively free to move with the lymph dissection; and 2) without sacrificing several cm of the small intestine, the distal jejunum from the jejunogastrostomy could be bent forward and lifted up to the cranial side. The jejunojejunostomy was placed a short distance from the caudal side of the esophagojejunostomy. Thereby, the jejunogastrostomy and jejunojejunostomy were placed at almost the same level as the remnant stomach. Our innovation was to create the anastomosis by twisting a small amount of the small intestine, bending the small intestine of the outflow path into a U-shaped appearance without sacrificing the small intestine. This enables food to enter the residual stomach more easily, because the bent distal jejunum reduces the

outflow of the contents. Hong et al. [30] reported a delta-shaped intracorporeal jejunogastrostomy in DTR in a procedure similar to ours. However, it was difficult to narrow the exit routes of the jejunum by twisting only the remnant stomach.

The second weak point is that the DTR needs more anastomotic sites than other reconstructive procedures for the proximal gastrectomy. However, every anastomosis for DTR used the same anastomotic procedure, which had been established as the intracorporeal reconstruction of laparoscopic gastrectomy. The esophagojejunostomy and jejunojejunostomy for DTR could be performed in the same procedure as Roux-en-Y reconstruction after laparoscopic total gastrectomy [7]. The jejunogastrostomy for DTR could be performed in the same fashion as the delta-anastomosis after laparoscopic distal gastrectomy [38]. In contrast, intracorporeal esophagogastronomy or jejunal interposed reconstruction are not well known even by experienced laparoscopic surgeons, because these procedures are not necessary for either total or distal gastrectomy. The total operation time was similar between TLPG with DTR and that with Roux-en-Y reconstruction in our department, although DTR requires one more reconstruction than Roux-en-Y reconstruction. The reconstruction time of DTR (median time: 108 minutes) was approximately 20 minutes longer than that of Roux-en-Y reconstruction (median time: 87.5 minutes) in our department when comparing the same periods (data not shown). As mentioned above, we consider that DTR is a better reconstructive procedure for TLPG, because the procedure can be more easily introduced.

The present study had several limitations. Subjective symptoms were not evaluated using a questionnaire, and long-term functional benefits were not assessed. Gastric emptying was not examined in all patients. Despite this being a retrospective study of a small number of cases, our technique appears to provide feasible results.

Conclusion

Our novel oblique jejunogastrostomy technique for DTR appeared to have the considerable advantage of passage of dietary intake to the remnant stomach, as well as reasonable complications rates without any anastomosis-related problems. Further studies to evaluate subjective function and symptoms are needed.

Authors' contribution

Study conception and design: Kimitaka Tanaka, Yuma Ebihara, Yo Kurashima, and Satoshi Hirano

Acquisition of data: Kimitaka Tanaka, Yuma Ebihara, and Yo Kurashima

Drafting of manuscript: Kimitaka, Tanaka, Yuma Ebihara, and Satoshi Hirano

Critical revision of manuscript: Yuma Ebihara, Yo Kurashima, Yoshitsugu Nakanishi, Toshimichi Asano, Takehiro

Noji, Soichi Murakami, Toru Nakamura, Takahiro Tsuchikawa, Keisuke Okamura, Toshiaki Shichinohe, and Satoshi Hirano

Compliance with ethical standards

Funding: This study received no funding.

Conflict of Interest: The authors declare that they have no conflicts of interest

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Reference

1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A (2015) Global cancer statistics, 2012. *CA Cancer J Clin* 65 (2):87-108. doi:10.3322/caac.21262
2. Sano T, Hollowood A (2006) Early gastric cancer: diagnosis and less invasive treatments. *Scand J Surg* 95 (4):249-255
3. Asaka M, Mabe K, Matsushima R, Tsuda M (2015) Helicobacter pylori Eradication to Eliminate Gastric Cancer: The Japanese Strategy. *Gastroenterology clinics of North America* 44 (3):639-648. doi:10.1016/j.gtc.2015.05.010
4. Jeong O, Park YK (2011) Clinicopathological features and surgical treatment of gastric cancer in South Korea: the results of 2009 nationwide survey on surgically treated gastric cancer patients. *J Gastric Cancer* 11 (2):69-77. doi:10.5230/jgc.2011.11.2.69
5. Kim YI, Kim SY, Cho SJ, Park JH, Choi IJ, Lee YJ, Lee EK, Kook MC, Kim CG, Ryu KW, Kim YW (2014) Long-term metformin use reduces gastric cancer risk in type 2 diabetics without insulin treatment: a nationwide cohort study. *Alimentary pharmacology & therapeutics* 39 (8):854-863. doi:10.1111/apt.12660
6. Kawamura H, Okada K, Isizu H, Masuko H, Yamagami H, Honma S, Ueki S, Noguchi K, Kondo Y (2008) Laparoscopic gastrectomy for early gastric cancer targeting as a less invasive procedure. *Surg Endosc* 22 (1):81-85. doi:10.1007/s00464-007-9373-y
7. Ebihara Y, Okushiba S, Kawarada Y, Kitashiro S, Katoh H (2013) Outcome of functional end-to-end esophagojejunostomy in totally laparoscopic total gastrectomy. *Langenbeck's archives of surgery / Deutsche Gesellschaft fur Chirurgie* 398 (3):475-479. doi:10.1007/s00423-013-1051-z
8. Takiguchi N, Takahashi M, Ikeda M, Inagawa S, Ueda S, Nobuoka T, Ota M, Iwasaki Y, Uchida N, Koderu Y, Nakada K (2015) Long-term quality-of-life comparison of total gastrectomy and proximal gastrectomy by postgastrectomy syndrome assessment scale (PGSAS-45): a nationwide multi-institutional study. *Gastric Cancer* 18 (2):407-416. doi:10.1007/s10120-014-0377-8
9. Nozaki I, Hato S, Kobatake T, Ohta K, Kubo Y, Kurita A (2013) Long-term outcome after proximal gastrectomy with jejunal interposition for gastric cancer compared with total gastrectomy. *World J Surg* 37 (3):558-564. doi:10.1007/s00268-012-1894-4
10. Ichikawa D, Komatsu S, Kubota T, Okamoto K, Shiozaki A, Fujiwara H, Otsuji E (2014) Long-term outcomes of patients who underwent limited proximal gastrectomy. *Gastric Cancer* 17 (1):141-145. doi:10.1007/s10120-013-0257-7
11. Papachristou DN, Fortner JG (1980) Adenocarcinoma of the gastric cardia. The choice of gastrectomy. *Ann Surg* 192 (1):58-64
12. Yoo CH, Sohn BH, Han WK, Pae WK (2004) Long-term results of proximal and total gastrectomy for adenocarcinoma of the upper third of the stomach. *Cancer Res Treat* 36 (1):50-55. doi:10.4143/crt.2004.36.1.50
13. Wen L, Chen XZ, Wu B, Chen XL, Wang L, Yang K, Zhang B, Chen ZX, Chen JP, Zhou ZG, Li CM, Hu JK (2012) Total vs. proximal gastrectomy for proximal gastric cancer: a systematic review and meta-analysis. *Hepato-gastroenterology* 59 (114):633-640. doi:10.5754/hge11834
14. Pu YW, Gong W, Wu YY, Chen Q, He TF, Xing CG (2013) Proximal gastrectomy versus total gastrectomy for proximal gastric carcinoma. A meta-analysis on postoperative complications, 5-year survival, and recurrence rate. *Saudi Med J* 34 (12):1223-1228
15. Uyama I, Ogiwara H, Takahara T, Kikuchi K, Iida S (1995) Laparoscopic and minilaparotomy proximal gastrectomy and esophagogastrostomy: technique and case report. *Surgical laparoscopy & endoscopy* 5 (6):487-491
16. Uyama I, Sugioka A, Fujita J, Komori Y, Matsui H, Hasumi A (2000) Completely laparoscopic proximal gastrectomy with jejunal interposition and lymphadenectomy. *J Am Coll Surg* 191 (1):114-119
17. Ahn SH, Jung do H, Son SY, Lee CM, Park do J, Kim HH (2014) Laparoscopic double-tract proximal gastrectomy for proximal early gastric cancer. *Gastric Cancer* 17 (3):562-570. doi:10.1007/s10120-013-0303-5
18. Aikou T, Natsugoe S, Shimazu H, Nishi M (1988) Antrum preserving double tract method for reconstruction following proximal gastrectomy. *Jpn J Surg* 18 (1):114-115
19. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240 (2):205-213
20. Nakajima T (2002) Gastric cancer treatment guidelines in Japan. *Gastric Cancer* 5 (1):1-5. doi:10.1007/s101200200000

21. Japanese Gastric Cancer A (1998) Japanese Classification of Gastric Carcinoma - 2nd English Edition. *Gastric Cancer* 1 (1):10-24. doi:10.1007/s101209800016
22. Matsui H, Uyama I, Sugioka A, Fujita J, Komori Y, Ochiai M, Hasumi A (2002) Linear stapling forms improved anastomoses during esophagojejunostomy after a total gastrectomy. *American journal of surgery* 184 (1):58-60
23. Kumagai K, Shimizu K, Yokoyama N, Aida S, Arima S, Aikou T, Japanese Society for the Study of Postoperative Morbidity after G (2012) Questionnaire survey regarding the current status and controversial issues concerning reconstruction after gastrectomy in Japan. *Surg Today* 42 (5):411-418. doi:10.1007/s00595-012-0159-z
24. Katsoulis IE, Robotis JF, Kouraklis G, Yannopoulos PA (2006) What is the difference between proximal and total gastrectomy regarding postoperative bile reflux into the oesophagus? *Digestive surgery* 23 (5-6):325-330. doi:10.1159/000097948
25. An JY, Youn HG, Choi MG, Noh JH, Sohn TS, Kim S (2008) The difficult choice between total and proximal gastrectomy in proximal early gastric cancer. *American journal of surgery* 196 (4):587-591. doi:10.1016/j.amjsurg.2007.09.040
26. Katai H, Morita S, Saka M, Taniguchi H, Fukagawa T (2010) Long-term outcome after proximal gastrectomy with jejunal interposition for suspected early cancer in the upper third of the stomach. *Br J Surg* 97 (4):558-562. doi:10.1002/bjs.6944
27. Kinoshita T, Gotohda N, Kato Y, Takahashi S, Konishi M, Kinoshita T (2013) Laparoscopic proximal gastrectomy with jejunal interposition for gastric cancer in the proximal third of the stomach: a retrospective comparison with open surgery. *Surg Endosc* 27 (1):146-153. doi:10.1007/s00464-012-2401-6
28. Nakamura M, Nakamori M, Ojima T, Katsuda M, Iida T, Hayata K, Matsumura S, Kato T, Kitadani J, Iwahashi M, Yamaue H (2014) Reconstruction after proximal gastrectomy for early gastric cancer in the upper third of the stomach: an analysis of our 13-year experience. *Surgery* 156 (1):57-63. doi:10.1016/j.surg.2014.02.015
29. Nomura E, Lee SW, Kawai M, Yamazaki M, Nabeshima K, Nakamura K, Uchiyama K (2014) Functional outcomes by reconstruction technique following laparoscopic proximal gastrectomy for gastric cancer: double tract versus jejunal interposition. *World J Surg Oncol* 12:20. doi:10.1186/1477-7819-12-20
30. Hong J, Qian L, Wang YP, Wang J, Hua LC, Hao HK (2015) A novel method of delta-shaped intracorporeal double-tract reconstruction in totally laparoscopic proximal gastrectomy. *Surg Endosc*. doi:10.1007/s00464-015-4490-5
31. Yang K, Bang HJ, Almadani ME, Dy-Abalajon DM, Kim YN, Roh KH, Lim SH, Son T, Kim HI, Noh SH, Hyung WJ (2016) Laparoscopic Proximal Gastrectomy with Double-Tract Reconstruction by Intracorporeal Anastomosis with Linear Staplers. *J Am Coll Surg*. doi:10.1016/j.jamcollsurg.2016.01.002
32. Davis JL, Selby LV, Chou JF, Schattner M, Ison DH, Capanu M, Brennan MF, Coit DG, Strong VE (2016) Patterns and Predictors of Weight Loss After Gastrectomy for Cancer. *Ann Surg Oncol* 23 (5):1639-1645. doi:10.1245/s10434-015-5065-3
33. Ohashi M, Morita S, Fukagawa T, Oda I, Kushima R, Katai H (2015) Functional Advantages of Proximal Gastrectomy with Jejunal Interposition Over Total Gastrectomy with Roux-en-Y Esophagojejunostomy for Early Gastric Cancer. *World J Surg* 39 (11):2726-2733. doi:10.1007/s00268-015-3180-8
34. Nishigori T, Okabe H, Tsunoda S, Shinohara H, Obama K, Hosogi H, Hisamori S, Miyazaki K, Nakayama T, Sakai Y (2017) Superiority of laparoscopic proximal gastrectomy with hand-sewn esophagogastronomy over total gastrectomy in improving postoperative body weight loss and quality of life. *Surg Endosc*. doi:10.1007/s00464-016-5403-y
35. Iwahashi M, Nakamori M, Nakamura M, Naka T, Ojima T, Iida T, Katsuda M, Ueda K, Yamaue H (2009) Evaluation of double tract reconstruction after total gastrectomy in patients with gastric cancer: prospective randomized controlled trial. *World J Surg* 33 (9):1882-1888. doi:10.1007/s00268-009-0109-0
36. Nomura E, Lee SW, Tokuhara T, Nitta T, Kawai M, Uchiyama K (2013) Functional outcomes according to the size of the gastric remnant and the type of reconstruction following distal gastrectomy for gastric cancer: an investigation including total gastrectomy. *Jpn J Clin Oncol* 43 (12):1195-1202. doi:10.1093/jjco/hyt141
37. Zhou D, Liu QX, Deng XF, Min JX, Dai JG (2015) Comparison of two different mechanical esophagogastric anastomosis in esophageal cancer patients: a meta-analysis. *J Cardiothorac Surg* 10:67. doi:10.1186/s13019-015-0271-4
38. Kanaya S, Gomi T, Momoi H, Tamaki N, Isobe H, Katayama T, Wada Y, Ohtoshi M (2002) Delta-shaped anastomosis in totally laparoscopic Billroth I gastrectomy: new technique of intraabdominal gastroduodenostomy. *J Am Coll Surg* 195 (2):284-287

Figure Legends

Figure 1

Positions of the surgical ports. Four 12-mm trocars are placed in the paraumbilical, bilateral abdominal, and epigastric regions. One 5-mm trocar is placed in the left hypochondral area.

Figure 2

A: A purple line is drawn on the antimesenteric border of the jejunum. C: Black squares show the entry holes of the linear stapler made on the greater curvature of the remnant stomach and the antimesenteric wall of the jejunum. The dotted line is drawn at the antimesenteric border of the jejunum. D: Both the stomach and jejunum are twisted posteriorly. The arrows indicate the twisting direction for the remnant stomach and the jejunum. Then, a side-to-side jejunogastrostomy is performed between the posterior wall of the remnant stomach and the jejunum. B: Intraoperative photo of the side-to-side jejunogastrostomy performed between the posterior wall of the remnant stomach and the jejunum. E: The jejunum rides on the remnant stomach by returning the torsion of the jejunum to the counter-clockwise direction, because the twisted jejunum with the support of the mesentery returns to the original position. The jejunum of the anastomotic site is lifted to the ventral side by the remnant stomach that enters behind.

ST: stomach, J: jejunum.

Figure 3

A: The distal jejunum from the jejunogastrostomy can be lifted up to the cranial side, because the esophagojejunosomy and jejunojejunosomy are placed a short distance from each other without sacrificing jejunum and dividing the mesentery. The white arrow shows the site of esophagojejunosomy. The white arrow head shows the jejunojejunosomy site. ST: stomach, J: jejunum. B: The schema shows an overview of the double-tract reconstruction before jejunojejunosomy is performed. The grey triangle shows the anastomotic sites. The position of the proximal jejunum (black arrow head) near the ligament of Treitz is able to be pulled toward the cranial side by not sacrificing part of the jejunum. C: The schema shows the overview of the double-tract reconstruction. The jejunum is bent toward the cranial side just distal from the gastrojejunosomy (black arrow) because it is lifted up by the jejunojejunosomy located in the cranial side.

Table 1. Patients' characteristics

	LPG (n = 10)
Age* (years)	70 (55 - 77)
Sex (male : female)	7 : 3
Body mass index* (BMI, kg/m ²)	23.7 (18.3 - 28.7)
ESD before operation	6
Previous abdominal operation	2
Comorbidity	
Hypertension	7
Diabetes mellitus	3
Cerebral infarction	2
ASA	
1	2
2	7
3	1

*Median (range)

ESD: endoscopic submucosal dissection

Table 2. Surgical and pathological findings

	LPG (n = 10)
Operation time* (minutes)	285 (146 - 440)
Reconstruction time* (minutes)	108 (77 - 194)
Estimated blood loss* (ml)	0 (0 - 25)
Tumor size* (cm)	3.0 (1.5 - 5.0)
Proximal resection margin* (cm)	3.4 (1.1 - 7.5)
distal resection margin* (cm)	4.4 (2.1 - 8.5)
pT1b	10
pN0	10
pStage	
Ia	10
Numbers of retrieved lymph node*	32 (22 - 56)

*Median (range)

POD: Postoperative day

Table 3. Operative outcomes and complications

	LPG (n = 10)
Time to fluid diet* (POD)	3 (3 - 5)
Postoperative hospital stays* (days)	13 (9 - 16)
Anastomotic complications	
Intraperitoneal hemorrhage	0
Leakage of anastomosis or duodenal stump	0
Bowel or anastomosis obstruction	
Anastomotic stenosis	0
	0
Late complications	
Anastomotic stenosis	0
Reflux symptoms	2
dumping syndrome	0
Re-operaton	0
Postoperative mortality	0
<u>Body weight loss (%), mean \pm SD</u>	<u>14.0 \pm 7.1</u>
<u>Decline in serum hemoglobin (%), mean \pm SD</u>	<u>5.4 \pm 10.4</u>
Recurrence	0
Adjuvant chemotherapy	0
Overall survival rate	100%
Median follow-up* (month)	10.7 (1 - 37.6)

*Median (range)