



Title	C1-triangulations of semialgebraic sets
Author(s)	Ohmoto, Toru; Shiota, Masahiro
Citation	Journal of topology, 10(3), 765-775 https://doi.org/10.1112/topo.12024
Issue Date	2017-09
Doc URL	http://hdl.handle.net/2115/71421
Rights	This is the peer reviewed version of the following article: Ohmoto, T. and Shiota, M. (2017), C1-triangulations of semialgebraic sets. Journal of Topology, 10: 765–775, which has been published in final form at doi:10.1112/topo.12024. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	article (author version)
File Information	C1_triangulation_JT.pdf



[Instructions for use](#)

C^1 -TRIANGULATIONS OF SEMIALGEBRAIC SETS

TORU OHMOTO AND MASAHIRO SHIOTA

ABSTRACT. We show that every semialgebraic set admits a semialgebraic triangulation such that each closed simplex is C^1 differentiable. As an application, we give a straightforward definition of the integration $\int_X \omega$ over a compact semialgebraic subset X of a differential form ω on an ambient semialgebraic manifold. This provides a significant simplification of the theory of semialgebraic singular chains and integrations without using geometric measure theory. Our results hold over every (possibly non-archimedean) real closed field.

1. INTRODUCTION

In this paper, we solve a fundamental question on the C^1 -regularity of triangulations for semialgebraic sets. Also an elementary application to the integration of differential forms is discussed, that would be useful, e.g. for the de Rham homotopy theory of semialgebraic sets introduced by Kontsevich-Soibelman [8]. For simplicity, we mainly work on the category of semialgebraic sets and maps over the real number field \mathbb{R} , although our argument goes well in a more general category over a general real closed field, e.g. an *o-minimal* category, the category of *subanalytic sets and maps* or more generally an \mathfrak{X} -category with Axiom (v) in [13, II, p.97].

To begin with, we explain a convention of terminologies which will be used throughout the present paper. Let $X \subset \mathbb{R}^m$ and $Y \subset \mathbb{R}^n$ be semialgebraic sets, *i.e.*, subsets defined by finitely many polynomial equations, inequalities and Boolean operations; a semialgebraic map $X \rightarrow Y$ is a map whose graph is a semialgebraic set in $\mathbb{R}^m \times \mathbb{R}^n$. A subset $X \subset \mathbb{R}^m$ is a *locally semialgebraic set* if for any compact m -disk D in \mathbb{R}^m , $X \cap D$ is semialgebraic; a map $X \rightarrow Y$ is called *locally semialgebraic* if for any compact semialgebraic subset X' of X , the restriction to X' is a semialgebraic map. Note that locally semialgebraic does not imply semialgebraic. Because we systematically deal with possibly non-compact spaces, *we use the notion semialgebraic to mean locally semialgebraic in the above sense*, unless specifically mentioned. Let k be an integer. A *semialgebraic C^k map* $X \rightarrow Y$ is the restriction of a semialgebraic map of class C^k from some semialgebraic open neighborhood of X in \mathbb{R}^m to \mathbb{R}^n ; in fact, we can extend it to $\mathbb{R}^m \rightarrow \mathbb{R}^n$ by multiplying a semialgebraic C^k function ϕ on \mathbb{R}^m such that $\phi = 1$ on X and $\phi = 0$ outside of a smaller closed neighborhood (cf. [13]).

We write (half-)open intervals like as $(0, 1) = \{t \in \mathbb{R} \mid 0 < t < 1\}$ and $[0, 1) = \{t \in \mathbb{R} \mid 0 \leq t < 1\}$ etc.

Date: June 8, 2017.

Key words and phrases. semialgebraic sets, subanalytic sets, \mathfrak{X} -sets, o-minimal category, triangulation, curve selection lemma, differential forms, piecewise algebraic differential forms, de Rham homotopy theory.

1.1. C^1 triangulation. In the present paper, a *semialgebraic triangulation* of a locally closed semialgebraic set X means the pair of a locally finite simplicial complex K and a semialgebraic homeomorphism $f : |K| \rightarrow X$, where $|K|$ is the polyhedron in some Euclidean space \mathbb{R}^ℓ obtained as the geometric realization of K . It is well known that such a semialgebraic triangulation always exists (see Theorem 2.2 below); moreover, it is possible to take (K, f) so that $f|_{\text{Int}(\sigma)}$ is a C^k diffeomorphism for each $\sigma \in K$, where $\text{Int}(\sigma)$ denotes the interior of σ and $k \geq 1$. However we here do not require any regularity of C^0 manifolds $f(\text{Int}(\sigma))$ in our definition of semialgebraic triangulation. Instead, we are concerned with the differentiability (the regularity) of $f(\sigma)$ along the boundary $\partial\sigma$. Our main theorem is the following:

Theorem 1.1. *For a locally closed semialgebraic set X , there exists a semialgebraic triangulation (K, f) so that the map $f : |K| \rightarrow X$ is of class C^1 .*

We call (K, f) in the theorem a *semialgebraic triangulation of X with a C^1 realization*. Then, it follows that for each $\sigma \in K$, $f(\sigma)$ is semialgebraically homeomorphic to the closed simplex $\sigma \subset |K|$ and the differential $d(f|_\sigma)$ exists and is continuous on σ (precisely saying, f is of class C^1 on a neighborhood U_σ of σ in \mathbb{R}^ℓ , and hence, taking the affine subspace H of the same dimension containing σ , $f|_{H \cap U_\sigma}$ is also C^1). However, as a cost for that, $d(f|_\sigma)$ may drop the rank along the boundary $\partial\sigma$ and some smaller dimensional semialgebraic subset of $\text{Int}(\sigma)$.

The proof is elementary, only using standard techniques such as triangulation, tube and the curve selection lemma (some basic facts are summarized in Section 2), so it would be readable even for non-experts. The proof is done in the case of semialgebraic sets over \mathbb{R} , but it is presented in such a way that it is valid in a more general context of \mathfrak{X} -category over a real closed field R in the sense of [13]; in particular, it works in the category of subanalytic sets and maps. Furthermore, our argument fits with *piecewise algebraic spaces* introduced in [8] (Remark 3.8).

There remain some open questions. For instance, we do not know whether or not f can be of class C^k ($2 \leq k \leq \infty$ or $k = \omega$). As known, any Whitney stratified set is triangulable, so it would be natural to ask if the realization map f can be C^1 , see e.g. [12].

1.2. Differential forms and integration. Let M be a semialgebraic manifold and X a compact semialgebraic subset of M ($p = \dim X$). Take a semialgebraic triangulation (K, f) of X with a C^1 realization. Here K is a finite simplicial complex. For a differential p -form $\omega \in \Omega^p(M)$ and a p -simplex $\sigma \in K$, the integral of ω on σ is defined in an obvious way by

$$\int_{f(\sigma)} \omega := \int_\sigma (f|_\sigma)^* \omega$$

because the map $f|_\sigma : \sigma \rightarrow M$ is of class C^1 (the integral is often written by $\int_\sigma \omega$ for short). Suppose that X has the \mathbb{Z} -fundamental class. Put

$$\int_X \omega := \sum \int_{f(\sigma)} \omega$$

where the sum runs over all p -simplices σ in K with suitable orientation so that they form the fundamental cycle. It is almost obvious that the value $\int_X \sigma$ does not depend on the choice of (K, f) of X (Theorem 4.1). Consequently, the theory of integration of differential forms over semialgebraic sets can be treated in entirely the same manner as in the case of manifolds – that simplifies the treatment of semialgebraic chains and integrations, without using geometric measure theory due to Hardt et al [3, 5, 6, 8] (see Remarks 4.3, 4.4). In fact, a traditional way using geometric currents involves the limiting process: $\int_\sigma \omega$ is defined by the integral of ω over the interior $\text{Int}(\sigma)$ which absolutely converges (that is shown by the fact that the induced volume is finite). On the other hand, our definition uses only the *definite integral* of a continuous function on a compact domain σ .

We remark that our definition of $\int_X \omega$ works over not only the real number field \mathbb{R} but also any possibly non-archimedean real closed field R . Then the value of the integral does not take values in the same field R but in a larger saturated field (see [11]). In particular, the volume of a bounded definable set in R^m is bounded within R (Remark 4.5).

Acknowledgement. This work was originally motivated by a question of Tatsuo Suwa about the definition of $\int_X \omega$ using triangulation, which was needed for rigorously establishing the Alexander duality for an embedded analytic subvariety in the Čech-de Rham context [14]. The authors would like to thank Tatsuo Suwa for guiding them to this problem and for useful discussions. During preparing the final version of the present paper, we were informed about two recent preprints [2] and [7]; we also thank G. Comte and M. Hanamura for letting us know of those papers, respectively. The first author is supported by JSPS KAKENHI Grants no.24340007 and no.15K13452, and the second author is supported by JSPS KAKENHI Grant no.26400084.

2. PRELIMINARY

We summarize some basic facts in semialgebraic geometry, which are also valid without any change for the *locally semialgebraic* setting in our sense. As noted in Introduction, we simply say semialgebraic to mean locally semialgebraic, unless specifically mentioned.

2.1. Triangulation. Let k be an integer and $X \subset \mathbb{R}^m$ a semialgebraic set. A *semialgebraic C^k stratification* of X is a family $\mathcal{S} = \{X_i\}$ of semialgebraic subsets of X (called *strata* of \mathcal{S}) so that each X_i is a locally closed C^k submanifold of \mathbb{R}^m , $X = \bigsqcup X_i$ (disjoint), $X_i \cap X_j = \emptyset$ ($i \neq j$), \mathcal{S} is locally finite, and \mathcal{S} satisfies the *frontier condition*, *i.e.*, for each X_i , the closure $\overline{X_i}$ is a union of X_i and some $X_j \in \mathcal{S}$ of smaller dimension. We say that \mathcal{S} is *compatible with* a family \mathcal{A} of subsets of X if for each stratum $V \in \mathcal{S}$ and each $A \in \mathcal{A}$, it holds either $V \subset A$ or $V \cap A = \emptyset$. A *semialgebraic C^k stratification of a semialgebraic C^0 map $f : X \rightarrow Y$* is the pair of semialgebraic C^k stratifications \mathcal{S} of X and \mathcal{T} of Y such that for each $V \in \mathcal{S}$, $f(V)$ is a stratum of \mathcal{T} and $f|_V : V \rightarrow f(V)$ is a semialgebraic C^k submersion. We denote it by $f : (X; \mathcal{S}) \rightarrow (Y; \mathcal{T})$. Such a semialgebraic stratification always exists (the statement can be more enhanced so that \mathcal{S} and \mathcal{T} satisfy the Whitney condition, but we do not need it later):

Lemma 2.1. (Canonical stratification of maps [4], [13, I. Lemmas 2.2, 2.6; 2.10, 2.14]) Let X be a semialgebraic set and $k \geq 1$. Let X' be a closed semialgebraic subset of X , \mathcal{S} a semialgebraic C^k stratification of $X - X'$, and \mathcal{A} a finite family of semialgebraic subsets of X' . Assume $\dim X' < \dim V$ for any $V \in \mathcal{S}$. Then there exists a semialgebraic C^k stratification \mathcal{S}' of X' compatible with \mathcal{A} such that $\mathcal{S} \sqcup \mathcal{S}'$ is a semialgebraic C^k stratification of X .

Moreover, let $f : X \rightarrow Y$ be a proper semialgebraic C^0 map such that $f(X') \cap f(X - X') = \emptyset$. Assume $f : (X - X'; \mathcal{S}) \rightarrow (f(X - X'); \mathcal{T})$, where \mathcal{T} consists of all $f(V)$, $V \in \mathcal{S}$, is a semialgebraic C^k stratification of $f|_{X - X'}$. Then we can choose \mathcal{S}' so that $f : \mathcal{S} \sqcup \mathcal{S}' \rightarrow \mathcal{T} \sqcup \mathcal{T}'$, where \mathcal{T}' consists of all $f(V')$, $V' \in \mathcal{S}'$, is a semialgebraic C^k stratification of $f : X \rightarrow f(X)$.

Theorem 2.2. (Łojasiewicz's triangulation theorem [9], [13, II.2.1]) Let X be a locally closed semialgebraic set in \mathbb{R}^m and \mathcal{A} a finite family of semialgebraic subsets of X . Then there is a pair of (K, h) of a locally finite simplicial complex K and a semialgebraic homeomorphism $h : |K| \rightarrow X$ so that $\{h(\text{Int}(\sigma)), \sigma \in K\}$ is a semialgebraic C^k stratification of X compatible with \mathcal{A} .

Those theorems have been extended to more general categories (o-minimal, subanalytic, \mathfrak{X}), e.g. see [13].

2.2. Curve selection lemma. We will frequently use the following version of the *curve selection lemma* (*wing lemma*) afterward. There is a number of literature; the origin goes back to Whitney and Milnor, and a recent one is, e.g. a version in o-minimal category [10, Lem.1.7]. For proving its \mathfrak{X} -version, it suffices to use the \mathfrak{X} -isotopy lemma [13, II.6].

Lemma 2.3. (Wing lemma) Let $V \subset \mathbb{R}^m$ be a non-empty semialgebraic open subset, and Z a semialgebraic subset of $\mathbb{R}^m \times \mathbb{R}^n$ so that its closure \bar{Z} contains $V \times \{0\}$ and $Z \cap (V \times \{0\}) = \emptyset$. Then there is a non-empty semialgebraic open subset $U \subset V$ which admits a semialgebraic C^0 map $c : U \times [0, a) \rightarrow \mathbb{R}^n$ with some $a > 0$ so that $c(x, 0) = 0$ for $x \in U$, $(x, c(x, t)) \in Z$ and $|c(x, t)| = t$ for any $t \neq 0$, and c is of class C^k on $U \times (0, a)$.

3. PROOF – PANEL BEATING

3.1. Panel beating. We will prove a bit more general statement.

Theorem 3.1. Let X be a locally closed semialgebraic set with a semialgebraic triangulation (K, f) , and $\varphi : X \rightarrow Y$ a semialgebraic C^0 map. Then there exists a semialgebraic homeomorphism χ of $|K|$ such that χ preserves any $\sigma \in K$ and the composed map

$$\varphi \circ f \circ \chi : |K| \xrightarrow{\cong} |K| \xrightarrow{\cong} X \rightarrow Y$$

is of class C^1 .

Theorem 1.1 is the case of $X = Y$ and $\varphi = id_X$.

Definition 3.2. We call χ a *panel beating* of the triangulation (K, f) of X with respect to $\varphi : X \rightarrow Y$.

Notice that in Theorem 3.1, χ can be chosen so that χ itself, $f \circ \chi$ and $\varphi \circ f \circ \chi$ are of class C^1 simultaneously. In fact, we have

Corollary 3.3. *Let X and (K, f) as above. Suppose that we are given semialgebraic C^0 maps $\varphi^{(j)} : X \rightarrow Y_j$ ($1 \leq j \leq s$). Then, there is a common panel beating χ such that each $\varphi^{(j)} \circ f \circ \chi$ is of class C^1 .*

Proof. Apply Theorem 3.1 to the map $\varphi = \prod \varphi^{(j)} : X \rightarrow Y = \prod Y_j$. □

Here is an obvious example. Consider $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ given by $\varphi(u) = \sqrt{|u|}$, and decompose \mathbb{R} into $(-\infty, 0)$, $\{0\}$ and $(0, \infty)$. Put

$$\chi_0(u) = \begin{cases} u^4 & (u \geq 0) \\ -u^4 & (u \leq 0) \end{cases}$$

then it is a panel beating; indeed, $\varphi \circ \chi_0(u) = u^2$. At a key step (Lemma 3.5) in the proof of Theorem 3.1, we glue together this kind of mappings χ_0 via a semialgebraic version of partition of unity to construct the desired semialgebraic homeomorphism $\chi : |K| \rightarrow |K|$.

3.2. Proof of Theorem 3.1. Let X be a locally closed semialgebraic set which admits a semialgebraic triangulation (K_1, f_1) with $|K_1| \subset \mathbb{R}^m$, and Y a semialgebraic set in \mathbb{R}^p . Let $\varphi : X \rightarrow Y$ be a semialgebraic C^0 map. By the semialgebraic Tietze theorem, e.g. [1, Prop.2.6.9] (or using a semialgebraic partition of unity and tubes [13]), we see that the map $\varphi \circ f_1$ is extended to a semialgebraic C^0 map $\mathbb{R}^m \rightarrow \mathbb{R}^p$. The graph map is denoted by $f : \mathbb{R}^m \rightarrow \mathbb{R}^n := \mathbb{R}^m \times \mathbb{R}^p$, which is semialgebraic C^0 and proper. Also we extend K_1 to a simplicial decomposition (locally closed infinite simplicial complex) K of \mathbb{R}^m . To prove Theorem 3.1 we find a semialgebraic homeomorphism $\chi : \mathbb{R}^m \rightarrow \mathbb{R}^m$ so that χ preserves any simplices of K and $f \circ \chi$ is of class C^1 .

By Lemma 2.1, there is a semialgebraic C^2 stratification $\mathcal{S} \rightarrow \mathcal{T}$ of f , such that each simplex in K is a finite union of some strata of \mathcal{S} . The restriction of f on each stratum is of class C^2 by definition, so in particular f is C^1 on the union of open strata. Now we use the downward induction on dimension d ($\leq m - 1$) of the ‘bad subset’ where f is not C^1 : We will construct a panel beating χ around the subset so that $f \circ \chi$ is C^1 off a smaller subset of less dimension. Clearly, it suffices to construct χ around each connected stratum of top dimension of the bad subset. Furthermore, we may assume that the stratum is semialgebraically diffeomorphic to \mathbb{R}^d (by subdividing \mathcal{S} and \mathcal{T} if necessary).

(Induction hypothesis) Suppose that there is a stratum $V \simeq \mathbb{R}^d$ of \mathcal{S} such that f is of class C^1 over the complement $\mathbb{R}^m - \bar{V}$.

Since V is a C^2 submanifold, the tangent bundle TV is of class C^1 . By using the orthogonal projection to $T_x V$, we can obtain a semialgebraic C^1 tube $T = (|T|, \pi, \rho)$ at V in \mathbb{R}^m as in [13, Lemma II.5.1]. Here $|T|$ is a semialgebraic neighborhood of V in \mathbb{R}^m together with semialgebraic C^1 maps $\pi : |T| \rightarrow V$ and $\rho : |T| \rightarrow \mathbb{R}$ so that

- $\rho(y) \geq 0$ ($y \in |T|$) and $V = \rho^{-1}(0)$,

- $\pi \times \rho : |T| - V \rightarrow V \times \mathbb{R}_{>0}$ is a proper submersion onto $V \times (0, 1]$,
- $d(y, V) = \inf\{\|x - y\|, x \in V\} \rightarrow 0$ ($y \in T$) as $\pi(y)$ tends to a point in $\bar{V} - V$.

We choose T small enough so that

- $|T| \cap \sigma = \emptyset$ for $\sigma \in K$ with $\sigma \cap V = \emptyset$.

Let D denote the unit closed disk in \mathbb{R}^{m-d} centered at 0. By the construction of T , there is a semialgebraic C^1 diffeomorphism

$$g : V \times D \rightarrow |T|, \quad g(x, u) = x + \sum_k u_k \mathbf{u}_k(x)$$

with $x = (x_1, \dots, x_d) \in V$ and $u = (u_1, \dots, u_{m-d}) \in D$ such that $\mathbf{u}_1(x), \dots, \mathbf{u}_{m-d}(x) \in T_x V^\perp$ form a semialgebraic C^1 vector frame so that

$$\pi \circ g(x, u) = x \quad \text{and} \quad \rho \circ g(x, u) = |u| = \left(\sum u_k^2 \right)^{1/2}.$$

For $u \neq 0$, set $\bar{u} := u/|u| \in \partial D$.

We construct a modification χ of $|T|$ as follow. Let $I = (0, 1) \subset \mathbb{R}$ an open interval. Let $\eta : V \times I \rightarrow I$ be a semialgebraic C^1 map such that $I \ni t \mapsto \eta(x, t) \in I$ is a diffeomorphism for each $x \in V$, $\eta(x, t) = t$ for $t \geq \frac{1}{2}$, and

$$\eta(x, t) \rightarrow 0 \quad \text{and} \quad \frac{\partial \eta}{\partial t}(x, t) \rightarrow 0 \quad \text{as} \quad t \rightarrow 0$$

(in fact, in Lemma 3.5 below, we will take η depending on the variables x in order to satisfy a nice property). We then define a semialgebraic homeomorphism $\chi : |T| \rightarrow |T|$ by

$$\chi(y) = \chi(g(x, u)) = g(x, \eta(x, |u|) \bar{u})$$

for $y = g(x, u) \in |T| - V$ and $\chi|_V = id_V$. We also write $\chi(x, u) = (x, \eta(x, |u|) \bar{u}) \in V \times D$. Since $\chi(y) = y$ if $\rho(y) \geq \frac{1}{2}$, χ is extended to $\chi : \mathbb{R}^m \rightarrow \mathbb{R}^m$ so that $\chi = id$ outside of $|T|$.

Lemma 3.4. *The semialgebraic homeomorphism χ preserves every $\sigma \in K$ with $\sigma \cap V \neq \emptyset$, and $f \circ \chi : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is a semialgebraic C^0 map and is of class C^1 on $\mathbb{R}^m - \bar{V}$.*

Proof. If $V \subset \tau$ and $\sigma \cap V \neq \emptyset$, then τ is a face of σ or $\tau = \sigma$; if $y \in |T| \cap \sigma$ and $x = \pi(y)$, then the segment \overline{xy} is in σ , hence $\chi(y) \in \sigma$. Since π, ρ, η are of class C^1 , χ is so over $|T| - V$. By the assumption that f is C^1 off \bar{V} , $f \circ \chi$ is also. \square

Lemma 3.5. *We can choose $\eta(x, t)$ such that for every $1 \leq k \leq m - d$ and for every $x \in V$, it holds that*

- $\frac{\partial (f \circ \chi)}{\partial u_k}(x, 0) = 0$
- $\frac{\partial (f \circ \chi)}{\partial u_k}(x', u') \rightarrow 0$, as $(x', u') \rightarrow (x, 0) \in V \times D$

Proof. For sufficiently small $t \geq 0$, we set

$$g(t) := \sup_{x \in V, \bar{u} \in \partial D} \|f(x, t \bar{u}) - f(x, 0)\|.$$

Note that $f(x, t\bar{u}) - f(x, 0) \rightarrow 0$ as x converges to a point in the boundary $\bar{V} - V$, because f is continuous and $d(g(x, t\bar{u}), V) \rightarrow 0$ by a condition of T , therefore $g(t)$ is attained as the maximum. It is easily checked that $g(t)$ is a semialgebraic C^0 -function; in particular, $g(t) \rightarrow g(0) = 0$ as $t \rightarrow 0$. By the (ordinary) curve selection lemma, we see $g(t) = O(t^\alpha)$ with some positive rational number α . Put $\xi(t) = t^{r_0}$ with $r_0 > \max\{\frac{1}{\alpha}, 1\}$, then $\frac{g(\xi(t))}{t} \rightarrow 0$ as $t \rightarrow 0$. Thus $\frac{1}{t}(f(x, \xi(t)\bar{u}) - f(x, 0)) \rightarrow 0$ for any x and \bar{u} . In particular, we can choose $\eta(x, t)$ so that $\eta(x, t) \leq \xi(t)$ for every $x \in V$; then every directional derivative of $f \circ \chi (= f(x, \eta(x, |u|)\bar{u}))$ at $(x, 0)$ exists, and it is actually zero. Thus the first claim is verified. Next, we find such a function $\eta(x, t)$ which also leads to the second claim. Since f and χ are C^1 on $T - V$, $f \circ \chi$ is C^1 and there are semialgebraic C^0 maps $A_k(x, u)$ and $B_k(x, u)$ for each k such that

$$\frac{\partial(f \circ \chi)}{\partial u_k}(x, u) = \eta(x, |u|)A_k(x, u) + \frac{\partial\eta}{\partial t}(x, |u|)B_k(x, u) \quad (x \in V, |u| \neq 0).$$

For $x \in V$ and $0 < t < 1$, set

$$a(x, t) := \min \left\{ 1, \frac{1}{|A_k(x, u)|}, \frac{1}{|B_k(x, u)|} \mid |u| = t, 1 \leq k \leq m - d \right\}.$$

Notice that $a(x, t)$ is a positive semialgebraic C^0 function on $V \times I$. By using a partition of unity on V (which is locally semialgebraic in our sense), we can construct $\eta(x, t)$ which satisfies $\eta(x, t) \leq \xi(t)$, $0 < \eta(x, t) < t \cdot a(x, t)$ and $0 < \frac{\partial\eta}{\partial t}(x, t) < t \cdot a(x, t)$ for $0 < t \ll 1$; then both $\eta \cdot |A_k|$ and $\frac{\partial\eta}{\partial t} \cdot |B_k|$ are less than $t = |u|$, thus we have $|\frac{\partial(f \circ \chi)}{\partial u_k}(x, u)| < 2|u|$, that implies the second claim. In fact, it is easy to find such a function $\eta(x, t)$ locally. For a compact semialgebraic subset $K \subset V$, which is the closure of an open set of V , put $a_K(t) := \min_{x \in K} a(x, t)$; by the curve selection lemma again, we may write $a_K(t) = O(t^\alpha)$ with some $\alpha \geq 0$; then we can take $\eta_K(t) = t^r$ ($r > r_0$) so that η_K and $\frac{d}{dt}\eta_K$ are smaller than $t \cdot a_K(t)$ near $t = 0$. By a semialgebraic bump function, it is extended to a semialgebraic diffeomorphism $\eta_K : I \rightarrow I$ with $\eta_K(t) = t$ for $t \geq 1/2$. Put $\eta_K(x, t) := \eta_K(t) : K \times I \rightarrow I$. To obtain the desired function $\eta(x, t)$ on $V \times I$, it suffices to gather together such η_K by mean of a semialgebraic partition of unity on V . \square

Here is an obvious claim in elementary calculus, which will be used in the proof of the next Lemma 3.7.

Lemma 3.6. *Let $\nu(s, t)$ be a C^1 function on $\mathbb{R} \times (0, 1)$ such that $|\frac{\partial\nu}{\partial s}(s, t)| > K$ for some fixed $K > 0$. If ν extends continuously to $\mathbb{R} \times \{0\}$, then $\nu(s, 0)$ is not constant.*

Proof. Assume $\frac{\partial\nu}{\partial s}(s, t) > K$. Use $\nu(s_1, t) - \nu(s_0, t) > K(s_1 - s_0)$ if $s_1 > s_0$, and take $t \rightarrow 0$. \square

Lemma 3.7. *Let η be as in Lemma 3.5. Then there is a semialgebraic subset S of V with $\dim S < d = \dim V$ satisfying that*

- $\frac{\partial(f \circ \chi)}{\partial x_j}(x, 0) = \frac{\partial f}{\partial x_j}(x, 0)$
- $\frac{\partial(f \circ \chi)}{\partial x_j}(x', u') \rightarrow \frac{\partial f}{\partial x_j}(x, 0)$, as $(x', u') \rightarrow (x, 0) \in (V - S) \times D$

for every $1 \leq j \leq d$ and for every $x \in V - S$.

Proof. The first claim follows from the condition that $\chi|_V = id_V$. For the second, we set

$$F(x, u) = f \circ \chi(x, u) - f(x, 0), \quad F = (F_1, \dots, F_n) \in \mathbb{R}^n,$$

which is continuous on $V \times D$ and of class C^1 on $V \times (D - \{0\})$. Note that

$$F(x, 0) = 0 \quad (x \in V).$$

Put

$$h(x, u) := \max_{i,j} \left| \frac{\partial F_i}{\partial x_j}(x, u) \right| \quad (u \neq 0),$$

which is a semialgebraic C^0 function on $V \times (D - \{0\})$. It holds that

$$\left| \frac{\partial(f_i \circ \chi)}{\partial x_j}(x', u') - \frac{\partial f_i}{\partial x_j}(x, 0) \right| \leq h(x', u') + \left| \frac{\partial f_i}{\partial x_j}(x', 0) - \frac{\partial f_i}{\partial x_j}(x, 0) \right|$$

and $\frac{\partial f_i}{\partial x_j}(x, 0)$ is continuous by the assumption that $f|_V$ is C^1 , hence it is enough to consider the condition that $h(x', u') \rightarrow 0$ as $(x', u') \rightarrow (x, 0)$.

Let S be the subset of V consisting of x such that there exists a sequence of points $(x', u') \in V \times D$ converging to $(x, 0)$ with $h(x', u')$ not converging to 0, that is,

$$S = \left\{ x \in V \mid \exists \epsilon > 0 \quad \forall \delta > 0 \quad \exists (x', u') \in V \times D \quad \text{s.t.} \quad \begin{array}{l} |x' - x| + |u'| < \delta, \\ h(x', u') > \epsilon \end{array} \right\}.$$

The ‘bad set’ S is semialgebraic; indeed, S is obtained by projecting a semialgebraic subset S' in $V \times \mathbb{R}_{>0}$ consisting of (x, ϵ) which satisfy the corresponding condition (*i.e.*, $\forall \delta > 0, \exists (x', u')$ such that $|x' - x| + |u'| < \delta$ and $h(x', u') > \epsilon$). We show that S is of dimension $< d = \dim V$.

Suppose that $\dim S = d$. Below we make five steps for deducing the contradiction. A major tool is the curve selection lemma (Lemma 2.3), which is used in each step from (1) to (4).

(1) There is a non-empty semialgebraic open subset $U_0 \subset S$ with a constant $\epsilon_1 > 0$ such that for each $x \in U_0$, there is a sequence of points $(x', u') \in V \times D$ converging to $(x, 0)$ with $h(x', u') > \epsilon_1$. This is easily verified by applying Lemma 2.3 to the above semialgebraic set $S' \subset S \times \mathbb{R}$; in fact, the closure $\overline{S'}$ contains $S \times 0$, hence there are U_0 and $\epsilon_1 > 0$ so that $U_0 \times (0, 2\epsilon_1) \subset S'$; in particular, $U_0 \times \epsilon_1 \subset S'$.

(2) Let $Z := \{(x, u) \in V \times D \mid h(x, u) > \epsilon_1\}$, then \overline{Z} contains $U_0 \times \{0\}$. By Lemma 2.3, there is a non-empty semialgebraic open subset U_1 of U_0 which admits a semialgebraic C^0 map

$$c : U_1 \times [0, a_1) \rightarrow D$$

for some $a_1 > 0$ so that $u = c(x, t)$ is C^1 for $t > 0$, and $c(x, 0) = 0$, $t = |c(x, t)|$ and

$$h(x, c(x, t)) > \epsilon_1 \quad (\forall (x, t) \in U_1 \times (0, a_1)).$$

(3) Put $c(x, t) = (c_1(x, t), \dots, c_{m-d}(x, t)) \in D$. We claim that all the derivatives $\frac{\partial c_k}{\partial x_j}(x, t)$ are bounded over some open subset $U \times (0, a)$ of $U_1 \times (0, a_1)$. Suppose that $v(x, t) := \left| \frac{\partial c_1}{\partial x_1}(x, t) \right|$ is

unbounded over any open subsets of the form $U \times (0, a)$. Take $M > 0$. Then for any $x \in U_1$, there is a sequence of points (x', u') converging to $(x, 0)$ with $v(x', u') > M$. Let

$$Z' := \{(x, u) \in U_1 \times (0, a_1) \mid v(x, u) > M\},$$

then $\overline{Z'}$ contains $U_1 \times \{0\}$, and Z' is semialgebraic, for $v(x, t)$ is semialgebraic. By Lemma 2.3, there is an open subset $U \times (0, a)$, over which $v(x, u) > M$. Take a small segment in U given by $\gamma(s) := b + (s, 0, \dots, 0)$ ($0 \leq s \leq \delta$, $b \in U$) and put $\nu(s, t) := c_1(\gamma(s), t)$. Since $c(x, 0) = 0$, we have $\nu(s, 0) = 0$, but it makes the contradiction by Lemma 3.6. Repeating the same argument for all j and k , the claim is proved.

(4) Without loss of generality, we may assume that the partial derivative of the first component function of F with respect to the first coordinate of V satisfies

$$\left| \frac{\partial F_1}{\partial x_1}(x, c(x, t)) \right| > \epsilon_1 \quad (\forall (x, t) \in U \times (0, a)).$$

In fact, if not so, then for each i, j , it holds that for any $x \in U$, there is a sequence $(x', t') \rightarrow (x, 0)$ with $\left| \frac{\partial F_i}{\partial x_j}(x', c(x', t')) \right| \leq \epsilon_1$. Hence by applying Lemma 2.3 again, we find a (common) small positive number a' so that $\left| \frac{\partial F_i}{\partial x_j}(x, c(x, t)) \right| \leq \epsilon_1$ on $U \times (0, a')$ for every i, j , but that contradicts the condition $h > \epsilon_1$ in the step (2).

(5) Consider a C^0 function on $U \times [0, a)$ given by

$$\phi(x, t) := F_1(x, c(x, t)),$$

which is C^1 off $t = 0$. By the chain rule, we have

$$\frac{\partial \phi}{\partial x_1}(x, t) = \frac{\partial F_1}{\partial x_1}(x, c(x, t)) + \sum_k \frac{\partial (f_1 \circ \chi)}{\partial u_k}(x, c(x, t)) \frac{\partial c_k}{\partial x_1}(x, t).$$

Since $\left| \frac{\partial c_k}{\partial x_j}(x, t) \right|$ is bounded by (3) above, the sum \sum_k in the right hand side converges to 0 as $t \rightarrow 0$ by Lemma 3.5. Therefore, by (4), we can take an open subset U' with $\overline{U'} \subset U$, and $0 < a' < a$ such that

$$\left| \frac{\partial \phi}{\partial x_1}(x, t) \right| > \frac{\epsilon_1}{2} \quad (\forall (x, t) \in U' \times (0, a')).$$

Take a segment in U' of the form $\gamma(s) := b + (s, 0, \dots, 0)$ and put $\nu(s, t) := \phi(\gamma(s), t)$. By Lemma 3.6, $\nu(s, 0)$ is not constant. However, for any $x \in U$,

$$\phi(x, 0) = F_1(x, c(x, 0)) = F_1(x, 0) = 0,$$

that causes the contradiction. This completes the proof. \square

Proof of Theorem 3.1: From the above Lemmas 3.5 and 3.7,

$$d(f \circ \chi)(y) \rightarrow d(f \circ \pi)(x) \quad \text{as } |T| \ni y \rightarrow x \in V - S,$$

thus $f \circ \chi$ is of class C^1 on the complement to $(\overline{V} - V) \sqcup S$. The bad subset is semialgebraic and of dimension less than $d = \dim V$. By Lemma 3.4, χ preserves K . By applying Lemma

2.1 again we obtain a stratification of $f \circ \chi : \mathbb{R}^m \rightarrow \mathbb{R}^n$ which subdivides K and the bad subset, hence the induction step is verified. This completes the proof of Theorem 3.1. \square

Remark 3.8. (PA spaces) In Theorems 1.1 X is assumed to be embedded in \mathbb{R}^m , but the global embeddability is not needed essentially, because key lemmas in the proof deal with local problems. Indeed, our argument fits with the framework of (possibly non-compact) *piecewise algebraic spaces* introduced by Kontsevich-Soibelman [8]; a PA space is a locally compact Hausdorff space $X = \bigcup X_\alpha$ made by glueing countably many compact semialgebraic sets X_α via semialgebraic homeomorphisms on constructible subsets $X_\alpha \supset X_{\alpha\beta} \xrightarrow{\cong} X_{\beta\alpha} \subset X_\beta$. In the same approach, we can see that any PA space admits a (PA) triangulation with a C^1 realization so that the (locally finite) simplicial complex is locally realized in some \mathbb{R}^m .

4. SEMIALGEBRAIC CHAINS AND INTEGRATIONS


4.1. Integrals of differential forms. Let M be a semialgebraic manifold, X a compact semialgebraic subset of M ($\dim X = p$) with a semialgebraic subset $\partial X \subset X$ so that ∂X is locally the boundary of the regular part X_{reg} around each regular point of ∂X . Assume that X has a fundamental class $\mu_X \in H_p(X, \partial X; \mathbb{Z})$, *i.e.*, $i_*\mu_X$ generates $H_p(X, X - x; \mathbb{Z}) \simeq \mathbb{Z}$ for any regular point $x \in X_{\text{reg}} - \partial X$ with inclusion $i : X - x \hookrightarrow X$.

Let (K, f) be a semialgebraic triangulation of X with a C^1 realization as in Theorem 1.1. Then for each simplex $\sigma \in K$, $f|_\sigma : \sigma \rightarrow X$ is a semialgebraic C^1 map. The chain $\sum \sigma$ of all p -simplices in K with suitable orientation represents μ_X ; we define the integral over X of a differential p -form ω on M by

$$\int_X \omega := \sum \int_\sigma (f|_\sigma)^* \omega.$$

We show that the integral does not depend on the choice of (K, f) .

Let (K_1, f_1) and (K_2, f_2) be semialgebraic triangulations of X with C^1 realizations. Obviously, there is a semialgebraic homeomorphism $\tau : |K_1| \rightarrow |K_2|$ with $f_1 = f_2 \circ \tau$. Applying Lemma 2.1, we have a stratification for τ , and let \mathcal{A} be the set of strata of the source stratification of τ , $\sigma \in K_1$ and $\tau^{-1}(\sigma')$ of $\sigma' \in K_2$. By Theorem 2.2, we can find a semialgebraic triangulation (K, ϕ_1) of $|K_1|$ compatible with \mathcal{A} . Set $\phi_2 = \tau \circ \phi_1 : |K| \rightarrow |K_2|$ and $g = f_1 \circ \phi_1 = f_2 \circ \phi_2 : |K| \rightarrow X$. Notice that any simplex of K_j ($j = 1, 2$) is the union of the image of finitely many simplices of K via ϕ_j . By Corollary 3.3, there is a common panel beating χ of $|K|$ such that $\phi_j \circ \chi$ ($j = 1, 2$) are of class C^1 . Put $f := g \circ \chi = f_j \circ (\phi_j \circ \chi)$, then f is C^1 , for f_j and $\phi_j \circ \chi$ are so. We call the obtained triangulation (K, f) of X a common C^1 refinement of (K_1, f_1) and (K_2, f_2) .



Theorem 4.1. *The integral $\int_X \omega$ is well-defined, i.e., it does not depend on the choice of semialgebraic triangulation of X with C^1 realization.*

Proof. Let $\sigma : \Delta^p \rightarrow X$ be a semialgebraic C^0 map. By a panel beating, we may assume that σ itself is of class C^1 . Let (K, h) be a semialgebraic triangulation of Δ^p with a C^1 realization. Then $h : \Delta^p = |K| \rightarrow \Delta^p$ is a C^1 map which is diffeomorphic off a measure zero subset, thus we have $\int_\sigma \omega = \int_\Delta g(x)dx = \int_\Delta g(h(u))|\frac{\partial h}{\partial u}|du = \int_{\sigma \circ h} \omega$ for a differential p -form $\sigma^* \omega = g(x)dx_1 \wedge \dots \wedge dx_p$. By using a common C^1 refinements of triangulations, the assertion immediately follows. \square

Proposition 4.2. *The Stokes formula holds, $\int_X d\omega = \int_{\partial X} \omega$, provided X and ∂X have fundamental cycles.*

Proof. It is the same as in the smooth case. \square

The above argument is also applied to the case of non-compact X and compact supported ω . We can also work on subanalytic sets or more generally \mathfrak{X} -sets (see Remarks below); in particular, the same treatment is available for complex analytic sets and integrations of (p, q) -differential forms. There is an alternative approach for the integration on subanalytic sets, e.g. [15].

Remark 4.3. (Semialgebraic currents) Our integral $\int_X \omega$ over a compact semialgebraic set X coincides with the integral as a *semialgebraic current* in geometric measure theory [3, 5, 6]. The key point in the theory is that X has a finite volume, hence the integral over the interior absolutely converges. Subanalytic currents are also available. However, the theory of currents is unclear for a more general setting such as the category of \mathfrak{X} -sets or o-minimal category over any real closed field, while our approach does not have any trouble in such a general context.

Remark 4.4. (de Rham homotopy theory of PA spaces) In [8] the foundation of *semialgebraic* differential forms on semialgebraic sets, more precisely *PA forms* on PA spaces, has been introduced for the use in proving the formality of little cubes operads; in particular, as an analogy to the de Rham algebra of differential forms on smooth manifolds, the algebra of PA forms encodes the information of the real homotopy type of the PA space. A large portion of the theory of PA forms has been clarified in [6] through a bit long discussion using semialgebraic currents. Our approach using panel beatings simplifies the arguments in

a straightforward way; it enables us to directly connect the de Rham algebra of PA forms with Sullivan's model (CDGA) with real coefficients. Another advantage is that our approach works over an *arbitrary* real closed field.

Remark 4.5. (O-minimal category) We may work over any o-minimal category over any general real closed field R . Let X be a bounded definable subset X of dimension d in R^m . Assume that $X \subset I^m$ where I is the closed interval $[0, L]$, $L \in R$. By subdividing I into n intervals $[L/n, (k+1)L/n]$, I^m breaks into m -dimensional small cubes U_i with $1 \leq i \leq n^m$. For each i , set $c_i = 1$ if $X \cap U_i \neq \emptyset$, $c_i = 0$ otherwise. Let $\delta_n := \sqrt{m} \cdot L/n$, the diameter of the small cube, and put $v_n^{(d)}(X) := \left(\sum_{i=1}^{n^m} c_i\right) (\delta_n)^d$. Then, by using Theorem 1.1, we can show that v_n is uniformly bounded in R , that is, there is $c \in R$ such that $v_n^{(d)}(X) < c$ for any $n \in \mathbb{N}$. Moreover, the d -th Hausdorff measure $\mathcal{H}^d(X)$ (or the volume) of X is given by $\lim_{n \rightarrow \infty} v_n^{(d)}(X)$, which exists in some saturated field containing R . However, the argument about the saturated field is quite complicated, so we do not discuss it here. See [11] for details.

REFERENCES

- [1] J. Bochnak, M. Coste and M.F Roy, *Real algebraic geometry*, Springer-Verlag (1998).
- [2] M. Czapla and W. Pawlucki, Strict C^1 -triangulations in o-minimal structures, preprint (2017).
- [3] H. Federer, *Geometric measure theory*, Springer-Verlag, Heidelberg and New York, 1969.
- [4] C. G. Gibson, K. Wirthmüller, A. A. Du Plessis, E. J. N. Looijenga, Topological stability of smooth mappings, Lecture Notes in Math., 552, Springer, 1976.
- [5] R.M. Hardt, Topological properties of subanalytic sets, Trans. Amer. Math. Soc., 211 (1975), 57–70.
- [6] R.M. Hardt, P. Lambrechts, V. Turchin and I. Volić, Real homotopy theory of semi-algebraic sets, Algebraic & Geometric Topology **11** (2011), 2477–2545, DOI: 10.2140/agt.2011.11.2477.
- [7] M. Hanamura, K. Kimura and T. Terasoma, Integrals of logarithmic forms on semi-algebraic sets and a generalized Cauchy formula, Part I: convergence theorems, preprint (2015), arXiv:1509.06950.
- [8] M. Kontsevich and Y. Soibelman, Deformations of algebras over operads and the Deligne conjecture, In Conference Moshé Flato 1999, Vol. I (Dijon), Math. Phys. Stud. **21**, Kluwer Acad. Publ., (2000), 255–307.
- [9] S. Lojasiewicz, Triangulation of semi-analytic sets, Ann. Scu. Norm. Sup. Pisa **18** (1964), 449–474.
- [10] T. L. Loi, Verdier and strict Thom stratifications in o-minimal structures, Illinois J. Math. **42** (1998), 347–356.
- [11] J. Maříková and M. Shiota, Measuring definable sets in o-minimal fields, Israel Journal of Math. **209** (2015), 687–714.
- [12] C. Murolo and D. Trotman, Horizontally- C^1 controlled stratified maps and Thom's first isotopy theorem, Comptes Rendus Acad. Sci. Ser. I - Math. **330** (8) (2000), 707–712.
- [13] M. Shiota, *Geometry of subanalytic and semialgebraic sets*, Progress in Math. **150**, Birkhäuser, 1997.
- [14] T. Suwa, Indices of vector fields and residues of singular holomorphic foliations, Actualités Mathématiques, Hermann, Paris (1998).
- [15] G. Valette, Stokes' formula for stratified forms, Ann. Polon. Math. **114** (2015), 197–206.

(T. Ohmoto) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN

E-mail address: ohmoto@math.sci.hokudai.ac.jp

(M. Shiota) GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, NAGOYA, 464-8602, JAPAN

E-mail address: shiota@math.nagoya-u.ac.jp