Title	Volume of high-risk intratumoral subregions at multi-parametric MR imaging predicts overall survival and complements molecular analysis of glioblastoma
Author(s)	Cui, Yi; Ren, Shangjie; Tha, Khin Khin; Wu, Jia; Shirato, Hiroki; Li, Ruijiang
Citation	European Radiology, 27(9), 3583-3592 https://doi.org/10.1007/s00330-017-4751-x
Issue Date	2017-09
Doc URL	http://hdl.handle.net/2115/71443
Rights	This is a post-peer-review, pre-copyedit version of an article published in European radiology. The final authenticated version is available online at: http://dx.doi.org/10.1007/s00330-017-4751-x
Type	article (author version)
File Information	EU_RADIO_CUI.pdf

[Instructions for use](#)

Manuscript title

Volume of high-risk intratumoral subregions at multi-parametric MR imaging predicts overall survival and complements molecular analysis of glioblastoma

Authors

Yi **Cui**, PhD^{1,2}, Shangjie **Ren**, PhD⁴, Khin Khin **Tha**, MD PhD^{2,3}, Jia **Wu**, PhD¹, Hiroki **Shirato** MD PhD^{2,3}, Ruijiang **Li**, PhD^{1,2}

Affiliations

1. Department of Radiation Oncology, Stanford University, USA
2. Global Station for Quantum Medical Science and Engineering, Global Institution for Collaborative Research and Education, Hokkaido University, Japan
3. Department of Radiology and Nuclear Medicine, Hokkaido University, Japan
4. School of Electrical Engineering and Automation, Tianjin University, China

Corresponding Author

Yi Cui

Department of Radiation Oncology, Stanford University

1070 Arastradero Rd., Palo Alto, CA 94304, USA

Tel: +1 (650) 724-9383

Email: cuiyi@stanford.edu

Volume of high-risk intratumoral subregions at multi-parametric MR imaging predicts overall survival and complements molecular analysis of glioblastoma

Abstract

Objective

To develop and validate a volume-based, quantitative imaging marker by integrating multi-parametric MR images for predicting glioblastoma survival, and to investigate its relations and synergy with molecular characteristics.

Materials and Methods

We retrospectively analyzed 108 patients with primary glioblastoma. The discovery cohort consisted of 62 patients from the cancer genome atlas (TCGA). Another 46 patients combining 30 from TCGA and 16 internally were used for independent validation. Based on integrated analyses of T1-weighted contrast-enhanced (T1-c) and diffusion-weighted MR images, we identified an intratumoral subregion with both high T1-c and low ADC, and accordingly defined a high-risk volume (HRV). We evaluated its prognostic value and biological significance with genomic data.

Results

On both discovery and validation cohorts, HRV predicted overall survival (OS) (concordance index: 0.642 and 0.653, $P<0.001$ and $P=0.038$, respectively). HRV stratified patients within the proneural molecular subtype (log-rank $P=0.040$, hazard ratio=2.787). We observed different OS among patients depending on their MGMT methylation status and HRV (log-rank $P=0.011$). Patients with unmethylated MGMT and high HRV had significantly shorter survival (median survival: 9.3 versus 18.4 months, log-rank $P=0.002$).

Conclusion

Volume of the high-risk intratumoral subregion identified on multi-parametric MRI predicts glioblastoma survival, and may provide complementary value to genomic information.

Keywords

Multi-parametric MRI, glioblastoma multiforme, high-risk tumor volume, overall survival, radiogenomics

Key Points

1. High-risk volume (HRV) defined on multi-parametric MRI predicted GBM survival.
2. The proneural molecular subtype tended to harbor smaller HRV than other subtypes.
3. Patients with unmethylated MGMT and high HRV had significantly shorter survival.
4. HRV complements genomic information in predicting GBM survival

Introduction

Glioblastoma (GBM) is the most deadly primary brain tumor in adults, with a median survival of 12-15 months despite aggressive treatment [1]. GBM is also a biologically heterogeneous disease, where four subtypes, i.e., the proneural, neural, classical, and mesenchymal subtypes, have been proposed based on molecular characteristics of the tumor [2].

Compared with the molecular approach, imaging provides a unique opportunity to noninvasively interrogate the anatomical and functional properties of the entire tumor. Given the routine use of imaging in GBM management, reliable imaging-based biomarkers would have tremendous value in precision medicine, by stratifying patients to guide individualized therapy.

There has been a significant interest in predicting survival of GBM patients based on multi-parametric magnetic resonance (MR) imaging that incorporates perfusion-weighted imaging [3-5], or diffusion-weighted (DW) imaging [6-12]. Most previous studies defined imaging prognosticators as a single point on the signal intensity histogram, e.g. maximum cerebral blood volume (CBV) [3], minimum apparent diffusion coefficient (ADC) [13], or simple quantiles [6]. On the other hand, volume-based imaging metrics that incorporate both intensity and volumetric information, may be more reliable indicators of tumor burden [14-16]. Furthermore, given the heterogeneous nature of GBM, detecting “high-risk” intratumoral subregions could potentially identify biological relevant, aggressive subclones within a tumor [17], and has therapeutic implications for intensified local therapy to improve survival [18].

Recent preliminary studies have shown promising results for predicting survival of GBM patients based on analysis of intratumoral subregions, using conventional T1-weighted contrast-enhanced (T1-c) and T2-weighted fluid-attenuated inversion recovery (FLAIR) MR images [19; 20]. However, a method to explicitly identify clinically relevant, high-risk tumor volume with robust and meaningful cutoffs has been lacking. Current MR imaging markers based on simple, predefined cutoffs such as median or quantile may not be optimal. In addition, individual cutoffs at the patient level can be sensitive to variations due to

1 differences in image acquisition protocols in different patients [21; 22]. Consequently, these
2 imaging markers are difficult to compare and reproduce across cohorts in multi-center
3 settings, which is a significant hurdle to their clinical translation.
4
5
6

7 In this study, we hypothesized that the volume of an intratumoral subregion associated with
8 abnormally high signal intensity on T1-weighted contrast-enhanced imaging and abnormally
9 low ADC on DW imaging can quantify the most aggressive disease burden within a tumor,
10 and thus may be a better predictor of prognosis of GBM patients compared with whole-tumor
11 imaging metrics. This is supported by recent studies showing that the tumor-enhancing
12 volume [14] and the volume of low ADC [16] were both prognostic of overall survival in GBM.
13
14 Instead of using predefined cutoffs for individual patients, we propose a novel method to
15 define robust cutoffs applicable to the entire study population and identify high-risk
16 intratumoral subregions by using a data-driven approach. Further, we evaluated the
17 biological significance of our imaging marker by associating with underlying molecular
18 features. Distinct from most previous radiogenomic studies [23-26], we further investigated
19 whether this imaging marker provides complementary value to the genomic counterparts.
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 The purpose of this study is two-fold: 1), to develop and validate a new volume-based,
35 quantitative imaging marker by integrating multi-parametric MR images for predicting
36 survival of GBM patients; 2), to investigate the relations and potential synergy between the
37 proposed imaging marker and underlying molecular characteristics of GBM.
38
39
40
41
42
43
44
45
46

47 **Materials and Methods**

48 *Study Population*

49
50 In this institutional-review-board approved study, a total of 108 patients were retrospectively
51 investigated. The inclusion criteria were: 1) pathologically confirmed diagnosis of GBM, 2)
52 availability of preoperative T1-weighted contrast enhanced (T1-c) and diffusion-weighted
53 (DW) images, and 3) availability of information about overall survival. The exclusion criteria
54
55
56
57
58
59
60
61
62
63
64
65

1 were prior surgery and other treatments. The majority of the study cohort, consisting of 92
2 patients from 1998 to 2011, was retrieved from the Cancer Imaging Archive (TCIA). We
3 initially identified 98 patients from TCIA whereas 6 of them were excluded due to poor image
4 quality (such as motion, metal artifacts, and RF inhomogeneity) as assessed by a
5 neuroradiologist with over 10 years' experience (KKT). Furthermore, we searched patient
6 records from 2004 to 2014 at the local institution using the same inclusion and exclusion
7 criteria and found 31 patients. Fifteen of them were excluded because they did not have
8 echo-planar T2-weighted images with zero diffusion weighting ($b = 0 \text{ sec/mm}^2$), leading to 16
9 additional patients eligible for this study. The median follow-up duration was 10.9 months for
10 the TCIA cohort and 9.6 months for the internal cohort.
11
12
13
14
15
16
17
18
19
20
21
22

23 We randomly split the TCIA cohort into two portions, where the first one containing
24 approximately two-thirds of the patients ($n = 62$) was used as a discovery cohort. The
25 remaining portion ($n = 30$) was combined with internal cohort ($n = 16$) to form the validation
26 cohort ($n = 46$). The overall study design and patient cohorts are illustrated in Fig. 1. Clinical
27 and demographic information of the study population is listed in Table 1.
28
29
30
31
32
33
34
35
36

37 *Image Acquisition*

38 Among the 108 patients, the magnetic fields used to acquire the MR images were 1T ($n = 1$),
39 1.5T ($n = 65$), 3T ($n = 35$), or unknown ($n = 7$). For the T1-c images, the sequence protocols
40 were spin-echo ($n = 77$), gradient echo ($n = 13$), or T1-weighted fluid-attenuated inversion
41 recovery ($n = 18$). The repetition time and echo time ranged respectively from 6 to 3189
42 msec, and from 3 to 20 msec. The intra-slice voxel resolution varied from 0.43 mm to 1.02
43 mm, the slice thickness was between 2.5 mm and 5 mm, and the inter-slice gap was
44 between 0 mm and 2.5 mm. For DW images, the b-values were 1000 sec/mm^2 ($n = 103$),
45 1500 sec/mm^2 ($n = 5$). Echo-planar T2-weighted images with zero diffusion-weighting ($b=0$
46 sec/mm^2) were acquired for all the patients ($n = 108$). The intra-slice voxel resolution, the
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 slice thickness, and the inter-slice gap of the DW images ranged from 0.86 mm to 1.8 mm, 3
2 mm to 7 mm, and 0 mm to 1 mm, respectively.

3 *Image Processing*

4
5
6
7
8
9
10 For each patient, we co-registered the MR images using the extensively validated software,
11 elastix [27]. Specifically, we used the T1-c image as the reference and rigidly transformed
12 and resliced the echo-planar T2-weighted image with zero diffusion weighting. However, for
13 those cases having noticeable geometric distortion by automatic registration, in-house
14 developed MATLAB software was used instead for manual registration. This process started
15 with visually inspecting the T1-c image and the echo-planar T2-weighted image with zero
16 diffusion weighting to identify slice pairs at the same locations. Then within each slice pair
17 we manually selected landmark points at salient anatomical structures (e.g., ventricles) in
18 both modalities and recorded their 3-dimensional coordinates. Given the corresponding
19 coordinates of the landmarks, we calculated an affine transformation function by least-
20 square estimation to register the echo-planar T2-weighted image with zero diffusion
21 weighting to the T1-c image. Finally, elastix was used to register the DW image to the T2-
22 weighted image with zero diffusion weighting with affine transformation in order to correct for
23 the eddy current distortion and motion effects [28]. The ADC maps were reconstructed from
24 the registered images. Any voxel with negative ADC values due to measurement noise was
25 set to zero and then imputed from its neighborhood.

26 *Image Normalization*

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 estimate the probability density function (PDF) underlying the observed data. It does not
2 assume a parametric form for the PDF to be estimated and therefore particularly suitable for
3 characterizing irregular (non-Gaussian) distributions. Specifically, for each T1-c or ADC
4 image, we applied the KDE to estimate the continuous PDF of the intensities for voxels
5 within the entire brain parenchyma. We then normalized the image by dividing each voxel
6 with the mode of this PDF (Fig. 2). The rationale for this approach is that the mode
7 represents the most frequently occurring voxel value in the image, which comes from the
8 normal-appearing white matter that constitutes the majority of brain tissues. For PDF
9 estimation we used the MATLAB code which used the Gaussian kernel and was able to
10 automatically choose the kernel bandwidth based on the “plug-in” KDE algorithm [29]. In
11 order to avoid estimating the background noise, we set the interval on which the density
12 estimate was constructed to be $[\max/50, \max]$ where $\max$ denoted the maximum intensity of
13 the image and used 10^8 meshes to discretize this interval. It should be noted that although
14 some studies reported similar ADC values among scanners when identical image acquisition
15 parameters were used [30-32], the imaging parameters in our study were not consistent.
16 Therefore, in order to minimize the effects of inter-scanner variations, we chose to normalize
17 the ADC maps. ADC normalization was also performed in other studies [33], where relative
18 ADC maps were used to correlate with genetic and cellular GBM features.

43 *High-Risk Volume Identification*

44 The gross tumor volume(GTV), including both the contrast-enhanced area as well as the
45 bounded non-enhancing and necrotic regions, was segmented semi-automatically on T1-c
46 images using MIPAV [34]. This process used the built-in level-set algorithm of MIPAV and
47 entailed the operator moving the cursor around the boundary of the tumor which was then
48 automatically captured. Manual correction was performed in 8 cases where automatic
49 segmentation failed. In order to assess the reproducibility of the segmentation as well as its
50 impact on subsequent analyses, all tumors were independently delineated by two observers

(SJR and YC). Dice and intra-class correlation coefficients were calculated for the two tumor segmentations.

After segmentation, we defined the high-risk volume (HRV) as the volume of the intratumoral subregion with both higher T1-c intensity and lower ADC. In order to obtain robust and meaningful thresholds to identify the HRV, we pooled the tumor voxels of all the patients in the discovery cohort, for the normalized T1-c and ADC images respectively. For each sequence we used the KDE to estimate their pooled PDFs and used the mode values as global cutoffs, hereafter denoted as t_1 and t_2 . The rationale for the use of mode is that statistically speaking, it is the most typical value of a certain population: any value above or below this value may be considered abnormally high or low. The HRV of an individual patient was defined as the volume of the tumor satisfying $T1-c > t_1$ and $ADC < t_2$. Fig. 2 illustrates the hierarchical flowchart for HRV identification. Once identified, the same thresholds were used to define the HRV for patients in the validation cohort. We examined whether HRV predicted overall survival (OS) in both discovery and validation cohorts. In addition to HRV, we computed the enhancing tumor volume (ETV), which we defined as the volume of the tumor satisfying $T1-c > 1$. The OS prediction performance of ETV was also evaluated.

Relations between high-risk volume and molecular features

We investigated the associations between the proposed imaging marker (HRV) and four molecular subtypes of GBM, i.e., the proneural, neural, classical, and mesenchymal subtypes, which was obtained for 88 patients in the TCIA cohort from the UCSC Cancer Genomics Browser [35; 36]. Given the established role of *MGMT* methylation status for prognosis, we evaluated whether imaging-based HRV could provide complementary value in predicting survival. Information about *MGMT* methylation status was obtained for 61 patients in the TCIA cohort from a previous study [2]. Finally, we evaluated the relationships between

HRV and the mutation status of 9 genes that are known to have important functions in GBM, including *TP53*, *RB1*, *IDH1*, *PIK3R1*, *PTEN*, *PDGFRA*, *NF1*, *EGFR*, and *PIK3CA* [2].

Statistical analysis

Survival prediction performances of the HRV were assessed by the concordance index (CI) [37] and Cox regression analysis. Survival differences among two or more patient groups were compared by the log-rank test as well as Kaplan-Meier analysis. One-way ANOVA was performed to analyze the correlation between a continuous variable (e.g., the HRV) and a nominal variable (e.g., molecular subtype). P values smaller than 0.05 were considered significant. All statistical analyses were done in the open-source statistical computing environment R.

Results

HRV predicted overall survival, independent of clinical factors and other imaging metrics in the overall cohort

In the discovery cohort, we determined the thresholds for defining the HRV to be $t_1=1.429$, $t_2=1.321$ for the normalized T1-c and ADC intensities, respectively (Fig. 2). Using this definition, HRV achieved a CI score of 0.642, and was significantly correlated with OS on univariate Cox regression analysis ($P<0.001$). HRV remained as a significant predictor of OS ($P<0.001$) when adjusted for clinical variables including age, Karnofsky performance status (KPS), eloquent brain involvement (EBI, encoded as a binary variable), and conventional imaging metrics including ETV and minimum ADC (Table 2). An optimal cutoff of 5.12 cm³ for HRV stratified the discovery cohort in terms of OS (Fig. 3A, log-rank $P=0.009$, hazard ratio=2.413).

1 In the validation cohort, HRV achieved a similar CI of 0.653, and was again significantly
2 correlated with OS ($P=0.038$). Further, using the same cutoff derived from the discovery
3 cohort, we stratified the validation cohort into short and long-survival groups, with a median
4 survival of 9.3 months 13.7 months (Fig. 3B, log-rank $P=0.009$, hazard ratio=2.718).
5
6 However, in the validation cohort alone, HRV was not significant ($P=0.20$) in multivariate
7 analysis, nor was any of the other analyzed risk factors ($P=0.13-0.74$), possibly due to the
8 smaller size of this cohort. Therefore, in order to increase the statistical power, we combined
9 the discovery and validation cohorts together and performed multivariate analysis on this
10 overall cohort again. The result showed that HRV was indeed a significant OS predictor in
11 the overall cohort, independent of clinical factors and other imaging metrics.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 *Reproducibility of tumor segmentation and definition of HRV*

26
27 The two independently delineated tumor volumes showed mostly high inter-observer
28 agreement, with the Dice indices ranging from 0.623 to 0.985 (median: 0.948). Importantly,
29 the HRVs computed from the two tumor segmentations were highly concordant
30 (Supplemental Material Fig. S1), with an intra-class correlation of 0.994 (95% confidence
31 interval = [0.990, 0.994], $P<1E-16$).
32
33
34
35
36
37
38
39
40
41
42

43 *HRV stratified patients within the proneural molecular subtype*

44
45 One-way ANOVA showed that HRV was not significantly correlated with the four GBM
46 subtypes ($P=0.1124$). However, we found that the tumors of the proneural subtype tended to
47 have the smallest HRV, which was confirmed by pair-wise comparison with the other three
48 subtypes (Fig. 4, $P=0.014-0.038$). Furthermore, HRV was significantly associated with OS
49 within the proneural group (CI=0.696, $P=0.003$). Using the median (2.29 cm^3) as a cutoff,
50 HRV stratified patients with proneural tumors into short and long-survival groups, with a
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 median survival of 6.4 months versus 12.3 months (Fig. 5A, log-rank $P=0.040$, hazard
2 ratio=2.787). HRV did not stratify patients among other molecular subtypes.
3
4
5
6

7 *HRV provided complementary information to MGMT methylation*

8
9

10 HRV was not correlated with MGMT methylation status ($P=0.1746$). Survival stratification
11 based on MGMT methylation status alone trended toward significance (log-rank $P=0.072$,
12 hazard ratio=1.762). However, by combining methylation status and HRV (using median as
13 the cutoff), we observed significantly different OS among the four groups (Fig. 5B, log-rank
14 $P=0.011$), i.e., methylated MGMT and low HRV ($n = 7$), methylated MGMT and high HRV (n
15 = 6), unmethylated MGMT and low HRV ($n = 21$), and unmethylated MGMT and high HRV (n
16 = 28). Of note, patients with unmethylated MGMT and high HRV had much shorter survival
17 compared with the others (median survival: 9.3 versus 18.4 months, log-rank $P=0.002$).
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 *Higher HRV was associated with NF1 and PIK3CA mutation*

33
34

35 HRV was significantly different between mutated and wide-type groups for *NF1* ($P=0.049$)
36 and *PIK3CA* ($P=0.028$). Tumors with mutation in either *NF1* or *PIK3CA* had higher HRV than
37 those of the wild type (Supplemental Material Fig. S2).
38
39
40
41
42
43
44

45 **Discussion**

46
47

48 In this study, we identified high-risk intratumoral subregions by using a data-driven approach
49 and defined a volume-based imaging marker by integrating multi-parametric MR images of
50 GBM. We found that HRV was prognostic for overall survival in two independent multi-
51 institutional cohorts, and remained a significant predictor after adjusting for clinical factors
52 and other imaging metrics. The predictive accuracy of HRV was higher than gross tumor
53 volume, suggesting that analysis of intratumoral subregions may afford more reliable
54
55
56
57
58
59
60
61
62
63
64
65

1 indicators of tumor burden compared with the whole tumor. HRV was also superior to
2 minimum ADC, which points to the benefits of volume-based metrics versus conventional
3 single-voxel approaches.
4
5

6
7 We used the kernel density estimation and mode approach under two scenarios: to obtain a
8 patient-specific background voxel value for image normalization, and to find a population-
9 level cutoff for defining the high-risk volume. In both scenarios, this approach has important
10 advantages in that the mode value is a more robust statistic of the pooled distribution
11 compared with other commonly used summary statistics (e.g., mean, median, or quantile
12 values) [3; 6; 13], which may be sensitive to variations in tumor segmentation [38; 39].
13
14

15 Compared with previous approaches that require manual selection of a region of interest for
16 image normalization [33], ours is fully automated and more robust.
17
18

19 Recent studies have used a quantitative radiomic approach to obtain comprehensive tumor
20 phenotypes such as shape and texture. While further validation is warranted, this approach
21 has showed promising results in identifying prognostic imaging markers in GBM and appears
22 to provide additional information beyond simple volume-based imaging metrics [20; 23; 40;
23 41]. Previously this approach has mostly been applied to the primary tumor to extract whole-
24 tumor aggregate characteristics. It would be interesting to apply radiomics to the high-risk
25 intratumoral subregions extracted in this work to derive further improvement in prognostic
26 value [19].
27
28

29 Our radiogenomic analysis revealed that the proposed imaging marker (HRV) was
30 associated with several important molecular features of GBM. We showed that HRV was
31 associated with overall survival and further stratified the proneural group. Compared with
32 other molecular subtypes, the proneural group tended to harbor smaller HRV, which was
33 correlated with longer survival. This is consistent with previous studies showing that the
34 proneural molecular subtype had a better prognosis than others [2].
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

We showed that higher HRV was associated with mutations in NF1 and PIK3CA, which are key genetic events driving the progression of GBM [2]. *NF1* is a tumor-suppressor gene and frequently inactivated in GBM [2; 42]. It has been shown that NF1 mutation is highly enriched in the mesenchymal molecular subtype, a known aggressive GBM subtype with poor outcomes [2]. The *PI3K* signaling pathway is frequently dysregulated in GBM, and plays a critical role in proliferation, cellular metabolism, and apoptosis [43]. Therapeutic agents inhibiting PI3K activity are under active development and have the potential for improvement in clinical outcome for GBM [44]. *IDH1* mutation has been shown to be an independent prognostic factor in patients diagnosed with glioma including GBM [45], but appeared to be not associated with HRV in our study (Fig. S2), suggesting that they may be driven by differing biological processes.

Importantly, we showed that HRV provided complementary information to *MGMT* methylation status for survival prediction. Patients with unmethylated *MGMT* and high HRV had much shorter survival compared with other), while *MGMT* methylation status alone was not prognostic within our study cohort. This is consistent with a recent study [6] showing worse prognosis for patients with unmethylated *MGMT* and lower mean ADC. Taken together, these data support that the imaging-based HRV recapitulates tumor biology of GBM and potentially could provide additional prognostic information beyond genomic analysis.

Limitations of our study include the retrospective design and relatively small validation cohorts. The image data came from multiple institutions and were acquired with different imaging protocols and parameters, which might have influenced the image quantification. Nevertheless, we used careful image standardization techniques and robust image analysis to minimize the potential biases. Our findings warrant further validation in larger prospective cohorts. Intra-tumor genetic heterogeneity in GBM [46; 47] may confound the radiogenomic analyses. Because imaging has the unique capability of sampling the entire tumor and surrounding tissue, it would be intriguing to prospectively test the idea that combines image-

guided stereotactic biopsy [48; 49] and the proposed method to identify high-risk intratumoral subregions, which might increase the likelihood of detecting the most aggressive part of a tumor.

Future studies would benefit from the incorporation of additional imaging modalities such as T2-weighted FLAIR and perfusion-weighted imaging for more comprehensive characterization of GBM such as surrounding edema/invasion [26] and blood volume/flow [3]. We also plan to test the ability of HRV to evaluate treatment response of GBM, in particular, to distinguish progression from pseudo-progression after chemoradiation therapy [50; 51].

In conclusion, volume of the high-risk intratumoral subregion on multi-parametric MRI predicts overall survival in GBM patients, and may provide complementary value to genomic information. We envision that the same approach could be applied to identify clinically and biologically relevant imaging markers in other cancer types.

Compliance with ethical standards:

Guarantor:

The scientific guarantor of this publication is Dr. Ruijiang Li.

Conflict of interest:

The authors of this manuscript declare no relationships with any companies, whose products or services may be related to the subject matter of the article.

Funding:

This research was partially funded by the NIH (grant number: R01 CA193730), and partially supported by the Global Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University, founded by the Ministry of Education, Culture, Sports, Science and

Technology MEXT, Japan. Part of the data used in this research was obtained from The Cancer Imaging Archive (TCIA) sponsored by the Cancer Imaging Program, DCTD/NCI/NIH.

Statistics and biometry:

One of the authors has significant statistical expertise.

Ethical approval:

Institutional Review Board approval was obtained.

Informed consent:

Written informed consent was waived by the Institutional Review Board.

Methodology:

- retrospective
- diagnostic or prognostic study
- multicenter study

References

- 1 Ostrom QT, Gittleman H, Fulop J et al (2015) CBTRUS Statistical Report: Primary Brain and Central Nervous System Tumors Diagnosed in the United States in 2008-2012. *Neuro Oncol* 17 Suppl 4:iv1-iv62
- 2 Verhaak RG, Hoadley KA, Purdom E et al (2010) Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer Cell* 17:98-110
- 3 Jain R, Poisson L, Narang J et al (2013) Genomic mapping and survival prediction in glioblastoma: molecular subclassification strengthened by hemodynamic imaging biomarkers. *Radiology* 267:212-220
- 4 Burth S, Kickingeder P, Eidel O et al (2016) Clinical parameters outweigh diffusion- and perfusion-derived MRI parameters in predicting survival in newly diagnosed glioblastoma. *Neuro Oncol*. 10.1093/neuonc/nov122
- 5 Schmainda KM, Zhang Z, Prah M et al (2015) Dynamic susceptibility contrast MRI measures of relative cerebral blood volume as a prognostic marker for overall survival in recurrent glioblastoma: results from the ACRIN 6677/RTOG 0625 multicenter trial. *Neuro Oncol* 17:1148-1156
- 6 Choi YS, Ahn SS, Kim DW et al (2016) Incremental Prognostic Value of ADC Histogram Analysis over MGMT Promoter Methylation Status in Patients with Glioblastoma. *Radiology*. 10.1148/radiol.2016151913:151913
- 7 Gupta A, Prager A, Young RJ, Shi W, Omuro AM, Graber JJ (2013) Diffusion-weighted MR imaging and MGMT methylation status in glioblastoma: a reappraisal of the role of preoperative quantitative ADC measurements. *AJNR Am J Neuroradiol* 34:E10-11
- 8 Moon WJ, Choi JW, Roh HG, Lim SD, Koh YC (2012) Imaging parameters of high grade gliomas in relation to the MGMT promoter methylation status: the CT, diffusion tensor imaging, and perfusion MR imaging. *Neuroradiology* 54:555-563

- 9 Pope WB, Lai A, Mehta R et al (2011) Apparent Diffusion Coefficient Histogram Analysis
Stratifies Progression-Free Survival in Newly Diagnosed Bevacizumab-Treated Glioblastoma.
American Journal of Neuroradiology 32:882-889
- 10 Romano A, Calabria LF, Tavanti F et al (2013) Apparent diffusion coefficient obtained by
magnetic resonance imaging as a prognostic marker in glioblastomas: correlation with
MGMT promoter methylation status. European Radiology 23:513-520
- 11 Saksena S, Jain R, Narang J et al (2010) Predicting Survival in Glioblastomas Using Diffusion
Tensor Imaging Metrics. Journal of Magnetic Resonance Imaging 32:788-795
- 12 Sunwoo L, Choi SH, Park CK et al (2013) Correlation of apparent diffusion coefficient values
measured by diffusion MRI and MGMT promoter methylation semiquantitatively analyzed
with MS-MLPA in patients with glioblastoma multiforme. Journal of Magnetic Resonance
Imaging 37:351-358
- 13 Higano S, Yun X, Kumabe T et al (2006) Malignant astrocytic tumors: Clinical importance of
apparent diffusion coefficient in prediction of grade and prognosis. Radiology 241:839-846
- 14 Wangaryattawanich P, Hatami M, Wang J et al (2015) Multicenter imaging outcomes study
of The Cancer Genome Atlas glioblastoma patient cohort: imaging predictors of overall and
progression-free survival. Neuro Oncol 17:1525-1537
- 15 Ellingson BM, Harris RJ, Woodworth DC et al (2016) Baseline pretreatment contrast
enhancing tumor volume including central necrosis is a prognostic factor in recurrent
glioblastoma: evidence from single- and multicenter trials. Neuro Oncol.
10.1093/neuonc/now187
- 16 Zhang M, Gulotta B, Thomas A et al (2016) Large-volume low apparent diffusion coefficient
lesions predict poor survival in bevacizumab-treated glioblastoma patients. Neuro Oncol
18:735-743
- 17 Gatenby RA, Grove O, Gillies RJ (2013) Quantitative imaging in cancer evolution and ecology.
Radiology 269:8-15

- 18 Ling CC, Humm J, Larson S et al (2000) Towards multidimensional radiotherapy (MD-CRT):
biological imaging and biological conformality. *Int J Radiat Oncol Biol Phys* 47:551-560
- 19 Cui Y, Tha KK, Terasaka S et al (2016) Prognostic Imaging Biomarkers in Glioblastoma:
Development and Independent Validation on the Basis of Multiregion and Quantitative
Analysis of MR Images. *Radiology* 278:546-553
- 20 Chang K, Zhang B, Guo X et al (2016) Multimodal imaging patterns predict survival in
recurrent glioblastoma patients treated with bevacizumab. *Neuro Oncol.*
10.1093/neuonc/now086
- 21 Chenevert TL, Malyarenko DI, Newitt D et al (2014) Errors in Quantitative Image Analysis due
to Platform-Dependent Image Scaling (vol 7, pg 65, 2014). *Translational Oncology* 7:523-523
- 22 Ellingson BM, Lai A, Nguyen HN, Nghiemphu PL, Pope WB, Cloughesy TF (2015)
Quantification of Nonenhancing Tumor Burden in Gliomas Using Effective T-2 Maps Derived
from Dual-Echo Turbo Spin-Echo MRI. *Clinical Cancer Research* 21:4373-4383
- 23 Gevaert O, Mitchell LA, Achrol AS et al (2014) Glioblastoma multiforme: exploratory
radiogenomic analysis by using quantitative image features. *Radiology* 273:168-174
- 24 Gutman DA, Cooper LA, Hwang SN et al (2013) MR imaging predictors of molecular profile
and survival: multi-institutional study of the TCGA glioblastoma data set. *Radiology* 267:560-
569
- 25 Jamshidi N, Diehn M, Bredel M, Kuo MD (2014) Illuminating Radiogenomic Characteristics of
Glioblastoma Multiforme through Integration of MR Imaging, Messenger RNA Expression,
and DNA Copy Number Variation. *Radiology* 270:212-222
- 26 Zinn PO, Mahajan B, Sathyan P et al (2011) Radiogenomic mapping of edema/cellular
invasion MRI-phenotypes in glioblastoma multiforme. *PLoS One* 6:e25451
- 27 Klein S, Staring M, Murphy K, Viergever MA, Pluim JPW (2010) elastix: A Toolbox for
Intensity-Based Medical Image Registration. *Ieee Transactions on Medical Imaging* 29:196-
205

- 28 Mohammadi S, Moller HE, Kugel H, Muller DK, Deppe M (2010) Correcting eddy current and motion effects by affine whole-brain registrations: evaluation of three-dimensional distortions and comparison with slice-wise correction. *Magn Reson Med* 64:1047-1056
- 29 Botev ZI, Grotowski JF, Kroese DP (2010) Kernel Density Estimation Via Diffusion. *Annals of Statistics* 38:2916-2957
- 30 Ogura A, Tamura T, Ozaki M et al (2015) Apparent Diffusion Coefficient Value Is Not Dependent on Magnetic Resonance Systems and Field Strength Under Fixed Imaging Parameters in Brain. *J Comput Assist Tomogr* 39:760-765
- 31 Lemkaddem A, Daducci A, Vullemoz S et al (2012) A multi-center study: intra-scan and inter-scan variability of diffusion spectrum imaging. *Neuroimage* 62:87-94
- 32 Grech-Sollars M, Hales PW, Miyazaki K et al (2015) Multi-centre reproducibility of diffusion MRI parameters for clinical sequences in the brain. *NMR Biomed* 28:468-485
- 33 Barajas RF, Hodgson JG, Chang JS et al (2010) Glioblastoma Multiforme Regional Genetic and Cellular Expression Patterns: Influence on Anatomic and Physiologic MR Imaging. *Radiology* 254:564-576
- 34 McAuliffe MJ, Lalonde FM, McGarry D, Gandler W, Csaky K, Trus BL (2001) Medical Image Processing, Analysis & Visualization in clinical research. Fourteenth IEEE Symposium on Computer-Based Medical Systems, Proceedings:381-386
- 35 Goldman M, Craft B, Swatloski T et al (2015) The UCSC Cancer Genomics Browser: update 2015. *Nucleic Acids Res* 43:D812-817
- 36 Zhu J, Sanborn JZ, Benz S et al (2009) The UCSC Cancer Genomics Browser. *Nat Methods* 6:239-240
- 37 Harrell FE (2001) Regression modeling strategies : with applications to linear models, logistic regression, and survival analysis. Springer, New York ; London
- 38 Leijenaar RT, Carvalho S, Velazquez ER et al (2013) Stability of FDG-PET Radiomics features: an integrated analysis of test-retest and inter-observer variability. *Acta Oncol* 52:1391-1397

- 39 Parmar C, Rios Velazquez E, Leijenaar R et al (2014) Robust Radiomics feature quantification using semiautomatic volumetric segmentation. PLoS One 9:e102107
- 40 Macyszyn L, Akbari H, Pisapia JM et al (2016) Imaging patterns predict patient survival and molecular subtype in glioblastoma via machine learning techniques. Neuro Oncol 18:417-425
- 41 Kickingeder P, Bonekamp D, Nowosielski M et al (2016) Radiogenomics of Glioblastoma: Machine Learning-based Classification of Molecular Characteristics by Using Multiparametric and Multiregional MR Imaging Features. Radiology. 10.1148/radiol.2016161382:161382
- 42 Cancer Genome Atlas Research N (2008) Comprehensive genomic characterization defines human glioblastoma genes and core pathways. Nature 455:1061-1068
- 43 Engelman JA (2009) Targeting PI3K signalling in cancer: opportunities, challenges and limitations. Nat Rev Cancer 9:550-562
- 44 Wen PY, Lee EQ, Reardon DA, Ligon KL, Alfred Yung WK (2012) Current clinical development of PI3K pathway inhibitors in glioblastoma. Neuro Oncol 14:819-829
- 45 Sanson M, Marie Y, Paris S et al (2009) Isocitrate dehydrogenase 1 codon 132 mutation is an important prognostic biomarker in gliomas. J Clin Oncol 27:4150-4154
- 46 Patel AP, Tirosh I, Trombetta JJ et al (2014) Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. Science 344:1396-1401
- 47 Sottoriva A, Spiteri I, Piccirillo SG et al (2013) Intratumor heterogeneity in human glioblastoma reflects cancer evolutionary dynamics. Proc Natl Acad Sci U S A 110:4009-4014
- 48 Barajas RF, Jr., Phillips JJ, Parvataneni R et al (2012) Regional variation in histopathologic features of tumor specimens from treatment-naïve glioblastoma correlates with anatomic and physiologic MR Imaging. Neuro Oncol 14:942-954
- 49 Hu LS, Ning S, Eschbacher JM et al (2016) Radiogenomics to characterize regional genetic heterogeneity in glioblastoma. Neuro Oncol. 10.1093/neuonc/now135

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
- Chu HH, Choi SH, Ryoo I et al (2013) Differentiation of true progression from pseudoprogression in glioblastoma treated with radiation therapy and concomitant temozolomide: comparison study of standard and high-b-value diffusion-weighted imaging. Radiology 269:831-840
- Park JE, Kim HS, Goh MJ, Kim SJ, Kim JH (2015) Pseudoprogression in Patients with Glioblastoma: Assessment by Using Volume-weighted Voxel-based Multiparametric Clustering of MR Imaging Data in an Independent Test Set. Radiology 275:792-802

Figure captions

Figure 1: Flow-chart of the proposed study design. The TCIA cohort was randomly split into the discovery cohort and a spin-off cohort which was further combined with an internal cohort to construct the validation cohort. The proposed imaging marker was developed on the discovery cohort and its performance for OS prediction was also evaluated on the validation cohort. The subtype, mutation, and methylation data associated with the TCIA cohort were used to correlate with the proposed imaging marker to show its biological relevance and complementarity to molecular-level information.

Figure 2: Diagram of the procedure of high-risk volume identification. In Step 1, the T1-c and ADC images for each patient were respectively normalized using the mode values of the intensity PDF estimates. In Step 2, the normalized intensities of the pixels within the segmented tumors (shaded in red) of all patients were pooled and two thresholds (t_1 , t_2) were respectively obtained as the mode values of the pooled intensity PDF estimates. In Step 3, HRV for each patient was defined as the volume of the tumor satisfying $T1-c > t_1$ and $ADC < t_2$.

Figure 3: Kaplan-Meier survival estimates for the discovery (A) and the validation (B) cohorts using HRV.

Figure 4: Boxplot shows that HRV was the smallest in the proneural subtype among the four GBM molecular subtypes. The P-value is for one-way ANOVA.

Figure 5: Radiogenomic analysis of HRV. (A) Kaplan-Meier survival estimates for proneural patients using HRV. (A) Kaplan-Meier survival estimates based on both methylation status and HRV.

Table 1: Demographic and clinical information of study population.

	Overall Cohort	TCIA Cohort	Local Cohort	Discovery Cohort	Validation Cohort
Number†	108 (23)	92 (15)	16 (7)	62 (9)	46 (14)
Age‡ (years)	58.1 ± 14.8	57.9 ± 14.5	58.9 ± 16.9	57.7 ± 14.1	58.5 ± 15.8
Karnofsky Performance Scale††	80 (60–100)*	80 (60–100)*	N/A	80 (60–100)*	80 (60–100)*
Overall Survival†† (days)	357 (16–1757)	362 (16–1757)	288 (37–713)	357 (16–1757)	426 (34–1561)
Gender					
Male	63	56	7	43	20
Female	45	36	9	19	26
Molecular Subtypes					
Classical	17	17	0	10	7
Proneural	24	24	0	16	8
Neural	21	21	0	14	7
Mesenchymal	26	26	0	18	8
Unknown	20	4	16	4	16
Methylation Status					
Methylated	12	12	0	6	6
Unmethylated	49	49	0	33	16
Unknown	47	31	16	23	24

Note. –Unless otherwise indicated, data are patient numbers.

† Data in parenthesis are censored patient numbers.

‡ Data are mean ± standard deviation

†† Data are median (range).

* Missing data were imputed with median value.

Table 2: Prognostic performances of high-risk volume in comparison with clinical and baseline indicators.

Cohort	Risk factor	Concordance Index	Univariate Analysis		Multivariate Analysis	
			P-Value	Hazard Ratio	P-Value	Hazard Ratio
Discovery	HRV	0.64 (0.45, 0.80)	3.9E-4***	1.63 (1.25, 2.14)	4.78E-3**	1.83 (1.20, 2.78)
	Age	0.64 (0.45, 0.80)	3.4E-3**	1.71 (1.20, 2.45)	2.15E-3**	1.77 (1.23, 2.56)
	KPS	0.37 (0.19, 0.59)	0.20	0.83 (0.63, 1.10)	0.73	0.95 (0.71, 1.27)
	ETV	0.60 (0.41, 0.76)	0.09	1.25 (0.96, 1.61)	0.98	1.01 (0.70, 1.43)
	EBI	0.52 (0.28, 0.76)	0.55	0.84 (0.48, 1.48)	0.28	0.71 (0.39, 1.31)
	Minimum ADC	0.43 (0.26, 0.62)	0.12	0.80 (0.61, 1.06)	0.94	1.01 (0.71, 1.44)
Validation	HRV	0.65 (0.42, 0.83)	0.04*	1.39 (1.01, 1.92)	0.20	1.97 (0.70, 5.56)
	Age	0.58 (0.35, 0.78)	0.23	1.32 (0.84, 2.09)	0.70	1.11 (0.66, 1.84)
	KPS	0.50 (0.25, 0.75)	0.38	0.86 (0.60, 1.21)	0.74	1.08 (0.69, 1.69)
	ETV	0.57 (0.34, 0.77)	0.20	1.23 (0.90, 1.68)	0.44	0.70 (0.28, 1.74)
	EBI	0.76 (0.43, 0.93)	0.02*	2.42 (1.16, 5.04)	0.13	2.06 (0.80, 5.30)
	Minimum ADC	0.53 (0.31, 0.74)	0.84	1.04 (0.70, 1.55)	0.47	1.21 (0.73, 2.00)
Discovery + Validation	HRV	0.64 (0.49, 0.77)	2.6E-4***	1.43 (1.18, 1.73)	0.03*	1.53 (1.04, 2.25)
	Age	0.61 (0.46, 0.74)	4.1E-3**	1.48 (1.13, 1.94)	4.8E-3**	1.50 (1.13, 1.99)
	KPS	0.41 (0.26, 0.58)	0.08	0.82 (0.66, 1.02)	0.23	0.87 (0.69, 1.09)
	ETV	0.59 (0.44, 0.73)	0.03	1.24 (1.02, 1.51)	0.72	0.94 (0.62, 1.35)
	EBI	0.61 (0.40, 0.78)	0.27	1.27 (0.83, 1.95)	0.57	0.87 (0.54, 1.41)
	Minimum ADC	0.46 (0.32, 0.61)	0.16	0.85 (0.68, 1.07)	0.60	0.93 (0.72, 1.21)

Note. –Unless otherwise indicated, data in parenthesis are 95% confidence intervals.

Abbreviations. –HRV: high-risk volume, KPS: Karnofsky performance status, ETV: enhancing tumor volume, ADC: apparent diffusion coefficient, EBI: eloquent brain involvement

* P<0.05

** P<0.005

*** P<0.0005

Step 1: T1-c & ADC image normalization by mode of patient-level distribution

Step 2: Thresholds determination by mode of population-level distribution

Step 3: HRV identification by thresholding T1-c & ADC images