Title	Fungal Decolorization of Microbially Treated Night Soils
Author(s)	HISHINUMA, DAISUKE; MAKABE, AKEMI; TSURUGA, HIDEO; OOI, TOSHIHIKO; KINOSHITA, SHINICHI; HIRAO, TOMOHIKO
Citation	日本水処理生物学会誌, 37(3), 123-128 https://doi.org/10.2521/jswtb.37.123
Issue Date	2001
Doc URL	http://hdl.handle.net/2115/71463
Rights	著作権は学会に帰属する
Type	article
File Information	water 37.3.pdf

[Instructions for use](#)

Fungal Decolorization of Microbially Treated Night Soils

DAISUKE HISHINUMA, AKEMI MAKABE, HIDEO TSURUGA, TOSHIHIKO OOI,
SHINICHI KINOSHITA, and TOMOHIKO HIRAO*

Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University
/Nishi-8, Kita-13, Kita-ku, Sapporo 060-8628, Japan

*Takuma Co. Ltd.,/2-2-23, Kinrakuji-cho, Amagasaki, 660-0806, Japan

Abstract

A fungal strain LM-12 was selected for the decolorization of microbially treated night soils {filtrate(iv) in Fig. 1} and it was identified as *Penicillium janthinellum*. Strain LM-12 decolorized 50% of the color of the medium with 5.0 g/l glucose and 0.6 g/l ammonium sulfate at 30°C for 24 hr. Over 50% decolorization was not attained in any culture conditions tested, but the increase of the mycelial amount enhanced the decolorization rate. So by using repeated replacement culture, the decolorization period was reduced to 2.0 hr. The amount of nutrient addition could be reduced in further replacement culture.

Key words : decolorization of night soils, night soil decolorization, *Penicillium janthinellum*, repeated replacement culture

INTRODUCTION

Recently night soil treatment in villages is carried out in a compact night soil treatment plant. Figure 1 shows a plant in Kamiukena, Matsuyama, and it is composed of following steps: (i) removal of solids such as sands, metals, and plastics by sedimentation and screening; (ii) treatment by aerobic microbes to remove BOD, COD, nitrogenous compounds, and SS (suspended solid); (iii) microbial denitrification; (iv) membrane filtration to remove micro-organisms and SS; (v) addition of chemicals and alkali to promote aggregation of phosphates and colored materials; (vi) membrane filtration to remove aggregates; (vii) decolorization with activated charcoal; (viii) disinfection with hypochlorite addition.

By the operation of this plant some color remains and so the effluent from this system is not discharged without adsorption on active charcoal.

In this study we tried to decolorize the

microbially treated night soil {filtrate (iv) in Fig. 1} for the cancellation or reduction of the use of activated charcoal. We found a fungal strain to decolorize it to half and optimized the reaction period to 2 h by the repeated replacement culture.

MATERIALS AND METHODS

Solution for decolorization The filtrate in step (vi) in Fig. 1 is used for the decolorization to save the amount of active charcoal. The solution after step (viii) can be used, but color of the solution is very faint (less than A_{400} 0.2), which is not easy to distinguish the color removal by naked eyes in screening of microorganisms. Also after decolorization the solution is expected to return to step (iii) in Fig. 1. The solution at step (ii) or (iii) can be used, but the cleared solution is better because most of micro-organisms is removed and we can keep it long during experiments.

The filtrate in step (iv) is obtained from Environmental Sanitation Center, Kami-

Fig. 1 Flow sheet of human night soil treatment system in a compact plant at Kamiukena, Ehime, Japan.

ukena, Matsuyama which is kept in a cold room. The analytical data of the filtrate: BOD, <1 mg/l; COD, 210 mg/l; SS, <10 mg/l; total P, 100 mg/l; total N, 18 mg/l; color unit, 400; pH, 6.9. As most of BOD and nitrogenous compounds are removed in the filtrate, so 5 g/l glucose and 0.6 g/l ammonium sulfate were tentatively added to the filtrate, which was used as a medium. The initial color of the medium is around 0.8 of absorption at 400 nm.

Decolorization experiments Strain LM12 was inoculated into 500-ml conical flask containing 100 ml of the medium from the agar slant and it was incubated at 30°C on a rotary shaker.

In a replacement culture mycelia are harvested by filtration apparatus which was sterilized before, and were suspended into the fresh medium. In this experiment we tried to operate under as aseptic conditions as possible. Also the pH of the preculture was decreased to lower than pH 3.0, which gives more advantage for the replacement culture against microbial contamination.

Measurements The decolorization was determined spectrophotometrically from the absorbance at 400 nm after filtration of the culture. The mycelial weight was determined after collecting the mycelia from the culture by filtration, washing repeatedly with water, and drying at 80°C until a constant weight.

RESULTS AND DISCUSSION

Absorption spectrum of the culture filtrate

The absorption spectrum (320 to 700 nm) of the medium gave a monotonous increasing curve toward shorter wave length (Fig. 2A), even before and after the microbial decolorization, which meant the absorption was composed of many colored materials. The absorption of the medium before and after the decolorization at 400 nm was not affected much by pH 2 to 8 (Fig. 2B), though a little increased in alkaline side. In the experiments we used the pH range between 2 and 6, where the absorption did not depend on pH, and so the color is measured without adjusting pH of the supernatant of medium.

Fig. 2 A Absorption spectra of the medium supernatant containing microbially treated night soil before and after the microbial decolorization. B Effect of pH on the absorption at 400 nm before and after the decolorization of the medium.

Screening of microorganisms for the decolorization

One loop from 300 soil samples was inoculated into a test tube containing 5 ml of the medium, and it was incubated on a reciprocal shaker at 30°C. After one week, the decolorization was judged by naked eyes, and one loop of which was seeded into the fresh medium. It was cultivated for 5 days, and 25 decolorized samples were seeded into 100-ml conical flasks containing 20 ml of fresh medium together with 15 stock cultures in our laboratory. They were incubated for 3 days on a rotary shaker and those absorbance at 400 nm of the supernatant were measured. A culture to show the highest decolorization, strain LM12, was selected, which had been isolated as a lignin decomposer before. It is used in this study.

The decolorizing activity was not located extracellularly, because the culture filtrate supplemented with the fresh medium was not decolorized at all.

The photo of strain LM12 is shown in Fig. 3. Taxonomical identification of strain LM-12 was carried out by NCIMB Japan (Shimizu, Japan) and identified as *Penicillium janthinellum*.(1, 2).

The fundamental properties are as follows: hypha has cepta; conidiophore branches irregularly; double-ring penicilli are composed of blanches, meturoe and fialides; ascus or sclerotium is not formed; colony is whity cotton fiber-like until conidiospore

10 μm

Fig. 3 The photo of strain LM-12.

formation and turns to green, then dark green; the reverse side of agar plate is dark orange; spores is mainly globe, rough in surface, and about 3 μm in diameter; size of colony is about 6 cm in diameter on a potato-dextrose agar plate after 10-day growth. It was quite rare that *P. janthinellum* was isolated as a decolorizing fungus, because white rot fungi such as *Phanerochaete chrysosporium* (3), *Geotrichum candidum* (4), and *Coriolus hirsutus* (5) were usually found, as decolorizers of azo dyes, molasses, and melanoidine, respectively. It was maintained on malt-extract agar medium.

Optimal culture condition The good decolorization was observed at 30–37°C in 2-day culture as shown in Fig. 4, and 30°C was used in further experiment.

The best decolorization was observed at initial pH of 7.0 as shown in Fig. 5, but in further experiments the initial pH of 6.0 was selected because glucose in the medium gave colored material at pH 7.0 after sterilization.

The decolorizations were high at 2.5–10.0 g/l glucose concentrations in the medium, the fastest at 5.0 g/l glucose (Fig. 6), and this concentration was used in further experiments. The strain could not decolorize the medium without the addition of glucose.

The decolorizations were high at 0.6–2.4 g/l ammonium sulfate concentration (Fig. 7), the fastest at 0.6 g/l ammonium sulfate, and this concentration was used in further

Fig. 4 Effect of incubation temperature on the decolorization of the medium using microbially treated night soils by strain LM12 for 2 days.

experiments. Strain LM-12 could not decolorize without the addition of ammonium sulfate.

The time course of the decolorization under the optimal condition was shown in Fig. 8. The growth became active after 12 h and continued to grow until 24 h. The pH began to decrease sharply after 12 h and reached to pH 3.0 at 16 h. The decolorization also started at 12 h, and after 24 h 50% of decolorization was attained.

Decolorization degree After the incubation for 24 h, 5.0 g/l glucose and 0.6 g/l ammonium sulfate were added to the culture and it was further incubated for more 24 h. The cell mass increased, but the

decolorization did not proceed any more (data were not shown).

The culture filtrate was recovered after 24 h, to which glucose and ammonium sulfate at 5.0 and 0.6 g/l, respectively, were added, LM12 was inoculated, and it was incubated for another 24 h. Strain LM-12 could grow as well as before, but the decolorization did not proceed any more (data were not shown).

To check the effect of amount of mycelia on the decolorization, the mycelia were harvested by filtration from one, two, or four flasks of 24-h culture, and they were suspended to fresh media to give different amounts of mycelia (1-4 folds), which were incubated for more 24 h, but the

Fig. 5 Effect of the initial pH on the decolorization of the medium by strain LM12 for 2 days.

Fig. 6 Time courses of the decolorization of the medium by the addition of different concentration of glucose by strain LM12.

Fig. 7 Time courses of the decolorization of the medium by the addition of different concentration of ammonium sulfate by strain LM12.

Fig. 8 The decolorization of the medium under the optimal culture condition.
Symbols: Δ , color; $\circ$, pH; $\bullet$, dry cell weight.

decolorization were about 50%, however, the decolorization periods were reduced to 3.5 h by the replacement culture (1 fold) from 24 h in the previous batch experiments. And also the increases of mycelial amount to 2 and 4 folds reduced the decolorization period to 2.0, and 1.5 h, respectively (Fig. 9). This meant that some part of colored components in the filtrate were resistant to further decolorization by strain LM-12. However, the maximum decolorization rates were 0.09, 0.15, and 0.26 $\Delta A_{400}/h$, with 1, 2, and 4 folds of mycelial amount, respectively, and so decolorization periods were also reduced in the same order. These results meant that the higher amount of mycelia promoted the decolorization rate very much and we should use the higher mycelial amount in the decolorization.

Decolorization by replacement culture

So to obtain higher mycelial amount, the strain was precultured for 48 h with intermittent additions of nutrients (5.0 glucose and 0.6 g/l ammonium sulfate) at 24 and 36 h, respectively. The mycelia were harvested by filtration from the preculture, suspended in the fresh medium, and incubated for 2 h. The decolorization attained to a steady level, then the mycelia were harvested, used for the further rounds of replacement cultures at 2 h-interval as

Fig. 9 Effect of cell mass in the replacement culture on the decolorization. The different amounts of mycelia were prepared by harvesting mycelia from different numbers of flasks precultured for 24 h. Symbols (relative call mass): ○, 1 fold; ■, 2 folds; △, 4 folds.

shown in Fig. 10. This experimental results showed that the replacement culture worked well for the reduction of decolorization period.

In the next experiment 24-h culture was used as a preculture without obtaining higher mycelial amount in the first round of replacement culture. So the culture was replaced when the decolorization attained to a steady level as shown in Fig. 11. The decolorization time in the first culture was 4.5 h, that in the second was 3.0 h, those in the third and fourth was 2.5 h, and those after the fifth were 2.0 h. This 2-h decolorization was attained for at least following 10 replacement cultures without using 48-h preculture. This reaction system would be very convenient for the scale-up, because one replacement culture takes only 2 h and replacement culture could be operated by natural sedimentation of mycelia, removal of supernatant, and addition of the fresh medium without filtration, and mixing of mycelia in the next decolorization process.

Though a constant amount of 5 g/l glucose and 0.6 g/l ammonium sulfate addition into the replacement culture was employed, but such much amounts of nutrient additions may not always be necessary and the amount of nutrient may be reduced in the replacement culture. So the replacement culture was carried out with addition of

Fig. 10 The replacement culture by using higher amounts of mycelia. The higher amounts of mycelia were prepared from the precultures, which was incubated for 48 h with twice additions of the nutrients.

different amounts of glucose and ammonium sulfate. The standard nutrient concentration is composing of 5.0 g/l glucose and 0.6 g/l ammonium sulfate. The first replacement cultures with addition of 10 and 20% of the standard nutrient did not give sufficient decolorization, and that with 40% nutrient addition gave a little less decolorization than that with the standard nutrient addition as shown in Fig. 12. But the second replacement culture with 40% nutrient addition gave the same decolorization as that with the standard nutrient addition. So it may be possible to reduce the addition of nutrient more in prolonged replacement cultures.

Fig. 11 Repeated replacement culture. The seed was precultured for 24-h and each replacement culture was carried out when the decolorization attained at a steady level (50% decolorization).

Fig. 12 Addition of reduced amounts of nutrients in replacement culture.
Symbols (nutrient addition) : ○, standard (100%, 5 g/l glucose and 0.6 g/l ammonium sulfate) ; ●, 40% ; △, 20% ; ■, 10% of the standard.

CONCLUSION

We selected *P. janthinellum* LM12 as a decolorizer of microbially treated night soil {filtrate in step (iv) in Fig. 1} with addition of 5 g/l glucose and 0.6 g/l ammonium sulfate, which decolorized 50% of the filtrate within 2-day culture. We optimized culture conditions for the decolorization, where the reaction period was reduced to 2 h by replacement culture but more than 50% of the decolorization was not attained. In a prolonged replacement culture the amount of glucose and ammonium sulfate addition is suggested to be reduced.

REFERENCES

- 1) Raper, K. B. and Thom, C. : A manual of the *Penicillia*, p. 120-125, p. 299-303, and p. 318-322, Williams & Wilkins (1949)
- 2) Udagawa, S. and Tsubaki, K. : Illustrated book of fungi (in Japanese), p. 1076-1081 and p. 1099-1100 (1978)
- 3) Spadaro, j. T., Gold, M. H., and Renganathan, V. : Degradation of azo dye by the lignin-degrading fungus *Phanerochaete chrysosporium*. Appl. Environ. Microbiol., 58, 2397-1401 (1992)
- 4) Kim, S. J. and Shoda, M. : Batch decolorization molasses by suspended and immobilized fungus of *Geotrichum candidum* Dec. 1. J. Biosci. Bioeng., 88, 577-581 (1999)
- 5) Miyata, N., Mori, T., Iwahori, K., and Fujita, M. : Microbial decolorization of melanoidine-containing wastewater: combined use of activated sludge and the fungus *Coriolus hirsutus*. J. Biosci. Bioneng., 89, 145-150 (2000)

(Submitted 2001. 5. 7)

(Accepted 2001. 6. 20)