Title	Selection and functional analysis of a <i>Pyropia yezoensis</i> ammonium transporter PyAMT1 in potassium deficiency
Author(s)	Adams, Eri; Mikami, Koji; Shin, Ryoung
Citation	Journal of Applied Phycology, 29(5), 2617-2626 https://doi.org/10.1007/s10811-017-1196-1
Issue Date	2017-10
Doc URL	http://hdl.handle.net/2115/71573
Rights	This is a post-peer-review, pre-copyedit version of an article published in Journal of Applied Phycology. The final authenticated version is available online at: http://dx.doi.org/10.1007/s10811-017-1196-1 .
Type	article (author version)
File Information	Mikami JAPhy.pdf

[Instructions for use](#)

Selection and functional analysis of a *Pyropia yezoensis* ammonium transporter PyAMT1 in potassium deficiency

Eri Adams¹, Koji Mikami², Ryoung Shin¹

¹RIKEN Center for Sustainable Resource Science, 1-7-22 Suehirocho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan

²Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate, Hokkaido 041-8611, Japan

Correspondence should be addressed to R.S. (ryoung.shin@riken.jp).

Abstract

Seaweeds are believed to have developed unique mechanisms to maintain optimal cellular potassium and sodium concentrations in order to survive in the saline marine environment. To gain a molecular understanding of underlying potassium/sodium homeostasis in seaweeds, full-length cDNA libraries from the multiple stages in the life cycle, including gametophytes, conchosporangia and sporophytes of a marine red alga *Pyropia yezoensis* were constructed. A large portion of genes from each library through the life cycle was revealed to be functionally unknown reconfirming the uniqueness of *P. yezoensis* genes in terms of evolutionary lineage. Genes that could potentially contribute to potassium deficiency tolerance were selected from the potassium uptake defective *Escherichia coli* strain expressing gametophytes and conchosporangia libraries under the low potassium conditions. Of those, an ammonium transporter gene, *PyAMT1*, was demonstrated to enhance potassium deficiency tolerance effectively when expressed in the *E. coli* strain. Potential roles of *PyAMT1* and other candidate components in this context are discussed.

Keywords

ammonium transporter, full-length cDNA library, potassium deficiency, *Pyropia*

31 *yezoensis*, salt tolerance

Introduction

Seaweeds have adapted to the extremely high salt environment in the ocean, an environment that most of land plants never encounter. High levels of sodium (Na^+) in the cell cause osmotic and ionic stress and disturb potassium (K^+) uptake and functions due to their similar physicochemical properties, often resulting in a K^+ deficiency response (Adams and Shin, 2014). Despite the high concentrations of Na^+ in seawater, cytosolic concentrations of Na^+ are generally maintained at low levels in marine algae, suggesting the existence of Na^+ extrusion mechanisms (Kirst, 1990; Karsten, 2012).

It has been long known that the marine red algae Bangiales (Rhodophyta) which include *Pyropia* and *Porphyra* (Sutherland et al., 2011) accumulate K^+ in the cytoplasm and either exclude or contain Na^+ preferentially in the vacuoles (Eppley, 1958; Wiencke et al., 1983). In order to maintain the appropriate cytosolic K^+/Na^+ ratios, active K^+ uptake mechanisms are considered essential. Unlike land plants and green algae (Chan et al., 2012; Pedersen et al., 2012), red algae such as *Pyropia yezoensis* and *Porphyridium purpureum* have been reported to possess animal-type Na^+/K^+ -ATPases which extrude three ions of Na^+ while taking up two ions of K^+ into the cell and they are predicted to provide the driving force for Na^+ -driven solute transporters (Barrero-Gil et al., 2005; Bhattacharya et al., 2013). There seems a tendency that freshwater algae and

land plants utilise H^+ gradient generated by H^+ -ATPases to energise secondary transporters whereas marine algae make use of Na^+ gradient albeit with some exceptions (Chan et al., 2012) and this notion is evolutionarily quite interesting.

Recently, the 43 Mb genome of *P. yezoensis* was sequenced, with more than 10,000 gene models predicted (Nakamura et al., 2013). In this alga, a gene encoding K^+ P-type ATPase, *PyKPA1*, was found to be phylogenetically related to the animal H^+/K^+ - and Na^+/K^+ -ATPases. Heterologous expression of *PyKPA1* in the *Escherichia coli* strain deficient in K^+ uptake demonstrated that PyKPA1 had a growth promoting effect in the K^+ -limited condition and that addition of Na^+ further enhanced the effect of PyKPA1 (Barrero-Gil et al., 2005), suggesting PyKPA1 to be a Na^+/K^+ -ATPase. Although gene expression of *PyKPA1* was not altered in response to salt stress in *P. yezoensis* (Uji et al., 2012a), ectopic overexpression of *PyKPA1* in rice plants increased salt tolerance by restoring growth (Kishimoto et al., 2013). Another P-type ATPase, PyKPA2, which shares a 65% sequence identity with PyKPA1 and Na^+/H^+ antiporters, PySOS1 and PyNhaD, have also been isolated from the genome of *P. yezoensis* (Barrero-Gil et al., 2005; Uji et al., 2012a; Uji et al., 2012b). These membrane proteins could potentially be the major machineries in ion homeostasis and Na^+ extrusion in *P. yezoensis*, however, ATPases may be too energetically costly to be the major K^+ uptake mechanism and

additional K⁺ transporters/channels are expected to exist. Although K⁺ uptake in land plants is commonly mediated by K⁺ channels and transporters such as AKT1, HAK5 and KUPs in a model land plant *Arabidopsis thaliana* (Adams and Shin, 2014), proteins with similar sequences and/or function have not been reported in *P. yezoensis*. In order to resolve the molecular mechanisms underlying the ability of red seaweeds to survive in the marine environment, investigation of regulatory components involved in K⁺/Na⁺ homeostasis in *P. yezoensis* needs to be performed.

P. yezoensis spends the winter in the form of gametophytes, the leafy structure commonly harvested as seaweed, and it turns into sporophytes, the filamentous structure during the summer. In autumn, sporophytes form conchosporangia from which conchospores are emitted to produce a new generation of gametophytes. There are several reports describing that different sets of genes are expressed in the extremely diverse structures observed throughout the life cycle of *P. yezoensis*: for instance, only 22.5% of ESTs and one out of 14 microRNAs are common among gametophytes and sporophytes (Asamizu et al., 2003; Shen et al., 2011; He et al., 2012). Indeed, phase-specific gene expression include genes encoding urea transporters (*PyDUR3s*), an alginate lyase (*PyAly*) and a bromoperoxidase (*PyBPOI*) (Inoue et al., 2015; Matsuda et al., 2015; Kakinuma et al., 2016b). Interestingly, *PyKPAI* has been reported to be

predominantly expressed in sporophytes while *PyKPA2* is expressed specifically in gametophytes (Uji et al., 2012a). These findings strongly suggest the existence of distinct regulatory mechanisms upon K^+/Na^+ homeostasis in each life stage.

To identify the genes playing roles in K^+ deficiency tolerance throughout the life cycle of *P. yezoensis*, we here constructed full-length cDNA libraries using three different stages, gametophytes, conchosporangia and sporophytes, and these libraries were then transferred into the *E. coli* expression vector system to isolate the genes involved in K^+ deficiency response. Candidate genes and possible mechanisms by which *P. yezoensis* tolerates K^+ deficiency are discussed.

Materials and methods

Plant material and growth conditions

The cultivation of the *P. yezoensis* strain U51 was performed as previously reported (Li et al., 2008) with a slight modification. Briefly, free-living sporophytes, free-living conchosporangia and gametophytes attached to polyvinyl alcohol (PVA) monofilaments were suspended in ESL (enriched SEALIFE) media, continuously aerated with filter-sterilised air and grown at 15°C in a 10 h light/14 h dark photocycle with a light intensity of $60 \mu\text{mol m}^{-2} \text{s}^{-1}$. The sterile ESL medium was made by

dissolving commercially available SEALIFE powder (Marintech Co. Ltd., Tokyo, Japan) in distilled water with added ESS₂ solution (Kitade et al., 2002) and this was exchanged weekly.

RNA extraction and cDNA library construction

An excess amount of sporophyte, gametophyte and conchosporangium samples were flash frozen in liquid N₂ and ground into fine powder using a mortar and a pestle. Total RNA was extracted using TRIzol reagent (Thermo Fisher Scientific, MA) and 75 µg of total RNA was used to isolate mRNA using Ambion Dynabeads mRNA Purification Kit (Thermo Fisher Scientific). Purified mRNA was concentrated by glycogen precipitation with 1 µL of glycogen, 0.5 volumes of NH₄OAc and 2.5 volumes of 100% ethanol. 0.88~3.31 µg of mRNA was used to create full-length cDNA libraries using CloneMiner II cDNA Library Construction Kit according to the manufacture's instruction (Thermo Fisher Scientific). In short, hybridisation of Biotin-*attB2*-Oligo(dT) Primer to the mRNA poly(A) tail and the first strand cDNA synthesis by SuperScript III Reverse Transcriptase were followed by the second strand cDNA synthesis by *E. coli* Polymerase I and ligation of *attB1* Adapter to the 5' end of the cDNA. The resultant double-stranded cDNA was size fractionated by a column to remove truncated cDNA

shorter than 500 bp and cloned into a Gateway entry vector pDONR222 through BP recombination reaction. The cDNA construct was then transformed into ElectroMAX DH10B T1 Phage Resistant Cells to create the final cDNA library. Titer was determined by spreading 1:10 serial dilutions (10^{-2} , 10^{-3} , 10^{-4}) of each library onto LB plates containing kanamycin. Titer was calculated as colony forming unit (cfu mL⁻¹) = colonies on plates × dilution factor / volume plated (mL) and total CFU (cfu) = average titer (cfu mL⁻¹) × total volume of cDNA library (mL). Single colonies were picked and plasmid DNAs (pDNAs) were prepared. Each pDNA was digested by *Bsr*I to determine the insert size and sequenced using M13 forward and reverse universal primers and the Sanger sequencing technique (HITACHI gene analysis system with ABI PRISM 3100-21 genetic analyser).

Selection of K⁺ deficiency tolerance-related genes

The pDONR222 entry libraries from gametophyte and conchosporangium samples were transferred into the pBAD-DEST49 Gateway destination vector according to the manufacture's instruction (Thermo Fisher Scientific). Plasmid DNA was prepared from the entry library culture grown till an OD₆₀₀ to be approximately 1.0. Polyethylene glycol (PEG) precipitation was performed to purify pDNA using 0.4 volumes of 30%

PEG/Mg solution. The entry library was transferred into the destination vector using Gateway LR Clonase II enzyme and transformed into ElectroMAX DH10B T1 Phage Resistant Cells. Plasmid DNA of the pBAD-DEST49 library prepared from the *E. coli* culture with an OD₆₀₀ of approximately 1.0 was transformed into an *E. coli* strain defective in K⁺ uptake, TK2463 (Epstein et al., 1993), and selected on minimal media (Ahn et al., 2004) containing 1-3 mM KCl, 0.1% arabinose and ampicillin. For functional analysis, overnight culture of TK2463 expressing pBAD-PyAMT1, PyβCA1 or PyHSP70 grown in KML media (10 g Bacto Tryptone, 10 g KCl, 5 g Bacto Yeast Extract in 1 L MilliQ water) containing ampicillin were pelleted, washed three times with autoclaved MilliQ water, resuspended in autoclaved MilliQ water and dropped onto minimal media containing 30, 1.5 or 1.25 mM KCl, 0.1% arabinose and ampicillin as five-fold serial dilutions.

Sequence analysis

Contig numbers were retrieved from the obtained sequences using the public *Pyropia* database (Nakamura et al., 2013). *Pyropia* genes were annotated using blastx function against the non-redundant protein sequences database at the NCBI search engine (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Amino acid sequences were aligned using

Vector NTI (Thermo Fisher Scientific).

Results

Construction and validation of full-length cDNA libraries of *Pyropia yezoensis*

In order to gain molecular information on each stage in the life of *P. yezoensis*, cDNA libraries derived from gametophyte, sporophyte and conchosporangium samples, whose diverse structures are shown in Fig 1 (a,b for gametophytes, c for sporophytes and d,e for conchosporangia), were constructed. Total CFU was greater than 10^7 for all the libraries with the greatest being $>10^8$ for the sporophytes library (Table 1). Twenty-four single colonies from each library were randomly picked to validate the diversity of the cDNA libraries and pDNA was digested with restriction enzyme *Bsr*G I to determine the insert size. The recombination efficiency was 100% for all libraries and the average size of inserts was approximately 1 kb (Table 1). Inserted cDNA from each colony was also sequenced and annotated against the public protein sequences database (Table 2). For the gametophytes library, one third of genes were either not annotated or annotated to encode proteins of unknown function and most of the other genes were predicted to encode rather ubiquitous proteins involved in the general biological processes such as protein synthesis, regulation and degradation (Table 2 and Fig. 2). By contrast, more

than half of the genes were annotated as unknown in the conchosporangia and sporophytes libraries. The annotated genes were for general functions such as protein synthesis and degradation.

Selection of genes potentially involved in K⁺ deficiency tolerance

In order to isolate genes responsible for efficient K⁺ utilisation and K⁺/Na⁺ balance in *P. yezoensis*, the entire entry clone libraries from gametophytes and conchosporangia were transferred into the *E. coli* expression vector system and transformed into TK2463, an *E. coli* strain defective in K⁺ uptake. Under the less stringent K⁺ deficiency conditions (2 or 3 mM KCl), 48 colonies were recovered and 45 genes were successfully sequenced from the gametophytes library (Table 3). Under the stringent condition (1 mM KCl), 23 genes from the gametophytes library and 16 genes from the conchosporangia library were revealed (Table 4). Although a large portion of genes could not be annotated for function as in the entry libraries (34.8% for gametophytes and 62.5% for conchosporangia), a higher number of the annotated genes was associated with specific functions in biological processes such as metabolism and signalling rather than general functions. Of these, three genes were annotated as β -carbonic anhydrase (β CA, two from 2 or 3 mM KCl, one from 1 mM KCl screening). All three were predicted to

194 represent the same gene model (*PyβCA1*, contig_16545_g4020). Upon sequence
195 alignment with known βCAs from *Chlamydomonas reinhardtii* (CrCAH4),
196 *Ostreococcus tauri* (OtβCA) and *Arabidopsis* (AtβCA5.1), *PyβCA1* was shown to
197 possess all three conserved zinc binding sites, two cysteine residues (C) and histidine
198 (H) (marked in blue in Fig. 3) (Provart et al., 1993; Bracey et al., 1994; Kimber and Pai,
199 2000), however, the rest of the sequence was fairly diverse among the species (20.1%,
200 22.0%, 23.1% identity with AtβCA5.1, CrCAH4, OtβCA, respectively). Two genes
201 recovered from stringent K⁺ deficiency screening were predicted to be a single
202 ammonium transporter (*PyAMT1*, contig_16335_g3953) (Kakinuma et al., 2016a).
203 Sequence alignment of *PyAMT1* with well-studied AtAMT1;2 (Yuan et al., 2007) and
204 algal OtAMT (Derelle et al., 2006) indicated that many conserved amino acids such as
205 those which form the ammonium binding site, tryptophan (W) and serine (S) (marked in
206 red in Fig. 4), phenylalanine (F) and aspartate (D) (marked in blue) were identical
207 among three whereas others, such as F and threonine (T) (marked in green) highlighted
208 the difference between algae and land plants (Pantoja, 2012). The TK2463 *E. coli*
209 strains expressing *PyβCA1* and *PyAMT1* were further analysed in the K⁺ deficient
210 conditions (1.25 and 1.5 mM KCl) and compared with the strain expressing *PyHSP70*
211 as a negative control. The strains expressing *PyAMT1* and, to a lesser extent, *PyβCA1*

grew well in K⁺ deficiency while the strain expressing *PyHSP70* could not survive (Fig. 5). Multiple ribosomal proteins of various sizes were also selected from the gametophyte library (Table 3 and 4).

Discussion

Full-length cDNA libraries from various life stages, including gametophytes, conchosporangia and sporophytes, were created for a model marine alga *P. yezoensis* with excellent recombination percentages and titer. The average insert size of approximately 1 kb corresponds with the predicted average coding sequence length in *P. yezoensis* (Nakamura et al., 2013). Analysis of the whole genome sequence of *P. yezoensis* has revealed that the function of 35% of the genes is unknown (Nakamura et al., 2013) and our results obtained from the gametophytes library was consistent with their report. It is intriguing to postulate why the conchosporangia and sporophytes libraries contain more than 50% of genes that are functionally unknown. Many of these genes do not even have any previously characterised conserved amino acid motif or domain, underlining the uniqueness of *P. yezoensis* genes, especially in the conchosporangia and sporophytes stages.

In the search of contributory factors in K⁺ deficiency tolerance in *P. yezoensis*, we

identified a series of genes that might be involved in such response as efficient uptake and use of K^+ . Gametophytes and conchosporangia cDNA libraries were expressed in the *E. coli* system and selected in two different stringency conditions of K^+ deficiency. More colonies were found in the less stringent condition (45 transformants from gametophytes) than in the more stringent condition (23 transformants from gametophytes and 16 transformants from conchosporangia). The ratios of unannotated genes were similar to the pattern in the entry libraries. Unlike the genes annotated in the entry libraries, selected genes were annotated as proteins with specific biological functions rather than ubiquitous proteins, suggesting specific pathways at work in K^+ deficiency response.

There were two types of proteins selected multiple times: β -carbonic anhydrase (Py β CA1) and ammonium transporter (PyAMT1). CA catalyses the reversible reaction between CO_2 and $HCO_3^- + H^+$ and is crucial for aquatic photosynthetic organisms which suffer in the low- CO_2 environment to concentrate CO_2 in the vicinity of ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) (Moroney et al., 2001). The existence of CA activity in marine macroalgae has been known for a long while (Bowes, 1969). A β CA has previously been cloned in *P. yezoensis* and its expression was reported to be the lowest in gametophytes, followed by sporophytes and conchospores

(Zhang et al., 2010). Although Py β CA1 selected in this screen is different from the one identified in the previous study, its expression is predicted to be low in gametophytes since the same expression pattern was also observed for *P. haitanensis* β CAs (Chen et al., 2016). Considering its lower abundance in the original gametophytes library and the fact that three independent transformants were recovered in the low K⁺ assays, it seems to point to the significance of Py β CA1 in K⁺ deficiency response. Furthermore, this particular β CA might be important in this response as only one gene was repeatedly isolated though multiple β CAs were expected to exist in the genome. As all three conserved amino acids which contribute to binding of the catalytic zinc ion are present in Py β CA1 (Fig. 3), it is predicted as a functional β CA. By contrast, overall sequence identity is not high among the species tested (approximately 20%) and it is possible that regulation and function of each β CA are distinct. This notion is also supported by the fact that CA is essential for *E. coli* growth under aerobic conditions probably due to HCO₃⁻ requirement for amino acid, nucleotide and fatty acid synthesis (Merlin et al., 2003), indicating that expression of *Py β CA1*, but not the innate CA activity of *E. coli*, could contribute to K⁺ deficiency tolerance (Fig. 5). Although a direct interaction between β CA and K⁺ has yet to be reported in seaweeds (Escassi et al., 2002), we speculate that increased carbon source and photosynthesis by Py β CA1 might

compensate the loss of K⁺.

Two independent transformants from the stringent K⁺ deficiency screen were found to carry a single *AMT* gene (*PyAMT1*). During the review process of the current paper, another group reported isolation of PyAMT1 as a functional ammonium transporter whose gene expression is dramatically induced in response to nitrogen deficiency (Kakinuma et al., 2016a). It is known that ammonium is preferentially taken up over nitrate by many algae and multiple *AMT* genes are present and expressed in the *Porphyra* species (Chan et al., 2012). Amino acid sequence alignment of PyAMT1 with *Arabidopsis* AtAMT1;2 and green alga *O. tauri* OtAMT indicated that many of the functional residues were conserved but some of the amino acids were the same among algae but this was not the case with *Arabidopsis*. Interestingly, substitution of the H125 residue identified from bean (*Phaseolus vulgaris*), which is generally replaced by proline (P) in other plant homologues (marked in orange in Fig. 4), for arginine (R) renders the transporter more active (Ortiz-Ramirez et al., 2011). Since the H125 position is R in PyAMT1, this might suggest it to be an active form. Expression of *PyAMT1* dramatically improved the ability of TK2463, an *E. coli* strain defective in K⁺ uptake, to survive in K⁺ deficiency compared to the negative control line expressing *PyHSP70* (Fig. 5). Although the negative control line showed somewhat compromised growth in the

sufficient K^+ condition, the degree of viability under K^+ deficiency between the lines expressing *PyHSP70* and *PyAMT1* was fairly clear. The effect of *PyAMT1* was stronger than that of *PyβCA1* (Fig. 5) and this point was consistent with the fact that *PyAMT1* was recovered solely from the stringent screen while *PyβCA1* from the mild screen as well. Interaction between K^+ and ammonium is known due to their chemical similarities such as charge and size. Replacement of a nitrogen source as nitrate to ammonium in tobacco (*Nicotiana tabacum*) was reported to cause growth retardation and a decrease in K^+ uptake (Lu et al., 2005). In *Arabidopsis*, ammonium has been shown to inhibit K^+ deficiency-induced expression of a high-affinity K^+ transporter gene, *AtHAK5* (Qi et al., 2008; Rubio et al., 2008). By contrast, in ammonium-tolerant rice species, ammonium inhibits high-affinity K^+ transport but promotes low-affinity K^+ uptake (Szczerba et al., 2008). Tomato (*Solanum lycopersicum*) *LeHAK5* expression is induced by ammonium although K^+ concentrations in roots are not altered, and K^+ uptake and accumulation are stimulated by ammonium in sorghum (*Sorghum bicolor*) (Alvarez-Pizarro et al., 2011). As shown in the examples from the previous reports, whether ammonium prevents or activates K^+ uptake depends on the plant species. There is no information available at present on the effect of ammonium on K^+ uptake in *P. yezoensis*, but it is possible that increased concentrations of ammonium due to *PyAMT1*,

directly or indirectly, help accumulate K^+ under K^+ starvation. It would be interesting to demonstrate the functions of PyAMT1 *in planta* and compare those with the functions of AMTs from land plants in terms of K^+ deficiency tolerance.

Taken together, our findings provided insight into the potential pathways involved in K^+ uptake and response in *P. yezoensis*, PyAMT1 and probably ammonium being strong candidate components, although further investigation is required to clarify the roles of the selected genes in the K^+ deficiency response. The cDNA libraries created will serve as a useful tool to understand the molecular mechanisms underlying K^+/Na^+ homeostasis in seaweeds.

Acknowledgements

We would like to thank Ms. Takae Miyazaki and Mr. Hajime Takiguchi for taking care of the seaweed cultures, Dr. Yuko Makita and Dr. Minami Matsui (RIKEN Center for Sustainable Resource Science) for creating the *P. yezoensis* BLAST search engine and Dr. Megumu Takahashi (Tokyo University of Agriculture) for providing a photo of conchosporangium generation. We appreciate the contribution of the Marine Resources Research Center, Aichi Fisheries Research Institute in kindly providing the *P. yezoensis* strain U51 and Dr. Nobuyuki Uozumi (Tohoku University) for providing the *E. coli*

TK2463 strain. Many thanks also go to Dr. Michael Adams for comments and discussion on the paper. This work was supported by funding from the RIKEN Incentive Research Projects.

References

Adams E, Shin R (2014) Transport, signaling, and homeostasis of potassium and sodium in plants. *J Integr Plant Biol* 56: 231-249.

Ahn SJ, Shin R, Schachtman DP (2004) Expression of *KT/KUP* genes in Arabidopsis and the role of root hairs in K⁺ uptake. *Plant Physiol* 134: 1135-1145.

Alvarez-Pizarro JC, Gomes E, Prisco JT, Grossi-De-Sa MF, Neto OBO (2011) NH₄⁺-stimulated low-K⁺ uptake is associated with the induction of H⁺ extrusion by the plasma membrane H⁺-ATPase in sorghum roots under K⁺ deficiency. *J Plant Physiol* 168: 1617-1626.

Asamizu E, Nakajima M, Kitade Y, Saga N, Nakamura Y, Tabata S (2003) Comparison of RNA expression profiles between the two generations of *Porphyra yezoensis* (Rhodophyta), based on expressed sequence tag frequency analysis. *J Phycol* 39: 923-930.

Barrero-Gil J, Garciadeblas B, Benito B (2005) Sodium, potassium-ATPases in algae

338 and oomycetes. J Bioenerg Biomembr 37: 269-278.

339 Bhattacharya D, Price DC, Chan CX, Qiu H, Rose N, Ball S, Weber APM, Arias MC,
340 Henrissat B, Coutinho PM, Krishnan A, Zauner S, Morath S, Hilliou F, Egizi A,
341 Perrineau MM, Yoon HS (2013) Genome of the red alga *Porphyridium*
342 *purpureum*. Nat Commun 4: 1941.

343 Bowes GW (1969) Carbonic anhydrase in marine algae. Plant Physiol 44: 726-732.

344 Bracey MH, Christiansen J, Tovar P, Cramer SP, Bartlett SG (1994) Spinach carbonic
345 anhydrase: Investigation of the zinc-binding ligands by site-directed
346 mutagenesis, elemental analysis, and EXAFS. Biochemistry 33: 13126-13131.

347 Chan CX, Zauner S, Wheeler G, Grossman AR, Prochnik SE, Blouin NA, Zhuang YY,
348 Benning C, Berg GM, Yarish C, Eriksen RL, Klein AS, Lin SJ, Levine I,
349 Brawley SH, Bhattacharya D (2012) Analysis of *Porphyra* membrane
350 transporters demonstrates gene transfer among photosynthetic eukaryotes and
351 numerous sodium-coupled transport systems. Plant Physiol 158: 2001-2012.

352 Chen CS, Dai ZZ, Xu Y, Ji DH, Xie CT (2016) Cloning, expression, and
353 characterization of carbonic anhydrase genes from *Pyropia haitanensis*
354 (Bangiales, Rhodophyta). J Appl Phycol 28: 1403-1417.

355 Derelle E, Ferraz C, Rombauts S, Rouze P, Worden AZ, Robbens S, Partensky F,

356 Degroeve S, Echeynie S, Cooke R, Saeys Y, Wuyts J, Jabbari K, Bowler C,
 357 Panaud O, Piegu B, Ball SG, Ral JP, Bouget FY, Piganeau G, De Baets B, Picard
 358 A, Delseny M, Demaille J, Van De Peer Y, Moreau H (2006) Genome analysis of
 359 the smallest free-living eukaryote *Ostreococcus tauri* unveils many unique
 360 features. Proc Natl Acad Sci USA 103: 11647-11652.

361 Eppley RW (1958) Sodium exclusion and potassium retention by the red marine alga,
 362 *Porphyra perforata*. J Gen Physiol 41: 901-911.

363 Epstein W, Buurman E, Mclaggan D, Naprstek J (1993) Multiple mechanisms, roles and
 364 controls of K⁺ transport in *Escherichia coli*. Biochem Soc T 21: 1006-1010.

365 Escassi L, Aguilera J, Figueroa FL, Fernandez JA (2002) Potassium drives daily
 366 reversible thallus enlargement in the marine red alga *Porphyra leucosticta*
 367 (Rhodophyta). Planta 214: 759-766.

368 He LW, Huang AY, Shen SD, Niu JF, Wang GC (2012) Comparative analysis of
 369 microRNAs between sporophyte and gametophyte of *Porphyra yezoensis*. Comp
 370 Funct Genom 2012: 912843.

371 Inoue A, Mashino C, Uji T, Saga N, Mikami K, Ojima T (2015) Characterization of an
 372 eukaryotic PL-7 alginate lyase in the marine red alga *Pyropia yezoensis*. Current
 373 Biotechnol 4: 240-248.

374 Kakinuma M, Nakamoto C, Kishi K, Coury DA, Amano H (2016a) Isolation and
 375 functional characterization of an ammonium transporter gene, *PyAMT1*, related
 376 to nitrogen assimilation in the marine macroalga *Pyropia yezoensis*
 377 (Rhodophyta). Mar Environ Res, In press.

378 Kakinuma M, Suzuki K, Iwata S, Coury DA, Iwade S, Mikami K (2016b) Isolation and
 379 characterization of a new *DUR3*-like gene, *PyDUR3.3*, from the marine
 380 macroalga *Pyropia yezoensis*. Fish Sci 82: 171-184.

381 Karsten U (2012) Seaweed acclimation to salinity and desiccation stress. In: Wiencke C,
 382 Bischof K (eds) Seaweed Biology. Springer, Heidelberg, pp 87-107.

383 Kimber MS, Pai EF (2000) The active site architecture of *Pisum sativum* b-carbonic
 384 anhydrase is a mirror image of that of a-carbonic anhydrases. Embo J 19:
 385 1407-1418.

386 Kirst GO (1990) Salinity tolerance of eukaryotic marine algae. Annu Rev Plant Phys 41:
 387 21-53.

388 Kishimoto M, Shimajiri Y, Oshima A, Hase A, Mikami K, Akama K (2013) Functional
 389 expression of an animal type- Na^+ -ATPase gene from a marine red seaweed
 390 *Porphyra yezoensis* increases salinity tolerance in rice plants. Plant Biotechnol
 391 30: 417-422.

392 Kitade Y, Fukuda S, Nakajima M, Watanabe T, Saga N (2002) Isolation of a cDNA
 393 encoding a homologue of actin from *Porphyra yezoensis* (Rhodophyta). J Appl
 394 Phycol 14: 135-141.

395 Li L, Saga N, Mikami K (2008) Phosphatidylinositol 3-kinase activity and asymmetrical
 396 accumulation of F-actin are necessary for establishment of cell polarity in the
 397 early development of monospores from the marine red alga *Porphyra yezoensis*.
 398 J Exp Bot 59: 3575-3586.

399 Lu YX, Li CJ, Zhang FS (2005) Transpiration, potassium uptake and flow in tobacco as
 400 affected by nitrogen forms and nutrient levels. Ann Bot 95: 991-998.

401 Matsuda R, Ozgur R, Higashi Y, Takechi K, Takano H, Takio S (2015) Preferential
 402 expression of a bromoperoxidase in sporophytes of a red alga, *Pyropia yezoensis*.
 403 Mar Biotechnol 17: 199-210.

404 Merlin C, Masters M, Mcateer S, Coulson A (2003) Why is carbonic anhydrase essential
 405 to *Escherichia coli*? J Bacteriol 185: 6415-6424.

406 Moroney JV, Bartlett SG, Samuelsson G (2001) Carbonic anhydrases in plants and algae.
 407 Plant Cell Environ 24: 141-153.

408 Nakamura Y, Sasaki N, Kobayashi M, Ojima N, Yasuike M, Shigenobu Y, Satomi M,
 409 Fukuma Y, Shiwaku K, Tsujimoto A, Kobayashi T, Nakayama I, Ito F, Nakajima

410 K, Sano M, Wada T, Kuhara S, Inouye K, Gojobori T, Ikeo K (2013) The first
 411 symbiont-free genome sequence of marine red alga, Susabi-nori (*Pyropia*
 412 *yezoensis*). PLoS One 8: e57122.

413 Ortiz-Ramirez C, Mora SI, Trejo J, Pantoja O (2011) PvAMT1;1, a highly selective
 414 ammonium transporter that functions as H⁺/NH₄⁺ Symporter. J Biol Chem 286:
 415 31113-31122.

416 Pantoja O (2012) High affinity ammonium transporters: molecular mechanism of action.
 417 Front Plant Sci 3: 34.

418 Pedersen CNS, Axelsen KB, Harper JF, Palmgren MG (2012) Evolution of plant P-type
 419 ATPases. Front Plant Sci 3.

420 Provart NJ, Majeau N, Coleman JR (1993) Characterization of pea chloroplastic
 421 carbonic anhydrase. Expression in *Escherichia coli* and site-directed
 422 mutagenesis. Plant Mol Biol 22: 937-943.

423 Qi Z, Hampton CR, Shin R, Barkla BJ, White PJ, Schachtman DP (2008) The high
 424 affinity K⁺ transporter AtHAK5 plays a physiological role in planta at very low
 425 K⁺ concentrations and provides a caesium uptake pathway in *Arabidopsis*. J Exp
 426 Bot 59: 595-607.

427 Rubio F, Nieves-Cordones M, Aleman F, Martinez V (2008) Relative contribution of

428 AtHAK5 and AtAKT1 to K⁺ uptake in the high-affinity range of concentrations.
 429 Physiol Plantarum 134: 598-608.

430 Shen S, Zhang G, Li Y, Wang L, Xu P, Yi L (2011) Comparison of RNA expression
 431 profiles on generations of *Porphyra yezoensis* (Rhodophyta), based on
 432 suppression subtractive hybridization (SSH). BMC Res Notes 4: 428.

433 Sutherland JE, Lindstrom SC, Nelson WA, Brodie J, Lynch MDJ, Hwang MS, Choi HG,
 434 Miyata M, Kikuchi N, Oliveira MC, Farr T, Neefus C, Mols-Mortensen A,
 435 Milstein D, Muller KM (2011) A new look at an ancient order: generic revision
 436 of the Bangiales (Rhodophyta). J Phycol 47: 1131-1151.

437 Szczerba MW, Britto DT, Ali SA, Balkos KD, Kronzucker HJ (2008) NH₄⁺-stimulated
 438 and -inhibited components of K⁺ transport in rice (*Oryza sativa* L.). J Exp Bot
 439 59: 3415-3423.

440 Uji T, Hirata R, Mikami K, Mizuta H, Saga N (2012a) Molecular characterization and
 441 expression analysis of sodium pump genes in the marine red alga *Porphyra*
 442 *yezoensis*. Mol Biol Rep 39: 7973-7980.

443 Uji T, Monma R, Mizuta H, Saga N (2012b) Molecular characterization and expression
 444 analysis of two Na⁺/H⁺ antiporter genes in the marine red alga *Porphyra*
 445 *yezoensis*. Fisheries Sci 78: 985-991.

446 Wiencke C, Stelzer R, Lauchli A (1983) Ion compartmentation in *Porphyra umbilicalis*
447 determined by electron-probe X-ray microanalysis. *Planta* 159: 336-341.

448 Yuan LX, Loque D, Kojima S, Rauch S, Ishiyama K, Inoue E, Takahashi H, Von Wiren
449 N (2007) The organization of high-affinity ammonium uptake in *Arabidopsis*
450 roots depends on the spatial arrangement and biochemical properties of
451 AMT1-type transporters. *Plant Cell* 19: 2636-2652.

452 Zhang BY, Yang F, Wang GC, Peng G (2010) Cloning and quantitative analysis of the
453 carbonic anhydrase gene from *Porphyra yezoensis*. *J Phycol* 46: 290-296.

454

455

456

Table 1 Titer, recombination % and average insert size of cDNA libraries for three life stages of the marine red alga *P. yezoensis*

cDNA library	Titer (cfu mL ⁻¹)	Total CFU	Recombination (%)	Average insert (kb)
Gametophytes	1.93×10^6	1.93×10^7	100	1.22
Conchosporangia	1.30×10^6	1.56×10^7	100	1.00
sporophytes	$>10^7$	$>10^8$	100	1.20

Table 2 Annotation of genes from cDNA libraries for three life stages of the marine red alga *P. yezoensis*

Number	Name	Involved in
Gametophytes		
1	ferritin	storage
1	nitrate reductase	metabolism
1	5-formyltetrahydrofolate cycloligase	metabolism
1	transmembrane 9 protein	transport
1	phosphate transporter	transport
1	mitochondrial substrate carrier family protein	transport
1	transcription initiation factor	transcription
1	ribosomal protein	protein synthesis
2	ribosomal RNA/hypothetical protein	protein synthesis
1	Ser/Thr protein phosphatase	protein regulation
1	Ser/Thr protein kinase	protein regulation
1	protein kinase	protein regulation
1	F-box protein	protein degradation
1	proteasome β subunit	protein degradation
8	unknown/no hit	
Conchosporangia		
1	thioredoxin	redox reaction
1	phosphotransferase	metabolism
1	actin	structure
1	bHLH DNA-binding superfamily protein	transcription
1	zinc finger transcription factor	transcription
1	ribosomal protein	protein synthesis
1	disulfide isomerase (thioredoxin superfamily)	protein regulation
1	transducin family protein/WD-40 repeat family protein	protein regulation
1	F-box family protein	protein degradation
1	RING/U-box superfamily protein/E3 ubiquitin-protein ligase	protein degradation
14	unknown/no hit	
Sporophytes		
1	catalase	redox reaction

1	alanine:glyoxylate aminotransferase	metabolism
1	kinesin	transport
1	ER membrane protein	Transport
3	ribosomal protein	protein synthesis
1	GTPase	protein synthesis
1	FKBP-type peptidyl-prolyl cis-transisomerase	protein regulation
1	peptidase	protein degradation
1	proteasome activator protein	protein degradation
2	YGGT family protein	unknown function
11	unknown/no hit	

465

466

Table 3 List of genes selected from the K⁺ tolerance screening of the gametophytes library expressed in the *E. coli* strain defective in K⁺ uptake under mild K⁺ deficiency (2 or 3 mM KCl)

Number	Name	Involved in
Gametophytes		
1	carrier superfamily protein	transport
1	glycyl-tRNA synthetase	metabolism
1	5' adenylyl phosphosulfate reductase	metabolism
1	GDP-D-mannose 3',5'-epimerase	metabolism
1	serine hydroxymethyl transferase	metabolism
1	carbohydrate binding protein	metabolism
1	alanine:glyoxylate transaminase	metabolism
1	glutamate-5-semialdehyde dehydrogenase	metabolism
2	β-carbonic anhydrase	metabolism
1	fructose/ketose-bisphosphate aldolase	metabolism
1	nicotinic acetylcholine receptor-like protein	signalling
1	calmodulin/centrin	signalling
1	chromosome associated-like protein	transcription
1	NAC transcription factor	transcription
1	histone superfamily protein	transcription
1	RNA-binding protein	RNA regulation
1	translational elongation factor EFG/EF2 protein	protein synthesis
10	ribosomal protein	protein synthesis
1	kinase-like protein	protein regulation
1	proteasome subunit	protein degradation
15	unknown	

Table 4 List of genes selected from the K⁺ tolerance screening of the gametophytes and conchosporangia libraries expressed in the *E. coli* strain defective in K⁺ uptake under severe K⁺ deficiency (1 mM KCl)

Number	Name	Involved in
Gametophytes		
2	ammonium transporter	transport
1	voltage-dependent anion channel	transport
1	valine-tRNA ligase/valyl trans synthase	metabolism
1	β-carbonic anhydrase	metabolism
1	phosphoglycerate mutase-like protein	metabolism
1	cytochrome c oxidase-like protein	respiration
1	calcium-binding EF-hand family protein/calcineurin	signalling
1	heat shock protein	defence
1	RNA-binding protein	RNA regulation
3	ribosomal protein	protein synthesis
1	Ser/Thr kinase/phototropin	protein regulation
1	F-box protein	protein degradation
8	unknown	
Conchosporangia		
1	ubiquinol-cytochrome c reductase	metabolism
1	senescence-associated protein	defence
1	DNA repair helicase	defence
2	GTP-binding protein	transcription
1	Ser protease-like protein	protein degradation
10	unknown	

Figure legends

Fig. 1 Images of the multiple stages in the life cycle of *Pyropia yezoensis*. **a** Gametophytes. **b** Vegetative cells of gametophytes. **c** Sporophytes. **d** Generation of a conchosporangium from a sporophyte. **e** Conchosporangia. Full-length cDNA libraries were constructed with RNA extracted from gametophytes, sporophytes and conchosporangia.

Fig. 2 Functional categories (%) of representative genes recovered from each cDNA library. Randomly selected 24 colonies from each of gametophytes, conchosporangia and sporophytes libraries were sequenced for the inserted genes and annotated based on the sequence similarities against the public protein sequences database.

Fig. 3 Amino acid sequence alignment of β CAs. *Pyropia yezoensis* Py β CA1 (contig_16545_g4020) was aligned with *Arabidopsis thaliana* At β CA5.1 (At4g33580), *Chlamydomonas reinhardtii* CrCAH4 (GI: 159475801) and *Ostreococcus tauri* Ot β CA (GI: 308799709). Identical amino acids among all four β CAs are highlighted as dark gray and identical amino acids between two or three among four are highlighted as light gray. The amino acids which form the conserved zinc binding site are marked in blue.

497

498 **Fig. 4** Amino acid sequence alignment of AMTs. *Pyropia yezoensis* PyAMT1
499 (contig_16335_g3953) was aligned with *Arabidopsis thaliana* AtAMT1;2 (At1g64780)
500 and *Ostreococcus tauri* OtAMT (GI:693496005). Identical amino acids among all three
501 AMTs are highlighted as dark gray and identical amino acids between two of three
502 AMTs are highlighted as light gray. The amino acids which form the ammonium
503 binding site are marked in red and other conserved amino acids reported are marked in
504 blue (identical among three), green (identical among algae) and orange (not identical).

505

506 **Fig. 5** Functional analysis of PyAMT1 and Py β CA1 in K⁺ deficiency. The *E. coli* strain
507 defective in K⁺ uptake expressing *PyAMT1* and *Py β CA1* were grown in the K⁺ sufficient
508 (30 mM KCl) and K⁺ deficient (1.5 and 1.25 mM KCl) conditions. *PyHSP70* was used
509 as a negative control.

Gametophytes

Conchosporangia

Sporophytes

PyβCA1 (1) -----MAAVASPTSVPATN
CrCAH4 (1) -----MSSRNVATALRMFATLGRSQAGEASAMMTGSALLAQRAAL
OtβCA (1) -----
AtβCA5 (1) MAATPTHFSVSHDPFSSTSLNLQTQAI FGNHSLKTTQLRIPASFRRKA

PyβCA1 (15) DEPLALLTGEC PAGDKVWASLLASNAQFATAGERPP-AEGVSVTHRGS LA
CrCAH4 (43) GGPQAVNKGCSRCRCGRVACMGACMPMRHLHAHPNPPSDPDQALEYLREGN
OtβCA (1) -----MSSPERAFERLLDGHRAFRRAHFAASDGAA-DVPRALRALSER-
AtβCA5 (51) TNLQVMASGKTFGLTQEANGVAIDRQNNTDVFDDMK-QRFLAFKKLKYMD

PyβCA1 (64) -----G-----GQSPASVVVTCADSRLSPELLFARG
CrCAH4 (93) KRFVNNKPHDSHPTRNLDRVKATAAGQKFFAAFLSCADSRVPVEIIFDQG
OtβCA (43) -----GQRPRALVVACSDSRADPAIVFDTA
AtβCA5 (100) -----DFEHYKNLADAQAPKFLVIACADSRVCPASVVLGFQ

PyβCA1 (90) LGELFVIRTAGNTTGDDS-----TVAS-VEYAVKNLSASLVVVLGHTK
CrCAH4 (143) FGDVFTVRVAGNIVTNEIT-----ASL--EFGTAVLGSKVLMVLGHS A
OtβCA (68) PGDVFTIRNVGSLVPAYAGLDGGHHGTCAATEYATVHLEVPVILVMGHTQ
AtβCA5 (135) PGDAFTVRNIANLVPPEY---SGPTETKAALFESVNTLNVENILVIGHSR

PyβCA1 (132) CGAVGA AVATEADPDAMAE-----QPRTLAA FVKEKLLAPVQAVKL
CrCAH4 (184) CGAVAATMNGAAVP-----GVISSLYYS-----IS-P
OtβCA (118) CGGAAAGLRKYGNP DADASVFGVNEATGEGF IGAWVAL---AEDAVRRV
AtβCA5 (182) CGGIQALMKMEDEG-----DSRSFIHNWVVVGKKAKESTKAV

PyβCA1 (173) RGDADSGFVAACEVENVHHA VRTLLTTSGLWLA KTRVGGGVKVVGAM YHL
CrCAH4 (210) ACKKAQAGD VDGAI AENVKVQMEQLKVSPVLQGLVKEGK-LKIVGGVYDL
OtβCA (165) CERHDPGVRARMLEYELVRQSVQNLLTFPFVKRRVDRGE-LVVKGAVFNV
AtβCA5 (219) ASNLHFDHQCHCEKASINHSLERLLGYPWIEEKVRQGS-LSLHGGYYNF

PyβCA1 (223) ETGVVEEC-----
CrCAH4 (259) ATGKVTEIA-----
OtβCA (214) WDG TLEVL RADGSFEQLDDDAEDGRGEAKRAKN-
AtβCA5 (268) VDCTFEKWTVDY AASRGKKKEGSGI AVKDRSVWS

1

PyAMT1 (1) -----MIATDMTAMAASPVGRQAVSEALAAALTDQVSRNSDS

AtAMT1;2 (1) MDTATTTCSAVDLSALLSSSSNSTSSLAAATFLCSQISNISKLSDTTYA

OtAMT (1) -----MSLTESGAETQSLYNN

51

PyAMT1 (37) MDVFFILVSGYLVFLMOTGFAMLTAGSVRSKNTKNVLLKNVLDACVGAIA

AtAMT1;2 (51) VDNTYLLFSAYLVFAMQLGFAMLCAGSVRAKNTMNIIMLTNVLDAAAGAIS

OtAMT (17) LDANFLLSSAYLVFFMQAGFAMLCAGSVRSKNTKNILIKNVLDACVGALA

101

PyAMT1 (87) YYLFGFAFAYGTEAN---SFLGHSDFALSGDR---TDFHFFFQWTFE

AtAMT1;2 (101) YYLFGFAFAFGTPSNG--FIGRHHSFALSSYPERPGSDFSFFLYQWAF

OtAMT (67) WFYFGYGFALGEASNGKLNSFIGSGNFAMKGVSGN--TGIAMYLFWSEFS

151

PyAMT1 (129) ATAATIVSGSVAERTSFYAYLGYAFFLSGFVYPIVSHWVWGG-GWLSTIF

AtAMT1;2 (149) IAAAGITSGSIAERTQFVAYLIYSTFLTGFVYPTVSHWFWS SDGWASASR

OtAMT (115) AAATTIVSGSVAERTKFEAYLGYSFFLCAFVYPVVHWGWSGQGWLG PWR

201

PyAMT1 (178) TVGAK-----DFAGDAVVHVMVGGFAGLAGATIVGPRLGRFDQ

AtAMT1;2 (199) SDNNLLFG-----SGAIDFAGSGVVHVMVGGIAGLCGALVEGPRI GRFDR

OtAMT (165) CEGSSNGCGPLLAGSGMLDFAGSGIVHMTGGVAGLVGAIIVGPRTGRFAP

251

PyAMT1 (215) DGRVVPMPGHSATLCTLGTFILWFGWYGFNPGSTLGISNTG----PDADY

AtAMT1;2 (243) SGRSVALRGHSASLVVLGTFLWFGWYGFNPGSFLTILKGYDKSRPYYGQ

OtAMT (215) DGRVNPMPGHSAPLVVLGTFILWLWFGWYGFNPGSQLAIVAFGG---AAADN

301

PyAMT1 (261) TVTAARCAVTTTIAAASAGVTTLIVIKLRDHIFDLLACLNGILAGLVAIT

AtAMT1;2 (293) WSAVGRTAVTTTTLTGCTAALTTLFSKRLLAGHWNVIDVCNGLLGGFAAIT

OtAMT (262) SRVIARTAVTTTTLAAGGGIMAMVLNYVLYHVWDLIAVCNGLILAGLVGIT

351

PyAMT1 (311) ASCAWVEVYAALVIGVIGALVYIGAAMLLLMFKIDDPLEAFPIHGAVGVW

AtAMT1;2 (343) SGC AVEPWAAIVCGFVASWVLIGFNLLAKKLKYDDPLEAAQLHGGCGAW

OtAMT (312) AGCSTTEPWAAIPICGALSALVIHASSKLLKLLKIDDPLEAAPMHGFCGAF

401

PyAMT1 (361) GAFVGLFARIELLTLSGYGNDNGWE--GVFYGGGGRLLAANCVMIASTIA

AtAMT1;2 (393) GLIFTGLFARKEYVNEIYSGDR----PYGLFMGGGGKLLAAQIVQIIVIV

OtAMT (362) GVLWVGFMKQSYVAEVFGTARNGYMPAGVFYGGNGKLLGAQIAGICVIT

451

PyAMT1 (409) GWTLMIVPLFVVLNLVGVLRLISP EMELIGNDVSKHGGAAYPDDVITTEE

AtAMT1;2 (439) GWVTVTMGPLFYGLHKMNLRLISAEDEMAGMDMTRHGGFAYAYNDEDDVS

OtAMT (412) AAVGATLGAFELLMKKLNLLRTSVEEETMGLDESKHGGSAAYAMELVAP

501 526

PyAMT1 (459) KQAGHTIDNLGVDDSLSRADDPTMV-

AtAMT1;2 (489) TKPWGHFAGRVEPTSRSSPTPTLTV

OtAMT (462) A-----