Title	Circulating insulin-like growth factor binding proteins in fish : their identities and physiological regulation
Author(s)	Shimizu, Munetaka; Dickhoff, Walton W.
Citation	General and Comparative Endocrinology, 252, 150-161 https://doi.org/10.1016/j.ygcen.2017.08.002
Issue Date	2017-10-01
Doc URL	http://hdl.handle.net/2115/71575
Rights	© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Shimizu17HUSCAP.pdf

[Instructions for use](#)

Title:

Circulating insulin-like growth factor binding proteins in fish: their identities and physiological regulation

Authors:

Munetaka Shimizu^{a*} and Walton W. Dickhoff^b

^aFaculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato, Hakodate, Hokkaido 041-8611, Japan.

^bNorthwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd. E., Seattle, WA 98112, USA.

*Corresponding author at: Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato, Hakodate, Hokkaido 041-8611, Japan.

Fax: +81 138 40 8897. E-mail address: mune@fish.hokudai.ac.jp (M. Shimizu)

Abstract

Insulin-like growth factor binding proteins (IGFBPs) play crucial roles in regulating the availability of IGFs to receptors and prolong the half-lives of IGFs. There are six IGFBPs present in the mammalian circulation with IGFBP-3 being most abundant. In mammals IGFBP-3 is the major carrier of circulating IGFs, facilitated by forming a ternary complex with IGF and an acid-labile subunit (ALS). IGFBP-1 is generally inhibitory to IGF action by preventing it from interacting with its receptors. In teleosts, the third-round of vertebrate whole genome duplication created paralogs of each IGFBP, except IGFBP-4. In the fish circulation, three major IGFBPs are typically detected at molecular ranges of 20-25, 28-32 and 40-50 kDa. However, their identities are not well established. Three major circulating IGFBPs in Chinook salmon have been identified through protein purification and cDNA cloning. Salmon 28- and 22-kDa IGFBPs are co-orthologs of IGFBP-1, termed IGFBP-1a and -1b, respectively. They are induced under catabolic conditions such as stress and fasting but their responses are somewhat different, with IGFBP-1b being the most sensitive of the two. Cortisol stimulates production and secretion of these IGFBP-1 subtypes while, unlike in mammals, insulin may not be a primary suppressor. Salmon 41-kDa IGFBP, a major carrier of IGF-I, is not IGFBP-3, as might be expected extrapolating from mammals, but is in fact IGFBP-2b. Salmon IGFBP-2b levels in plasma are high when fish are fed, and GH treatment increases its circulating levels similar to mammalian IGFBP-3. These findings suggest that salmon IGFBP-2b acquired the role and regulation similar to mammalian IGFBP-3. Multiple replication of fish IGFBPs offer a unique opportunity to investigate molecular evolution of IGFBPs.

Keywords: insulin-like growth factor binding protein, circulation, fish, identification, hormone, environment

1. Introduction

Insulin-like growth factor (IGF)-I plays a pivotal role in growth regulation in vertebrates and exerts its action through endocrine, paracrine and autocrine modes (Daughaday and Rotwein, 1989; LeRoith et al., 2001; Ohlsson et al., 2009). Endocrine IGF-I is produced mainly by the liver in response to stimulation by growth hormone (GH), and it mediates many of the GH actions in target tissues (Salmon and Daughaday, 1957; Daughaday and Rotwein, 1989). IGF-I is also expressed in virtually all tissues and acts as a local growth factor (D'Ercole et al., 1980; LeRoith et al., 2001). IGF-I is structurally related to insulin arising from a common ancestor, and the ancestral IGF molecule further diverged into IGF-I and -II by gene duplication (Collet et al., 1998; Reinecke and Collet, 1998; Wood et al., 2005a). In contrast to insulin, circulating levels of IGFs are relatively high for hormones averaging around 200 ng/ml for IGF-I and 500 ng/ml for IGF-II in humans (Daughaday and Rotwein, 1989). These high circulating levels are due to stabilization by IGF-binding proteins (IGFBPs). IGFBPs prolong the half-lives of IGFs by protecting them from glomerular filtration and enzymatic degradation (Guler et al., 1987, 1989). In the mammalian circulation, there are three pools of IGFs around 150-, 50- and 7.5-kDa (Fig. 1; Guler et al., 1987, 1989; Zapf, 1995; Rajaram et al., 1997). In humans, 75-80% of circulating IGFs are in the 150-kDa pool consisting of IGF, IGFBP-3/-5 and an acid-labile subunit (ALS) (Baxter and Martin, 1989). ALS does not bind IGFs but binds IGFBP-3 to form a ternary complex. As this complex is too large to cross the capillary barrier or to be filtered at the kidney, it constitutes a large pool of IGFs in the circulation (Binoux and Hossenlopp, 1988; Hasegawa et al., 1992). The 50-kDa pool consists of IGFBP and IGF and carries about 20% of circulating IGFs (Guler et al., 1987, 1989). Less than 1% of circulating IGFs is in the free form (Frystyk et al., 1994).

IGFBPs also regulate the availability of IGFs to target tissues. There are six IGFBPs in the mammalian circulation, and they modulate the activity of IGFs in various ways (Rajaram et al., 1997). IGFBP-1 is the first member of the IGFBP family to be identified and generally inhibits IGF action by preventing it from interacting with its receptor (Lee et al., 1993, 1997; Kajimura and Duan, 2007; Wheatcroft and Kearney, 2009). Unlike other IGFBPs, circulating levels of IGFBP-1 show dynamic fluctuations in response to meals and increase under catabolic conditions such as fasting (Yeoh and Baxter, 1988). In mammals insulin is the primary suppressor of IGFBP-1 whereas cortisol increases its production and secretion (Unterman et al., 1991; Katz et al., 1998). IGFBP-2 is also considered to be an inhibitor of IGF-I action and IGFBP-2 levels are generally high under catabolic conditions, although its levels are relatively

stable and no clear hormonal control has been reported (Rajaram et al., 1997; Wheatcroft and Kearney, 2009). IGFBP-3, as mentioned above, is a major carrier of circulating IGF-I in mammals by forming a ternary complex with IGF and ALS (Baxter and Martin, 1989; Martin and Baxter, 1992). The major site of production of IGFBP-3 and ALS is the liver, but different cell types within the liver produce them. IGFBP-3 is produced by Kupffer and endothelial cells, and ALS is produced by hepatocytes (Chin et al., 1994; Villafuerte et al., 1994). All components of the ternary complex are induced by GH. IGFBP-3 can be a stimulator of IGF-I action in terms of protecting IGF-I in the circulation and releasing it to target tissues when the binding protein is partly degraded by specific enzymes (Martin and Baxter, 1992; Rajaram et al., 1997; Firth and Baxter, 2002).

In fish blood, three major IGFBPs around 20-25, 28-32 and 40-50 kDa have been detected. They have molecular weights similar to mammalian IGFBP-4, -1/-2 and -3, respectively (Kelley et al., 1992, 2001). Their circulating levels are regulated by hormones, nutrition, stress and other factors as is the case for mammalian IGFBPs, suggesting that fish IGFBPs are also important for modulating activity of circulating IGFs (Duan, 1997; Kelley et al., 2000, 2001, 2002, 2006; Wood et al., 2005a). However, the identities of the three IGFBPs in fish are not well established, making comparisons with mammalian IGFBPs difficult. This review summarizes identities of the three major IGFBPs in the fish circulation with special reference to salmon and their physiological regulation. As this review focuses only on selected numbers of IGFBPs in the fish circulation, readers should refer to other excellent reviews/references dealing with molecular and functional aspects of fish IGFBPs (Wood et al., 2005a; Rodgers et al., 2008; Daza et al., 2011; Maqueen et al., 2013; Lappin et al., 2016).

2. Detection of IGFBPs in fish blood

IGFBPs in plasma/serum are usually detected by ligand blotting using ¹²⁵I-labeled human IGF-I developed by Hossenlopp et al. (1986). This method visualizes IGFBP bands using a labeled IGF-I instead of a primary antibody and is based on the ability of proteins to bind the ligand. Modified ligand blotting using non-radio labeled ligands such as biotinylated and digoxigenin-labeled IGF-Is have been developed (Grulich-Henn et al., 1998; Shimizu et al., 2000). Normal human serum consistently exhibits five IGFBP bands at 41.5, 38.5, 34, 30 and 24 kDa, which correspond to IGFBP-3, -3, -2, -1 and -4, respectively (Fig. 2). The doublet bands at 41.5 and 38.5 kDa are of IGFBP-3 with different degrees of N-glycosylation (Firth and Baxter, 1999). IGFBP-5 and -6 are probably difficult to detect in normal human serum/plasma since their

concentrations are low in the circulation, they are diffuse due to different degrees of glycosylation, and/or they are unable to bind IGFs when electroblotted onto a nitrocellulose membrane (Rajaram et al., 1997). Since band intensity is a reflection of both relative abundance and affinity to IGF used as a ligand, care should be taken when comparing different types of IGFBPs.

The presence of fish IGFBPs was first reported by Kelley et al. (1992) in the circulation of four teleost species (coho salmon, *Oncorhynchus kisutch*; striped bass, *Morone saxatilis*; tilapia, *Oreochromis mossambicus*; longjawed mudsucker (goby), *Gillichthys mirabilis*) by ligand blotting using ¹²⁵I-labeled human IGF-I (Fig. 3). Niu and Le Bail (1993) also utilized this technique and detected a major IGFBP band at 41.5 kDa and minor bands at 47.7 kDa and 30-34 kDa in serum of rainbow trout (*O. mykiss*) (Fig. 3). The presence of fish IGFBP was also confirmed by IGF-binding assays (Anderson et al., 1993; Niu et al., 1993). The detection of IGFBPs both by ligand blotting and binding assay in plasma of the lamprey (*Geotria australis*) (Fig. 3) led researchers to hypothesize that IGFBP is an ancient protein family that emerged in the early history of vertebrates (Upton et al., 1993).

Multiple IGFBP bands have since been detected in several fish species and they are within a molecular range of 20-50 kDa. Since the molecular weights of these IGFBPs are close to each other, there is some inconsistency in reporting their sizes even in the same species. This inconsistency is probably due to differences in gel concentration for electrophoresis and molecular markers used. High-molecular-weight bands around 70 kDa are often detected in fish plasma on ligand blotting (Fig. 2; Fukuzawa et al., 1996; Shimizu et al., 2000), but these are unlikely to be specific IGFBP bands since the displacement of the binding by unlabelled IGF-I in ligand blotting is often difficult. However, a possibility that these bands are aggregates of IGFBP(s) cannot be ruled out.

3. Purification of fish IGFBPs

Bauchat et al. (2001) were the first to purify a 30-kDa IGFBP from conditioned medium of the rainbow trout hepatoma cell (RTH-149) culture. The medium was first fractionated by hydrophobic chromatography using a Phenyl-Sepharose column. Fractions containing IGF-binding activity were loaded onto an IGF-I affinity column, and IGFBP was eluted with 1 M acetic acid. The IGFBP fraction was further purified by reversed-phase HPLC using a Vydac C-4 column. Approximately 70 µg of purified IGFBP was obtained from 600 ml conditioned medium.

Chinook salmon serum contains three major IGFBPs at 41, 28 and 22 kDa (Fig. 2), and they were purified from serum of spawning males (Shimizu et al., 2003, 2005, 2011b). The purification procedures were based on those for mammalian and trout IGFBPs (Walton et al., 1989; Bauchat et al., 2001) with some modifications. The 41- and 28-kDa IGFBPs were co-purified by the same procedure (Shimizu et al., 2003b, 2011a). Typically, one liter of salmon serum was mixed with proteinase inhibitors and acidified with 2 M acetic acid to dissociate endogenous IGFs from IGFBPs. IGFs were removed by adsorbing to a SP-Sephadex C-25 followed by centrifugation. The supernatant containing IGFBPs was neutralized and the precipitate was removed. The supernatant was loaded onto an IGF-I affinity column and IGFBPs were eluted with 1 M acetic acid. IGFBPs were purified by reversed-phase HPLC using a Vydac C-4 column as described in Bauchat et al. (2001). The 41- and 28-kDa IGFBPs were eluted at 32% and 33% acetonitrile, respectively (Shimizu et al., 2003b, 2011a). The recovery of purified proteins from 1 L serum was 70 µg and 11-24 µg for 41- and 28-kDa IGFBPs, respectively. It is worth noting that purified IGFBPs were "sticky" being easily adsorbed to tubes when lyophilized (Shimizu personal communication). They should be stored in a low adsorption tube and not be completely lyophilized. Since the 22-kDa IGFBP was found to be labile under acidic conditions and when unoccupied by endogenous IGFs, the acidification step was omitted from its purification (Shimizu et al., 2005). Instead, salmon serum was first fractionated by ammonium sulfate precipitation and loaded onto an IGF-I column. The 22-kDa IGFBP was further purified by reversed-phase HPLC and eluted at 37% acetonitrile. Final yield of the purified protein per 1 L serum was 45 µg (Shimizu et al., 2005).

When band patterns of purified salmon IGFBPs on SDS-PAGE were compared between non-reducing and reducing conditions, the molecular weight of the 41-kDa IGFBP decreased by 2 kDa while those of the 28- and 22-kDa IGFBPs increased by approximately 5 kDa (Shimizu personal communication). These changes should be attributed to differential conformational changes by the dissociation of intra-peptide disulfide bonds (Shimizu, personal communication). Purified 41-kDa IGFBP was shown to be N-glycosylated (Shimizu et al., 2003b). Purified fish IGFBPs were also analyzed for their partial N-terminal amino acid sequences (Bauchat et al., 2001; Shimizu et al., 2003b, 2005, 2011a). Four of five N-terminal amino acid sequences of rainbow trout and Chinook salmon 28-30-kDa IGFBPs were identical and similar to those of human IGFBP-1 and -4 (Shimizu et al., 2011a). Salmon 22-kDa IGFBP also showed relatively high sequence homologies to human IGFBP-1 and -4 (Shimizu et al., 2006). On the other hand, the partial N-terminal amino acid sequence of salmon 41-kDa, which

had a molecular weight similar to human IGFBP-3, had the highest sequence homology with IGFBP-2 (Shimizu et al., 2003b). The discrepancy between its molecular weight and N-terminal amino acid sequence raised questions about the identity of salmon 41-kDa IGFBP.

4. cDNA cloning of circulating salmon IGFBPs

cDNAs for the three circulating salmon IGFBPs were cloned from the liver of Chinook salmon by reverse-transcribed (RT)-PCR followed by 5'- and 3'-rapid amplification of cDNA ends (RACE) (Shimizu et al., 2005, 2011a,b). IGFBPs were first amplified by RT-PCR using degenerate primers based on the partial N-terminal amino acid sequences of purified proteins. Forward primers were specific to each IGFBP and a reverse primer was universal to the IGFBP family designed based on a conserved C-terminal region. These combinations of primers successfully amplified different IGFBP cDNAs and partial cDNA sequences were used to further design gene specific primers for RACE. As results, full-length cDNAs for the 22-, 28- and 41-kDa IGFBPs were obtained (Shimizu et al., 2005, 2011a,b). An additional IGFBP sequence was obtained during the cloning of the 41-kDa IGFBP (Shimizu et al., 2011b). These cDNA sequences were compared with those of human and zebrafish IGFBPs. Sequence comparison and phylogenetic analysis revealed that salmon 22-, 28- and 41-kDa IGFBP were IGFBP-1b, -1a and -2b, respectively (Shimizu et al., 2011a,b). The additional IGFBP was identified as IGFBP-2a (Shimizu et al., 2011b). The presence of two subtypes of each IGFBP is common in teleosts except for IGFBP-4 (Daza et al., 2011). These two subtypes are due to the teleost-specific third round of whole genome duplication (Daza et al., 2011). In addition, since salmonids underwent a fourth round of whole genome duplication, they possess up to four copies of each IGFBP (Macqueen et al., 2013). The cloned circulating salmon IGFBPs were assigned to be IGFBP-1a1, -1b1, -2a and -2b1, respectively (Macqueen et al., 2013). However, in this review, they are simply called "a" or "b" to avoid confusion about exact correspondence to IGFBP subtypes of other fishes.

The finding that salmon 41-kDa IGFBP was IGFBP-2b was controversial to what was understood for this protein. Salmon 41-kDa IGFBP had a molecular weight similar to that of mammalian IGFBP-3. Salmon IGFBPs often appeared as doublet bands of 41- and 43-kDa and they were N-glycosylated as was the case of mammalian IGFBP-3 (Shimizu et al., 2003b). Salmon 41-kDa IGFBP was also similar to mammalian IGFBP-3 in terms of its physiological regulation; it was up-regulated under anabolic states and induced by GH treatment (Shimizu et al., 1999, 2003a). These biochemical and physiological data suggested that the salmon 41-kDa

IGFBP corresponded functionally to mammalian IGFBP-3 (Shimizu et al., 2003b). However, when a cDNA for salmon IGFBP-3 was cloned, its N-terminal amino acid sequence did not match that of purified 41-kDa IGFBP (Shimizu et al., 2011b). Moreover, amino acids obtained after digesting purified 41-kDa IGFBP by cyanogen bromide were assigned to the internal regions of the deduced IGFBP-2b sequence. These results indicated that circulating salmon 41-kDa IGFBP was not IGFBP-3 but was instead IGFBP-2b (Shimizu et al., 2011b). Fish 40-50 kDa IGFBP had been assumed to be an ortholog of mammalian IGFBP-3, and the structure and function of IGFBP-3 were assumed to be "conserved". However, it was IGFBP-2b that physiologically corresponded to mammalian IGFBP-3. Our findings suggested that there is a functional convergence between mammalian IGFBP-3 and salmon IGFBP-2b. On the other hand, despite a relatively high sequence homology between human and salmon IGFBP-3, their functions appeared to have diverged.

5. Structural features of circulating salmon IGFBPs

IGFBPs are a single chain polypeptide consisting of three functional domains: N-, C- and L-domains (Fig. 4; Shimasaki and Ling, 1991; Hwa et al., 1999; Firth and Baxter, 2002; Forbes et al., 2012). The N-terminal domains of IGFBP-1-5 have 12 cysteine residues to form six disulfide bonds within the domain, which is necessary for IGF binding. A GCGCCXXC motif is well conserved in the IGFBP family except for IGFBP-6. IGFBP-6 lacks two cysteine residues resulting in five disulfide bonds. There are several IGFBP-related proteins (IGFBPrPs) having sequence homology to the IGFBP N-terminus but not to other domains (Kim et al., 1997; Hwa et al., 1999). It is of note that IGFBPrPs have the ability to bind IGFs and insulin but with much lower affinity compared to IGFBPs due to lack of the IGFBP C-terminus (Hwa et al., 1999). The C-terminal domain of IGFBP has 6 cysteine residues with a conserved CWCV motif, which are also required for high affinity binding to IGFs (Forbes et al., 2012). There are some motifs in the C-terminus important for IGFBPs to associate with the cell surface or translocate into the cell. Such motifs include an Arg-Gly-Asp (RGD) sequence found in IGFBP-1 and -2 (Firth and Baxter, 2002) and a nuclear localization signal (NLS) found in IGFBP-3, -5 and -6 (Forbes et al., 2012). The middle linker (L) domain of IGFBP is less conserved among IGFBPs and contains several motifs specific to each type such as sites for N-glycosylation, proteolytic cleavage, phosphorylation, nuclear localization and heparin binding (Shimazaki and Ling, 1991; Hwa et al., 1999; Firth and Baxter, 2002; Forbes et al., 2012).

The deduced amino acid sequences of cloned salmon IGFBPs have relatively high

sequence homologies to mammalian counterparts, especially for the N- and C-terminal domains, but some motifs were absent (Fig. 4). The RGD motif is an integrin recognition site important for IGFBP interaction with cell surface $\alpha 5 \beta 1$ integrin, and is present in mammalian IGFBP-1 and -2 (Jones et al., 1993). This motif is conserved in fish IGFBP-2s but substituted in fish IGFBP-1 (Fig. 4; Maures and Duan, 2002; Kamei et al., 2008; Shimizu et al., 2011a). NLS is an important motif for IGFBP-3 to express IGF-independent action by translocating to the nucleus and acting as a transcription factor (Schedlich et al., 2000). Zebrafish (*Danio rerio*) IGFBP-3 possesses NLS and has been shown to exert its ligand-independent action by antagonizing bone morphogenetic protein (BMP) (Zhong et al., 2011). This motif is conserved in salmon IGFBP-3 but its functionality has not been confirmed (Shimizu et al., 2011b).

Post-translational modification is an important step for proteins to exert their functions. Phosphorylation and N-glycosylation are two major modifications for IGFBPs. Mammalian IGFBP-1 is highly phosphorylated, which may be important for high IGF binding since a non-phosphorylated form had a lower IGF binding ability and failed to suppress IGF action (Jones et al., 1991). A Pro-Glu-Ser-Thr (PEST)-rich domain is responsible for phosphorylation of mammalian IGFBP-1 (Julkunen et al., 1988) and also present in salmon IGFBP-1s (Shimizu et al., 2005; 2011a), suggesting they are phosphorylated although it has not been directly confirmed by phosphoprotein staining. N-glycosylation is characteristic of mammalian IGFBP-3 (Firth and Baxter, 1999). There are three N-glycosylation sites in human IGFBP-3 whereas two sites are found in salmon IGFBP-3 (Fig. 4; Shimizu et al., 2011b). It is of note that human IGFBP-2 has no N-glycosylation sites while salmon IGFBP-2b has three (Fig. 4; Shimizu et al., 2011b). These multiple glycosylations are probably why salmon IGFBP-2b appears as doublet bands at 41- and 43-kDa similar to the size of mammalian IGFBP-3 on electrophoresis gels. Glycosylation of these two bands was confirmed by digestion with endoglycopeptidase of purified salmon IGFBP-2b (Shimizu et al., 2003b). The exact role of N-glycosylation of mammalian IGFBP-3 and salmon IGFBP-2b is unknown, but it may affect interaction with the cell surface (Firth and Baxter, 1999). No N-glycosylation sites are found in salmon IGFBP-1s, but zebrafish IGFBP-1a has one site (Maures and Duan, 2002; Kamei et al., 2008; Shimizu et al., 2011a).

6. Molecular distribution of circulating IGF-I in fish

As mentioned earlier, the ternary complex of IGFBP-3, IGF and ALS is critical for maintaining high concentrations of IGFs in the mammalian circulation. However, there is so far no evidence

of the presence of the ternary complex in the fish circulation. When lamprey plasma was separated by size-exclusion chromatography, IGF-binding activity was detected around 50 kDa (Upton et al., 1993), which was much smaller than the mammalian 150-kDa complex. A similar result was obtained with barramundi (*Lates calcarifer*; Degger et al., 2000). However, it should be noted that under unequilibrated conditions the IGF-binding assay detects mainly unoccupied IGFBPs but may not do so with IGFBPs saturated with endogenous IGFs. Shimizu et al. (1999) fractionated coho salmon serum by size-exclusion chromatography under neutral conditions and measured IGF-I in the eluted fractions. As a result, IGF-I immunoreactivity was detected at approximately 50-kDa (binary complex) and 7.5 kDa (free form), but not at 150 kDa (ternary complex) (Fig. 5). These attempts were unable to detect a high-molecular-weight pool of IGF-I in the fish circulation. However, sequences of ALS are present in fish genome databases and appear to be similar to those of mammalian counterparts (Shimizu, personal communication). The apparent lack of the ternary complex is probably due to the extremely low hepatic expression of *igfbp-3* at least in salmon (Shimizu et al., 2011b). The liver is the major site of production of IGFBP-3 in mammals. Shimizu et al. (2011b) cloned a cDNA of salmon *igfbp-3* from the heart since its expression was very low in the liver. Macqeen et al. (2013) comprehensively analyzed tissue expression levels of 19 salmon *igfbps* and found four *igfbp-3* paralogs had low expression in the liver and also other tissues examined. Such low hepatic expression makes it unlikely that IGFBP-3 is a major circulating IGFBP in salmon. It is not known if ALS circulates in fish blood, but the extremely low levels of production of IGFBP-3 by the liver may be a major reason for the lack of ternary complex in salmon circulation.

The significance and/or reason of the apparent lack of ternary complex in the salmon circulation are not known at present, but the following is a possible explanation why the ternary complex is not formed. An advantage of forming the high-molecular-weight complex of 150 kDa is to prevent IGF from being filtered out by the kidney and from leaving the capillary barrier (Zapf, 1995; Rajaram et al., 1997). The capillary barrier in mammals has a molecular cutoff around 60 kDa, which does not allow the 150 kDa ternary complex to pass through, although the 50 kDa binary complex may be filtered (Hasegawa et al., 1992). For this reason, the ternary complex is critical to form large pools of IGFs in mammalian circulation. In contrast, fish capillaries are relatively "leaky" lacking a clear molecular cutoff (Hargens et al., 1974), so that even if the ternary complex is formed, IGFs may not be sequestered in the circulation. Furthermore, renal glomerular filtration in fish is at least an order of magnitude lower than in mammals, so renal loss of IGF may not be as great in fish as in mammals

(Hickman and Trump, 1969). In addition, mammalian IGFBP-3 is produced by the Kupffer and endothelial cells in the liver. Fish apparently lack Kupffer cells (Robertson and Bradley, 1992; Bruslé and Anadon, 1995), which may be a reason for the low *igfbp-3* expression in the liver. Supporting this, cultured striped bass liver pieces did not secrete a 35-39-kDa IGFBP, but did produce 28-30-kDa and 23-24-kDa IGFBPs (Fukazawa et al., 1995; Siharath et al., 1996). Based on this evidence, it is hypothesized that the apparent lack of the ternary complex in fish may be due to the leaky nature of the vascular system and low renal filtration in fish, which does not provide a selective advantage for the formation of a ternary complex along with the apparent lack of the Kupffer cells in the liver, all of which might prevent IGFBP-3 from being a major circulating IGFBP. The significance of the lack of the ternary complex needs to be addressed by taking account of the difference in the vascular system between mammals and fishes.

7. Hormonal regulation of circulating fish IGFBPs

Kelley et al. (1992) detected IGFBPs in fish blood and demonstrated that they were under control of hormones. Fukazawa et al. (1995) examined in vitro effects of several hormones (GH, insulin, prolactin, glucagon, triiodothyronine, thyroxine, testosterone and estradiol) and growth factors (epidermal growth factor and IGF-I) on the secretion of the two low-molecular-weight IGFBPs from liver pieces of striped bass. These results indicate that many hormones are involved in the regulation of fish IGFBPs.

7.1. Insulin

In mammals, insulin is a potent inhibitor of IGFBP-1 (Unterman et al., 1991). Circulating levels of IGFBP-1 are generally inversely related to those of insulin after meals, and insulin treatment inhibits IGFBP-1 both at the mRNA and protein levels (Lee et al., 1993, 1997). In the goby, an experimental model to create insulin-dependent diabetes mellitus (IDDM) by surgical removal of pancreatic islets (isletectomy) was established (Kelley et al., 1993). The isletectomized goby showed induction of 24- and 30-kDa IGFBPs in plasma, and insulin treatment restored levels of these IGFBPs to basal conditions (Kelley et al., 2001), suggesting that negative regulation by insulin of putative fish IGFBP-1 is conserved at least in this species. In contrast, an in vitro experiment using primary cultured salmon hepatocytes found that insulin had no effect on reducing *igfbp-1b* mRNA and protein (Pierce et al., 2006). On the other hand, there was a weak negative correlation between circulating IGFBP-1b and insulin levels in coho salmon (Shimizu

et al., personal communication). Thus, the inhibitory effect of insulin may be species-specific or an indirect effect in fish.

The effect of insulin on the 40-50-kDa IGFBP is not clear. A study using isletectomized goby showed that a high dose of insulin (1U/kg) increased the intensity of the 40-50-kDa IGFBP band (Kelley et al., 1992), although no statistical analysis was conducted.

7.2. Glucocorticoids

Cortisol is known to induce IGFBP-1 in mammals although its positive effect is secondary to the inhibitory effect of insulin (Katz et al., 1998). In fish, the involvement of cortisol in inducing the low-molecular-weight IGFBPs in blood has been suggested based on the findings that IGFBP levels were increased under catabolic conditions such as fasting, isletectomy, or handling stress when cortisol is also elevated (Kelley et al., 2001; 2002; Peterson and Small, 2004; Davis and Peterson, 2005, 2006). Several studies also provided evidence of a direct induction by cortisol of these smaller IGFBPs in fish (Kajimura et al., 2003; Peterson and Small, 2005; Pierce et al., 2006; Shimizu et al., 2011a). A single cortisol injection of tilapia resulted in a clear induction of 24- and 30-kDa IGFBPs within two hours and its effect lasted for eight hours, but levels returned to baseline by 24 hours (Kajimura et al., 2003). A long-term (four weeks) treatment of channel catfish with dietary cortisol also induced a 20-kDa IGFBP in plasma (Peterson and Small, 2005). A glucocorticoid agonist, dexamethasone, directly stimulated *igfbp-1b* mRNA in salmon hepatocytes in vitro (Pierce et al., 2006). In rainbow trout, IGFBP-1a as well as IGFBP-1b were inducible by exogenous cortisol (Shimizu et al., 2011a). These findings well support the notion that cortisol induces salmon IGFBP-1s and the low-molecular-weight IGFBPs of other fishes. However, Kelley et al. (2000) reported that a 30-kDa IGFBP in the goby was induced by isletectomy but cortisol treatment reduced it in a dose dependent manner. This result suggests an interactive regulation by cortisol and pancreatic hormone(s) of the 30-kDa IGFBP in the goby. Thus, cortisol may have divergent effects on multiple IGFBPs in various tissues.

Cortisol has been found to reduce the plasma 40-50-kDa IGFBP in tilapia (Kajimura et al., 2003). However, it is not known if the same regulation exists in other fish species.

7.3. Growth hormone (GH)

In mammals GH is the major regulator of the components of the ternary complex (i.e. IGF-I,

IGFBP-3 and ALS) (Martin and Baxter, 1992). The liver is the primary site of production of all three components, and GH treatment induces their production, although cellular localization of the components in the liver is different as mentioned earlier. Induction by GH of fish IGFBPs was first reported in coho salmon (Kelley et al., 1992). In striped bass, hypophysectomy was effective in reducing a 35-kDa IGFBP and injecting ovine GH restored its levels (Siharath et al., 1995b). The same approach was used to assess the effect of GH on tilapia IGFBPs (Park et al., 2000). In that study, fish were first hypophysectomized and given homologous GH. Exogenous GH increased a 40-kDa IGFBP, which had been reduced by hypophysectomy (Park et al., 2000). However, the GH effect in pituitary-intact fish was less clear. In a study of tilapia by Breves et al. (2014) injection of GH increased hepatic expression of *igfbp-2b*, suggesting that GH induces hepatic production of the 40-kDa IGFBP in intact tilapia. GH treatment of intact coho salmon increased 41-kDa IGFBP (IGFBP-2b) levels as measured by radioimmunoassay (RIA) (Shimizu et al., 2003a). In channel catfish (*Ictalurus punctatus*), GH injection reduced plasma levels of 44- and 47-kDa IGFBPs, which goes along with other atypical responses to GH treatment in channel catfish such as increased body fat and reduced body protein (Johnson et al., 2003). However, long-term GH treatment of the same species had no significant effect on a 45-kDa IGFBP (Peterson et al., 2004). It should be noted that in none of these fish species has 40-50-kDa IGFBPs been proven to be IGFBP-3 orthologs. On the contrary, salmon 41-kDa IGFBP was identified as IGFBP-2b (Shimizu et al., 2011b). In humans, GH is weakly inhibitory to IGFBP-2 (Blum et al., 1993), which contrasts to the GH effect on salmon IGFBP-2b. In this respect, a major target of GH has diversified between mammals and salmon.

The effect of GH on other IGFBPs in fish is not consistent. GH injection had no effect on the striped bass 23-24-kDa IGFBP, reduced the tilapia 20-kDa IGFBP and the catfish 35-kDa IGFBP, or increased the tilapia 29- and 32-kDa IGFBPs (Siharath et al., 1995b; Park et al., 2000; Johnson et al., 2003). These results suggest that GH does not act directly on the lower-molecular-weight IGFBPs, and its effect is secondary through modulating other hormones such as insulin and/or cortisol. In contrast, in zebrafish GH injection decreased whole body mRNA levels of *igfbp-2*, which corresponds to IGFBP-2a with molecular weight of 31 kDa when expressed in CHO-K1 cells (Duan et al., 1999).

7.4. Sex steroids

Only a few studies examined effects of sex steroids on circulating IGFBPs in fish (Fukuzawa et al., 1995; Larsen et al., 2004), but the IGFBP responses at the mRNA level have been

comprehensively evaluated in rainbow trout (Cleveland and Weber, 2015). Fukuzawa et al. (1995) examined in vitro effects of testosterone and estradiol-17 β on the secretion of the low-molecular-weight IGFBPs from the liver pieces of striped bass and found that estradiol-17 β at 5 ng/ml significantly increased the secretion of two IGFBPs, whereas testosterone at the same concentration had no effect. The striped bass 35-39 kDa IGFBP was undetectable in the cultured media (Fukazawa et al., 1995; Siharath et al., 1995a). On the other hand, in vivo injection of testosterone or 11-ketotestosterone significantly increased circulating IGFBP-2b levels in postsmolt coho salmon (Larsen et al., 2004), although responses of IGFBP-1a or -1b were not examined in that study.

8. Environmental and developmental regulation of circulating fish IGFBPs

Circulating fish IGFBPs are presumably important for adjusting growth under changing environments through regulating the availability of IGFs to target tissues. Accordingly, circulating levels of IGFBPs are controlled by environmental factors (Picha et al., 2008a).

8.1. Feeding and nutrition

Circulating levels of fish IGFBPs are affected by nutritional status, but their responses differ among the IGFBP types in terms of direction and/or magnitude of change. The 20-25 kDa IGFBP including salmon IGFBP-1b is relatively sensitive to food deprivation or feeding ration. Fasting of striped bass for 30 days induced a 25-kDa IGFBP in plasma, and refeeding for two weeks reduced it to undetectable levels (Siharath et al., 1996). Similar responses of the 20-25-kDa IGFBP have been reported using ligand blotting in coho salmon, goby, channel catfish and Atlantic salmon (*Salmo salar*) (Shimizu et al., 1999; Kelley et al., 2001; Peterson and Small, 2004; Hevrøy et al., 2011). The availability of an RIA for salmon 22-kDa IGFBP (IGFBP-1b) made it possible to process a large number of samples with greater measurement precision (Shimizu et al., 2006). When postsmolt coho salmon were reared under different feeding rations (1.75, 1.0, 0.5 and 0% body weight/day), IGFBP-1b levels as measured by RIA were graded by the ration being highest in the lowest ration (Shimizu et al., 2006). Changes in feeding ration influenced circulating IGFBP-1b levels within two weeks (Shimizu et al., 2006). Moreover, a significant reduction in IGFBP-1b was observed by four hours after a meal in postsmolt coho salmon (Shimizu et al., 2009). In mammals, IGFBP-1 shows a dynamic change within a day in response to a meal (Yeoh and Baxter, 1988). Salmon IGFBP-1b is similar to the mammalian counterpart by showing a significant fluctuation within a day (Shimizu et al., 2009).

The 28-32-kDa IGFBP, including salmon IGFBP-1a, is also inducible by fasting in some species, but its response is not as sensitive as the 20-25-kDa form. In the goby, a 30-kDa IGFBP was induced by fasting and reduced by re-feeding, which was in parallel with a 24-kDa form (Kelley et al., 2001). On the other hand, fasting of channel catfish up to 45 days had no effect on a 35-kDa IGFBP (Peterson and Small, 2004). In Chinook salmon, IGFBP-1a was not induced after six weeks of fasting (Shimizu et al., 2011a), whereas fasting of masu salmon (*O. masou*) for four weeks resulted in increases of both IGFBP-1a and -1b (Fig. 6; Kawaguchi et al., 2013). These findings suggest that IGFBP-1 sensitivity to fasting may be species-specific.

The response of the 40-50-kDa IGFBP including salmon IGFBP-2b appears to be opposite to what is seen for the 20-24-kDa IGFBP; the 40-50-kDa levels are high when fish are fed, and fasting decreases its levels. However, its response to fasting is sometimes unclear partly because this IGFBP often appears as diffused bands on ligand blotting, making it difficult to quantify. For example, fasting of striped bass for 60 days tended to decrease a 35-39-kDa IGFBP, but the response was not significant (Siharath et al., 1996). On the other hand, in a hybrid striped bass (*M. chrysops* X *M. saxatilis*) a 40-kDa IGFBP, presumably corresponding to the 35-39 kDa form in striped bass, significantly decreased in fasted fish in 20 days; it recovered to fed control levels after re-feeding for 20 days (Picha et al., 2008b). With the development of an RIA for salmon 41-kDa IGFBP (IGFBP-2b), it became clear in salmon that IGFBP-2b is sensitive to nutritional conditions including fasting and feeding ration (Shimizu et al., 2003a, 2009; Beckman et al., 2004a,b). IGFBP-2b responded to fasting as early as four days in Chinook salmon and a single meal increased it within several hours in fasted coho salmon (Pierce et al., 2005; Shimizu et al., 2009). In addition, the trout 47-kDa IGFBP increased when fish were fed diets with increasing percentage of plant proteins (0, 50, 75 or 100%) (Gomez-Requeni et al., 2005). In juvenile olive flounder (*Paralichthys olivaceus*), dietary supplementation of glycoprotein extracts from the sea mustard (*Hizikia fusiformis*) increased a plasma 43-kDa IGFBP while decreasing a 34-kDa IGFBP (Choi et al., 2014). It is of note that in some fish species the 40-50-kDa IGFBP is undetectable, which could partly account for considerably lower IGF-I levels in fish compared to mammals (Kelley et al., 2000, 2001, 2002, 2006).

There are a number of studies looking at responses of multiple *igfbp* mRNAs in the liver and/or white muscle to fasting and refeeding in fish (Gabillard et al., 2006; Bower et al., 2008; Pedroso et al., 2009; Bower and Johnston, 2010; Safian et al., 2012; Cleveland and Weber, 2014; Garcia de la serrana et al., 2017). Fasting of adult zebrafish increases whole body

mRNA levels of an *igfbp-2a* (Duan et al., 1999). Fasting and refeeding of rainbow trout showed significant changes in *igfbp* expression, including in liver increases in *igfbp-2*, *-4*, and *-6* after refeeding, and in muscle increases in *igfbp-2*, *-4*, and *-5* (Gabillard et al. 2006). An in vitro approach to a fasting and refeeding experiment is removal of amino acids and subsequent addition in cultured Atlantic salmon myotubes (Bower and Johnston, 2010). This approach showed an increase in *igfbp-5* expression in myotube cells due to amino acid addition alone and an increase in *igfbp-4* expression when amino acid addition was combined with IGF or insulin addition (Bower and Johnston, 2010). These studies indicate that hepatic and muscle IGFBPs are regulated by nutrition and emphasize that local IGFBPs play significant roles in regulating muscle growth in response to changes in nutritional status.

8.2. Stress

Stress is a strong inducer of the two low-molecular-weight IGFBPs in the fish circulation (Kelley et al., 2000, 2001, 2002, 2006). Hypophysectomy of tilapia induced a 20-kDa IGFBP in the circulation, and levels of a 29-kDa IGFBP were increased (Park et al., 2000). These increases should be due, at least in part, to the stress associated with surgery. Kelley et al. (2001) pointed out that although both 24- and 30-kDa IGFBPs of jack mackerel (*Trachurus symmetricus*) were induced by handling stress, the 30-kDa form was more sensitive than the 24-kDa form. In contrast, a direct transfer of Chinook salmon parr, which had low hypoosmoregulatory capacity, to full-strength seawater resulted in a strong induction of IGFBP-1b, but not IGFBP-1a, at six hours after transfer (Shimizu et al., 2011a). In the same experiment, IGFBP-1a was induced at 12 h after transfer (Shimizu et al., 2011a). A 15-min low-water stress at 25°C caused increases in both 24- and 30-kDa IGFBPs in sunshine bass, a hybrid between female white bass (*M. chrysops*) and male striped bass (Davis and Peterson, 2006). These studies suggest that the relative sensitivity of the two low-molecular-weight IGFBPs vary among species or type of stress employed. Supporting this, in rainbow trout an acute handling stress (five min handling) had no effect on 21- and 32-kDa IGFBPs (Shepherd et al., 2011).

The 40-50-kDa IGFBP appears to be less sensitive to stress. An acute handling stress tended to increase circulating 42- and 50-kDa IGFBPs in rainbow trout (Shepherd et al., 2011). An osmotic stress to Chinook salmon parr had no effect on circulating IGFBP-2b levels when assessed by RIA (Shimizu et al., 2011a). More work need to be done to reveal stress effects on the 40-50-kDa IGFBP.

8.3. Temperature

Temperature affects all biological processes as well as specific components of the GH/IGF/IGFBP system in poikilothermic fish (Gabillard et al., 2005). Several studies examined effects of decreased or increased water temperature on IGFBPs and suggested that responses were different among IGFBP types. When postsmolt coho salmon were reared under two water temperatures (11°C or 7°C) in combination with feeding ration (1.8% or 1.0%/body weight), a drop in water temperature from 11°C to 7°C decreased plasma IGFBP-1b levels in postsmolt coho salmon regardless of feeding ration in the first two weeks (Shimizu et al., 2006). However, four weeks after a water temperature change, IGFBP-1b levels in fish at 11°C became lower than those at 7°C and the effect of the water temperature change disappeared by the ninth week (Shimizu et al., 2006). In the same experiment, however, the change in water temperature alone had no effect on IGFBP-2b levels but masked the effect of decreased feeding ration (Beckman et al., 2004a). These findings suggest that IGFBP-1b is more sensitive than IGFBP-2b to decreasing water temperature. Hevrøy et al. (2013) examined effects of elevated water temperatures in Atlantic salmon in seawater and found that a 32-kDa IGFBP showed the highest levels at 17°C but further increase in water temperature to 19°C restored it to the basal levels. Another study by the same group showed that an elevated water temperature at 19°C decreased plasma IGFBP-1b in Atlantic salmon whereas higher levels of IGFBP-1b were seen in rainbow trout, suggesting these two species handled elevated temperature differently (Hevrøy et al., 2015). In channel catfish, a higher water temperature (26°C versus 20°C) increased a 19-kDa IGFBP (Johnson et al., 2003). In sunshine bass a decrease in water temperature from 23°C to 5°C suppressed plasma 33-kDa IGFBP levels whereas 24- and 28-kDa IGFBPs were unchanged (Davis and Peterson, 2006). These studies indicate contrasting regulation by water temperature change for different types of IGFBPs in fish, although pathways by which temperature affects IGFBPs are unknown. Since fish growth is largely affected by water temperature, its direct and indirect effects on circulating IGFBPs need to be comprehensively explored.

8.4. Salinity

Salinity influences the GH-IGF-I system in euryhaline fishes such as salmon. However, relatively little is known about the effect of salinity on fish IGFBPs. Shepherd et al. (2005) were the first to examine salinity effects on IGFBPs. A gradual acclimation of rainbow trout from freshwater to 66‰ seawater (22 ppt) resulted in elevated IGFBPs at 21, 32, 42 and 50 kDa sizes (Shepherd et al., 2005). When postsmolt coho salmon were transferred from freshwater to 50‰

seawater (15 ppt), IGFBP-2b transiently increased one day after transfer and then returned to levels similar to freshwater controls (Shimizu et al., 2007). In the same study, GH treatment was found to be more effective in fish in 50% seawater presumably due to a lowered glomerular filtration rate so that exogenous GH was retained longer in the circulation and stimulated IGFBP-2b to a greater extent (Shimizu et al., 2007).

8.5. Developmental/seasonal effects (smoltification)

Smoltification is a transitional process for juvenile salmon by which river-dwelling parr become ocean-type smolt (Hoar, 1988; Stefansson et al., 2008; Björnsson et al., 2011; McCormick, 2013). Smolt acquire hypoosmoregulatory ability during smoltification through activating gill Na^+, K^+ -ATPase (NKA) prior to the seawater entry. Smoltification is a seasonal event generally occurring in spring, although some strains undergo smoltification in autumn. Many endocrine axes including the GH-IGF-I system are activated during smoltification (Dickhoff et al., 1997; Stefansson et al., 2008; Björnsson et al., 2011; McCormick, 2013). It is thus not surprising that IGFBPs change during this period. However, a limited number of studies examined changes in circulating IGFBPs in salmon. In coho salmon, plasma IGFBP-1b levels showed a peak at the end of April when condition factor was decreasing (Shimizu et al., 2006). A negative relationship between plasma IGFBP-1b and condition factor is generally found in salmon, suggesting that this form is involved in catabolism (Shimizu et al., 2006). In addition, serum IGFBP-1b levels in smolting masu salmon were positively correlated with gill NKA activity (Fukuda et al., 2015). During smoltification, circulating IGF-I in smolting salmon are generally high and both growth and hypo-osmoregulatory ability are promoted (Dickhoff et al., 1997; Stefansson et al., 2008; Björnsson et al., 2011; McCormick, 2013). Thus, circulating IGF-I is most likely adequately distributed to different organs involved in growth or osmoregulation. Circulating IGFBP-1b may be involved in delivering IGF-I to the gills (Fukuda et al., 2015). Activity of the endocrine IGF-I at the gills in turn may be regulated by local IGFBPs. A recent work by Breves et al. (2017) reported changes in local *igfbp-6bs* in the gills and suggested their importance in the development of hypoosmoregulatory ability.

Plasma IGFBP-2b levels also changed during smoltification of coho salmon reaching a peak at the end of March, one month earlier than the peak of IGFBP-1b (Shimizu et al., 2003a, 2006). In the same fish, plasma IGF-I showed two peaks in late March and late April corresponding to peaks of IGFBP-2b and -1b, respectively (Shimizu et al., 2003a, 2006). These results suggest that IGFBPs play different roles during smoltification, although unraveling the

role of each IGFBP is subject to future study.

9. Perspective

Our better understanding of the fish IGFBPs invites speculation on how this system evolved and how it compares functionally to the well-characterized system in mammals. The evolution of IGFBP genes has been elegantly described in a number of studies proposing that an ancestral IGFBP gene was duplicated in chordates and was followed by a number of whole genome duplications; two whole-genome duplications in ancestral vertebrates, followed by a third whole-genome duplication leading to teleosts, and a fourth whole-genome duplication in some fish species, e.g., salmonids (Daza et al., 2011, Macqueen et al., 2013). Thus, the Atlantic salmon has 19 IGFBP genes, which are 13 more than found in humans (Macqueen et al., 2013). Therefore we should be cautious in extrapolating the functions of IGFBPs in mammals to those in fish because there is an approximately 500 million year span involving a lot of genetic change since their evolutionary divergence. However, this complexity in IGFBP evolution offers a great opportunity for future study of the range of functions (subfunctional partitioning) of IGFBPs in fish and how they compare with mammals. Subfunction partitioning is one of the fates of duplicated copies of a gene where ancestral regulatory and structural subfunctions are preserved by gene duplicates (Postlethwait et al., 2004). Subfunction partitioning in turn provides gene duplicates opportunities to acquire new function and/or regulation. Interestingly, the earliest IGFBP function may have been independent of binding IGF (Zhou et al., 2013; Zhong and Duan, 2017).

Studies of IGFBP functions in fish have made good use of the zebrafish model. Zebrafish IGFBPs are generally inhibitory to IGF-induced cell proliferation (Duan et al., 1999). Knocking down IGFBPs resulted in abnormal organ formation (Li et al., 2005; Wood et al., 2005b). In addition, while IGFBP-1 delayed the speed of embryonic development under hypoxic conditions (Kajimura et al., 2005), such a response limits oxygen consumption due to IGF-induced anabolism and may be adaptive to increase embryo survival. Functions of duplicated zebrafish IGFBPs are overlapping, but there are certain differences such as affinity for IGFs, site of production, developmental changes and responses to fasting (Kamei et al., 2008; Zhou et al., 2008; Wang et al., 2009; Dai et al., 2010). The fish circulation appears to contain duplicated IGFBP-1s where they may play overlapping yet distinct roles in regulating postnatal growth. However, there are few attempts examining functions of circulating fish IGFBPs in the context of gene duplication. This is most likely due to the lack of purified

proteins. Plasma/serum is a source for protein purification, however levels of circulating IGFBPs are low, e.g., < 300 ng/ml (Shimizu et al., 2003a, 2006). Thus, producing recombinant IGFBPs is desirable for functional analyses of fish IGFBPs. Recombinant salmon IGFBP-1a, -1b, -2a and -2b are currently being produced by using bacterial expression systems (Tanaka et al., in press). Although recombinant proteins produced in bacteria are not glycosylated or phosphorylated, the availability of recombinant fish IGFBPs should promote functional analyses.

A series of studies of circulating salmon IGFBPs suggest that IGFBP-3 plays little role in regulating endocrine IGFs (Shimizu et al., 1999, 2003a, 2009, 2011b). However, whether or not the finding on salmon IGFBP-3 applies to other fishes is unknown. In zebrafish, since IGFBP-3 plays a crucial role in embryonic development and exhibits an IGF-independent action (Li et al., 2005; Zhong et al., 2011), it is important for normal development at least in this species. In tilapia and yellowtail (*Seriola quinqueradiata*), GH treatment increased *igfbp-3* mRNA in the liver (Cheng et al., 2002; Pedroso et al., 2009), suggesting it is secreted into the bloodstream and modulates IGF action. Local action of IGFBP-3 is also suggested in fine flounder (*Paralichthys adspersus*; Safian, et al., 2012). The stage-, tissue- or species-specific roles of fish IGFBP-3 need to be examined in future studies.

In view of the large number of IGFBP genes that have been retained in fish, the fish circulation probably contains more than three IGFBPs. There is some evidence for this to be the case. For example, in coho salmon there are two additional bands often detected at 37 and 31 kDa, and the 31-kDa IGFBP appeared to be decreased by fasting (Fig. 7). In order to compare regulation and function of fish IGFBPs with mammalian counterparts, the fish proteins need to be identified. One strategy to clone cDNAs of unidentified IGFBPs is to use degenerate primer(s) designed from partial amino acid sequences of purified proteins. However, as mentioned above, purifying each IGFBP from serum/plasma is not practical. Instead, partial N-terminal amino acid sequence could be obtained by IGF-affinity chromatography of serum/plasma followed by electrophoresis and MALDI-TOF MS/MS (Matrix Assisted Laser Desorption/Ionization-Time of Flight Mass Spectrometry/Mass Spectrometry). Once a cDNA is cloned, identification and recombinant production of an IGFBP are possible. Moreover, IGFBPs can be identified from genome sequences when they are available, thus avoiding the need to clone cDNAs. In the case of salmonids, protein-coding sequences of all 19 *igfbp* paralogs in 10 species are now available (Macqueen et al., 2013; Lappin et al., 2016), which provides a very useful reference to assign unidentified circulating salmon IGFBPs and accelerate functional

studies on IGFBP paralogs.

The evidence for salmon IGFBPs suggests that salmon IGFBP-2 diverged in its function from that of its mammalian counterpart, and convergently acquired characters similar to mammalian IGFBP-3 (Fig. 8). Roch et al. (2009) stated that studies on duplicate hormones and receptors were vulnerable to misidentification if only structural similarity was used. Salmon IGFBP-2b is a good example for such case and highlights the importance of actually testing the function to unravel the evolutionary fate of duplicated IGFBPs. Moreover, both salmon IGFBP-1a and -1b are increased under catabolic conditions but may have different sensitivity, suggesting they underwent subfunction partitioning (Fig. 8). Thus, salmon as well as other fish species provide a unique opportunity to investigate how functional divergence, convergence and subfunction partitioning of IGFBPs occurred during vertebrate evolution.

Acknowledgments

Much of the work on circulating salmon IGFBPs reviewed here were conducted at Northwest Fisheries Science Center, NOAA Fisheries, Seattle WA. We thank our collaborators in Seattle for their great help and stimulating discussions: Brian R. Beckman, Penny Swanson, Donald A. Larsen, Jon Dickey, Andrew L. Pierce, and Haruhisa Fukada. We also thank Akihiko Hara for his help in protein analyses and antibody production. Our research on salmon IGFBPs was supported by funding from US Department of Agriculture, NOAA Fisheries, and Japan Society for the Promotion of Science.

References

- Anderson, T.A., Bennett, L.R., Conlon, M.A., Owens, P.C., 1993. Immunoreactive and receptor-active insulin-like growth factor-I (IGF-I) and IGF-binding protein in blood plasma from the freshwater fish *Macquaria ambigua* (golden perch). *J. Endocrinol.* 136, 191-198.
- Bauchat, J.R., Busby, W.H., Jr., Garmong, A., Swanson, P., Moore, J., Lin, M., Duan, C., 2001. Biochemical and functional analysis of a conserved IGF-binding protein isolated from rainbow trout (*Oncorhynchus mykiss*) hepatoma cells. *J. Endocrinol.* 170, 619-628.
- Baxter, R.C., Martin, J.L., 1989. Structure of the Mr 140,000 growth hormone-dependent insulin-like growth factor binding protein complex: determination by reconstitution and affinity-labeling. *Proc. Natl. Acad. Sci. USA* 86, 6898-6902.
- Beckman, B.R., Shimizu, M., Gadberry, B.A., Cooper, K.A., 2004a. Response of the

somatotropic axis of juvenile coho salmon to alterations in plane of nutrition with an analysis of the relationships among growth rate and circulating IGF-I and 41 kDa IGFBP. Gen. Comp. Endocrinol. 135, 334-344.

Beckman, B.R., Shimizu, M., Gadberry, B.A., J., P.P., Cooper, K.A., 2004b. The effect of temperature change on the relations among plasma IGF-I, 41-kDa IGFBP, and growth rate in postsmolt coho salmon. Aquaculture 241, 601-619.

Binoux, M., Hossenlopp, P., 1988. Insulin-like growth factor (IGF) and IGF-binding proteins: Comparison of human serum and lymph. J. Clin. Endocrinol. Metab. 67, 509-514.

Björnsson, B.T., Stefánsson, S.O., McCormick, S.D., 2011. Environmental endocrinology of salmon smoltification. Gen. Comp. Endocrinol. 170, 290-298.

Blum, W.F., Horn, N., Kratzsch, J., Jorgensen, J.O., Juul, A., Teale, D., Mohnike, K., Ranke, M.B., 1993. Clinical studies of IGFBP-2 by radioimmunoassay. Growth Regul. 3, 100-104.

Bower, N.I., Jonnston, I.A., 2010. Transcriptional regulation of the IGF signaling pathway by amino acids and insulin-like growth factors during myogenesis in Atlantic salmon. PLoS ONE 5, 1-14.

Bower, N.I., Li, X., Taylor, R., Johnston, I.A., 2008. Switching to fast growth: the insulin-like growth factor (IGF) system in skeletal muscle of Atlantic salmon. J. Exp. Biol. 211, 3859-3870.

Breves, J.P., Tipsmark, C.K., Stough, B.A., Seale, A.P., Flack, B.R., Moorman, B.P., Laerner, D.T., Grau, E.G., 2014. Nutritional status and growth hormone regulate insulin-like growth factor binding protein (igfbp) transcripts in Mozambique tilapia. Gen. Comp. Endocrinol. 207, 66-73.

Breves, J.P., Fujimoto, C.K., Phipps-Costin, S.K., Einarsdottir, I.E., Björnsson, B.T., McCormick, S.D., 2017. Variation in branchial expression among insulin-like growth-factor binding proteins (igfbps) during Atlantic salmon smoltification and seawater exposure. BMC physiol. 17, 2.

Bruslé, J., Anadon, G.I., 1995. The structure and function of fish liver, in: Munshi, J.S.D., Dutta, H.M. (Eds.), Fish Morphology. Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi, pp. 77-93.

Cleveland, B.M., Weber, G.M., 2014. Ploidy effects on genes regulating growth mechanisms during fasting and refeeding in juvenile rainbow trout (*Oncorhynchus mykiss*). Mole. Cell. Endocrinol., 382, 139-149.

702 Cleveland, B.M., Weber, G.M., 2015. Effects of sex steroids on expression of genes regulating
 703 growth-related mechanisms in rainbow trout (*Oncorhynchus mykiss*). Gen. Comp.
 704 Endocrinol. 216, 103-115.

705 Cheng, R., Chang, K.-M., Wu, J.-L., 2002. Different temporal expressions of tilapia
 706 (*Oreochromis mossambicus*) insulin-like growth factor-I and IGF binding protein-3 after
 707 growth hormone induction. Mar. Biotechnol. 4, 218-225.

708 Chin, E., Zhou, J., Dai, J., Baxter, R.C., Bondy, C.A., 1994. Cellular localization and regulation
 709 of gene expression for components of the insulin-like growth factor ternary binding
 710 protein complex. Endocrinology 134, 2498-2504.

711 Choi, Y.H., Kim, K.W., Han, H.S., Nam, T.J., Lee, B.J., 2014. Dietary Hizikia fusiformis
 712 glycoprotein-induced IGF-I and IGFBP-3 associated to somatic growth, polyunsaturated
 713 fatty acid metabolism, and immunity in juvenile olive flounder *Paralichthys olivaceus*.
 714 Comp. Biochem. Physiol. A 167, 1-6.

715 Collet, C., Candy, J., Sara, V., 1998. Evolutionary aspects of the IGF system, in: Takano, K.,
 716 Hizuka, N., Takahashi, S.-I. (Eds.), Molecular Mechanism to Regulate the Activities of
 717 Insulin-like Growth Factors. Elsevier Science B.V., Amsterdam, Netherland, pp. 215-
 718 223.

719 Dai, W., Kamei, H., Zhao, Y., Ding, J., Du, Z., Duan, C., 2010. Duplicated zebrafish insulin-
 720 like growth factor binding protein-5 genes with split functional domains: evidence for
 721 evolutionarily conserved IGF binding, nuclear localization, and transactivation activity.
 722 FASEB J. 24, 2020-2029.

723 Daughaday, W.H., Rotwein, P., 1989. Insulin-like growth factors I and II. Peptide, messenger
 724 ribonucleic acid and gene structures, serum, and tissue concentrations. Endocr. Rev. 10,
 725 68-91.

726 Davis, K.B., Peterson, B.C., 2005. Comparison of insulin-like growth factor-I and insulin-like
 727 growth factor binding protein concentrations of the palmetto and sunshine bass and the
 728 effects of gender and stress. J. World Aquacult. Soc. 36, 384-392.

729 Davis, K.B., Peterson, B.C., 2006. The effect of temperature, stress, and cortisol on plasma
 730 IGF-I and IGFBPs in sunshine bass. Gen. Comp. Endocrinol. 149, 219-225.

731 Daza, D.O., Sundstrom, G., Bergqvist, C.A., Duan, C.M., Larhammar, D., 2011. Evolution of
 732 the insulin-like growth factor binding protein (IGFBP) family. Endocrinology 152, 2278-
 733 2289.

734 Degger, B., Upton, Z., Soole, K., Collet, C., Richardson, N., 2000. Comparison of recombinant

735 barramundi and human insulin-like growth factor (IGF)-I in juvenile barramundi (*Lates*
736 *calcarifer*): in vivo metabolic effects, association with circulating IGF-binding proteins,
737 and tissue localisation. Gen. Comp. Endocrinol. 117, 395-403.

738 D'Ercole, A.J., Applewhite, G.T., Underwood, L.E., 1980. Evidence that somatomedin is
739 synthesized by multiple tissues in the fetus. Dev. Biol. 75, 315-328.

740 Dickhoff, W.W., Beckman, B.R., Larsen, D.A., Duan, C., Moriyama, S., 1997. The role of
741 growth in endocrine regulation of salmon smoltification. Fish. Physiol. Biochem. 17, 231-
742 236.

743 Duan, C., 1997. The insulin-like growth factor system and its biological actions in fish. Am.
744 Zool. 37, 491-503.

745 Duan, C., Ding, J., Li, Q., Tsai, W., Pozios, K., 1999. Insulin-like growth factor binding protein
746 2 is a growth inhibitory protein conserved in zebrafish. Proc. Natl. Acad. Sci. USA 96,
747 15274-15279.

748 Firth, S.M., Baxter, R.C., 1999. Characterisation of recombinant glycosylation variants of
749 insulin-like growth factor binding protein-3. J. Endocrinol. 160, 379-387.

750 Firth, S.M., Baxter, R.C., 2002. Cellular actions of the insulin-like growth factor binding
751 proteins. Endocr. Rev. 23, 824-854.

752 Forbes, B.E., McCarthy, P., Norton, R.S., 2012. Insulin-like growth factor binding proteins: a
753 structural perspective. Front. Endocrinol. 3, 38.

754 Frystyk, J., Skjaerbaek, C., Dinesen, B., Orskov, H., 1994. Free insulin-like growth factors
755 (IGF-I and IGF-II) in human serum. FEBS Lett. 348, 185-191.

756 Fukazawa, Y., Siharath, K., Iguchi, T., Bern, H.A., 1995. In vitro secretion of insulin-like
757 growth factor-binding proteins from liver of striped bass, *Morone saxatilis*. Gen. Comp.
758 Endocrinol. 99, 239-247.

759 Fukuda, M., Kaneko, N., Kawaguchi, K., Hevrøy, E.M., Hara, A., Shimizu, M., 2015.
760 Development of a time-resolved fluoroimmunoassay for salmon insulin-like growth
761 factor binding protein-1b. Comp. Biochem. Physiol. A 187, 66-73.

762 Gabillard, J.C., Kamanger, B.B., Montserrat, N. 2006. Coordinated regulation of the GH/IGF
763 system genes during refeeding in rainbow trout (*Oncorhynchus mykiss*). J. Endocrinol.
764 191, 15-24.

765 Gabillard, J.C., Weil, C., Rescan, P.Y., Navarro, I., Gutierrez, J., Le Bail, P.Y., 2005. Does the
766 GH/IGF system mediate the effect of water temperature on fish growth? A review.
767 Cybium 29, 107-117.

768 Garcia de la serrana, D., Fuentes, E.N., Martin, A.M.M., Johnston, I.A., Macqueen, D.J., 2017.
 769 Divergent regulation of insulin-like growth factor binding protein genes in cultured
 770 Atlantic salmon myotubes under different models of catabolism and anabolism. Gen.
 771 Comp. Endocrinol. 247, 53-65.

772 Gomez-Requeni, P., Calduch-Giner, J., Vega-Rubin de Celis, S., Medale, F., Kaushik, S.J.,
 773 Perez-Sanchez, J., 2005. Regulation of the somatotrophic axis by dietary factors in
 774 rainbow trout (*Oncorhynchus mykiss*). Br. J. Nutr. 94, 353-361.

775 Grulich-Henn, J., Spiess, S., Heinrich, U., Schonberg, D., Bettendorf, M., 1998. Ligand blot
 776 analysis of insulin-like growth factor-binding proteins using biotinylated insulin-like
 777 growth factor-I. Horm. Res. 49, 1-7.

778 Guler, H.P., Zapf, J., Froesch, E.R., 1987. Short-Term Metabolic Effects of Recombinant
 779 Human Insulin-Like Growth Factor-I in Healthy-Adults. New Engl. J. Med. 317, 137-
 780 140.

781 Guler, H.P., Zapf, J., Schmid, C., Froesch, E.R., 1989. Insulin-like growth factors I and II in
 782 healthy man. Estimations of half-lives and production rates. Acta Endocrinologica 121,
 783 753-758.

784 Hargens, A.R., Millard, R.W., Johansen, K., 1974. High capillary permeability in fishes. Comp.
 785 Biochem. Physiol. 48A, 675-680.

786 Hasegawa, Y., Cohen, P., Yorgin, P., Rosenfeld, R.G., 1992. Characterization of urinary
 787 insulin-like growth-factor binding-proteins. J. Clin. Endocrinol. Metab. 74, 830-835.

788 Hevrøy, E.M., Azpeleta, C., Shimizu, M., Lanzen, A., Kaiya, H., Espe, M., Olsvik, P.A., 2011.
 789 Effects of short-term starvation on ghrelin, GH-IGF system, and IGF-binding proteins in
 790 Atlantic salmon. Fish Physiol. Biochem. 37, 217-232.

791 Hevrøy, E.M., Hunsaker, C., de Gelder, S., Shimizu, M., Waagbo, R., Breck, O., Takle, H.,
 792 Sussort, S., Hansen, T., 2013. GH-IGF system regulation of attenuated muscle growth
 793 and lipolysis in Atlantic salmon reared at elevated sea temperatures. J. Comp. Physiol. B
 794 183, 243-259.

795 Hevrøy, E.M., Tipsmark, C.K., Remo, S.C., Hansen, T., Fukuda, M., Torgersen, T., Vikesa, V.,
 796 Olsvik, P.A., Waagbo, R., Shimizu, M., 2015. Role of the GH-IGF-1 system in Atlantic
 797 salmon and rainbow trout postsmolts at elevated water temperature. Comp. Biochem.
 798 Physiol. A 188, 127-138.

799 Hickman, C.P., Trump, B.F. 1969. The Kidney, in: Hoar, W.S., Randall, D.J. (eds). Fish
 800 Physiology, Academic Press NY, pp.91-239.

801 Hoar, W.S., 1988. The physiology of smolting salmonids, *Fish Physiology*, pp. 275-343.

802 Hossenlopp, P., Seurin, D., Segovia-Quinson, B., Hardouin, S., Binoux, M., 1986. Analysis of
803 serum insulin-like growth factor binding proteins using western blotting: use of the
804 method for titration of the binding proteins and competitive binding studies. *Anal.*
805 *Biochem.* 154, 138-143.

806 Hwa, V., Oh, Y., Rosenfeld, R.G., 1999. The insulin-like growth factor-binding protein
807 (IGFBP) superfamily. *Endocr. Rev.* 20, 761-787.

808 Johnson, J., Silverstein, J., Wolters, W.R., Shimizu, M., Dickhoff, W.W., Shepherd, B.S., 2003.
809 Disparate regulation of insulin-like growth factor-binding proteins in a primitive,
810 ictalurid, teleost (*Ictalurus punctatus*). *Gen. Comp. Endocrinol.* 134, 122-130.

811 Jones, J.I., D'Ercole, A.J., Camacho-Hubner, C., Clemmons, D.R., 1991. Phosphorylation of
812 insulin-like growth factor (IGF)-binding protein 1 in cell culture and in vivo: effects on
813 affinity for IGF-I. *Proc. Natl. Acad. Sci. USA* 88, 7481-7485.

814 Jones, J.I., Gockerman, A., Busby, W.H., Jr., Wright, G., Clemmons, D.R., 1993. Insulin-like
815 growth factor binding protein 1 stimulates cell migration and binds to the alpha 5 beta 1
816 integrin by means of its Arg-Gly-Asp sequence. *Proc. Natl. Acad. Sci. USA* 90, 10553-
817 10557.

818 Julkunen, M., Koistinen, R., Aalto-Setälä, K., Seppälä, M., Janne, O.A., Kontula, K., 1988.
819 Primary structure of human insulin-like growth factor-binding protein/placental protein
820 12 and tissue-specific expression of its mRNA. *FEBS Lett.* 236, 295-302.

821 Kajimura, S., Duan, C., 2007. Insulin-like growth factor-binding protein-1: an evolutionarily
822 conserved fine tuner of insulin-like growth factor action under catabolic and stressful
823 conditions. *J. Fish Biol.* 71, 309-325.

824 Kajimura, S., Hirano, T., Visitacion, N., Moriyama, S., Aida, K., Grau, E.G., 2003. Dual mode
825 of cortisol action on GH/IGF-I/IGF binding proteins in the tilapia, *Oreochromis*
826 *mossambicus*. *J. Endocrinol.* 178, 91-99.

827 Kajimura, S., Aida, K., Duan, C., 2005. Insulin-like growth factor-binding protein-1 (IGFBP-1)
828 mediates hypoxia-induced embryonic growth and developmental retardation. *Proc. Natl.*
829 *Acad. Sci. USA* 102, 1240-1245.

830 Kamei, H., Lu, L., Jiao, S., Li, Y., Gyrup, C., Laursen, L.S., Oxvig, C., Zhou, J., Duan, C.,
831 2008. Duplication and diversification of the hypoxia-inducible IGFBP-1 gene in
832 zebrafish. *PLoS One* 3, e3091.

833 Katz, L.E., Satin-Smith, M.S., Collett-Solberg, P., Baker, L., Stanley, C.A., Cohen, P., 1998.

834 Dual regulation of insulin-like growth factor binding protein-1 levels by insulin and
835 cortisol during fasting. J. Clin. Endocrinol. Metab. 83, 4426-4430.

836 Kawaguchi, K., Kaneko, N., Fukuda, M., Nakano, Y., Kimura, S., Hara, A., Shimizu, M., 2013.
837 Responses of insulin-like growth factor (IGF)-I and two IGF-binding protein-1 subtypes
838 to fasting and re-feeding, and their relationships with individual growth rates in yearling
839 masu salmon (*Oncorhynchus masou*). Comp. Biochem. Physiol. A 165, 191-198.

840 Kelley, K.M., Siharath, K., Bern, H.A., 1992. Identification of insulin-like growth factor-
841 binding proteins in the circulation of four teleost fish species. J Exp. Zool 263, 220-224.

842 Kelley, K.M., 1993. Experimental diabetes mellitus in a teleost fish. I. Effect of complete
843 isletectomy and subsequent hormonal treatment on metabolism in the goby, *Gillichthys*
844 *mirabilis*. Endocrinology 132, 2689-2695.

845 Kelley, K.M., Desai, P., Roth, J.T., Haigwood, J.T., Arope, S.A., Flores, R.M., Schmidt, K.E.,
846 Perez, M., Nicholson, G.S., Song, W.W., 2000. Evolution of endocrine growth
847 regulation: the insulin like growth factors (IGFs), their regulatory binding proteins
848 (IGFBPs), and IGF receptors in fishes and other ectothermic vertebrates, in: Fingerman,
849 M., Nagabhushanam, R. (Eds.), Recent Advances in Marine Biotechnology, Aquaculture,
850 Part B Fishes 4. Science Publishers, Plymouth, UK, pp. 189-228.

851 Kelley, K.M., Haigwood, J.T., Perez, M., Galima, M.M., 2001. Serum insulin-like growth factor
852 binding proteins (IGFBPs) as markers for anabolic/catabolic condition in fishes. Comp.
853 Biochem. Physiol. B 129, 229-236.

854 Kelley, K.M., Schmidt, K.E., Berg, L., Sak, K., Galima, M.M., Gillespie, C., Balogh, L.,
855 Hawayek, A., Reyes, J.A., Jamison, M., 2002. Comparative endocrinology of the insulin-
856 like growth factor-binding protein. J. Endocrinol. 175, 3-18.

857 Kelley, K.M., Price, T.D., Galima, M.M., Sak, K., Reyes, J.A., Zepeda, O., Hagstrom, R.,
858 Truong, T.A., Lowe, C.G., 2006. Insulin-like growth factor-binding proteins (IGFBPs) in
859 fish: beacons for (disrupted) growth endocrine physiology, in: Reinecke, M., Zacccone, G.,
860 Kapoor, B.G. (Eds.), Fish Endocrinology. Science Publishers, Enfield, New Hampshire,
861 pp. 167-195.

862 Kim, H.S., Nagalla, S.R., Oh, Y., Wilson, E., Roberts, C.T., Rosenfeld, R.G., 1997.
863 Identification of a family of low-affinity insulin-like growth factor binding proteins
864 (IGFBPs): Characterization of connective tissue growth factor as a member of the IGFBP
865 superfamily. Proc. Natl. Acad. Sci. USA 94, 12981-12986.

866 Lappin, F.M., Shaw, R.L., Macqueen, D.J., 2016. Targeted sequencing for high-resolution

867 evolutionary analyses following genome duplication in salmonid fish: Proof of concept
 868 for key components of the insulin-like growth factor axis. *Mar. Genom.* 30, 15-26.

869 Larsen, D.A., Shimizu, M., Cooper, K.A., Swanson, P., Dickhoff, W.W., 2004. Androgen
 870 effects on plasma GH, IGF-I, and 41-kDa IGFBP in coho salmon (*Oncorhynchus*
 871 *kisutch*). *Gen. Comp. Endocrinol.* 139, 29-37.

872 Lee, P.D., Conover, C.A., Powell, D.R., 1993. Regulation and function of insulin-like growth
 873 factor-binding protein-1. *Proc. Soc. Exp. Biol. Med.* 204, 4-29.

874 Lee, P.D., Giudice, L.C., Conover, C.A., Powell, D.R., 1997. Insulin-like growth factor binding
 875 protein-1: recent findings and new directions. *Proc. Soc. Exp. Biol. Med.* 216, 319-357.

876 LeRoith, D., Bondy, C., Yakar, S., Liu, J.L., Butler, A., 2001. The somatomedin hypothesis:
 877 2001. *Endocr. Rev.* 22, 53-74.

878 Li, Y., Xiang, J., Duan, C., 2005. Insulin-like growth factor-binding protein-3 plays an
 879 important role in regulating pharyngeal skeleton and inner ear formation and
 880 differentiation. *J. Biol. Chem.* 280, 3613-3620.

881 Macqueen, D.J., de la Serrana, D.G., Johnston, I.A., 2013. Evolution of ancient functions in the
 882 vertebrate insulin-Like growth factor system iuncovered by study of duplicated salmonid
 883 fish genomes. *Mol. Biol. Evol.* 30, 1060-1076.

884 Martin, J.L., Baxter, R.C., 1992. Insulin-like growth factor binding protein-3: biochemistry and
 885 physiology. *Growth Regul.* 2, 88-99.

886 Maures, T.J., Duan, C., 2002. Structure, developmental expression, and physiological regulation
 887 of zebrafish IGF binding protein-1. *Endocrinology* 143, 2722-2731.

888 McCormick, S.D., 2013. Smolt physiology and endocrinology, in: McCormick, S.D., Farrell,
 889 A.P., Brauner, C.J. (Eds.), *Euryhaline Fishes*. Academic Press, Oxford, UK, pp. 199-251.

890 Niu, P.-D., Le Bail, P.-Y., 1993. Presence of insulin-like growth factor binding protein (IGF-
 891 BP) in rainbow trout (*Oncorhynchus mykiss*) serum. *J. Exp. Zool.* 265, 627-636.

892 Niu, P.-D., Perez-Sanchez, J., Le Bail, P.-Y., 1993. Development of a protein binding assay for
 893 teleost insulin-like growth factor (IGF)-like: relationships between growth hormone (GH)
 894 and IGF-like in the blood of rainbow trout (*Oncorhynchus mykiss*). *Fish Physiol.*
 895 *Biochem.* 11, 381-391.

896 Ohlsson, C., Mohan, S., Sjogren, K., Tivesten, A., Isgaard, J., Isaksson, O., Jansson, J.O.,
 897 Svensson, J., 2009. The role of liver-derived insulin-like growth factor-I. *Endocr. Rev.*
 898 30, 494-535.

899 Park, R., Shepherd, B.S., Nishioka, R.S., Grau, E.G., Bern, H.A., 2000. Effects of homologous

900 pituitary hormone treatment on serum insulin-like growth-factor-binding proteins
 901 (IGFBPs) in hypophysectomized tilapia, *Oreochromis mossambicus*, with special
 902 reference to a novel 20-kDa IGFBP. Gen. Comp. Endocrinol. 117, 404-412.

903 Pedroso, F.L., Fukada, H., Masumoto, T., 2009. In vivo and in vitro effect of recombinant
 904 salmon growth hormone treatment on IGF-I and IGFBPs in yellowtail *Seriola*
 905 *quinqueradiata*. Fish. Sci. 75, 887-894.

906 Peterson, B.C., Small, B.C., 2004. Effects of fasting on circulating IGF-binding proteins,
 907 glucose, and cortisol in channel catfish (*Ictalurus punctatus*). Domest. Anim. Endocrinol.
 908 26, 231-240.

909 Peterson, B.C., Small, B.C., 2005. Effects of exogenous cortisol on the GH/IGF-I/IGFBP
 910 network in channel catfish. Domest. Anim. Endocrinol. 28, 391-404.

911 Peterson, B.C., Small, B.C., Bosworth, B.G., 2004. Effects of bovine growth hormone (Posilac
 912 (R)) on growth performance, body composition, and IGFBPs in two strains of channel
 913 catfish. Aquaculture 232, 651-663.

914 Picha, M.E., Turano, M.J., Beckman, B.R., Borski, R.J., 2008a. Endocrine biomarkers of
 915 growth and applications to aquaculture: A minireview of growth hormone, insulin-like
 916 growth factor (IGF)-I, and IGF-Binding proteins as potential growth indicators in fish. N.
 917 Am. J. Aquacult. 70, 196-211.

918 Picha, M.E., Turano, M.J., Tipsmark, C.K., Borski, R.J., 2008b. Regulation of endocrine and
 919 paracrine sources of Igfs and Gh receptor during compensatory growth in hybrid striped
 920 bass (*Morone chrysops* X *Morone saxatilis*). J. Endocrinol. 199, 81-94.

921 Pierce, A.L., Shimizu, M., Beckman, B.R., Baker, D.M., Dickhoff, W.W., 2005. Time course of
 922 the GH/IGF axis response to fasting and increased ration in chinook salmon
 923 (*Oncorhynchus tshawytscha*). Gen. Comp. Endocrinol. 140, 192-202.

924 Pierce, A.L., Shimizu, M., Felli, L., Swanson, P., Dickhoff, W.W., 2006. Metabolic hormones
 925 regulate insulin-like growth factor binding protein-1 mRNA levels in primary cultured
 926 salmon hepatocytes; lack of inhibition by insulin. J. Endocrinol. 191, 379-386.

927 Postlethwait, J., Amores, A., Cresko, W., Singer, A., Yan, Y.L., 2004. Subfunction partitioning,
 928 the teleost radiation and the annotation of the human genome. Trends Genet. 20, 481-490.

929 Rajaram, S., Baylink, D.J., Mohan, S., 1997. Insulin-like growth factor-binding proteins in
 930 serum and other biological fluids: regulation and functions. Endocr. Rev. 18, 801-831.

931 Reinecke, M., Collet, C., 1998. The phylogeny of the insulin-like growth factors. Int. Rev.
 932 Cytol. 183, 1-94.

933 Robertson, J.C., Bradley, T.M., 1992. Liver ultrastructure of juvenile Atlantic salmon (*Salmo*
 934 *salar*). J. Morph. 211, 41-54.
 935 Roch, G.J., Wu, S., Sherwood, N.M., 2009. Hormones and receptors in fish: do duplicates
 936 matter? Gen. Comp. Endocrinol. 161, 3-12.
 937 Rodgers, B.D., Roalson, E.H., Thompson, C., 2008. Phylogenetic analysis of the insulin-like
 938 growth factor binding protein (IGFBP) and IGFBP-related protein gene families. Gen.
 939 Comp. Endocrinol. 155, 201-207.
 940 Safian, D., Fuentes, E.N., Valdes, J.A., Molina, A., 2012. Dynamic transcriptional regulation of
 941 autocrine/paracrine igfbp1, 2, 3, 4, 5, and 6 in the skeletal muscle of the fine flounder
 942 during different nutritional statuses. J. Endocrinol. 214, 95-108.
 943 Salmon, W.D., Jr., Daughaday, W.H., 1957. A hormonally controlled serum factor which
 944 stimulates sulfate incorporation by cartilage in vitro. J. Lab. Clin. Med. 49, 825-836.
 945 Schedlich, L.J., Le Page, S.L., Firth, S.M., Briggs, L.J., Jans, D.A., Baxter, R.C., 2000. Nuclear
 946 import of insulin-like growth factor-binding protein-3 and -5 is mediated by the importin
 947 beta subunit. J. Biol. Chem. 275, 23462-23470.
 948 Schoen, T.J., Beebe, D.C., Clemmons, D.R., Chader, G.J., Waldbillig, R.J., 1992. Local
 949 synthesis and developmental regulation of avian vitreal insulin-like growth factor-binding
 950 proteins-a model for independent regulation in extravascular and vascular compartments.
 951 Endocrinology 131, 2846-2854.
 952 Shepherd, B.S., Aluru, N., Vijayan, M.M., 2011. Acute handling disturbance modulates plasma
 953 insulin-like growth factor binding proteins in rainbow trout (*Oncorhynchus mykiss*).
 954 Domest. Anim. Endocrinol. 40, 129-138.
 955 Shepherd, B.S., Drennon, K., Johnson, J., Nichols, J.W., Playle, R.C., Singer, T.D., Vijayan,
 956 M.M., 2005. Salinity acclimation affects the somatotropic axis in rainbow trout. Am. J.
 957 Physiol. Regul. Integr. Comp. Physiol. 288, R1385-1395.
 958 Shimasaki, S., Ling, N., 1991. Identification and molecular characterization of insulin-like
 959 growth factor binding proteins (IGFBP-1, -2, -3, -4, -5 and -6). Prog. Growth Factor Res.
 960 3, 243-266.
 961 Shimizu, M., Swanson, P., Dickhoff, W.W., 1999. Free and protein-bound insulin-like growth
 962 factor-I (IGF-I) and IGF-binding proteins in plasma of coho salmon, *Oncorhynchus*
 963 *kisutch*. Gen. Comp. Endocrinol. 115, 398-405.
 964 Shimizu, M., Swanson, P., Fukada, H., Hara, A., Dickhoff, W.W., 2000. Comparison of
 965 extraction methods and assay validation for salmon insulin-like growth factor-I using

commercially available components. Gen. Comp. Endocrinol. 119, 26-36.

Shimizu, M., Hara, A., Dickhoff, W.W., 2003a. Development of an RIA for salmon 41 kDa IGF-binding protein. J. Endocrinol. 178, 275-283.

Shimizu, M., Swanson, P., Hara, A., Dickhoff, W.W., 2003b. Purification of a 41-kDa insulin-like growth factor binding protein from serum of chinook salmon, *Oncorhynchus tshawytscha*. Gen. Comp. Endocrinol. 132, 103-111.

Shimizu, M., Dickey, J.T., Fukada, H., Dickhoff, W.W., 2005. Salmon serum 22 kDa insulin-like growth factor-binding protein (IGFBP) is IGFBP-1. J. Endocrinol. 184, 267-276.

Shimizu, M., Beckman, B.R., Hara, A., Dickhoff, W.W., 2006. Measurement of circulating salmon IGF binding protein-1: assay development, response to feeding ration and temperature, and relation to growth parameters. J. Endocrinol. 188, 101-110.

Shimizu, M., Fukada, H., Hara, A., Dickhoff, W.W., 2007. Response of the salmon somatotrophic axis to growth hormone administration under two different salinities. Aquaculture 273, 320-328.

Shimizu, M., Cooper, K.A., Dickhoff, W.W., Beckman, B.R., 2009. Postprandial changes in plasma growth hormone, insulin, insulin-like growth factor (IGF)-I, and IGF-binding proteins in coho salmon fasted for varying periods. Am. J. Physiol. Regul. Integr. Comp. Physiol. 297, R352-361.

Shimizu, M., Kishimoto, K., Yamaguchi, T., Nakano, Y., Hara, A., Dickhoff, W.W., 2011a. Circulating salmon 28- and 22-kDa insulin-like growth factor binding proteins (IGFBPs) are co-orthologs of IGFBP-1. Gen. Comp. Endocrinol. 174, 97-106.

Shimizu, M., Suzuki, S., Horikoshi, M., Hara, A., Dickhoff, W.W., 2011b. Circulating salmon 41-kDa insulin-like growth factor binding protein (IGFBP) is not IGFBP-3 but an IGFBP-2 subtype. Gen. Comp. Endocrinol. 171, 326-331.

Siharath, K., Kelley, K.M., Bern, H.A., 1996. A low-molecular-weight (25-kDa) IGF-binding protein is increased with growth inhibition in the fasting striped bass, *Morone saxatilis*. Gen. Comp. Endocrinol. 102, 307-316.

Siharath, K., Nishioka, R.S., Bern, H.A., 1995a. In vitro production of IGF-binding proteins (IGFBP) by various organs of the striped bass, *Morone saxatilis*. Aquaculture 135, 195-202.

Siharath, K., Nishioka, R.S., Madsen, S.S., Bern, H.A., 1995b. Regulation of IGF-binding proteins by growth hormone in the striped bass, *Morone saxatilis*. Mol. Mar. Biol. Biotech. 4, 171-178.

999 Stefansson, S.O., Björnsson, B.T., Ebbesson, L.O.E., McCormick, S.D., 2008. Smoltification,
 1000 in: Finn, R.N., Kapoor, B.G. (Eds.), Fish Larval Physiology. Science Publishers, Enfield,
 1001 NH, pp. 639-681.

1002 Tanaka, H., Oishi, G., Nakano, Y., Mizuta, H., Nagano, Y., Hiramatsu, N., Ando, H., Shimizu,
 1003 M. Production of recombinant salmon insulin-like growth factor binding protein-1
 1004 subtypes. Gen. Comp. Endocrinol. (in press).

1005 Unterman, T.G., Oehler, D.T., Murphy, L.J., Lacson, R.G., 1991. Multihormonal regulation of
 1006 insulin-like growth factor-binding protein-1 in rat H4IIE hepatoma cells: the dominant
 1007 role of insulin. Endocrinology 128, 2693-2701.

1008 Upton, Z., Chan, S.J., Steiner, D.F., Wallace, J.C., Ballard, F.J., 1993. Evolution of insulin-like
 1009 growth factor binding proteins. Growth Regul. 3, 29-32.

1010 Villafuerte, B.C., Koop, B.L., Pao, C.I., Gu, L., Birdsong, G.G., Phillips, L.S., 1994. Coculture
 1011 of primary rat hepatocytes and nonparenchymal cells permits expression of insulin-like
 1012 growth factor binding protein-3 in vitro. Endocrinology 134, 2044-2050.

1013 Walton, P.E., Baxter, R.C., Burleigh, B.D., Etherton, T.D., 1989. Purification of the serum acid-
 1014 stable insulin-like growth factor binding protein from the pig (*Sus scrofa*). Comp.
 1015 Biochem. Physiol. 92B, 561-567.

1016 Wang, X., Lu, L., Li, Y., Li, M., Chen, C., Feng, Q., Zhang, C., Duan, C., 2009. Molecular and
 1017 functional characterization of two distinct IGF binding protein-6 genes in zebrafish. Am.
 1018 J. Physiol. Regul. Integr. Comp. Physiol. 296, R1348-1357.

1019 Wheatcroft, S.B., Kearney, M.T., 2009. IGF-dependent and IGF-independent actions of IGF-
 1020 binding protein-1 and-2: implications for metabolic homeostasis. Trends Endocrinol.
 1021 Metab. 20, 153-162.

1022 Wood, A.W., Duan, C., Bern, H.A., 2005a. Insulin-like growth factor signaling in fish. Int. Rev.
 1023 Cytol. 243, 215-285.

1024 Wood, A.W., Schlueter, P.J., Duan, C., 2005b. Targeted knockdown of insulin-like growth
 1025 factor binding protein-2 (IGFBP-2) disrupts cardiovascular development in zebrafish
 1026 embryos. Mol. Endocrinol. 19, 1024-1034.

1027 Yeoh, S.I., Baxter, R.C., 1988. Metabolic regulation of the growth hormone independent
 1028 insulin-like growth factor binding protein in human plasma. Acta Endocrinologica 119,
 1029 465-473.

1030 Zapf, J., 1995. Physiological role of the insulin-like growth factor binding proteins. Eur. J.
 1031 Endocrinol. 132, 645-654.

1032 Zapf, J., Jagars, G., Sand, I., Froesch, E.R., 1978. Evidence for the existence in human serum of
 1033 large molecular weight nonsuppressible insulin-like activity (NSILA) different from the
 1034 small molecular weight forms. *FEBS Lett.* 90, 135-140.
 1035 Zhong, Y., Duan, C., 2017. Lamprey IGF-binding protein-3 has IGF-dependent and -
 1036 independent actions. *Front. Endocrinol.* 7:174.
 1037 Zhong, Y., Lu, L., Zhou, J., Li, Y., Liu, Y., Clemmons, D.R., Duan, C., 2011. IGF binding
 1038 protein 3 exerts its ligand-independent action by antagonizing BMP in zebrafish embryos.
 1039 *J. Cell Sci.* 124, 1925-1935.
 1040 Zhou, J., Li, W., Kamei, H., Duan, C., 2008. Duplication of the IGFBP-2 gene in teleost fish:
 1041 protein structure and functionality conservation and gene expression divergence. *PLoS*
 1042 *One* 3, e3926.
 1043 Zhou, J., Xiang, J., Zhang, S., Duan, C., 2013. Structural and functional analysis of the
 1044 amphioxus IGFBP gene uncovers ancient origin of IGF-independent functions.
 1045 *Endocrinology* 154, 3753-3763.

Figure legends

Fig. 1. Three forms of circulating IGFs in humans. The majority of circulating IGF is complexed with IGFBP-3 and an acid-labile subunit (ALS). IGFBP-5 also participates in the formation of the ternary complex. This ternary complex is too large to leave the vascular system so that a large circulating pool of IGF is formed. IGF is also bound to other IGFBPs to form binary complexes. The binary complexes can cross the capillary barrier, but the availability of IGF to the receptor is limited by IGFBP. Less than 1% of IGF is in the free form and biologically active.

Fig. 2. Ligand blotting using labeled IGF-I of human and Chinook salmon sera. IGFBP bands were visualized by using digoxigenin (DIG)-labeled human IGF-I and anti-DIG conjugated with horseradish peroxidase. Migration positions of human and Chinook salmon IGFBPs are indicated by arrows on the left and right, respectively. Molecular weights (kDa) of Chinook salmon IGFBPs are indicated. NS: non-specific bands.

Fig. 3. Schematic patterns of IGFBP bands in human and fish sera/plasma on ligand blotting using labeled IGF-I. Migration positions of circulating IGFBPs in human, chicken, Chinook salmon, rainbow trout, catfish, striped bass, tilapia, goby and lamprey are reproduced from Hossenlopp et al. (1986), Schoen et al. (1992), Niu et al. (1993), Johnson et al. (2003), Siharath et al. (1995), Park et al. (2000), Kelley et al. (1992) and Upton et al. (1993), respectively. Types of IGFBP are indicated by numbers and alphabets when identity is known.

Fig. 4. Alignment of deduced amino acid sequences of Chinook salmon IGFBP-1a, -1b, -2a, -2b and -3 with those of human counterparts. Sequences are from NP_000587, NP_000588, NP_000589, AEO18300.1, AAV83995.1, AEC33109.1, AEC33110.1 and AEC33113.1. Asterisks indicate cysteine residues conserved among the IGFBP family. Underlines and boxes indicate a Arg-Gly-Asp (RGD) integrin recognition motif and potential N-glycosylation sites, respectively. The starting positions of the N-, L- and C-domains of human IGFBPs are shown by triangles.

Fig. 5. Elution profiles of serum IGF-I in coho salmon on gel filtration under neutral conditions. Arrows indicate expected elution positions of ternary, binary and free forms of IGF-I. Modified from Shimizu et al. (1999) with permission.

1080

1081 Fig. 6. IGFBP patterns in sera of fed, fasted and refed masu salmon. Serum samples were the
1082 same as used in Kawaguchi et al. (2013). Postsmolt masu salmon were fed or fasted for 6
1083 weeks. An additional group (Re-fed) was fasted for 4 weeks followed by 2 weeks of refeeding.
1084 Arrows indicate migration positions of salmon IGFBPs.

1085

1086 Fig. 7. IGFBP patterns in plasma of fed and fasted coho salmon. Postsmolt coho salmon were
1087 fed or fasted for 3 weeks. Arrows indicate migration positions of identified and unidentified
1088 (New?) IGFBPs. NS: non-specific bands.

1089

1090 Fig. 8. Hypothetical functional relationships between human and salmon IGFBPs in the
1091 circulation.

Shimizu, Fig. 1

Shimizu, Fig. 2

Human BP1	-----MS-----EVPVARVWLVLVLLLTQVVG-VTAGAPWQCAPCSAEKLALC--PP	43
Salmon BP1a	-----MSGLFHRNVLVAAAVCCSVLVRSVLQEPVLAQEPVIRCAPCSPEKLSEC--PA	50
Salmon BP1b	-----MLGLYKK-LTLAAMSLSLLTTLAQSSPVVGPPIRCAPCTQEKLDEC--PA	49
Human BP2	MLPRVGC PALPLPPPPLPLPLLLLLLGGSGGGGARA EVLFRCPPCTPERLAACGPPR	60
Salmon BP2a	MTRRS-----TPRMISYSGCSLLLLS-VAFVGASFAEMVFRCPCTAERQAAC--PK	49
Salmon BP2b	-----MVLYFSCGLFLLTLLVLPGLLLGDLVFYCPKCTAERQTAC--PK	42
Human BP3	-----MQRARPTLWAAALTLVLLRGPPVARAGASSGGLGPVVRCEPCDARALAQCAPPP	55
Salmon BP3	-----MPGLCVLCLTAVLAA----FTRFAET---VGPVVRCEPCDAGALMECKPLP	44
	* * *	
Human BP1	VS-----ASCSEVTRSAGCGCCPMCALPLGAACGVATARCARGLS CRALPGE	90
Salmon BP1a	VA-----PGCAEVLREPGCGCLACALKTGDLGCIYTAPCGSGLRCTPRPGD	97
Salmon BP1b	IS-----PDCKQVLRPGCGCCMACALEKGASCGVYTAHCAQGLKCS PRAGD	96
Human BP2	VAPPAVA AVAGGARMPCAE LVREPGCGCCSVCARLEGEACGVYTPRCGQGLRCYPHPGS	120
Salmon BP2a	LT-----ETCAEIVREPGCGCCPV CARQEGELCGVYTPRCSSGLRCYPKPD	96
Salmon BP2b	LA-----TNC TEIVREPACGCPVCARLEGEFCGVYTPRCSTGLRCYPTVDS	89
Human BP3	AV-----CAELVREPGCGCLTCALSEGQPCGIYTERCGSGLRCQPSDE	100
Salmon BP3	KD-----CAERVREPGCGCLSCALAEQACGVYTGRCGSGLICQFQPG	89
	* * * * *	
Human BP1	QQPLHALTRGQACVQESD---ASAPH-----AAEAGSPES-----	123
Salmon BP1a	LRPLHSLTRGQAVCTEIPVSSSVSQ-----NPDQGAADNA-----	134
Salmon BP1b	PRPLHSLTRGQAICTE-----DQG-----	115
Human BP2	ELPLQALVMGEGTCEKRRDAEY GASPEQVAD-----NGDDHSEGLVENHVDST	169
Salmon BP2a	DLPLEQLVQGLGLCGHKVVTPTGSGE-----HREKLSGEVVDVLDTS	139
Salmon BP2b	KLPLEQLVQGLGRCSQKVDTVFNRTTEE-----HRDT-SGELPG-----	126
Human BP3	ARPLQALLDGRGLCVNNA SAVSRLRAYLLPAP---PAPGNAS-ESEEDRSAGSVESPSVSS	156
Salmon BP3	TRPLQALLEGRGACS-SAASKLNTFLFPVQKQETTSGEHSGAGDERANNGTVTTTKTV	148
	*	
Human BP1	-----PESTEITEEELLDNFHLMAPSEEDHSILWDAISTYDGS KALHV	166
Salmon BP1a	-----ETENTAMVSDSGSSLYLHGHSKPFDPRAAADALESMK-AKVNAI	177
Salmon BP1b	-----QEKVEGVDPDHSSLAYFLGLNTPFDTKNEG-AQESIK-AKVNTI	156
Human BP2	MN-----MLGGGSAGRKPLKSGMKELAVFREKVTEQHRQMGKGKHHGLGLEPKK	220
Salmon BP2a	L-----TEIPPLRKATKDN-PWLGPKENAMRQHRREMKTMKMSNK-PEDPKT	184
Salmon BP2b	-----TEGPTMKKPTKDVRIWIWSKDMAPKQAQNELKTKMKTNNCPEEPKT	172
Human BP3	-----THRVS DPKFHPLHSKII I IKKGHAKDSQRYKVDYESQ---STDTQNFSS	203
Salmon BP3	GGAVGVEGGGGGHRGAIEAKPPLHTKLDVIKKEQNKKSQSYKVESVSGGVSSDMHNFS	208
	* * *	
Human BP1	► C-domain	
Human BP1	TNIKKWKEPCRIELYRVVESLAKA----QETS GEEISKFYLPNCNKNNGFYHSRQCETSMD	222
Salmon BP1a	RKKLVEQGPCHVELQRALEKIAKS----QQLGDKLIRFYLPNC DKHGLYKAKQCESSLD	233
Salmon BP1b	RKKLVEQGPCHIELHAALDKITSS----QQLGEKFTNFYLPNC DKHGFYKAKQCESSLV	212
Human BP2	LRPPPARTPCQQELDQVLERISTMRLPDERGPLEHLYSLHIPNC DKHGLYNLQCKMSLN	280
Salmon BP2a	PRG--KQIQCQQELDQVLERISKMPFRDNRGPLEDLALHIPNC DMRGQYNLQCKMSLN	242
Salmon BP2b	QQP--MKGPCAQELEKVMEEISKMSFHDNRGHVDNLYQLKFPNCEKIGQYNLQCKMSTH	230
Human BP3	SKRETEYGPCCRREMEDTLNHLKFLN-----VLSPRGVHIPNC DKKGFYKKKQCRPSKG	256
Salmon BP3	NKRETEYGPCCRREME SILNSLKISN-----VLNPRGFRIPNC DKKGFYKKKQCRPSKG	261
	* * *	
Human BP1	GEAGLCWCVPWNGKRIPGSPEIRGDPNCQIYFNVQN-----	259
Salmon BP1a	GQKGRWCWVSFWNGKKILGSTDLGDAECAYEINH-----	268
Salmon BP1b	GPHARCWCVSSWNGKKILGSNYLPG-LECQLEL-----	244
Human BP2	GQRGECWCVPNPTGKLIQGA PTIRGDP ECHLFYNEQQEARGVDTQRMQ-----	328
Salmon BP2a	GQRGECWCVPNPHXGRPI SAPTVRGDPNCSQYLRGPEMDTLVSAQK-----	288
Salmon BP2b	GQRGECWCVPNPTGVQIAQSTKVRGDPNCSQYVEEQEMETGTQSTAVLQMAEI	283
Human BP3	RKRGFWCVDKYGQPLPGYT TTKGKEDVHCYSMQSK-----	291
Salmon BP3	RKRGYCWCVDKYGQPLPGYDGKERGESQCNNLENK-----	296
	* *

Shimizu, Fig. 7

Shimizu and Dickhoff, Fig. 8