



Title	ON OBSTACLE PROBLEM FOR MEAN CURVATURE FLOW WITH DRIVING FORCE
Author(s)	GIGA, YOSHIKAZU; TRAN, HUNG V.; ZHANG, LONGJIE
Citation	Hokkaido University Preprint Series in Mathematics, 1116, 1-26
Issue Date	2018-10-11
DOI	10.14943/85798
Doc URL	http://hdl.handle.net/2115/71653
Type	bulletin (article)
File Information	OnObstacleProblem.pdf



[Instructions for use](#)

ON OBSTACLE PROBLEM FOR MEAN CURVATURE FLOW WITH DRIVING FORCE

YOSHIKAZU GIGA, HUNG V. TRAN, AND LONGJIE ZHANG

ABSTRACT. In this paper, we study an obstacle problem associated with the mean curvature flow with constant driving force. Our first main result concerns interior and boundary regularity of the solution. We then study in details the large time behavior of the solution and obtain the convergence result. In particular, we give full characterization of the limiting profiles in the radially symmetric setting.

1. INTRODUCTION

In this paper, we study an obstacle problem for level-set forced mean curvature flow equation. We assume further that the surface evolution is described by the mean curvature with constant driving force A . Under the assumption, the equation is

$$(1.1) \quad u_t = |Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) + A|Du|, \quad \text{in } \mathbb{R}^N \times (0, T),$$

$$(1.2) \quad u(x, 0) = u_0(x), \quad \text{on } \mathbb{R}^N,$$

$$(1.3) \quad \psi^-(x) \leq u(x, t) \leq \psi^+(x), \quad \text{on } \mathbb{R}^N \times [0, T].$$

Here $T > 0$, $\operatorname{supp} \psi^\pm \subset \bar{\Omega}$ for some open, bounded, smooth domain $\Omega \subset \mathbb{R}^N$. The given positive constant A is called driving force. Moreover we assume $u_0 \in C^{1,1}(\mathbb{R}^N)$, and $\psi^- \leq u_0 \leq \psi^+$ on \mathbb{R}^N , where u_0, ψ^\pm are all L -Lipschitz continuous for some fixed $L > 0$. In this paper, by $C^{1,1}(\mathbb{R}^N)$ we mean the space of all functions on \mathbb{R}^N whose first derivatives are Lipschitz continuous. For $U \subset \mathbb{R}^N$, we say that a function $a : U \rightarrow \mathbb{R}$ is L -Lipschitz continuous if

$$|a(x) - a(y)| \leq L|x - y| \quad \text{for all } x, y \in U.$$

We postpone the definition of viscosity solutions of (1.1), (1.2), and (1.3) until Section 2.

1.1. Main results. Here we give our main results.

2010 *Mathematics Subject Classification.* 35A01, 35A02, 35K55, 53C44.

Key words and phrases. mean curvature flow; driving force; obstacle problem; large time behavior; limiting profiles; radially symmetric setting.

The work of the first author is partly supported by the Japan Society for the Promotion of Science (JSPS) through the grants No. 26220702 (Kiban S), No. 17H01091 (Kiban A), No. 16H03948 (Kiban B) and No. 18H05323 (Kaitaku). The work of the second author is partly supported by NSF grant DMS-1615944. The third author is the Research Fellow of Japan Society for the Promotion of Science, Number: 17J05160. Corresponding author: Longjie Zhang.

Theorem 1.1. *Let u be the viscosity solution of (1.1), (1.2), and (1.3). Then*

$$|u(x, t) - u(y, t)| \leq L|x - y|,$$

and

$$|u(x, t) - u(x, s)| \leq M|t - s|,$$

for all $x, y \in \mathbb{R}^N$, $t, s \in (0, T)$. Here M is a constant such that

$$M > 2 \left\| D^2 u_0 \right\|_{L^\infty(\mathbb{R}^N)} + AL.$$

As we mention in Section 2 more precisely, the existence of a unique viscosity solution is known by [8]. However, such a global estimate is new although the proof for Lipschitz bound is an adjustment of the proof without obstacles; see [3] for the spatial bound.

Theorem 1.2. *Let u be the viscosity solution of (1.1), (1.2), and (1.3). There exists a L -Lipschitz function v in \mathbb{R}^N such that*

$$u(\cdot, t) \rightarrow v, \quad \text{as } t \rightarrow \infty,$$

uniformly in \mathbb{R}^N . Here v is a stationary solution of (1.1), and (1.3).

A similar result is proved for the Neumann problem in a convex domain for the level-set mean curvature flow equation [7] without obstacles. We shall adjust their proof for our setting.

We aim at characterizing the limiting profile v in term of given initial condition u_0 , and obstacles ψ^\pm . This is a challenging task, and at this moment, we are able to get full characterization in the radially symmetric case. This is done by a careful study of radial solution of (1.1)–(1.3). A key observation is that (1.1) becomes a first order equation with singularity at the center of radial symmetricity. Here is our statement.

Theorem 1.3. *Let $\Omega = B_R(O)$, and*

$$\psi^+(x) = \begin{cases} \lambda \text{dist}(x, \partial\Omega) = \lambda(R - |x|), & x \in \Omega, \\ 0, & x \in \mathbb{R}^N \setminus \Omega, \end{cases}$$

where $\lambda > 0$ is given.

Assume $\psi^- < 0$ in Ω , and $\psi^-(x)$ is radial, that is, it depends only on $|x|$. Moreover, assume that u_0 is radial, and $\psi^- \leq u_0 \leq \psi^+$. The following holds.

(1) If $R \leq (N - 1)/A$, then $u(\cdot, t) \rightarrow 0$ uniformly in \mathbb{R}^N , as $t \rightarrow \infty$.

(2) If $R > (N-1)/A$, then for $B = \max_{(N-1)/A \leq |x| \leq R} u_0 \geq 0$,

$u(\cdot, t) \rightarrow \psi_B$ uniformly in \mathbb{R}^N , as $t \rightarrow \infty$, where, for $C \in \mathbb{R}$,

$$\psi_C := \psi^+ \wedge C = \min\{\psi^+, C\}.$$




FIGURE 1. Case (2) of Theorem 1.3

Remark 1.4. We have the following observations.

(1) Obviously, $\psi^+(x)$ is radial symmetric.

(2) In Theorem 1.3, noting that $u_0 = 0$ on $\partial B_R(O)$, and $u_0 \leq \psi^+$, we have $0 \leq B \leq \lambda(N-1)/A$.

(3) We will see that, for all $0 \leq C \leq \lambda(N-1)/A$, ψ_C are all (viscosity) subsolutions of (1.1) (without obstacle). Moreover, ψ_C are all (viscosity) radially symmetric stationary solutions of (1.1), and (1.3).

(4) We consider

$$(1.1e) \quad 0 = |Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) + A|Du|, \quad \text{in } \Omega,$$

the corresponding elliptic problem of (1.1). It is not hard to see that the comparison principle for (1.1e) does not hold as the equation (1.1e) is not monotone in u . Indeed, obviously 0 is a solution of (1.1e). As mentioned in (3), for all $0 \leq C \leq \lambda(N-1)/A$, ψ_C are (viscosity) subsolutions of (1.1e) with $\psi_C = 0$ on $\partial\Omega$.

1.2. Motivation and Background. In 1994, Sternberg and Ziemer [13] consider the following problem

$$\begin{cases} u_t - |Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) = 0, & (x, t) \in \Omega \times (0, T), \\ u(x, 0) = u_0(x), & x \in \bar{\Omega}, \\ u(x, t) = g(x), & (x, t) \in \partial\Omega \times [0, T]. \end{cases}$$

Under the assumption that domain Ω is mean convex, they show that the solution exists globally in time, is unique, and

$$(1.4) \quad \|u\|_{C^{0,1}(\bar{\Omega} \times [0, \infty))} \leq C.$$

Moreover, they also obtain large time behavior result of $u(x, t)$ as $t \rightarrow \infty$. We are tempting to derive similar results for generalized motion by mean curvature with driving force. However, global estimate (1.4) may fail for the solution of

$$(*) \quad \begin{cases} u_t - |Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) - A|Du| = 0, & (x, t) \in \Omega \times (0, T), \\ u(x, 0) = u_0(x), & x \in \bar{\Omega}, \\ u(x, t) = g(x), & (x, t) \in \partial\Omega \times [0, T], \end{cases}$$

if the boundary condition is fulfilled in classical sense. For instance, let $\Omega = B_2(O) \subset \mathbb{R}^2$, $g(x) = 0$ on $\partial\Omega$, and $A = 1$. Consider

$$\underline{\psi}(x, t) = \begin{cases} 1, & |x| \leq 2 - \frac{1}{2}e^{-\frac{1}{6}t}, \\ 2e^{\frac{1}{6}t}(2 - |x|), & 2 - \frac{1}{2}e^{-\frac{1}{6}t} < |x| \leq 2. \end{cases}$$

As we will see in Appendix, $\underline{\psi}$ is a subsolution of (*), and satisfies

$$\|\underline{\psi}\|_{C^{0,1}(\bar{\Omega} \times [0, \infty))} = \infty.$$

Therefore, as long as $u_0 \geq \underline{\psi}(\cdot, 0)$, by the comparison principle, the solution u also satisfies

$$\|u\|_{C^{0,1}(\bar{\Omega} \times [0, \infty))} = \infty,$$

provided that the boundary value of u agrees with g .

For these reasons, we study the problem with obstacle instead of Dirichlet problem. In 2014, Mercier and Novaga [9] study the mean curvature flow with obstacle in classical sense. In 2016, Mercier [8] gives the well-posed result for the problem (1.1)–(1.3) in the viscosity sense. In this research, they prove the comparison principle and give the existence, uniqueness results. We introduce them in Section 2.

Background. It is expected that a proper understanding of the Dirichlet problem is an obstacle formulation. Consider a curve evolving by the forced curvature flow equation $V = -\kappa + A$, where V is the normal velocity and $-\kappa$ is the curvature in the direction of the normal. Suppose both ends are fixed at the points P_1 and P_2 . We take an obstacle functions $\psi^+ \geq 0 \geq \psi^-$ such that it vanishes only on P_1 and P_2 . Let us explain naively. We denote the curve a part of the boundary of $\{u > 0\}$ which consists of at least two connected component “front” and “back”. Then the front level set of solution of (1.1) is expected to give a solution of Dirichlet problem for $V = -\kappa + 1$. The difficulty of the Dirichlet problem is that the curve does not divide into two parts. This is a reason we mention “front” and “back” of the level-set. Such a problem has been arisen when one discusses spiral growths. In [10, 11], a spiral growth by $V = -\kappa + 1$ is discussed for the Neumann boundary condition by using a modified level-set method. It seems to be possible to discuss the Dirichlet problem by using this obstacle approach.

The level set method for mean curvature flow in viscosity solution frame work was developed independently by Chen, Giga and Goto [1], and Evans, Spruck [2]. They prove the viscosity solution for level set method exists and is unique. Recently, there are some researches considering the mean curvature flow with driving force. In 2016, Giga, Mitake, and Tran [6] consider a crystal growth phenomenon in both vertical and horizontal directions. Indeed, the horizontal direction growth is our mean curvature flow with driving force; see also [4], [5] and [6] for a survey and more developments. In our case here, there is no source term, hence, no vertical growth. In 2017, Zhang consider the mean curvature flow with driving force by level set method and give some criteria to judge whether the zero set is fattening or not (see [14, 15]).

This paper is organized as following. In Section 2, we give the notion of viscosity solutions to the obstacle problem and some basic results. In Section 3, we prove the gradient estimates and give the large time behavior result. In Section 4, we give a full characterization of the limiting profile in the radially symmetric setting. To the best of our knowledge, this result is new in the literature. It is still an open problem on analyzing the limit in general setting.

2. PRELIMINARIES

In this section, we introduce the notion of viscosity solution to the obstacle problem (1.1)–(1.3) and give some related results.

Let $F : \mathbb{R}^N \setminus \{0\} \times \mathcal{S}_N \rightarrow \mathbb{R}$ (\mathcal{S}_N is the set of square symmetric matrices of size N) be such that

$$F(p, X) = \text{trace} \left(\left(I - \frac{p \otimes p}{|p|^2} \right) X \right) + A|p|.$$

Denote by, for $(p, X) \in \mathbb{R}^N \times \mathcal{S}_N$,

$$F_*(p, X) = \liminf_{(q, Y) \rightarrow (p, X)} F(q, Y),$$

and

$$F^*(p, X) = \limsup_{(q, Y) \rightarrow (p, X)} F(q, Y).$$

The above limits are $(q, Y) \rightarrow (p, X)$ in $\mathbb{R}^N \times \mathbb{R}^{N^2}$.

Definition 2.1. A function $u : \mathbb{R}^N \times [0, \infty) \rightarrow \mathbb{R}$ is said to be a (viscosity) subsolution of the problem (1.1)–(1.3) if

- u is upper semicontinuous (usc),
- for all $(x, t) \in \mathbb{R}^N \times [0, T)$, $\psi^-(x) \leq u(x, t) \leq \psi^+(x)$,
- for all $x \in \mathbb{R}^N$, $u(x, 0) \leq u_0(x)$,
- for smooth function φ , if $(\hat{x}, \hat{t}) \in \mathbb{R}^N \times (0, T)$ is a maximizer of $u - \varphi$, and $u(\hat{x}, \hat{t}) > \psi^-(\hat{x})$, then, at (\hat{x}, \hat{t}) ,

$$\varphi_t - F^*(D\varphi, D^2\varphi) \leq 0.$$

Similarly, a function $u : \mathbb{R}^N \times [0, \infty) \rightarrow \mathbb{R}$ is said to be a (viscosity) supersolution of the problem (1.1)–(1.3) if

- u is lower semicontinuous (lsc),
- for all $(x, t) \in \mathbb{R}^N \times [0, T)$, $\psi^-(x) \leq u(x, t) \leq \psi^+(x)$,
- for all $x \in \mathbb{R}^N$, $u(x, 0) \geq u_0(x)$,
- for smooth function φ , if $(\hat{x}, \hat{t}) \in \mathbb{R}^N \times (0, T)$ is a minimizer of $u - \varphi$, and $u(\hat{x}, \hat{t}) < \psi^+(\hat{x})$, then, at (\hat{x}, \hat{t}) ,

$$\varphi_t - F_*(D\varphi, D^2\varphi) \geq 0.$$

Finally, u is said to be a (viscosity) solution of (1.1)–(1.3) if u is both a viscosity subsolution and a supersolution.

Proposition 2.2 (Comparison principle). *We assume u is a subsolution and v is a supersolution of (1.1)–(1.3), respectively. Then $u \leq v$ in $\mathbb{R}^N \times (0, T)$.*

Theorem 2.3 (Well-posedness). *Under our assumptions, (1.1)–(1.3) has a unique solution.*

The comparison principle and well-posedness of (1.1)–(1.3) are quite standard. We refer to [8]. To derive convergence results, it is convenient to consider the approximate problem of (1.1)–(1.3) by considering, for $\varepsilon > 0$, $T > 0$,

$$(1.1\varepsilon) \quad u_t^\varepsilon = \sqrt{\varepsilon^2 + |Du^\varepsilon|^2} \operatorname{div} \left(\frac{Du^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} \right) + A\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}, \quad \text{in } \mathbb{R}^N \times (0, T),$$

$$(1.2\varepsilon) \quad u^\varepsilon(x, 0) = u_0(x), \quad \text{on } \mathbb{R}^N,$$

$$(1.3\varepsilon) \quad (\psi^\varepsilon)^-(x) \leq u^\varepsilon(x, t) \leq (\psi^\varepsilon)^+(x), \quad \text{on } \mathbb{R}^N \times [0, T].$$

Here $(\psi^\varepsilon)^\pm$ are smooth, $\operatorname{supp}(\psi^\varepsilon)^\pm \subset \Omega^\varepsilon$, and $\psi^{\varepsilon^\pm} \rightarrow \psi^\pm$ uniformly in \mathbb{R}^N , where

$$\Omega^\varepsilon = \{x \in \mathbb{R}^N \mid d(x, \Omega) < \varepsilon\}.$$

Moreover, assume $(\psi^\varepsilon)^\pm$ are L_ε -Lipschitz continuous, and $L_\varepsilon \leq K$ for some constant K .

Let $v^\varepsilon = u^\varepsilon/\varepsilon$. Then v^ε satisfies

$$(2.1) \quad v_t^\varepsilon = \sqrt{1 + |Dv^\varepsilon|^2} \operatorname{div} \left(\frac{Dv^\varepsilon}{\sqrt{1 + |Dv^\varepsilon|^2}} \right) + A\sqrt{1 + |Dv^\varepsilon|^2}, \quad \text{in } \mathbb{R}^N \times (0, T),$$

$$(2.2) \quad v^\varepsilon(x, 0) = u_0(x)/\varepsilon, \quad \text{on } \mathbb{R}^N,$$

$$(2.3) \quad (\psi^\varepsilon)^-(x)/\varepsilon \leq v^\varepsilon(x, t) \leq (\psi^\varepsilon)^+(x)/\varepsilon, \quad \text{in } \mathbb{R}^N \times [0, T].$$

Proposition 2.4. *We assume u is a subsolution and v is a supersolution of (2.1)–(2.3), respectively, and $u(\cdot, 0) \leq v(\cdot, 0)$ in \mathbb{R}^N . Then $u \leq v$ in $\mathbb{R}^N \times (0, T)$.*

Proposition 2.5. *Problem (2.1)–(2.3) has a unique continuous solution v^ε , and furthermore, $v^\varepsilon \in C(\mathbb{R}^N \times [0, T]) \cap C^{1,1}(\mathbb{R}^N \times (0, T))$.*

In [9], Mercier and Novaga give the well-posedness for problem (2.1)–(2.3) with $A = 0$ in the viscosity sense by Perron's method and the usual comparison principle. By repeating these standard arguments, the well-posed result for problem (2.1)–(2.3) with $A > 0$ holds. For the regularity, using [12, Theorem 4.1], we deduce that $v^\varepsilon \in C^{1,1}(\mathbb{R}^N \times (0, T))$. We omit the details here.

Propositions 2.4 and 2.5 give us the following results immediately.

Theorem 2.6. *We assume u is a subsolution and v is a supersolution of (1.1 ε)–(1.3 ε), respectively. Then $u \leq v$ in $\mathbb{R}^N \times (0, T)$.*

Theorem 2.7. *Problem (1.1 ε)–(1.3 ε) has a unique solution $u^\varepsilon \in C(\mathbb{R}^N \times [0, T]) \cap C^{1,1}(\mathbb{R}^N \times (0, T))$.*

Lemma 2.8. *Assume u^ε is the unique solution of (1.1 ε)–(1.3 ε) for each $\varepsilon > 0$, and there exists u such that*

$$u^\varepsilon \rightarrow u, \quad \text{as } \varepsilon \rightarrow 0,$$

uniformly in $\mathbb{R}^N \times [0, T)$.

Then u is the unique viscosity solution of (1.1)–(1.3).

Proof. We first show that u is a supersolution. Let φ be a smooth test function such that, at some point $(\hat{x}, \hat{t}) \in \mathbb{R}^N \times (0, T)$, $u(\hat{x}, \hat{t}) < \psi^+(\hat{x})$, and

$$(u - \varphi)(\hat{x}, \hat{t}) < (u - \varphi)(x, t), \quad (x, t) \in \mathbb{R}^N \times (0, T) \setminus \{(\hat{x}, \hat{t})\}.$$

Then there exists a neighborhood $V \subset \mathbb{R}^N \times [0, T)$ of (\hat{x}, \hat{t}) such that, for $(x, t) \in V$,

$$(2.4) \quad u^\varepsilon(x, t) < (\psi^\varepsilon)^+(x).$$

Assume

$$\min_{\mathbb{R}^N \times (0, T)} (u^\varepsilon - \varphi) = (u^\varepsilon - \varphi)(x^\varepsilon, t^\varepsilon).$$

By a standard argument ([3, Lemma 2.2.5]), $(x^\varepsilon, t^\varepsilon) \rightarrow (\hat{x}, \hat{t})$, as $\varepsilon \rightarrow 0$, by passing to a subsequence if necessary. Thanks to (2.4), for $\varepsilon > 0$ small enough, the viscosity supersolution test gives that

$$\left(\varphi_t - \sqrt{\varepsilon^2 + |D\varphi|^2} \operatorname{div} \left(\frac{D\varphi}{\sqrt{\varepsilon^2 + |D\varphi|^2}} \right) + A\sqrt{\varepsilon^2 + |D\varphi|^2} \right) (x^\varepsilon, t^\varepsilon) \geq 0.$$

Letting $\varepsilon \rightarrow 0$, we have

$$(\varphi_t - F_*(D\varphi, D^2\varphi))(\hat{x}, \hat{t}) \geq 0.$$

The proof of subsolution property is similar to the above, and hence, is omitted. \square

3. LIPSCHITZ BOUNDS AND LARGE TIME PROFILES

In this section, we prove Theorems 1.1 and 1.2.

Proof of Theorem 1.1. The proof of spatial Lipschitz bounds is a simple adjustment of that without obstacle; see e.g., [3].

Denote

$$u_z^+(x, t) = (u(x + z, t) + L|z|) \wedge \psi^+(x),$$

$$u_z^-(x, t) = (u(x + z, t) - L|z|) \vee \psi^-(x),$$

for $x, z \in \mathbb{R}^N$, $t \in [0, T)$. We claim u_z^+ is a supersolution and u_z^- is a subsolution of (1.1)–(1.3), respectively. Once we have this claim, Proposition 2.2 shows that

$$u_z^-(x, t) \leq u(x, t) \leq u_z^+(x, t), \quad \text{for all } (x, t) \in \mathbb{R}^N \times (0, T).$$

Thus,

$$u(x + z, t) - L|z| \leq u(x, t) \leq u(x + z, t) + L|z|.$$

Consequently, for every $x, y \in \mathbb{R}^N$, $t \in [0, T)$,

$$|u(x, t) - u(y, t)| \leq L|x - y|.$$

We only prove the claim for u_z^+ . First, we note

$$u(x + z, t) + L|z| \geq \psi^-(x + z) + L|z| \geq \psi^-(x).$$

Consequently,

$$(3.1) \quad \psi^-(x) \leq u_z^+(x, t) \leq \psi^+(x), \quad \text{for all } (x, t) \in \mathbb{R}^N \times [0, T).$$

Since the initial data u_0 is L -Lipschitz,

$$(3.2) \quad u(x + z, 0) + L|z| = u_0(x + z) + L|z| \geq u_0(x).$$

Then we get $u_z^+(\cdot, 0) \geq u_0$ in \mathbb{R}^N . Obviously, $w_z^+(x, t) := u(x + z, t) + L|z|$ satisfies equation (1.1). Combining (3.1), (3.2), and the definition of viscosity supersolution, u_z^+ is a viscosity supersolution of (1.1)–(1.3).

Next we consider

$$u_s^+(x, t) = (u(x, t + s) + Ms) \wedge \psi^+,$$

$$u_s^-(x, t) = (u(x, t + s) - Ms) \vee \psi^-,$$

$(x, t) \in \mathbb{R}^N \times [0, s]$, for $0 \leq s < T$. We claim u_s^+ is a supersolution and u_s^- is a subsolution of (1.1)–(1.3), respectively. If we have this claim, then by the comparison principle,

$$u_s^- \leq u \leq u_s^+.$$

Consequently,

$$u(x, t + s) - Ms \leq u(x, t) \leq u(x, t + s) + Ms.$$

Therefore,

$$|u(x, t) - u(x, r)| \leq M|t - r|,$$

for all $x \in \mathbb{R}^N$, $t, r \in (0, T)$.

Next we only prove the claim for u_s^+ . Seeing the choice of M , it is easy to prove $u_1(x, t) = (u_0(x) - Mt) \vee \psi^-$ is a subsolution of (1.1)–(1.3). By the comparison principle,

$$u(x, t) \geq u_1(x, t) \geq u_0(x) - Mt.$$

This implies

$$(3.3) \quad u_s^+(x, 0) = (u(x, s) + Ms) \wedge \psi^+ \geq u_0(x).$$

Obviously,

$$(3.4) \quad \psi^+ \geq u_s^+(x, t) = (u(x, t + s) + Ms) \wedge \psi^+ \geq \psi^-.$$

Note $h_s^+(x, t) := u(x, t + s) + Ms$ satisfies equation (1.1). Combining (3.3), (3.4), and the definition of viscosity supersolution, u_s^+ is a viscosity supersolution of (1.1)–(1.3). \square

Next we prove Theorem 1.2.

Proof of Theorem 1.2. We divide the proof into four steps.

Step 1. u^ε are Lipschitz continuous for all $\varepsilon \in (0, 1)$. Moreover,

$$|u^\varepsilon(x, t) - u^\varepsilon(y, t)| \leq L_\varepsilon|x - y| \leq K|x - y|,$$

and

$$|u^\varepsilon(x, t) - u^\varepsilon(x, r)| \leq M|t - r|,$$

for $x, y \in \mathbb{R}^N$, $t, r \in (0, T)$.

By constructing subsolution and supersolution as in Theorem 1.1, we can prove these results easily. We leave the details to the readers.

Step 2. There exists constant $C > 0$ independent of ε and T such that

$$(3.5) \quad \int_0^T \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dx dt \leq C.$$

We consider the following Lyapunov function (see e.g., [7])

$$I^\varepsilon(t) = \int_{\mathbb{R}^N} \sqrt{\varepsilon^2 + |Du^\varepsilon|^2} dx.$$

By calculation,

$$\frac{d}{dt} \int_{\mathbb{R}^N} \sqrt{\varepsilon^2 + |Du^\varepsilon|^2} dx = \int_{\mathbb{R}^N} \frac{Du^\varepsilon \cdot Du_t^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} dx = - \int_{\mathbb{R}^N} u_t^\varepsilon \operatorname{div} \left(\frac{Du^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} \right) dx.$$

If $u^\varepsilon(x_0, t_0) = (\psi^\varepsilon)^+(x_0)$ at some $(x_0, t_0) \in \mathbb{R}^N \times (0, T)$, it also means

$$\max_{(x,t) \in \mathbb{R}^N \times (0,T)} (u^\varepsilon(x, t) - (\psi^\varepsilon)^+(x)) = u^\varepsilon(x_0, t_0) - (\psi^\varepsilon)^+(x_0) = 0.$$

then there holds $u_t^\varepsilon(x_0, t_0) = 0$. Same claim holds if $u^\varepsilon(x_0, t_0) = (\psi^\varepsilon)^-(x_0)$. Consequently,

$$F^\pm(t) = \{x \in \mathbb{R}^N \mid u^\varepsilon(x, t) = (\psi^\varepsilon)^\pm(x), u_t^\varepsilon(x, t) \neq 0\} = \emptyset,$$

for $t \in (0, T)$. Let

$$Q^\pm(t) = \{x \in \mathbb{R}^N \mid u^\varepsilon(x, t) \neq (\psi^\varepsilon)^\pm(x)\}, \quad t \in (0, T).$$

Note the fact that for $x \in Q^+(t) \cap Q^-(t)$, we have

$$u_t^\varepsilon = \sqrt{\varepsilon^2 + |Du^\varepsilon|^2} \operatorname{div} \left(\frac{Du^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} \right) + A\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}$$

at (x, t) . Therefore,

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^N} \sqrt{\varepsilon^2 + |Du^\varepsilon|^2} dx &= - \int_{\mathbb{R}^N \setminus (F^+ \cup F^-)} u_t^\varepsilon \operatorname{div} \left(\frac{Du^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} \right) dx \\ &= - \int_{Q^+ \cap Q^-} u_t^\varepsilon \operatorname{div} \left(\frac{Du^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} \right) dx \\ &= - \int_{Q^+ \cap Q^-} \left(\frac{(u_t^\varepsilon)^2}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} - Au_t^\varepsilon \right) dx \\ &= - \int_{\mathbb{R}^N} \left(\frac{(u_t^\varepsilon)^2}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} - Au_t^\varepsilon \right) dx \\ &\leq - \frac{1}{\sqrt{\varepsilon^2 + K^2}} \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dx + A \int_{\mathbb{R}^N} u_t^\varepsilon dx. \end{aligned}$$

Then,

$$\frac{d}{dt} \left(\int_{\mathbb{R}^N} \sqrt{\varepsilon^2 + |Du^\varepsilon|^2} dx - A \int_{\mathbb{R}^N} u^\varepsilon dx \right) \leq - \frac{1}{\sqrt{\varepsilon^2 + K^2}} \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dx.$$

Integrating the inequality above, we have

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dx dt &\leq A\sqrt{\varepsilon^2 + K^2} \int_{\mathbb{R}^N} (u^\varepsilon(x, T) - u^\varepsilon(x, 0)) dx \\ &\quad + \sqrt{\varepsilon^2 + K^2} \int_{\mathbb{R}^N} \left(\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}(x, 0) - \sqrt{\varepsilon^2 + |Du^\varepsilon|^2}(x, T) \right) dx. \end{aligned}$$

The assumptions for $(\psi^\varepsilon)^\pm$ show that we can find a constant C_1 independent of $\varepsilon \in (0, 1)$ such that

$$\int_{\mathbb{R}^N} |(\psi^\varepsilon)^\pm|(x) dx \leq C_1.$$

Then, for $\varepsilon \in (0, 1)$,

$$\int_0^T \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dxdt \leq 2AC_1 \sqrt{1+K^2} + \sqrt{1+K^2} \int_{\mathbb{R}^N} \sqrt{1+|Du_0^\varepsilon|^2(x)} dx := C.$$

Thus,

$$\int_0^T \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dxdt \leq C.$$

Step 3. $u_t^\varepsilon \rightharpoonup u_t$ weakly in $L^2(\mathbb{R}^N \times [0, \infty))$, as $\varepsilon \rightarrow 0$.

Step 2 shows that u_t^ε is bounded in $L^2(\mathbb{R}^N \times [0, \infty))$. Since $L^2(\mathbb{R}^N \times [0, \infty))$ is a Hilbert space, there exists $g \in L^2(\mathbb{R}^N \times [0, \infty))$ such that, by passing to a subsequence if needed,

$$u_t^\varepsilon \rightharpoonup g,$$

weakly in $L^2(\mathbb{R}^N \times [0, \infty))$, as $\varepsilon \rightarrow 0$. For every $\phi \in C_0^\infty(\mathbb{R}^N \times (0, T))$,

$$\int_0^T \int_{\mathbb{R}^N} u_t^\varepsilon \phi dxdt \rightarrow \int_0^T \int_{\mathbb{R}^N} g \phi dxdt,$$

as $\varepsilon \rightarrow 0$, for every $T > 0$. On the other hand, Step 1 and Lemma 2.8 show that $u^\varepsilon \rightarrow u$ uniformly on $\mathbb{R}^N \times [0, T]$ for every $T > 0$. Therefore,

$$\int_0^T \int_{\mathbb{R}^N} u_t^\varepsilon \phi dxdt = - \int_0^T \int_{\mathbb{R}^N} u^\varepsilon \phi_t dxdt \rightarrow - \int_0^T \int_{\mathbb{R}^N} u \phi_t dxdt.$$

Then

$$\int_0^T \int_{\mathbb{R}^N} g \phi dxdt = - \int_0^T \int_{\mathbb{R}^N} u \phi_t dxdt$$

which shows that $g = u_t$ in $\mathbb{R}^N \times [0, \infty)$. Thus, $u_t^\varepsilon \rightharpoonup u_t$ weakly in $L^2(\mathbb{R}^N \times [0, \infty))$, as $\varepsilon \rightarrow 0$ (whole sequence).

Step 4. We complete the proof in this step.

By weakly lower semi-continuity,

$$(3.6) \quad \int_0^\infty \int_{\mathbb{R}^N} (u_t)^2 dxdt \leq \liminf_{\varepsilon \rightarrow 0} \int_0^\infty \int_{\mathbb{R}^N} (u_t^\varepsilon)^2 dxdt \leq C.$$

For every $\{t_k\} \rightarrow \infty$, by the Arzelà-Ascoli theorem, there exist a subsequence $\{t_{k_j}\}$ and a Lipschitz continuous function v such that

$$u_{k_j}(x, t) = u(x, t + t_{k_j}) \rightarrow v(x, t),$$

locally uniformly on $\mathbb{R}^N \times [0, \infty)$. As $u(\cdot, t)$ is compactly supported on $\bar{\Omega}$,

$$(3.7) \quad u_{k_j}(x, t) = u(x, t + t_{k_j}) \rightarrow v(x, t),$$

uniformly on $\mathbb{R}^N \times [0, T]$, for every $T > 0$. By stability results of viscosity solutions, v satisfies

$$(1.1a) \quad v_t = |Dv| \operatorname{div} \left(\frac{Dv}{|Dv|} \right) + A|Dv|, \quad (x, t) \in \mathbb{R}^N \times (0, \infty),$$

$$(1.3b) \quad \psi^-(x) \leq v(x, t) \leq \psi^+(x), \quad (x, t) \in \mathbb{R}^N \times [0, \infty).$$

Thanks to the fact that

$$\int_0^\infty \int_{\mathbb{R}^N} (u_t)^2 dx dt \leq C < \infty,$$

we have

$$\int_0^1 \int_{\mathbb{R}^N} (u_{k_j})_t^2 dx dt = \int_{t_{k_j}}^{1+t_{k_j}} \int_{\mathbb{R}^N} (u_t)^2 dx dt \rightarrow 0,$$

as $j \rightarrow \infty$. This shows that

$$(u_{k_j})_t \rightharpoonup 0,$$

weakly in $L^2(\mathbb{R}^N \times [0, 1])$. On the other hand, (3.7) implies that, by passing to a further subsequence if necessary

$$(u_{k_j})_t \rightharpoonup v_t,$$

weakly in $L^2(\mathbb{R}^N \times [0, 1])$. Consequently, $v_t = 0$ on $\mathbb{R}^N \times [0, 1]$. Thus, v is a solution of

$$(1.1e) \quad 0 = |Dv| \operatorname{div} \left(\frac{Dv}{|Dv|} \right) + A|Dv|, \quad x \in \mathbb{R}^N,$$

$$(1.3) \quad \psi^-(x) \leq v(x) \leq \psi^+(x), \quad x \in \mathbb{R}^N.$$

As mentioned in Remark 1.4, the solution of (1.1e), (1.3) is not necessary unique. Therefore, v may depend on the choice of subsequence of $\{t_k\}_k$.

At last, we prove that v is independent of the choice of subsequence of $\{t_k\}_k$. Since u_{k_j} converges uniformly to v on $\mathbb{R}^N \times [0, 1]$, for every $\varepsilon > 0$ there exists j large enough such that

$$|u_{k_j}(x, t) - v(x)| < \varepsilon, \quad \text{for all } (x, t) \in \mathbb{R}^N \times [0, 1].$$

Setting $t = 0$, $v(x) - \varepsilon < u_{k_j}(x, 0) < v(x) + \varepsilon$ in \mathbb{R}^N . By the comparison principle,

$$v(x) - \varepsilon \leq (v(x) - \varepsilon) \vee \psi^- \leq u(x, t) \leq (v(x) + \varepsilon) \wedge \psi^+ \leq v(x) + \varepsilon,$$

for $(x, t) \in \mathbb{R}^N \times [t_{k_j}, \infty)$. This implies that $u(\cdot, t)$ converges uniformly to v in \mathbb{R}^N without taking a subsequence.

□

Remark 3.1. The proof of large time behavior is quite standard. Nevertheless, as we do not have uniqueness of solutions to (1.1e), (1.3), it is not easy to analyze what is the limiting profile given the initial data u_0 , and obstacles ψ^\pm . In the following, we are able to characterize this in the radially symmetric setting.

4. THE RADIALLY SYMMETRIC SETTING

In this section, we prove Theorem 1.3. First, we find radially symmetric solutions of (1.1e), and (1.3). To establish this, we need the following lemma ([6, Lemma 8.1]).

Lemma 4.1. *Let $\psi : [0, \infty) \rightarrow \mathbb{R}$ be a continuous function, which is C^2 in $(0, R) \cup (R, R_1)$, and $\psi = 0$ in $[R_1, \infty)$, for some given $R, R_1 > 0$ with $R_1 > R$. Assume further that*

$$\psi'(R-) = a \quad \text{and} \quad \psi'(R+) = b.$$

The following holds.

(i) *If $a < b$, then for any $\phi \in C^2(\mathbb{R}^N)$ such that $\psi(|x|) - \phi(x)$ has a strict minimum at $x_0 \in \partial B(0, R)$, then for some $s \in [a, b]$,*

$$D\phi(x_0) = s \frac{x_0}{R} \quad \text{and} \quad \text{tr} \left[\left(I - \frac{Du \otimes Du}{|Du|^2} \right) D^2 u \right] \leq \frac{(N-1)s}{R}.$$

(ii) *If $a > b$, then for any $\phi \in C^2(\mathbb{R}^N)$ such that $\psi(|x|) - \phi(x)$ has a strict maximum at $x_0 \in \partial B(0, R)$, then for some $s \in [b, a]$,*

$$D\phi(x_0) = s \frac{x_0}{R} \quad \text{and} \quad \text{tr} \left[\left(I - \frac{Du \otimes Du}{|Du|^2} \right) D^2 u \right] \geq \frac{(N-1)s}{R}.$$

We always assume that the hypotheses of Theorem 1.3 are in force in this section. Let us recall them here for clarity

(1) $\Omega = B_R(O)$ for given $R > 0$;

(2)

$$\psi^+(x) = \begin{cases} \lambda \text{dist}(x, \partial\Omega) = \lambda(R - |x|), & x \in \Omega, \\ 0, & x \in \mathbb{R}^N \setminus \Omega, \end{cases}$$

where $\lambda > 0$ is given;

(3) ψ^- is radial, and $\psi^- < 0$ in Ω . The initial data u_0 is radial, and $\psi^- \leq u_0 \leq \psi^+$.

Proposition 4.2. *Assume $R \geq (N-1)/A$. The solution v of (1.1e), (1.3) satisfies*

$$0 \leq v \leq \psi^+ \wedge \lambda \left(R - \frac{N-1}{A} \right).$$

Proof. Let

$$u_* = \varphi_* \vee \psi^-,$$

where

$$\varphi_*(x, t) = \begin{cases} -Le^{-\gamma t}(R - |x|), & |x| \leq R, t \geq 0, \\ 0, & |x| > R, t \geq 0, \end{cases}$$

where γ is chosen such that

$$0 < \gamma < \frac{1}{R} \left(\frac{N-1}{R} + A \right).$$

At (x_0, t_0) , for $0 < |x_0| < R$, and $t_0 > 0$,

$$\begin{aligned} \left((\varphi_*)_t - |D\varphi_*| \operatorname{div} \left(\frac{D\varphi_*}{|D\varphi_*|} \right) - A|D\varphi_*| \right) (x_0, t_0) &= Le^{-\gamma t_0} \left(\gamma(R - |x_0|) - \frac{N-1}{|x_0|} - A \right) \\ &\leq Le^{-\gamma t_0} \left(\gamma R - \frac{N-1}{R} - A \right) < 0. \end{aligned}$$

Obviously, $u_*(\cdot, 0) = \psi^- \leq v$ and $\psi^- \leq u_* \leq \psi^+$. Thus, u_* is subsolution of (1.1), (1.2), and (1.3). By the comparison principle,

$$u_*(x, t) \leq v(x), \quad \text{for all } (x, t) \in \mathbb{R}^N \times [0, \infty).$$

Since $u_*(\cdot, t) \rightarrow 0$, as $t \rightarrow \infty$, we conclude that $v \geq 0$.

For small $\varepsilon > 0$, let

$$u^{*\varepsilon}(x, t) = \begin{cases} \lambda e^{-\mu t} \left(\frac{N-1}{A} - \varepsilon - |x| \right) + \lambda \left(R - \frac{N-1}{A} + \varepsilon \right), & |x| \leq \frac{N-1}{A} - \varepsilon, t \geq 0, \\ \psi^+, & |x| > \frac{N-1}{A} - \varepsilon, t \geq 0. \end{cases}$$

Here, $0 < \mu < \frac{A}{N-1} \left(\frac{N-1}{(N-1)/A - \varepsilon} - A \right)$.




FIGURE 2. Supersolution $u^{*\varepsilon}$

At (x_0, t_0) , where $0 < |x_0| < \frac{N-1}{A} - \varepsilon$, and $t_0 > 0$, we compute

$$\begin{aligned} & \left(u_t^{*\varepsilon} - |Du^{*\varepsilon}| \operatorname{div} \left(\frac{Du^{*\varepsilon}}{|Du^{*\varepsilon}|} \right) - A|Du^{*\varepsilon}| \right) (x_0, t_0) \\ &= \lambda e^{-\mu t_0} \left(-\mu \left(\frac{N-1}{A} - \varepsilon - |x_0| \right) + \frac{N-1}{|x_0|} - A \right) \\ &\geq \lambda e^{-\mu t_0} \left(-\mu \frac{N-1}{A} + \frac{N-1}{(N-1)/A - \varepsilon} - A \right) > 0. \end{aligned}$$

Thus, $u^{*\varepsilon}$ is supersolution of (1.1), (1.2), and (1.3).

Obviously, $u^{*\varepsilon}(\cdot, 0) = \psi^+ \geq v$, and $\psi^- \leq u^{*\varepsilon} \leq \psi^+$. By the comparison principle,

$$v(x) \leq u^{*\varepsilon}, \quad \text{for all } (x, t) \in \mathbb{R}^N \times [0, \infty).$$

Since

$$u^{*\varepsilon}(\cdot, t) \rightarrow \psi^+ \wedge \lambda \left(R - \frac{N-1}{A} + \varepsilon \right),$$

we deduce, as $t \rightarrow \infty$,

$$v \leq \psi^+ \wedge \lambda \left(R - \frac{N-1}{A} + \varepsilon \right).$$

Letting $\varepsilon \rightarrow 0$ in the above to imply

$$v \leq \psi^+ \wedge \lambda \left(R - \frac{N-1}{A} \right).$$

The proof is now complete. \square

Proposition 4.3. *Assume $R \geq (N-1)/A$. Then*

$$\psi_C := \psi^+ \wedge C, \quad \text{for } 0 \leq C \leq \lambda \left(R - \frac{N-1}{A} \right),$$

are all radially symmetric and Lipschitz continuous solutions to (1.1e), and (1.3) satisfying that

$$0 \leq v \leq \psi^+ \wedge \lambda \left(R - \frac{N-1}{A} \right).$$

Proof. Let v be a radially symmetric solution to (1.1e), and (1.3). By abuse of notion, we write $v(x) = v(|x|)$ for $x \in \mathbb{R}^N$. Then, $v(r)$ satisfies following ODE

$$(1.1r) \quad -\frac{(N-1)v'}{r} - A|v'| = 0, \quad r > 0,$$

$$(1.3r) \quad \psi^-(r) \leq v(r) \leq \psi^+(r), \quad r \geq 0,$$

in the viscosity sense.

Denote by

$$E = \{r \in (0, R) \mid v(r) = \psi^+(r) \text{ or } v(r) = \psi^-(r)\}.$$




FIGURE 3. Stationary solution of (1.1e), (1.3)

Obviously, E is a closed set. Then we can find at most a countable number of intervals (a_i, b_i) , $i \in \mathbb{N}$, satisfying

$$b_i \leq a_{i+1},$$

and

$$(0, R) \setminus E = \bigcup_{i=1}^{\infty} (a_i, b_i).$$

Under our assumption, $v(0) < \psi^+(0)$. Thus $a_1 = 0$.

In each interval (a_i, b_i) , v satisfies (1.1r) in the viscosity sense. If v is differentiable at $r_0 \in (a_i, b_i)$, and $r_0 \neq (N-1)/A$, there holds

$$-\frac{(N-1)v'(r_0)}{r_0} - A|v'(r_0)| = 0,$$

which implies $v'(r_0) = 0$. Since v is Lipschitz continuous, v is differentiable in (a_i, b_i) almost everywhere. Thus

$$v' = 0, \quad \text{a.e. in } (a_i, b_i),$$

which gives that $v \equiv c_i$ in (a_i, b_i) as v is Lipschitz continuous. We have

$$(a_i, b_i) = \emptyset,$$

provided that $\psi^+(a_i) > \psi^+(b_i)$, $i \geq 2$. Therefore, $v = c_1$ in $[a_1, b_1) = [0, b_1)$, and $v = \psi^+$ in $[b_1, \infty)$. In other words,

$$v = \psi^+ \wedge c_1.$$

Under our assumption, $0 \leq c_1 \leq \lambda(R - (N-1)/A)$.

At last we prove that for all $0 \leq C \leq \lambda(R - (N - 1)/A)$, $\psi^+ \wedge C$ is the solution of problem (1.1e), (1.3). It is easy to see $\psi^+ \wedge C$ is a supersolution. We only show ψ_C is a subsolution.

This claim is clear for $|x| < R - \frac{C}{\lambda}$ or $|x| \geq R$. We only check carefully where $R - \frac{C}{\lambda} \leq |x_0| < R$.

For $R - \frac{C}{\lambda} < |x_0| < R$, we have $\psi_C(x_0) = \psi^+(x_0) > \psi^-(x_0)$, and

$$\left(-|D\psi^+| \operatorname{div} \left(\frac{D\psi^+}{|D\psi^+|} \right) - A|D\psi^+| \right) (x_0) = \lambda \left(\frac{N-1}{|x_0|} - A \right) < 0.$$

Here we use the fact $|x_0| > R - \frac{C}{\lambda} \geq (N-1)/A$.

For $|x_0| = R - \frac{C}{\lambda}$, assume that there is a test function $\varphi \in C^2(\mathbb{R}^N)$ such that $\varphi - \psi_C$ has a strict maximum at x_0 . By Lemma 4.1, there exists $s \in [-\lambda, 0]$ such that

$$D\varphi(x_0) = \frac{s x_0}{|x_0|} \quad \text{and} \quad \operatorname{tr} \left[\left(I - \frac{Du \otimes Du}{|Du|^2} \right) D^2 u \right] \geq \frac{(N-1)s}{|x_0|}.$$

Since $|x_0| = R - \frac{C}{\lambda} \geq (N-1)/A$, and $s \leq 0$, we have

$$\left(-|D\varphi| \operatorname{div} \left(\frac{D\varphi}{|D\varphi|} \right) - A|D\varphi| \right) (x_0) \leq -\frac{(N-1)s}{|x_0|} - A|s| = -\frac{(N-1)s}{|x_0|} + As \leq 0.$$

Thus, ψ_C is subsolution. The proof is complete. □

Proof of Theorem 1.3. If $R = (N-1)/A$, Proposition 4.3 shows that 0 is the unique solution of (1.1e), (1.3). The result is obvious by Theorem 1.2.

If $R < (N-1)/A$, let

$$u^*(x, t) = \begin{cases} \lambda e^{-\nu_1 t} (R - |x|), & |x| < R, \\ 0, & |x| \geq R. \end{cases}$$

Here ν_1 is chosen such that

$$0 < \nu_1 < \frac{1}{R} \left(\frac{N-1}{R} - A \right).$$

At (x_0, t_0) , where $0 < |x_0| < R$, and $t_0 > 0$,

$$\begin{aligned} \left(u_t^* - |\nabla u^*| \operatorname{div} \left(\frac{\nabla u^*}{|\nabla u^*|} \right) - A|\nabla u^*| \right) (x_0, t_0) &= \lambda e^{-\nu_1 t_0} \left(-\nu_1 (R - |x_0|) + \frac{N-1}{|x_0|} - A \right) \\ &\geq \lambda e^{-\nu_1 t_0} \left(-\nu_1 R + \frac{N-1}{R} - A \right) > 0. \end{aligned}$$

Then u^* is a supersolution. Besides, u_* constructed in Proposition 4.2 is a subsolution. By the comparison principle,

$$u_* \leq u \leq u^*.$$

Therefore, $u(\cdot, t) \rightarrow 0$ uniformly in \mathbb{R}^N , as $t \rightarrow 0$.

We consider the final case where $R > (N-1)/A$. First we construct a supersolution

$$\bar{u}^\varepsilon(x, t) = \begin{cases} (B + \varepsilon) \wedge \psi^+, & |x| > \frac{N-1}{A} - \frac{\varepsilon}{L}, \\ Le^{-\nu_2 t} \left(\frac{N-1}{A} - \frac{\varepsilon}{L} - |x| \right) + B + \varepsilon, & |x| \leq \frac{N-1}{A} - \frac{\varepsilon}{L}. \end{cases}$$

Here ν_2 is chosen so that

$$0 < \nu_2 < \frac{A}{N-1} \left(\frac{N-1}{A} - \frac{\varepsilon}{L} - A \right).$$




FIGURE 4. Supersolution \bar{u}^ε

Recall $B = \max_{(N-1)/A \leq |x| \leq R} u_0(x) \leq \lambda(R - (N-1)/A)$, and u_0 is L -Lipschitz continuous. It is easy to check

$$\psi^- \leq \bar{u}^\varepsilon \leq \psi^+ \quad \text{and} \quad \bar{u}^\varepsilon(\cdot, 0) \geq u_0.$$

For $|x_0| > \frac{N-1}{A} - \frac{\varepsilon}{L}$, and $t_0 > 0$, obviously \bar{u}^ε is supersolution in the viscosity sense.

For $|x_0| < \frac{N-1}{A} - \frac{\varepsilon}{L}$, and $t_0 > 0$,

$$\begin{aligned} & \left(\bar{u}_t^\varepsilon - |D\bar{u}^\varepsilon| \operatorname{div} \left(\frac{D\bar{u}^\varepsilon}{|D\bar{u}^\varepsilon|} \right) - A|D\bar{u}^\varepsilon| \right) (x_0, t_0) \\ &= Le^{-\nu_2 t_0} \left(-\nu_2 \left(\frac{N-1}{A} - \frac{\varepsilon}{L} - |x_0| \right) + \frac{N-1}{|x_0|} - A \right) \\ &> Le^{-\nu_2 t_0} \left(-\nu_2 \left(\frac{N-1}{A} - \frac{\varepsilon}{L} \right) + \frac{N-1}{\frac{N-1}{A} - \frac{\varepsilon}{L}} - A \right) > 0. \end{aligned}$$

For $|x_0| = \frac{N-1}{A} - \frac{\varepsilon}{L}$, $t_0 > 0$, and test function $\varphi \in C^2(\mathbb{R}^N \times (0, \infty))$ satisfies

$$(\varphi - \bar{u}^\varepsilon)(x_0, t_0) < (\varphi - \bar{u}^\varepsilon)(x, t), \quad (x, t) \neq (x_0, t_0).$$

Then

$$\varphi_t(x_0, t_0) \geq 0.$$

By abuse of notions, we write $\bar{u}^\varepsilon(x, t) = \bar{u}^\varepsilon(r, t)$ for $r = |x|$. It is easy to see

$$\lim_{r \rightarrow (\frac{N-1}{A} - \frac{\varepsilon}{L})_-} \bar{u}_r^\varepsilon(r, t) = -Le^{-\nu_2 t},$$

and

$$\lim_{r \rightarrow (\frac{N-1}{A} - \frac{\varepsilon}{L})_+} \bar{u}_r^\varepsilon(r, t) = 0 \quad \text{or} \quad -\lambda.$$

Then by Lemma 4.1, there exists $s \leq 0$ such that

$$D\varphi(x_0, t_0) = \frac{sx_0}{|x_0|} \quad \text{and} \quad \operatorname{tr} \left[\left(I - \frac{Du \otimes Du}{|Du|^2} \right) D^2u \right] (x_0, t_0) \leq \frac{(N-1)s}{|x_0|}.$$

Since $|x_0| = \frac{N-1}{A} - \frac{\varepsilon}{L} < (N-1)/A$, and $s \leq 0$, we have

$$\left(\varphi_t - |D\varphi| \operatorname{div} \left(\frac{D\varphi}{|D\varphi|} \right) - A|D\varphi| \right) (x_0, t_0) \geq -\frac{(N-1)s}{|x_0|} - A|s| = -\frac{(N-1)s}{|x_0|} + As > 0.$$


Thus, \bar{u}^ε is a supersolution of (1.1), (1.2), and (1.3).

Next, we construct a subsolution $\underline{u} = \varphi \vee \psi^-$ to equation (1.1), (1.2), and (1.3), where

$$\underline{\varphi}(x, t) = \begin{cases} -Le^{-\nu_3 t}(r_0 - |x|) + B, & |x| \leq r_0, \\ (L(r_0 - |x|) + B) \vee (Le^{-\nu_4 t}(|x| - R)), & r_0 < |x| \leq R, \\ 0, & |x| > R. \end{cases}$$

Here $r_0 \in [(N-1)/A, R]$ satisfies $B = u_0(r_0)$,

$$0 < \nu_3 < \frac{N-1 + Ar_0}{r_0^2},$$

FIGURE 5. Subsolution $\underline{\varphi}$

and

$$0 < \nu_4 < \frac{1}{R} \left(\frac{N-1}{R} + A \right).$$

It is easy to check $u_0 \geq \underline{u}(\cdot, 0)$, and

$$\psi^- \leq \underline{u} \leq \psi^+.$$

Clearly, \underline{u} is a subsolution for $|x| \geq R$, $t > 0$. We show the claim for $|x| < R$, $t > 0$. For $r_0 < |x_0| < R$, $t_0 > 0$, if $\underline{\varphi} = Le^{-\nu_4 t_0}(|x| - R)$, we compute

$$\begin{aligned} \left(\underline{\varphi}_t - |D\underline{\varphi}| \operatorname{div} \left(\frac{D\underline{\varphi}}{|D\underline{\varphi}|} \right) - A|D\underline{\varphi}| \right) (x_0, t_0) &= Le^{-\nu_4 t_0} \left(\nu_4(R - |x_0|) - \frac{N-1}{|x_0|} - A \right) \\ &\leq Le^{-\nu_4 t_0} \left(\nu_4 R - \frac{N-1}{R} - A \right) < 0. \end{aligned}$$

For $r_0 < |x_0| < R$, $t_0 > 0$, if $\underline{\varphi} = L(r_0 - |x|) + B$, we compute

$$\left(\underline{\varphi}_t - |D\underline{\varphi}| \operatorname{div} \left(\frac{D\underline{\varphi}}{|D\underline{\varphi}|} \right) - A|D\underline{\varphi}| \right) (x_0, t_0) = L \left(\frac{N-1}{|x_0|} - A \right) < L \left(\frac{N-1}{r_0} - A \right) \leq 0.$$

For $|x_0| < r_0$, $t_0 > 0$,

$$\begin{aligned} \left(\underline{\varphi}_t - |D\underline{\varphi}| \operatorname{div} \left(\frac{D\underline{\varphi}}{|D\underline{\varphi}|} \right) - A|D\underline{\varphi}| \right) (x_0, t_0) &= Le^{-\nu_3 t_0} \left(\nu_3(r_0 - |x_0|) - \frac{N-1}{|x_0|} - A \right) \\ &< Le^{-\nu_3 t_0} \left(\nu_3 r_0 - \frac{N-1}{r_0} - A \right) < 0. \end{aligned}$$

Finally, we need to check the case where $|x_0| = r_0$, $t_0 > 0$. Take a test function $\varphi \in C^2(\mathbb{R}^N \times (0, \infty))$ such that $\underline{\varphi} - \varphi$ has a strict maximum at (x_0, t_0) . Then

$$\varphi_t(x_0, t_0) \leq 0.$$

By Lemma 4.1, there exists $s \in [-L, Le^{-\nu_3 t_0}]$ such that

$$D\varphi(x_0, t_0) = \frac{sx_0}{|x_0|} \quad \text{and} \quad \text{tr} \left[\left(I - \frac{Du \otimes Du}{|Du|^2} \right) D^2 u \right] \geq \frac{(N-1)s}{|x_0|}.$$

We use the above and the fact that $|x_0| = r_0 \geq (N-1)/A$ to imply

$$\left(\varphi_t - |D\varphi| \operatorname{div} \left(\frac{D\varphi}{|D\varphi|} \right) - A|D\varphi| \right) (x_0, t_0) \leq -\frac{(N-1)s}{|x_0|} - A|s| \leq \frac{(N-1)|s|}{|x_0|} - A|s| \leq 0,$$

Therefore, \underline{u} is a subsolution. By the comparison principle,

$$\underline{u} \leq u \leq \bar{u}^\varepsilon.$$

Letting $t \rightarrow \infty$ and $\varepsilon \rightarrow 0$ in this order,

$$\underline{\psi} \leq v \leq B \wedge \psi^+ = \psi_B.$$

Here

$$\underline{\psi}(x) = \begin{cases} B, & |x| \leq r_0, \\ L(r_0 - |x|) + B, & r_0 < |x| \leq r_0 + \frac{B}{L}, \\ 0, & r_0 + \frac{B}{L} < |x|. \end{cases}$$

Proposition 4.3 shows that $v = \psi_B$. □

Remark 4.4. Theorem 1.3 also holds for $\psi^- \leq 0$ in \mathbb{R}^N by using approximation arguments. We leave the proof to the readers.


Remark 4.5. We give another idea to prove Theorem 1.3 as following. Recall

$$u(x, t) = u(|x|, t) = u(r, t).$$

For $r > (N-1)/A$ such that $\psi^-(r) < u(r, t) < \psi^+(r)$, we have the following equation

$$u_t(r, t) = \frac{N-1}{r} u_r + A|u_r| \geq 0.$$

Therefore, at r_0 , we yield that $t \mapsto u(r_0, t)$ is non-decreasing. This immediately gives us the desired conclusion for $r > (N-1)/A$.

FIGURE 6. $\underline{\psi}$ and ψ_B

5. APPENDIX

Lemma 5.1. *Let $N = 2$. The function $\underline{\psi}$ is a subsolution of problem*

$$(5.1) \quad \begin{cases} u_t - |Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) - |Du| = 0, & (x, t) \in B_2(O) \times (0, \infty), \\ u(x, 0) = u_0(x), & x \in \overline{B_2(O)}, \\ u(x, t) = 0, & (x, t) \in \partial B_2(O) \times [0, \infty), \end{cases}$$

and satisfy

$$\|\underline{\psi}\|_{C^{0,1}(\overline{B_2(O)} \times [0, \infty))} = \infty,$$

where $u_0 \geq \underline{\psi}(\cdot, 0)$ in $\overline{B_2(O)}$, and

$$\underline{\psi}(x, t) = \begin{cases} 1, & |x| \leq 2 - \frac{1}{2}e^{-\frac{1}{6}t}, \\ 2e^{\frac{1}{6}t}(2 - |x|), & 2 - \frac{1}{2}e^{-\frac{1}{6}t} < |x| \leq 2. \end{cases}$$

Proof. Obviously, $\underline{\psi}(\cdot, t) = 0$ on $\partial B_2(O)$. There is nothing to check if $|x_0| < 2 - \frac{1}{2}e^{-\frac{1}{6}t_0}$, and $t_0 > 0$.

If $2 - \frac{1}{2}e^{-\frac{1}{6}t_0} < |x_0| < 2$, and $t_0 > 0$, noting $|x_0| > \frac{3}{2}$, we compute

$$\begin{aligned} \left(\underline{\psi}_t - |D\underline{\psi}| \operatorname{div} \left(\frac{D\underline{\psi}}{|D\underline{\psi}|} \right) - |D\underline{\psi}| \right) (x_0, t_0) &= 2e^{\frac{1}{6}t_0} \left(\frac{1}{6}(2 - |x_0|) + \frac{1}{|x_0|} - 1 \right) \\ &\leq 2e^{\frac{1}{6}t_0} \left(\frac{1}{6}(2 - |x_0|) + \frac{1}{|x_0|} - 1 \right) \leq 2e^{\frac{1}{6}t_0} \left(\frac{1}{12} + \frac{2}{3} - 1 \right) < 0. \end{aligned}$$

Finally, we check the case where $|x_0| = 2 - \frac{1}{2}e^{-\frac{1}{6}t_0}$, and $t_0 > 0$. Choose a test function $\varphi \in C^2(\mathbb{R}^2 \times (0, \infty))$ such that $\underline{\psi} - \varphi$ has a strict maximum at (x_0, t_0) . By Lemma 4.1, there exists $s \in [-\frac{1}{2}e^{\frac{1}{6}t_0}, 0]$ such that

$$D\varphi(x_0, t_0) = \frac{sx_0}{|x_0|} \quad \text{and} \quad \text{tr} \left[\left(I - \frac{Du \otimes Du}{|Du|^2} \right) D^2u \right] \geq \frac{s}{|x_0|}.$$

Let $x(t) = (2 - \frac{1}{2}e^{-\frac{1}{6}t})\frac{x_0}{|x_0|}$. For $t < t_0$, $1 = \underline{\psi}(x(t), t) < \varphi(x(t), t)$, and $1 = \varphi(x(t_0), t_0)$. Therefore,

$$0 \geq \frac{d}{dt}\varphi(x(t), t) \Big|_{t=t_0} = \frac{1}{12}e^{-\frac{1}{6}t_0} \frac{x_0}{|x_0|} \cdot D\varphi(x_0, t_0) + \varphi_t(x_0, t_0) = \frac{1}{12}se^{-\frac{1}{6}t_0} + \varphi_t(x_0, t_0).$$

Thus, for $s \leq 0$,

$$\begin{aligned} \left(\varphi_t - |D\varphi| \operatorname{div} \left(\frac{D\varphi}{|D\varphi|} \right) - |D\varphi| \right) (x_0, t_0) &\leq -\frac{1}{12}se^{-\frac{1}{6}t_0} - \frac{s}{|x_0|} - |s| \\ &\leq -\frac{s}{12}e^{-\frac{1}{6}t_0} - \frac{s}{2} + s \leq -\frac{s}{12} - \frac{s}{2} + s < 0. \end{aligned}$$

Hence, $\underline{\psi}$ is a subsolution.

Next we compute

$$\frac{\partial}{\partial \mathbf{n}} \underline{\psi}(x, t), \quad \text{for } x \in \partial B_2(O),$$

where \mathbf{n} denotes the outward normal vector on $\partial B_2(O)$. For $x_0 \in \partial B_2(O)$, $t > 0$,

$$\frac{\partial}{\partial \mathbf{n}} \underline{\psi}(x_0, t) = -2e^{\frac{1}{6}t}.$$

Therefore,

$$\lim_{t \rightarrow \infty} \left| \frac{\partial}{\partial \mathbf{n}} \underline{\psi}(x_0, t) \right| = \infty.$$

We complete the proof. \square

Corollary 5.2. *Assume u is a solution of problem (5.1) satisfies $u = 0$ on the boundary in the classical sense. If $u_0 \geq \underline{\psi}(\cdot, 0)$ on $\partial B_2(O)$, then*

$$\|u\|_{C^{0,1}(\overline{B_2(O)} \times [0, \infty))} = \infty.$$

6. OPEN PROBLEMS

In this paper, we get precise behaviors of the limiting profile for special obstacles, and initial data in Theorem 1.3. The following problems are open, and of great interests.

Problem A. Assume that ψ^\pm , and u_0 are radially symmetric (but not of the precise forms of Theorem 1.3). Study the limiting profile and its dependence on ψ^\pm , and u_0 .

Problem B. Let ψ^\pm be as in Theorem 1.3. Study the limiting profile and its dependence on ψ^\pm , and u_0 .

It is important noting that we do not assume u_0 is radially symmetric in Problem B, and therefore, u needs not be radially symmetric. In order to understand clearly the behavior of $v(x) = \lim_{t \rightarrow \infty} u(x, t)$, we need to characterize all stationary solutions of (1.1e), and (1.3), which is not yet known in the literature.

Finally, a most general, and most challenging problem is as following.

Problem C. Characterize all stationary solutions of (1.1e), and (1.3) in the general setting. Use this to study the limiting profile.

REFERENCES

- [1] **Y. G. Chen, Y. Giga, and S. Goto**, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Diff. Geom.*, **33**, 1991, 749-786.
- [2] **L. C. Evans, and J. Spruck**, Motion of level sets by mean curvature. I. *J. Diff. Geom.* **33**, 1991, No. 3, 635-681.
- [3] **Y. Giga**, Surface evolution equations-A level set approach, *Monographs in Mathematics*, Birkhäuser(2006).
- [4] **Y. Giga**, On large time behavior of growth by birth and spread, *Proc. Int. Cong. of Math. 2018 Rio de Janeiro* (2018).
- [5] **Y. Giga, H. Mitake, T. Ohtsuka and Hung V. Tran**, Existence of asymptotic speed of solutions to birth and spread type nonlinear partial differential equations, <https://arxiv.org/abs/1808.06312>.
- [6] **Y. Giga, H. Mitake, and Hung V. Tran**, On Asymptotic Speed of Solutions to Level-Set Mean Curvature Flow Equations with Driving and Source Terms, *SIAM Journal on Mathematical Analysis*, 2016, Vol. 48, No. 5, p3515-3546.
- [7] **Y. Giga, M. Ohnuma, and M. Sato**, On the Strong Maximum Principle and the Large Time Behavior of Generalized Mean Curvature Flow with the Neumann Boundary Condition, *Journal of Differential Equations*, Vol. 154, Issue 1, 1999, Pages 107-131.
- [8] **G. Mercier**, Mean curvature flow with obstacles: a viscosity approach, <https://arxiv.org/abs/1409.7657>.
- [9] **G. Mercier, and M. Novaga**, Mean curvature flow with obstacles: existence, uniqueness and regularity of solutions, *Interfaces Free Bound.* 17 (2015), No. 3, 399-426.
- [10] **T. Ohtsuka, T.-H. R. Tsai, and Y. Giga**, A level set approach reflecting sheet structure with single auxiliary function for evolving spirals on crystal surfaces, *Journal of Scientific Computing* 62 (2015), 831-874.
- [11] **T. Ohtsuka, Y.-H. R. Tsai, and Y. Giga**, Growth rate of crystal surfaces with several dislocation centers, *Cryst. Growth Des.* 18 (2018), 1917-1929.
- [12] **A. Petrosyan, and H. Shahgholian**, Parabolic obstacle problems applied to finance. In *Recent developments in nonlinear partial differential equations*, Vol. 439 of *Contemp. Math.*, 2007, p117-133. Amer. Math. Soc., Providence, RI.
- [13] **P. Sternberg, and W. P. Ziemer**, Generalized motion by curvature with a Dirichlet condition. *Journal of Differential Equation*, Vol. 114, 1994, 580-600.
- [14] **L. J. Zhang**, On curvature flow with driving force starting as singular initial curve in the plane, *J. Geom. Anal.*, <https://doi.org/10.1007/s12220-017-9937-6>.
- [15] **R. Mori, and L. J. Zhang**, On mean curvature flow with driving force starting as singular initial hypersurface, [arXiv:1712.09590](https://arxiv.org/abs/1712.09590).

(Y. Giga) GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN

E-mail address: labgiga@ms.u-tokyo.ac.jp

(H. V. Tran) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN MADISON, VAN VLECK HALL, 480 LINCOLN DRIVE, MADISON, WISCONSIN 53706, USA

E-mail address: hung@math.wisc.edu

(L. Zhang) GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN

E-mail address: zhanglj@ms.u-tokyo.ac.jp