



Title	Equilibrium Study on Interactions between Proteins and Bile-Salt Micelles by Micellar Electrokinetic Chromatography
Author(s)	SAITOH, Tohru; FUKUDA, Teruyuki; TANI, Hirofumi; KAMIDATE, Tamio; WATANABE, Hiroto
Citation	Analytical Sciences, 12(4), 569-573 https://doi.org/10.2116/analsci.12.569
Issue Date	1996-08-10
Doc URL	http://hdl.handle.net/2115/71708
Type	article
File Information	Anal.sci.12-569.pdf



[Instructions for use](#)

Equilibrium Study on Interactions between Proteins and Bile-Salt Micelles by Micellar Electrokinetic Chromatography

Tohru SAITOH, Teruyuki FUKUDA, Hirofumi TANI, Tamio KAMIDATE
and Hiroto WATANABE

Faculty of Engineering, Hokkaido University, Sapporo 060, Japan

Interactions between proteins and bile-salt micelles were evaluated on the basis of the binding constants, which were obtained from the migration of proteins in micellar electrokinetic chromatography. The binding constants, $K_b = [P_b]/([P_w][B])$, where $[P_b]$ is the concentration of a protein binding with a bile-salt, $[P_w]$ free protein concentration, and $[B]$ the concentration of bile salt present in the micelles, were successfully determined from the slope of a linear curve of the capacity factor (k') of the protein against $[B]$. The binding constants were almost identical to those obtained by a gel-filtration method. The value of K_b increased with increasing the hydrophobicity of the bile salt or that of the protein.

Keywords Protein, bile-salt micelle, binding constant, micellar electrokinetic chromatography

Bile salts have been extensively used in several steps of protein purification, such as the selective solubilization of membranes, the chromatographic separation of proteins, and the reconstitution of proteins.¹⁻⁵ A predominant factor controlling the separation efficiency of protein purification is the interactive nature between the proteins and the bile-salt micelles. Therefore, a quantitative description of protein-micelle interactions is necessary for designing micellar-mediated separation methods.

Tanford *et al.*^{6,7} determined the binding constants for interactions between some proteins and bile-salt molecules using equilibrium dialysis and gel-filtration methods. However, these methods are both tedious and time consuming. In addition, few quantitative data describing the exact nature of the protein-micelle interactions have yet been obtained. Thus, a rapid and simple method is desirable for evaluating the equilibrium constants for the interactions.

In this study, we tested the use of micellar electrokinetic chromatography (MEKC) for determining the binding constants between the proteins and bile salts in aqueous micellar solutions. The constants agreed well with those obtained by the gel-filtration method. Furthermore, we discuss the relationship between the protein-micelle interaction and the hydrophobic properties of a bile salt or protein, based on the binding constants.

Experimental

Apparatus

An MEKC instrument was made in a manner similar to that reported by Terabe *et al.*⁸⁻¹⁰ The instrument comprised a fused silica capillary tubing with an internal diameter of 50 μm and a total length of 65 cm; detection was made 50 cm downstream. The detector was a JASCO UVIDEC-100 II type UV detector with a modified flow-cell compartment. A high voltage (0–20 kV) was supplied by Matsusada Precision Devices (HCZE-30PN0.25 high voltage power supply). The temperature was maintained at $25 \pm 1^\circ\text{C}$.

Materials

Bile salts and proteins were obtained from Sigma Chemical Co. They were sodium cholate (SC), sodium deoxycholate (SDC), sodium taurocholate (STC), sodium taurodeoxycholate (STDC), albumin (bovin serum), ovalbumin (chicken egg), myoglobin (horse heart), trypsin (bovin pancreas), lysozyme (chicken egg white), cytochrome *c* (horse heart), and β -lactoglobulin B (bovin milk). Cytochrome *b₅* and P450 were purified by Kamataki's method.³ 3-(Cyclohexyl)propanesulfonic acid (CAPS) was used as a buffer component. Sucrose and Sudan III were used as marker reagents for the solvent and micelles, respectively.

MEKC separation of proteins

The solution used for protein separation was an aqueous CAPS (0.01 M, pH 11.0) buffer solution containing bile salt (0–50 mM). Electroinjection (10 kV for 10 s) was performed in order to introduce a sample

T. S. present address: Tokyo University of Pharmacy & Life Science, 1432-1 Horinouchi, Hachioji, Tokyo 192-03, Japan.

solution. A typical voltage for migration was 20.0 ± 0.1 kV.

Gel-filtration

An eluting solution was loaded with a Shimadzu LC-6A type HPLC pump, and passed through a Sephadex G-75 column (1.5 cm i.d. \times 50 cm). The solution was then fractionated with a Toyo SF-100G fraction collector. Bile-salt in the respective fraction was determined by measuring the fluorescence intensity of the product caused from a 3α -hydroxy steroid dehydrogenase-diaphorase combined enzymatic reaction in which the substrates were NAD and resazurin, respectively.¹¹

Measurement of critical micelle concentration

The critical micelle concentration (cmc) of the bile-salt was determined by measuring the enhanced fluorescence intensity of 8-anilino-1-naphthalenesulfonate sodium salt in an aqueous buffer solution of 0.02 M CAPS containing prescribed amounts of bile salt.¹²

Results and Discussion

Estimation of binding constants

The variation of the chromatograms with and without micelles are clearly illustrated in Fig. 1, where different migration patterns can be observed. The migration times of four proteins with the micelles are greater than those without micelles, obviously indicating protein binding with the bile-salt micelles. The difference in the migration time suggests that the chemical natures of the protein-micelle interactions are dependent on the kind of proteins.

A qualitative description of the interactions of proteins with surfactants (or micelles) was made by Tanford *et al.*^{6,7,13} For example, SDC molecules strongly bind to specific sites onto albumin. However, additional SDC molecules do not cooperatively bind to the surface of albumin. In contrast, cytochrome *b*₅ has no specific sites for SDC, but can interact with SDC aggregates in a higher SDC concentration region. In some cases, one protein molecule may interact with a number of SDC aggregates. Tanford's study suggests that the interaction of surfactant molecules (or micelles) with proteins is largely dependent on the kind of proteins or micelles, or their combination. Therefore, a quite complicate model is required for representing the interactions between proteins and bile-salt micelles.

To a first approximation, we simplify the interaction as the binding of a negatively charged protein with a bile-salt micelle. The protein-micelle interaction can be evaluated in the same manner as that for the interactions of small charged molecules with micelles.¹⁰ An expanded view of a capillary column of MEKC, depicting mass-transport phenomena, is illustrated in Fig. 2. When a high voltage is applied to both ends of the column, an electroosmotic flow occurred from the positive to the negative ends, while negatively charged




Fig. 1 MEKC chromatogram of proteins. [SDC]: (A) 0 mM, (B) 58 mM. Protein: (a) lysozyme, (b) cytochrome *c*, (c) ovalbumin, (d) albumin.




Fig. 2 Expanded view of the column dynamics.

micelles were dynamically immobilized, due to electrophoretic migration in the opposite direction. The migration velocity of a negatively charged protein (P) is influenced simultaneously by the electrophoretic force and the interaction with the micelles.

The capacity factor (k') of a negatively charged protein can be represented by

$$k' = (v_{eo} + v_{ep} - v_p) / (v_p - v_{mc}) \quad (1)$$

Here, v_{eo} is the electroosmotic velocity, v_{ep} the electrophoretic velocity of the protein, v_p the apparent velocity of the protein, and v_{mc} the velocity of micelle. They can be calculated by

$$v_{eo} = L/t_{eo}; v_{ep} = L/t_{p(cmc)} - v_{eo}; v_p = L/t_p; v_{mc} = L/t_{mc}$$

where L is the capillary length from the inlet to the detector; t_{eo} , t_{mc} and t_p are the migration times of the solvent, micelles and protein, respectively. The subscript (cmc) means the absence of micelles. The values of cmc for the respective bile-salts obtained in this study were 12.5 mM for SC, 7.2 mM for SDC, 12.5 mM for STC, and 4.6 mM for STDC, respectively.

The shape and size of the micelles are expected to be largely dependent on the kind of bile salt. Furthermore, a protein has a larger size than does a bile-salt micelle, and, hence, a protein is hardly incorporated into a micelle. Thus, we use a binding constant for representing the protein-micelle interaction rather than a distribution constant.

Since the bile salt is in large excess, the binding constant is defined by

$$K_b = [P_b]/([P_w][B]), \quad (2)$$

where $[P_b]$ is the concentration of protein binding with the bile salt, $[P_w]$ the free protein concentration, and $[B]$ the concentration of the bile salt exceeding the cmc.¹⁴ The distribution constant (K) of a protein between the micellar and aqueous phases is given by

$$K = [P]_b/[P]_w, \quad (3)$$

if $[P]_b$ and $[P]_w$ are defined as the concentration of the protein in the micellar and aqueous phases. The capacity factor (k') can be related to the distribution constant (K) through

$$k' = KV_{mc}/V_{aq} = K\nu[B]/(1 - \nu[B]), \quad (4)$$

where V_{mc} and V_{aq} are the volumes of the micellar and aqueous phases, and ν is the partial specific volume of the micelle.⁹ Under these experimental conditions, since the volume fraction of the micellar phase is very small, the denominator on the right side of Eq. (4) can be approximated to be equal to unity, *i.e.*

$$k' \doteq K\nu[B]. \quad (5)$$

Similarly, $[P]_w$ can be approximate to be $[P_w]$.




Fig. 3 Relationship between k' and $[B]$. Protein: lysozyme (○), cytochrome c (□), ovalbumin (●), albumin (△).

$$[P]_w = [P_w]/(1 - \nu[B]) \doteq [P_w] \quad (6)$$

Since $[P]_b$ is defined as the concentration of the protein in the micellar phase, it can be represented using $[P_b]$,

$$[P]_b = [P_b]/V_{mc} = [P_b]/(\nu[B]). \quad (7)$$

When Eqs. (6) and (7) are substituted to Eq. (3), the distribution constant can be related to the binding constant,

$$K \doteq [P_b]/([P_w]\nu[B]) = K_b/\nu. \quad (8)$$

From Eqs. (5) and (8), we obtain

$$k' \doteq K_b[B]. \quad (9)$$

In Fig. 3, linear relationships were obtained between k' and $[B]$ for all of the proteins tested. From the slopes of the curves, the values of K_b were obtained. They are

Table 1 Binding constants of protein to bile acid

Protein	SDC	STDC	SC	STC	SDC ^a
Albumin	13.9±0.1	21.6±0.3	—	—	14.5±1.3
Ovalbumin	3.9±0.3	3.9±1.0	—	—	2.7±1.1
Myoglobin	33.7±1.3	17.8±1.0	—	—	—
Cytochrome c	16.8±1.0	13.6±0.4	4.9±1.0	0.0±0.2	—
Lysozyme	47.8±0.3	34.0±0.2	16.8±0.9	6.9±1.1	42.4±7.2
Hemoglobin α	25.6±1.0	20.5±0.5	—	—	—
Hemoglobin β	19.8±2.3	18.1±0.6	—	—	—
Cytochrome P450	41.5±0.9	34.1±0.6	—	—	—
Cytochrome b_5	73.4±1.3	136.8±2.6	—	—	—

a. Data obtained by gel filtration method.




Fig. 4 Gel-chromatogram of lysozyme (dotted line) and SDC. The Y-axis is the fluorescence intensity of the product formed from resazurine by a combined enzymatic reaction of 3α -hydroxysteroid dehydrogenase and diaphorase.

summarized in Table 1, together with K_b for the other bile salts.

Comparison with K_b determined by the gel-filtration method

The K_b values obtained in the present study may include errors, because a high electric field (typically 20 kV/65 cm) can influence the orientation or shape of large protein molecules. Indeed, it is well known that the orientation of large DNA molecules is seriously influenced by the electric field in conventional gel electrophoresis.^{15,16} The change in the mobility of the micelle bound with relatively large protein molecules is another source of errors. Such errors can not be prevented when K_b is to be estimated from the retention time of the protein.

An attractive alternative is a gel-filtration method of Hummel and Dreyer¹⁷, which has usually been employed for determining the binding constant of small molecules to proteins.^{7,13,17-22} Since the binding constant is calculated from the peak area of the chromatogram, it should be free from experimental errors, as described above. Figure 4 shows the gel-chromatogram of lysozyme (dotted line) when an aqueous buffer solution of 50 mM SDC was used for the mobile phase. In the chromatogram of SDC, a positive peak appears at the eluting position of lysozyme, and a negative peak follows the peak. This fact is indicative of the specific interaction between lysozyme and SDC.

From the area of a negative peak at different bile salt concentrations, we can calculate the concentration ratio of a protein bound with a bile salt in micelles to the total proteins, $\nu = [P_b]/([P_w] + [P_b])$. If K_b is represented by ν , we obtain

$$\nu/[B] = K_b(1 - \nu). \quad (10)$$




Fig. 5 Scatchard plot for the binding of lysozyme by the SDC micelle.

The values of K_b obtained from the slope of $\nu/[B]$ against ν (Scatchard plot)²³, as shown in Fig. 5, are listed in Table 1. They are almost the same as the values obtained by MEKC, indicating the validity of K_b determined by MEKC.

Influence of hydrophobic properties to K_b

Table 1 apparently shows that the magnitudes of K_b for cytochrome *c* and lysozyme with SDC are greater than those with SC. SDC, having two hydroxyl groups, is more hydrophobic than SC, having three hydroxyl groups. Indeed, the values of the hydrophilic-lipophilic balance (HLB) were 11.9 for SC and 9.5 for SDC, respectively, when they were calculated on the basis of Oda's method.²⁴ Similarly, the K_b values with STDC are greater than those with STC (HLB: 16.2 for STC, 14.0 for STDC). These facts indicate that the hydrophobic interaction plays an important role in the protein-micelle interaction. In contrast, K_b of a certain protein with SDC is not always greater than that with STDC, although the free bile acid is more hydrophobic than the tauro-conjugated one. The hydrophobicity of only the bile-acid moiety is important for the protein-micelle interaction.

The hydrophobic property of a protein can be represented by the hydrophathy index. The index was calculated from the hydrophobicities of amino acids while taking into account the probability of their surface location.²⁵ In Fig. 6, there is a rough, but apparent, correlation between logarithmic K_b and the hydrophathy index. A similar tendency was also observed when a comparison was performed in the STDC system. These facts strongly support the idea that the predominating factor of the protein-micelle interaction is a hydrophobic interaction.

However, $\log K_b$ of albumin is less than that expected from the correlation. As mentioned above, bile salts do not cooperatively interact with the surface of albumin,




Fig. 6 Relationship between the logarithmic binding constant and the hydrophathy index of protein. Protein: (a) albumin, (b) hemoglobin β , (c) hemoglobin α , (d) cytochrome c, (e) myoglobin, (f) lysozyme, (g) cytochrome P450, (h) cytochrome b_5 .

although some bile-salt molecules strongly bind to its specific sites. Such binding may weaken the interaction between the bile salt and albumin. On the other hand, $\log K_b$ of cytochrome b_5 is much greater than that expected from the correlation. Since cytochrome b_5 has a large hydrophobic domain, cooperative binding of the surfactant aggregates with the hydrophobic domain enhances the interaction between cytochrome b_5 and the bile-salt micelle. The fuzzy correlation between $\log K_b$ and the hydrophathy index is also ascribable to the difference in the electrostatic repulsive forces between negatively charged proteins and micelles.

Finally, it has to be mentioned that SDC is a useful surfactant for solubilizing cytochromes P450 and b_5 from microsomal membranes.^{3,26} SDC has also been employed for eluting these proteins in a column chromatographic separation.^{3,26} The relatively great binding constants of cytochromes P450 and b_5 obtained in this study suggest that SDC is a good choice for utilization in the membrane solubilization or column chromatographic elution of these proteins. Quantitative description of the protein-micelle interaction, as proposed in the present study, would provide useful insight concerning micellar-mediated separation systems.

This work was supported by a Grant-in-Aid for Scientific Research (No. 06453108) from Ministry of Education, Science and Culture, for which the authors are greatly thankful.

References

- O. T. Jones, J. P. Earnest, M. G. McNamee, in "Biological Membranes: Practical Approach", ed. J. B. C. Findlay and W. H. Evans, p. 139, IRL Press, Oxford, 1987.
- Y. Banno, Y. Yada and Y. Nozawa, *J. Biol. Chem.*, **263**, 11459 (1987).
- T. Kamataki, K. Maeda, Y. Yamazoe, T. Nagai and R. Kato, *Biochem. Biophys. Res. Commun.*, **103**, 1 (1981).
- D. L. Foster and R. H. Fillingame, *J. Biol. Chem.*, **254**, 8230 (1979).
- H. Hirata, N. Sone, M. Yoshida and K. Kagawa, *Biochem. Biophys. Res. Commun.*, **69**, 665 (1976).
- N. C. Robinson and C. Tanford, *Biochemistry*, **14**, 369 (1975).
- S. Makino, J. A. Reynolds and C. Tanford, *J. Biol. Chem.*, **248**, 4926 (1973).
- S. Terabe, K. Otsuka, K. Ichikawa, A. Tsuchiya and T. Ando, *Anal. Chem.*, **56**, 111 (1984).
- S. Terabe, K. Otsuka and T. Ando, *Anal. Chem.*, **57**, 834 (1985).
- K. Otsuka, S. Terabe and T. Ando, *J. Chromatogr.*, **348**, 39 (1985).
- F. Mashige, K. Imai and T. Osuga, *Clin. Chim. Acta*, **70**, 79 (1976).
- E. D. Vendittis, G. Palumbo, G. Parlato and V. Bocchini, *Anal. Biochem.*, **115**, 278 (1981).
- J. A. Reynolds and R. H. Simon, *J. Biol. Chem.*, **249**, 3937 (1974).
- E. Pramauro, G. Saini and E. Pelizzetti, *Anal. Chim. Acta*, **166**, 233 (1984).
- D. C. Swartz and C. R. Cantor, *Cell*, **37**, 67 (1984).
- T. Y. Zhang, C. L. Smith and C. R. Cantor, *Nucleic Acids Res.*, **19**, 1291 (1991).
- J. P. Hummel and W. J. Dreyer, *Biochem. Biophys. Acta*, **63**, 530 (1962).
- G. F. Fairclough, Jr. and J. S. Fruton, *Biochemistry*, **5**, 673 (1966).
- G. C. Wood and P. F. Cooper, *Chromatogr. Rev.*, **12**, 88 (1970).
- W. Hoffman and U. Westphal, *Anal. Biochem.*, **32**, 48 (1969).
- J. Clausen, *J. Pharmacol. Exp. Therap.*, **153**, 167 (1966).
- P. Acad, D. M. Brown, T. L. Hardy and K. R. L. Mansford, *Nature [London]*, **199**, 758 (1963).
- G. Scatchard, *Ann. N. Y. Acad. Sci.*, **51**, 660 (1949).
- R. Oda, *Teijin Times*, **22**, 10 (1952).
- J. Kyte and R. F. Doolittle, *J. Mol. Biol.*, **157**, 105 (1982).
- Y. Funae and S. Imaoka, *Biophys. Biochim. Acta*, **842**, 119 (1985).

(Received February 7, 1996)

(Accepted May 13, 1996)